Brazilian Journal of Physical Therapy

<https://www.journals.elsevier.com/brazilian-journal-of-physical-therapy>

ORIGINAL RESEARCH

Influence of posture, sex, and age on breathing pattern and chest wall motion in healthy subjects

Liliane Patrícia De Souza Mendes^a, Danielle Soares Rocha Vieira^b,
Leticia Silva Gabriel^a, Giane Amorim Ribeiro-Samora^a,
Armêle Dornelas De Andrade^c, Daniella Cunha Brandão^c, Maria Clara Goes^d,
Guilherme Augusto Freitas Fregonezi^d, Raquel Rodrigues Britto^e,
Verônica Franco Parreira^{e,*}

^a Rehabilitation Sciences Program, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, MG, Brazil

^b Health Science Department, Universidade Federal de Santa Catarina (UFSC), Araranguá, SC, Brazil

^c Department of Physical Therapy, Universidade Federal de Pernambuco (UFPE), Recife, PE, Brazil

^d PneumoCardioVascular Lab, Hospital Universitário Onofre Lopes – Empresa Brasileira de Serviços Hospitalares (HUOL/EBSERH), Department of Physical Therapy, Universidade Federal do Rio Grande do Norte (UFRN), Natal, RN, Brazil

^e Department of Physical Therapy, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, MG, Brazil

Received 10 April 2018; received in revised form 12 February 2019; accepted 18 February 2019

Available online 1 March 2019

KEYWORDS

Aging;
Optoelectronic
plethysmography;
Physical therapy;
Position

Abstract

Objective: We evaluated the effects of posture, sex, and age on breathing pattern and chest wall motion during quiet breathing in healthy participants.

Methods: Eighty-three participants aged 42.72 (SD = 21.74) years presenting normal pulmonary function were evaluated by optoelectronic plethysmography in the seated, inclined (with 45° of trunk inclination), and supine positions. This method allowed to assess the chest wall in a three dimensional way considering the chest wall as three compartments: pulmonary rib cage, abdominal rib cage and abdomen.

Results: Posture influenced all variables of breathing pattern and chest wall motion, except respiratory rate and duty cycle. Chest wall tidal volume and minute ventilation were reduced ($p < 0.05$) in both sexes from seated to inclined and from seated to supine positions, mainly in males. Moreover, moving from seated to supine position significantly increased the percentage contribution of the abdomen to the tidal volume in both sexes ($p < 0.0001$). Regarding sex, women showed higher contribution of thoracic compartment compared to men ($p = 0.008$). Aging provided reductions on rib cage contributions to tidal volume that were compensated by increases of abdomen contributions ($p < 0.0001$). In addition, increases in end-inspiratory and end-expiratory volumes over the years were observed.

* Corresponding author at: Departamento de Fisioterapia, Avenida Antônio Carlos, 6627, Pampulha, CEP: 31270-901, Belo Horizonte, MG, Brazil.

E-mails: veronicaparreira@yahoo.com.br, veronica.parreira@pq.cnpq.br (V.F. Parreira).

Conclusion: The degree of contribution of chest wall compartments is dependent on posture, sex, and age. Therefore, verticalization increases expansion of pulmonary rib cage as well as horizontalization increases abdominal displacement. Women presented higher thoracic contribution to tidal volume than men. Aging reduces rib cage contributions to tidal volume that were compensated by increases of abdomen contributions.

© 2019 Published by Elsevier Editora Ltda. on behalf of Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia.

Introduction

The assessment of breathing pattern and chest wall motion provide important information about respiratory function.¹ It is known that these parameters are influenced by different factors, including posture, sex, and age.^{1–9} The influence of posture on chest wall motion has been intensively studied.^{1,2,6,9} Nevertheless, its influence on breathing pattern as well as the effect of sex^{7,9} has been less investigated.^{3,8,9} Moreover, the influence of sex on chest wall kinematics is still controversial.^{1,5,7,9} Regarding age, the literature suggests that aging does not change the breathing pattern and chest wall motion.^{1,4,7}

Although several studies have evaluated the influence of posture, sex, and age on breathing pattern and/or chest wall motion of healthy individuals, they focused on the isolated effects of these variables.^{1–9} In addition, respiratory inductive plethysmography was the main method used for the assessment.^{1,4,7,10,11}

As advances had been made to elucidate the mechanisms involved in the control of breathing pattern and chest wall motion, improvements in instrumentation used for its evaluation have also been developed.¹² In this context, optoelectronic plethysmography (OEP) has emerged as a noninvasive,² valid,² and reliable method¹⁵ capable of accurately providing an indirect measure of chest wall volume changes and its three compartments (pulmonary rib cage, abdominal rib cage, and abdomen) in different positions and situations.^{2,12–14} OEP allows analyzing volume variations, in a tricompartimental way, with no need to pre-establish degrees of freedom for the chest wall. This enables a more detailed study of these parameters in the different compartments of the chest wall, as well as the evaluations of end-inspiratory and end-expiratory volumes.^{2,12}

Taking into account the relevance of breathing pattern and chest wall motion assessment and considering the great differential of evaluating the influence of posture, sex, and age both on breathing pattern and chest wall motion using OEP on healthy individuals, it is important to study the influence of these factors on chest wall compartments. The aim of this study was to assess the influence of posture, sex, and age, as well as their interaction on breathing pattern and chest wall motion at rest in healthy individuals using OEP.

Methods

Participants

This was a multicenter cross-sectional study with 83 participants who met the following inclusion criteria: age between 20 and 99 years; body mass index (BMI) between 18.5 and 29.99 kg/m²¹⁶; normal lung function¹⁷; self-reported absence of neuromuscular diseases and/or congestive heart failure; and preserved cognitive function assessed by the Mini-Mental State Examination (MMSE).¹⁸ The exclusion criteria were the inability to understand and/or perform any of the procedures and OEP data acquisition problems. The study was approved by the Ethics Committee of The Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil (ETIC 0194.0.2036000-11). All participants signed a written consent form.

Measurement instrument

Optoelectronic plethysmography (OEP)

The OEP (BTS Bioengineering, Milan, Italy) was used to evaluate the breathing pattern and chest wall motion, during quiet breathing, in three positions: seated, inclined (with 45° of trunk inclination), and supine. This method assesses the chest wall in a three dimensional way considering the chest wall as three compartments: pulmonary rib cage, abdominal rib cage and abdomen. Technical details including marker positions, calibration processes and clinical applicability of the OEP system were previously published.¹² In this study, 89 markers were used for the seated position and 52 markers for the inclined and supine positions, based on anatomical reference points.^{19,20}

Procedures

Data were collected in one or two days, based on the availability of the participant. Initially, clinical and demographic data were registered. The participants then answered the MMSE administered by a trained investigator. Next, they immediately underwent the pulmonary function test (*Koko*®, PFT type; *nSpireHealth Inc.*, CO, USA or Micro Medical Microloop MK8, UK) and subsequently answered the Human Activity Profile (HAP) to register their physical activity level.

The HAP adjusted activity score (AAS) was used in this study to classify the participants as inactive (AAS < 53), moderately active (AAS between 53 and 74) and active (AAS > 74).²¹

Next, the examiner placed 89 markers on the participants' chest wall, and the OEP was calibrated according to the established protocol.^{12,20} Quiet breathing – defined as the participants' own breathing pattern – was assessed for 5 min in each of the three positions. Breathing pattern and chest wall motion were registered for 5 min in seated position, followed by 5 min in inclined and 5 min in supine positions. Calibration of OEP was performed before changing each position.

Variables analyzed

The following variables were analyzed: chest wall tidal volume (V_{cw}); respiratory rate (RR); minute ventilation (VE); duty cycle (Ti/Ttot); end-inspiratory chest wall volume ($V_{ei_{cw}}$); end-expiratory chest wall volume ($V_{ee_{cw}}$); pulmonary rib cage percentage contribution ($V_{RCp\%}$); abdominal rib cage percentage contribution ($V_{RCa\%}$); and abdomen percentage contribution ($V_{AB\%}$). More details regarding the description of the variables were previously published.¹²

Data reduction

To determine the breathing pattern and chest wall motion variables, the middle 100 s from the 5 min registered were used.

Sample size calculation

Sample size calculation was determined for mixed factorial analysis of variance (ANOVA) and linear regression model. For ANOVA, GPower[®] version 3.1 was used. Considering an effect size of 0.25, an alpha error probability of 0.05 and a power of 0.80, the estimated sample size was 44 participants. For regression analysis, considering the 5 independent variables in the model (age, sex, delta seated-inclined, delta inclined-supine, and delta seated-supine), the equation $n \geq 30 + 10k$ ²² was used, which resulted in a sample of at least 80 participants. Therefore, the higher number of individuals required for regression analysis was used for the recruitment of the individuals.

Statistical analysis

Data were presented as measures of central tendency and dispersion, and the normality was verified by the Shapiro–Wilk test. The comparisons between sexes for demographic, spirometric, and clinical variables were performed by Student *t* test for independent samples, and Mann–Whitney or chi-square test, according to the characteristic and/or variable distribution.

To verify the influence of position and sex on breathing pattern and chest wall motion, we performed mixed factorial ANOVA with repeated measures in a split-plot design 3 × 2 (positions and sex). The results regarding the comparisons of breathing pattern and chest wall motion variables in different positions and between sexes were

presented as deltas. This allows a dynamic analysis of the results, as opposed to comparing the absolute values found in each position in a static way. Post hoc analysis of the differences among positions (delta seated-inclined, delta inclined-supine, delta seated-supine) were performed by Bonferroni test.

A stepwise multiple linear regression model was used for assessing the influence of age and was analyzed together with position and sex. For bivariate analysis, we used the Spearman correlation test. The final model was determined from the adjusted coefficient of determination (R^2) and by the statistical significance. To determine the statistical quality of the model, the absence of multicollinearity was verified by the variance inflation factor as well as the presence of homogeneity and normal distribution of residuals, by graphic visual analysis. The level of significance was set at 5%. The Statistical Package for the Social Sciences v 15.0 (Chicago, IL, USA) was used for analyses.

Results

Sample

Initially, 161 individuals were approached to participate in the study in the three centers, of whom 7 refused to participate and 14 did not attend the data collection. From the 140 remaining participants, 57 were not included: 13 presented BMI over 29.99 kg/m² and 2 had BMI below 18.50 kg/m²; 20 showed abnormal pulmonary function test results; two reported respiratory or neurological disease; and 20 were excluded due to problems with OEP data acquisition. Thus, data of 83 individuals were included in the study.

Table 1 presents demographic, anthropometric, spirometric, and clinical data of the individuals by sex. The sample comprised mostly women and compared with men, women were older. There was a significant difference in the HAP score between men and women, but it did not influence the classification as both were considered active.

Influence of posture and sex

Table 2 presents breathing pattern and chest wall motion data in men and women in the three positions. We observed significant differences between the sexes for the variables $V_{RCp\%}$ and $V_{AB\%}$ and among positions (seated, inclined, and supine) for the variables $V_{RCp\%}$, $V_{RCa\%}$, and $V_{AB\%}$. In addition, there was an interaction effect (sex × position) for the variables V_{cw} , VE, $V_{ei_{cw}}$, and $V_{ee_{cw}}$. Neither sex nor position significantly influenced RR or Ti/Ttot.

Fig. 1 presents the results for the influence of posture and sex on breathing pattern. There were interaction effects for all variables, except for RR and Ti/Ttot. The V_{cw} and VE were significantly reduced in both sexes for positions seated to inclined and for seated to supine, with a greater reduction mainly in males. The variables $V_{ei_{cw}}$ and $V_{ee_{cw}}$ were significantly decreased from seated to inclined positions in both sexes, mainly in males and from seated to the supine position only in males. The variables RR and Ti/Ttot were not influenced by sex or position.

Fig. 2 presents the influence of posture and sex on chest wall motion variables. From seated to supine and from

Table 1 Demographic, anthropometric, spirometric, and clinical data of participants by sex.

Variables	Men	Women	p value
Sex	31 (37%)	52 (63%)	0.028
Age (years)	36 (19) [21–81]	47 (22) [21–85]	0.034
BMI (kg/m ²)	25.21 (2.50)	24.19 (2.99)	0.122
FEV ₁ (% predicted)	94.05 (11.60)	93.58 (9.25)	0.846
FVC (% predicted)	94.23 (10.54)	92.40 (8.71)	0.409
FEV ₁ /FVC	0.83 (0.06)	0.83 (0.07)	0.993
HAP score	92.00 (2.91)	80.30 (14.15)	<0.0001
Inactive	0%	7.5%	
Moderately active	0%	22.5%	
Active	100%	70%	
MMSE	28 (2)	29 (2)	0.613

Values are expressed as mean (standard deviation), except sex (%). Minimal and maximal values for age are shown in brackets. BMI, body mass index; FEV₁, forced expiratory volume in first second; FVC, forced vital capacity; FEV₁/FVC, ratio of FEV₁ to FVC; HAP, Human Activity Profile; MMSE, Mini-Mental State Examination.

Figure 1 Influence of posture and sex on breathing pattern.

Data are presented as mean values. *Abbreviations:* V_{cw} , chest wall tidal volume; VE, minute ventilation; $V_{ei_{cw}}$, end-inspiratory chest wall volume; $V_{ee_{cw}}$, end-expiratory chest wall volume. *, significant difference for men between posture; **, significant difference for women; §, significant difference between sexes.

inclined to supine positions $V_{RCp\%}$ was significantly reduced in both sexes, whereas from seated to the inclined position, we observed a reduction of $V_{RCp\%}$ only in men. From the inclined to the supine position, women presented a greater reduction in $V_{RCp\%}$ than men. Regarding $V_{RCa\%}$, a significant reduction was observed from seated to inclined as well as from seated to supine positions, in both sexes without other changes. The $V_{AB\%}$ significantly increased with trunk horizontalization (from seated to inclined and to supine) in both sexes.

Besides the groups were significantly different for HAP score, there was no correlation between the physical

activity level and most of the analyzed variables, except for $V_{RCa\%}$ in the inclined position ($r=0.294$; $p=0.028$), $V_{AB\%}$ in the supine position ($r=-0.281$; $p=0.032$), RR in the supine position ($r=-0.285$; $p=0.032$) and VE in the seated position ($r=0.271$; $p=0.042$), where significant correlations, however, weak, were observed.

Influence of age

Table 3 presents the influence of age adjusted for sex and position, on breathing pattern and chest wall motion. Age

Table 2 Analysis of breathing pattern and chest wall motion variables by sex in three corporal positions.

Variables	Men			Women			ANOVA position ($F_{df} = 2; p$)	ANOVA sex ($F_{df} = 1; p$)	ANOVA interaction ($F_{df} = 2; p$)
	Seated	Inclined	Supine	Seated	Inclined	Supine			
V_{cw} (L)	0.84 (0.45)	0.59 (0.26)	0.58 (0.28)	0.53 (0.23)	0.44 (0.16)	0.41 (0.16)	15.48; <0.0001	40.17; <0.0001	6.20; 0.009
$V_{rcp}\%$	37.41 (8.44)	30.97 (10.63)	19.50 (9.85)	43.02 (11.38)	39.74 (14.31)	23.01 (10.51)	118.08; <0.0001	7.34; 0.008	2.13; 0.122
$V_{rca}\%$	21.93 (5.81)	14.25 (3.92)	11.50 (4.36)	19.51 (5.35)	13.13 (4.95)	11.90 (4.63)	103.65; <0.0001	1.52; 0.221	2.28; 0.118
$V_{ab}\%$	40.83 (10.52)	55.40 (13.26)	69.77 (13.36)	37.42 (12.23)	46.49 (15.89)	65.41 (13.26)	183.91; <0.0001	4.66; 0.034	1.95; 0.148
VE (L/min)	11.67 (4.89)	8.21 (2.55)	8.32 (2.78)	7.98 (1.95)	6.71 (1.83)	6.36 (1.47)	66.77; <0.0001	19.96; <0.0001	11.38; <0.0001
$V_{ee_{cw}}$ (L)	25.13 (3.77)	20.90 (3.59)	22.38 (4.48)	19.79 (3.77)	17.75 (4.03)	19.53 (3.91)	82.45; <0.0001	19.11; <0.0001	15.48; <0.0001
$V_{ei_{cw}}$ (L)	25.96 (3.80)	21.47 (3.63)	22.96 (4.50)	20.32 (3.84)	18.19 (4.06)	19.94 (3.95)	92.99; <0.0001	20.63; <0.0001	17.61; <0.0001
RR (bpm)	15.66 (5.49)	15.69 (4.59)	16.15 (4.72)	16.72 (4.67)	16.86 (5.22)	17.35 (5.49)	0.82; 0.423	1.16; 0.285	0.01; 0.978
Ti/Ttot	0.41 (0.06)	0.41 (0.05)	0.42 (0.05)	0.41 (0.04)	0.40 (0.04)	0.42 (0.09)	1.87, 0.170	0.13; 0.715	0.001; 0.994

Values are expressed as mean (standard deviation). *Abbreviations:* V_{cw} , chest wall volume; $V_{rcp}\%$, percentage contribution of the pulmonary rib cage; $V_{rca}\%$, percentage contribution of the abdominal rib cage; $V_{ab}\%$, percentage contribution of the abdomen; VE, minute ventilation; $V_{ee_{cw}}$, end-expiratory chest wall volume; $V_{ei_{cw}}$, end-inspiratory chest wall volume; RR, respiratory rate; Ti/Ttot, duty cycle.

Figure 2 Influence of posture and sex on chest wall motion. Data are presented as mean values. *Abbreviations:* $V_{RCP\%}$, pulmonary rib cage percent contribution; $V_{RCa\%}$, abdominal rib cage percent contribution; $V_{AB\%}$, abdomen percent contribution. *, significant difference for men; **, significant difference for women; §, significant difference between sexes.

significantly influenced $V_{RCP\%}$, $V_{RCa\%}$, $V_{AB\%}$, $V_{ei\text{cw}}$, and $V_{ee\text{cw}}$. Between 21 and 85 years, for each year of increase in age, we observed an average reduction of 0.20% in $V_{RCP\%}$ and 0.08% in $V_{RCa\%}$ which was associated with an increase of 0.29% in $V_{AB\%}$. With regard to the values for $V_{ei\text{cw}}$ and $V_{ee\text{cw}}$, we observed an increase of 0.08% for each year of increase in age.

Discussion

The main results of this study were: (1) Aging caused reductions in rib cage contributions to tidal volume that were compensated by increases in abdomen contributions. In addition, increases in $V_{ei\text{cw}}$ and $V_{ee\text{cw}}$ with increasing

age were observed. (2) Posture influenced all variables of breathing pattern and chest wall motion, except RR and Ti/T_{tot} . (3) Women presented higher thoracic contribution to tidal volume than men.

Currently, the literature is consistent in suggesting that age does not cause changes in the breathing pattern and chest wall motion of healthy subjects at rest,^{1,4,7} despite the structural and physiological changes that occur in the respiratory system with aging.^{23,24} Contrary to results presently available in the literature,^{1,4,7} our study suggests that chest wall motion is influenced by age. The main alteration of the rib cage related to aging is the reduction of its compliance. This reduction is possibly related to the calcification of costal cartilage and costovertebral joints, as well as the narrowing of the intervertebral spaces.^{23,25} These changes can explain the average volume reduction observed in the two rib cage compartments – 0.20% in the $V_{RCP\%}$ and 0.08% in the $V_{RCa\%}$ – for each year of age increase. However, this reduction is compensated by an average increase in 0.29% in $V_{AB\%}$, which explains the absence of influence of age on the variable V_{cw} found in other studies.^{1,4,7}

In the senile lung, structural changes in connective tissue due to changes in the proportions of collagen and elastin, result in loss of pulmonary elastic recoil, leading to progressive air retention.^{23,25} These changes may explain the mean 0.08% increase in $V_{ei\text{cw}}$ and in $V_{ee\text{cw}}$ for each year of age observed in our study. To our knowledge, this is the first study to provide data on the influence of age in $V_{ei\text{cw}}$ and $V_{ee\text{cw}}$. Other authors have investigated the impact of age on the variables of breathing pattern; however, those studies were performed by inductive plethysmography that does not allow the analysis of these volumes.^{1,4,7}

Previous studies have shown that moving from seated to the supine position reduces the lung compliance and increases static and dynamic resistance and pulmonary elastance.^{3,26} Barnas et al.¹¹ demonstrated that the change from seated to supine position increases pulmonary elastance by about 24% and lung resistance and total resistance of the respiratory system (defined as resistance of lungs and chest wall components) by around 40–50%. These findings can explain the reduction of V_{cw} when changing from seated to inclined and from seated to supine positions. These results were similar to the ones reported by Romei et al.⁹ concerning the reduction of trunk verticalization. Our results are also in agreement with this study which reported a significantly higher reduction of the V_{cw} in men with trunk horizontalization compared to women. According to some authors,^{27,28} lung compliance between men and women is similar. However, to our knowledge, there are no studies that evaluated the influence of posture on the lung compliance of men and women.

Considering the reduction of V_{cw} from seated to inclined and from seated to supine, a similar VE response was expected, since RR is not influenced by posture and sex. Our results were similar to those found by Romei et al.⁹ who also observed a greater reduction of VE in men than in women. Kilbride et al.²⁹ showed that at rest the VE in women is lower than in men, and this is entirely due to the lower tidal volume in women.

The RR was not influenced by sex or change of position. Our results are in agreement with previous studies that also did not observe any influence of posture^{8,9} or of sex^{7,9} on

Table 3 Models obtained by regression analysis for the influence of age on breathing pattern and chest wall motion variables.

Dependent variables	Independent variables	Coefficients	95% CI	R^2 partial	R^2 model	p-value
$V_{rcp}\%$	Constant	60.50	54.68, 66.33		0.45	<0.0001
	Age	-0.20	-0.26, -0.14	0.14		<0.0001
	Seated-supine	-19.06	-22.34, -15.78	0.35		<0.0001
	Seated-inclined	-4.79	-8.07, -1.51	0.03		0.004
	Sex	-8.01	-10.87, -5.15	0.11		<0.0001
$V_{rca}\%$	Constant	23.93	22.39, 25.46		0.45	<0.0001
	Age	-0.08	-0.11, -0.05	0.13		<0.0001
	Seated-supine	-8.76	-10.19, -7.34	0.38		<0.0001
	Seated-inclined	-6.89	-8.32, -5.46	0.27		<0.0001
$V_{ab}\%$	Constant	14.88	8.31, 21.44		0.56	<0.0001
	Age	0.29	0.22, 0.36	0.21		<0.0001
	Seated-supine	28.34	24.65, 32.03	0.49		<0.0001
	Seated-inclined	11.31	7.61, 15.01	0.13		<0.0001
	Sex	8.16	4.95, 11.38	0.09		<0.0001
$V_{ee_{cw}}$ (L)	Constant	11.69	9.76, 13.62		0.40	<0.0001
	Age	0.08	0.06, 0.10	0.20		<0.0001
	Seated-supine	-1.29	-2.38, -0.21	0.02		0.02
	Seated-inclined	-2.90	-3.99, -1.82	0.10		<0.0001
	Sex	4.78	3.84, 5.73	0.29		<0.0001
$V_{ei_{cw}}$ (L)	Constant	12.07	10.12, 14.03		0.41	<0.0001
	Age	0.08	0.06, 0.10	0.20		<0.0001
	Seated-supine	-1.46	-2.56, -0.36	0.03		0.009
	Seated-inclined	-3.06	-4.16, -1.96	0.11		<0.0001
	Sex	4.98	4.02, 5.94	0.30		<0.0001

Values are coefficients, confidence intervals, and determination coefficients (R^2). $V_{RCp}\%$, percentage contribution of the pulmonary rib cage; $V_{RCa}\%$, percentage contribution of the abdominal rib cage; $V_{AB}\%$, percentage contribution of the abdomen; $V_{ee_{cw}}$, end-expiratory chest wall volume; $V_{ei_{cw}}$, end-inspiratory volume chest wall volume; 95% CI, confidence interval.

RR. Concerning Ti/T_{tot} , no literature was found concerning the influence of posture on this variable. However, the values observed in the present study are similar to the ones reported by Tobin et al.³⁰ (0.42 ± 0.03) in the supine position, as well as by Parreira et al.⁷ (average of 0.39) in healthy individuals aged 20–80 years.

The variables $V_{ei_{cw}}$ and $V_{ee_{cw}}$ presented a significant reduction when moving from the seated to inclined position in both sexes and from seated to the supine position, but only in males. The reduction for both positions was more pronounced in males. The decrease of $V_{ei_{cw}}$ could be associated with the reduction observed for V_{cw} from the seated to supine position which was more pronounced in males. In addition, the lower $V_{ei_{cw}}$ observed in the supine position could be explained by the increase in pulmonary elastance and lung resistance.¹¹ On the other hand, we can hypothesize that the reduction of $V_{ee_{cw}}$ from the seated to supine position may reflect the smaller residual functional capacity in the supine position.³¹

Regarding chest wall motion, our results are also in agreement with previous studies, which showed by different methods that in the upright or seated position, the relative contribution of the rib cage to tidal volume is higher than in the supine position, and consequently the contribution of the abdomen to tidal volume is lower in the most upright position of the trunk.^{1,3,5,8–10} The increase in $V_{AB}\%$ when moving from seated to inclined, from seated to supine, and from inclined to supine positions in both men and women is explained by the elastic properties of the rib cage and

abdomen. During quiet breathing in the standing posture, the abdomen is as compliant as the rib cage. In the seated position, the weight of the abdominal contents distends the abdominal wall, and consequently the elastance of the diaphragm and the abdomen becomes larger, thus reducing the contribution of this compartment to the tidal volume. In the supine position, only the abdomen varies its static characteristic by the increase of its compliance.³²

Regarding the effect of sex, women presented a higher thoracic contribution than men, similar to the results found by Romei et al.⁹ Binazzi et al.³³ by considering esophageal and gastric pressures demonstrated that women have a greater contribution of rib cage muscles than men. By using chest X-rays, these authors observed that women have a lower radial dimension of RC in relation to height than men and a greater inclination of the ribs. The reduction of $V_{RCp}\%$ with trunk horizontalization occurred between all positions, except in women during the transition from seated to inclined. Consequently, probably due to this factor women showed a greater reduction of this variable in the change of position from inclined to supine positions than men.

The $V_{RCa}\%$ was also reduced with trunk horizontalization, except in the change from the inclined posture to supine. Priori et al.⁸ and Magalhães et al.³⁴ also observed a significant reduction of this variable in healthy individuals, when comparing the sitting position with the supine. However, in these studies, the authors evaluated the responses between supine and seated positions only, thereby preventing a comparison with the results obtained in the present

study regarding inclined positions. On the other hand, Romei et al.⁹ analyzed the rib cage response as a whole ($V_{RCp\%} + V_{RCa\%}$) and results for $V_{RCa\%}$ with the trunk horizontalization were not presented.

The action of the diaphragm on the rib cage has two components. The first is related to the apposition zone. During inspiration, the increase in intra-abdominal pressure is transmitted through the attached diaphragm to expand the lower part of the rib cage. Its magnitude depends on the size of the apposition zone and on the magnitude of intra-abdominal pressure. The second component is related to the insertion of the muscles in the lower ribs. When the diaphragm contracts, it exerts a cranially oriented force on the lower ribs that has the effect of raising them and turning them outwards. However, for this mechanism to work correctly, the muscle fibers of the diaphragm must be oriented cranially, and the abdominal contents should effectively oppose diaphragm descent. If the resistance is small, the dome of the diaphragm descends easily during inspiration; at the same time, the apposition zone decreases in size, and the abdominal pressure increases only slightly, causing the two action components of the diaphragm to have a small action on the rib cage.^{35,36} Priori et al.⁸ demonstrated by means of ultrasonography that the movement of the diaphragm apposition zone of healthy individuals does not differ between the seated and supine positions. Nonetheless, in the supine position, the resistance offered by the abdominal contents to the descent of the diaphragm dome is smaller than that in the seated position. In this way, the diaphragm has a small action on the rib cage, thus reducing the $V_{RCp\%}$ and the $V_{RCa\%}$.

One limitation of the present study is the fact that volume values were obtained indirectly, without association with a direct measurement by a pneumotachograph. Therefore, they cannot be used as absolute values. In addition, the sample consisted mostly of women, and the imbalance of the number of participants in the age groups did not allow a more detailed study of the effect of age on variables of breathing pattern and chest wall motion. However, the sample size had sufficient power to study the average influence of age between 21 and 85 years. Another limitation is the fact that the positions were not randomized in order to optimize data collection time regarding to the process to place the markers and to calibrate the system.

In summary, the results of the present study suggest that posture, sex, and age influence the breathing pattern and the chest wall motion of healthy individuals at rest. The sitting position can benefit patients who present with reduced tidal volume for different reasons, such as pain, postoperative lung volume changes, pulmonary collapse or any other restriction. Regarding respiratory rate and duty cycle, any of the studied positions produced a significant reduction or increase of these parameters.

The rib cage contribution to tidal volume is greater in vertical postures, whereas the abdomen contribution is greater in horizontal postures. The demonstration that the degree of contribution of the compartments is dependent on posture could be useful for orientation of positioning in patients with specific expansion reduction in some compartments, such as cardiac and bariatric postoperative surgery.

The reduction of the rib cage contribution to the tidal volume with aging may encourage the development of

strategies to minimize the volume loss of this compartment. Future studies to investigate whether interventions aiming to increase rib cage expansion would minimize this loss may be relevant.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

The work was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brazil (CAPES), Finance Code 001; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Grant 309990/2017-3 and Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG). The funding sources had no role in the design, conduct or reporting of this study.

References

1. Verschakelen JA, Demedts MG. Normal thoracoabdominal motions influence of sex, age, posture, and breath size. *Am J Respir Crit Care Med*. 1995;151(2 Pt 1):399–405.
2. Aliverti A, Dellacà R, Pelosi P, Chiumello D, Gattinoni L, Pedotti A. Compartmental analysis of breathing in the supine and prone positions by optoelectronic plethysmography. *Ann Biomed Eng*. 2001;29(1):60–70.
3. Baydur A, Behrakis PK, Zin WA, Jaeger MJ, Weiner JM, Milic-Emili J. Effect of posture on ventilation and breathing pattern during room air breathing at rest. *Lung*. 1987;165(6):341–351.
4. Britto RR, Zampa CC, de Oliveira TA, Prado LF, Parreira VF. Effects of the aging process on respiratory function. *Gerontology*. 2009;55(5):505–510.
5. Fugl-Meyer AR. Relative respiratory contribution of the rib cage and the abdomen in males and females with special regard to posture. *Respiration*. 1974;31(3):240–251.
6. Nozoe M, Mase K, Takashima S, et al. Measurements of chest wall volume variation during tidal breathing in the supine and lateral positions in healthy subjects. *Respir Physiol Neurobiol*. 2014;193(3):38–42.
7. Parreira VF, Bueno CJ, França DC, Vieira DS, Pereira DR, Britto RR. Breathing pattern and thoracoabdominal motion in healthy individuals: influence of age and sex. *Rev Bras Fisioter*. 2010;14(5):411–416.
8. Priori R, Aliverti A, Albuquerque AL, Quaranta M, Albert P, Calverley PM. The effect of posture on asynchronous chest wall movement in COPD. *J Appl Physiol* (1985). 2013;114(8):1066–1075.
9. Romei M, Mauro AL, D'Angelo MG, et al. Effects of gender and posture on thoraco-abdominal kinematics during quiet breathing in healthy adults. *Respir Physiol Neurobiol*. 2010;172(3):184–191.
10. Sharp JT, Goldberg NB, Druz WS, Danon J. Relative contributions of rib cage and abdomen to breathing in normal subjects. *J Appl Physiol*. 1975;39(4):608–618.
11. Barnas GM, Green MD, Mackenzie CF, et al. Effect of posture on lung and regional chest wall mechanics. *Anesthesiology*. 1993;78(2):251–259.
12. Parreira VF, Vieira DS, Myrrha MA, Pessoa IM, Lage SM, Britto RR. Optoelectronic plethysmography: a review of the literature. *Rev Bras Fisioter*. 2012;16(6):439–453.

13. Cabral EEA, Resqueti VR, Lima INDF, Gualdi LP, Aliverti A, Fregonezi AFG. Effects of positive expiratory pressure on chest wall volumes in subjects with stroke compared to healthy controls: a case-control study. *Braz J Phys Ther.* 2017;21(6):416–424.
14. Lage SM, Britto RR, Brandão DC, Pereira DAG, De Andrade A, Parreira VF. Can diaphragmatic breathing modify chest wall volumes during inspiratory loaded breathing in patients with heart failure? *Braz J Phys Ther.* 2018;22(6):452–458.
15. Vieira DS, Hoffman M, Pereira DA, Britto RR, Parreira VF. Optoelectronic plethysmography: intra-rater and inter-rater reliability in healthy subjects. *Respir Physiol Neurobiol.* 2013;189(3):473–476.
16. WHO Organization. *Global database on body mass index*; 2014.
17. Pereira CA, Sato T, Rodrigues SC. New reference values for forced spirometry in white adults in Brazil. *J Bras Pneumol.* 2007;33(4):397–406.
18. Bertolucci PH, Brucki SM, Campacci SR, Juliano Y. O mini-exame do estado mental em uma população geral: impacto da escolaridade. *Arq Neuropsiquiatr.* 1994;52(1):1–7.
19. Aliverti A. Opto-electronic plethysmography: new findings in chronic obstructive pulmonary disease. *Int J Respir Care.* 2008;4(2):45–50.
20. Aliverti A, Pedotti A. Opto-electronic plethysmography. *Monaldi Arch Chest Dis.* 2003;59(1):12–16.
21. Daughton DM, Fix AJ, Kass I, Bell CW, Patil KD. Maximum oxygen consumption and the ADAPT quality-of-life scale. *Arch Phys Med Rehabil.* 1982;63(12):620–622.
22. Knapp TR, Campbell-Heider N. Numbers of observations and variables in multivariate analyses. *West J Nurs Res.* 1989;11(5):634–641.
23. Chan ED, Welsh CH. Geriatric respiratory medicine. *Chest.* 1998;114(6):1704–1733.
24. Zaugg M, Lucchinetti E. Respiratory function in the elderly. *Anesthesiol Clin North Am.* 2000;18(1):47–58.
25. Janssens JP, Pache JC, Nicod LP. Physiological changes in respiratory function associated with ageing. *Eur Respir J.* 1999;13(1):197–205.
26. Behrakis PK, Baydur A, Jaeger MJ, Milic-Emili J. Lung mechanics in sitting and horizontal body positions. *Chest.* 1983;83(4):643–646.
27. Gillissen A, Höltnann B, Schött D, Ulmer WT. Static compliance in subjects with intact lungs. *Respiration.* 1989;55(3):176–180.
28. Colebatch HJ, Greaves IA, Ng CK. Exponential analysis of elastic recoil and aging in healthy males and females. *J Appl Physiol Respir Environ Exerc Physiol.* 1979;47(4):683–691.
29. Kilbride E, McLoughlin P, Gallagher CG, Harty HR. Do gender differences exist in the ventilatory response to progressive exercise in males and females of average fitness? *Eur J Appl Physiol.* 2003;89(6):595–602.
30. Tobin MJ, Chadha TS, Jenouri G, Birch SJ, Gazeroglu HB, Sackner MA. Breathing patterns. 1. Normal subjects. *Chest.* 1983;84(2):202–205.
31. Lumb AB, Nunn JF. Respiratory function and ribcage contribution to ventilation in body positions commonly used during anesthesia. *Anesth Analg.* 1991;73(4):422–426.
32. Agostoni E, Rahn H. Abdominal and thoracic pressures at different lung volumes. *J Appl Physiol.* 1960;15:1087–1092.
33. Binazzi B, Lanini B, Bianchi R, et al. Breathing pattern and kinematics in normal subjects during speech, singing and loud whispering. *Acta Physiol (Oxf).* 2006;186(3):233–246.
34. Magalhaes CM, Fregonezi GA, Vidigal-Lopes M, Vieira BS, Vieira DS, Parreira VF. Effects of non-invasive ventilation and posture on chest wall volumes and motion in patients with amyotrophic lateral sclerosis: a case series. *Braz J Phys Ther.* 2016;20(4):336–344.
35. De Troyer A, Estenne M. Functional anatomy of the respiratory muscles. *Clin Chest Med.* 1988;9(2):175–193.
36. D'Angelo E, Sant'Ambrogio G. Direct action of contracting diaphragm on the rib cage in rabbits and dogs. *J Appl Physiol.* 1974;36(6):715–719.