Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

RT-PCR for SARS-CoV-2: quantitative versus qualitative

We read the Article by Lescure and colleagues¹ with great interest. During the ongoing coronavirus disease 2019 (COVID-19) pandemic, monitoring patients infected with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) using viral kinetics or viral loads in various sample types by real-time RT-PCR has become essential. However, understanding whether the RT-PCR test results are interpreted as quantitative, qualitative, or semi-quantitative is important. Since the cycle threshold (Ct) values from RT-PCR can be affected by batch effect, variations among different runs need to be closely monitored by laboratories—especially for quality control in quantitative RT-PCR. Unfortunately, several papers on COVID-19 use the naive Ct values from qualitative RT-PCR as a quantitation unit or use the Δ Ct values with incorrect quantitation unit.^{2,3} Quantitative RT-PCR is entirely different from qualitative RT-PCR. Ct value itself cannot be directly interpreted as viral load without a standard curve using reference materials. Thorough evaluation of the reliability and robustness of the standard curve is the key to accurately quantify the expected viral copy number.

There is wide heterogeneity and inconsistency of the standard curves calculated from studies that provided Ct values from serial dilution samples and the estimated viral loads (figure).^{2,4,5} An appropriate standard curve with adequate limit of detection is required for viral load quantification to correctly track the viral titre kinetics. A two-step approach using qualitative RT-PCR (for detection) and quantitative RT-PCR (for viral load quantification) is highly recommended for studies focusing on viral loads, as clearly

Figure: Standard curves drawn from papers providing serial dilution factors and corresponding Ct in patient samples

All Ct values were derived from clinical samples targeting RdRp/Orf1b sequence of SARS-CoV-2. Ct=cycle threshold. RdRp=RNA-dependent RNA polymerase. SARS-CoV-2=severe acute respiratory syndrome coronavirus 2. *Ct values provided in the figure 1 legend were used. †Ct values of Orf1b in patient 2 provided in table 1 were used. ‡Ct values of RdRp in test 2 of clinical specimen were used.

presented by Lescure and colleagues.¹ Furthermore, using appropriate quantification units according to different sample types—ie, copies per 1000 cells (for respiratory samples), copies per mL (for plasma), and copies per g (for stool)—should be followed by clinicians, as outstandingly shown by Lescure and colleagues.¹

When interpreting the results of SARS-CoV-2 RT-PCR, the validity of the standard curve using reference materials or in-house plasmid controls with known viral copy numbers should be confirmed first to interpret Ct values as viral loads. In conclusion, precautions are needed when interpreting the Ct values of SARS-CoV-2 RT-PCR results shown in COVID-19 publications to avoid misunderstanding of viral load kinetics for comparison across different studies.

We declare no competing interests.

Mi Seon Han, Jung-Hyun Byun, Yonggeun Cho, *John Hoon Rim
johnhoon1@yuhs.ac

MSH and JHR contributed to study design, data collection, data analysis, data interpretation, literature search, and writing of the Correspondence. J-HB and YC contributed to data collection, data analysis, and data interpretation. All authors reviewed and approved the final version of the Correspondence.

Department of Pediatrics, Seoul Metropolitan Government-Seoul National University Boramae Medical Center, Seoul, Korea (MSH); Department

of Laboratory Medicine, Gyeongsang National University Hospital, Gyeongsang National University College of Medicine, Jinju, Korea (J-HB); Department of Laboratory Medicine, Hallym University Sacred Heart Hospital, Anyang, Korea (YC); Department of Pharmacology, Yonsei University College of Medicine, Seoul 03722, Korea (JHR); Department of Medicine, Physician-Scientist Program, Yonsei University Graduate School of Medicine, Seoul, Korea (JHR); and Department of Laboratory Medicine, Yonsei University College of Medicine, Seoul, Korea (JHR).

- 1 Lescure FX, Bouadma L, Nguyen D, et al. Clinical and virological data of the first cases of COVID-19 in Europe: a case series. *Lancet Infect Dis* 2020; published online March 27. [https://doi.org/10.1016/S1473-3099\(20\)30200-0](https://doi.org/10.1016/S1473-3099(20)30200-0).
- 2 Zou L, Ruan F, Huang M, et al. SARS-CoV-2 viral load in upper respiratory specimens of infected patients. *N Engl J Med* 2020; **382**: 1177–79.
- 3 Shen C, Wang Z, Zhao F, et al. Treatment of 5 critically ill patients with COVID-19 with convalescent plasma. *JAMA* 2020; published online March 27. DOI:10.1001/jama.2020.4783.
- 4 Chu DKW, Pan Y, Cheng SMS, et al. Molecular diagnosis of a novel coronavirus (2019-nCoV) causing an outbreak of pneumonia. *Clin Chem* 2020; **66**: 549–55.
- 5 Chan JF, Yip CC, To KK, et al. Improved molecular diagnosis of COVID-19 by the novel, highly sensitive and specific COVID-19-RdRp/Hel real-time reverse transcription-polymerase chain reaction assay validated in vitro and with clinical specimens. *J Clin Microbiol* 2020; **58**: e00310–20.

Ratio, rate, or risk?

In epidemiology, the terms ratio, rate, and risk have clear definitions.¹ In the emerging publications related to coronavirus disease 2019 (COVID-19), the phrase case fatality rate is being

Published Online
May 20, 2020
[https://doi.org/10.1016/S1473-3099\(20\)30424-2](https://doi.org/10.1016/S1473-3099(20)30424-2)

Published Online
May 27, 2020
[https://doi.org/10.1016/S1473-3099\(20\)30439-4](https://doi.org/10.1016/S1473-3099(20)30439-4)

Published Online

May 28, 2020

[https://doi.org/10.1016/S1473-3099\(20\)30435-7](https://doi.org/10.1016/S1473-3099(20)30435-7)

used instead of case fatality ratio.^{2,3} A ratio is a comparison of two similar quantities. Ratios have no dimensions and can take any value; a ratio of 1 means the two quantities being compared are equal to each other. The case fatality ratio is the ratio of deaths (numerator) to identified cases (denominator), and is usually expressed as percentage.^{2,3} As clearly described by Kelly and Cowling,¹ a rate has a time dimension (ie, time⁻¹); it expresses changes in one quantity over a time period. Risk, however, is the probability associated with an adverse outcome that is likely to occur in the future during follow-up. Like ratios, risk has no dimensions but, unlike ratios, risk is confined to values between 0 and 1.

From a scientific standpoint, these terms acquire different meanings depending on the variables being measured in the relevant study designs. In the context of the COVID-19 pandemic, case fatality ratios are obtained from cross-sectional studies, whereas risk estimates are obtained from prospective cohort studies. The infection rate is calculated by counting new infections in equally spaced time intervals (eg, daily or weekly). This growth rate reflects the speed and scale of new cases and can be used to monitor the effects of preventive interventions.

Thus, in scientific research, these terms do not always align with their usual English language semantics and should not be used interchangeably. Ratio, rate, and risk have different meanings that are based on the variable being measured in the relevant study designs. It is hoped that future publications will use the terms appropriately, in line with the context.

I declare no competing interests

Srinivas Mantha

smantha@srinivasmantha.com

Mantha Heart Clinic, Barkatpura, Hyderabad 500027, India

1 Kelly H, Cowling BJ. Case fatality: rate, ratio, or risk? *Epidemiology* 2013; **24**: 622–23.

- 2 Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020; published online Feb 28. DOI:10.1056/NEJMoa2002032.
- 3 Rajgor DD, Lee MH, Archuleta S, Bagdasarian N, Quek SC. The many estimates of the COVID-19 case fatality rate. *Lancet Infect Dis* 2020; published online March 27. [https://doi.org/10.1016/S1473-3099\(20\)30244-9](https://doi.org/10.1016/S1473-3099(20)30244-9).

Implication of SARS-CoV-2 evolution in the sensitivity of RT-qPCR diagnostic assays

Since the initial outbreak of COVID-19 and identification of a novel enveloped RNA betacoronavirus, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), reports suggest the virus might be evolving, albeit at a lower rate than influenza viruses.^{1–3} Nevertheless, in light of the rapid and pandemic-scale spread of COVID-19, mutations in SARS-CoV-2 raise new diagnostic challenges, including the redesign of the oligonucleotide sequences in use in RT-qPCR assays to circumvent potential primer-sample mismatches.

We report an analysis of all high-coverage SARS-CoV-2 genome sequences (1825 in total) deposited in the Global Initiative on Sharing All Influenza Data (GISAID) database⁴ as of March 30, 2020. We aligned the sequences against the reference sequence obtained from the Wuhan seafood market pneumonia virus isolate, Wuhan-Hu-1 (NC_045512). Subsequently, we annotated in the alignments the binding sites of 33 oligonucleotides developed by different centres and shared by WHO⁵ for use in the RT-qPCR detection of SARS-CoV-2 from human samples. We then calculated the nucleotide diversity (π)⁶ in the binding region of each oligonucleotide (figure).

The analysis revealed that 79% (26 of 33) of the primer binding

sites used in the RT-qPCR assays were mutated in at least one genome. The substitution of three nucleotides (GGG substituted to AAC) at the beginning of the binding site of the forward primer designed by the Chinese National Institute for Viral Disease Control and Prevention in the gene encoding for the nucleocapsid phosphoprotein was of relevance. The AAC variant was in 14% (258 of 1825) of the genomes isolated and sequenced in 24 different countries (Australia, Belgium, Brazil, Chile, China, Czech Republic, Denmark, England, Finland, France, Germany, Greece, Hungary, Iceland, Italy, Mexico, the Netherlands, Peru, Portugal, Spain, Switzerland, the USA, Vietnam, and Wales).

Notwithstanding the possibility of sequencing errors, the fact that some variants were consistently found in different sequencing experiments at independent test sites suggests that these genetic variants are true variants. Moreover, the observation that at least one of the previously designed primers is now likely to be ineffective at detecting up to 14% of the virus variants in circulation strengthens the need to continue optimising the oligonucleotides in use in assays being developed. Oligonucleotide optimisation will be facilitated by global sharing of SARS-CoV-2 genomes and the frequently updated reports on sequence analysis that are available on the GISAID⁴ website.

We declare no competing interests.

We acknowledge all authors originating and submitting the sequences used in this Correspondence (appendix).

***Nuno Sampaio Osório,**
Margarida Correia-Neves
nosorio@med.uminho.pt

Life and Health Sciences Research Institute, School of Medicine, University of Minho, Braga 4710-057, Portugal

1 Lai A, Bergna A, Acciarri C, Galli M, Zehender G. Early phylogenetic estimate of the effective reproduction number of SARS-CoV-2. *J Med Virol* 2020; published online Feb 25. DOI:10.1002/jmv.25723.

See Online for appendix