Published in final edited form as:

J Am Chem Soc. 2019 May 15; 141(19): 7797–7806. doi:10.1021/jacs.9b01262.

Catalytic C–H Amination Mediated by Dipyrin Cobalt Imidos

Yunjung Baek, Theodore A. Betley*

Department of Chemistry and Chemical Biology, Harvard University, 12 Oxford Street, Cambridge, Massachusetts 02138, United States

Abstract

Reduction of ($^{\text{ArL}}$)Co^{II}Br ($^{\text{ArL}}$ = 5-mesityl-1,9-(2,4,6-Ph₃C₆H₂)dipyrin) with potassium graphite afforded the novel Co^I synthon ($^{\text{ArL}}$)Co^I. Treatment of ($^{\text{ArL}}$)Co^I with a stoichiometric amount of various alkyl azides (N₃R) furnished three-coordinate Co^{III} alkyl imidos ($^{\text{ArL}}$)Co(NR), as confirmed by single-crystal X-ray diffraction (R: CMe₂Bu, CMe₂(CH₂)₂CHMe₂). The exclusive formation of four-coordinate cobalt tetrazido complexes ($^{\text{ArL}}$)Co(κ^2 -N₄R₂) was observed upon addition of excess azide, inhibiting any subsequent C–H amination. However, when a weak C–H bond is appended to the imido moiety, as in the case of (4-azido-4-methylpentyl)benzene, intramolecular C–H amination kinetically outcompetes formation of the corresponding tetrazene species to generate 2,2-dimethyl-5-phenylpyrrolidine in a catalytic fashion without requiring product sequestration. The imido ($^{\text{ArL}}$)Co(NAd) exists in equilibrium in the presence of pyridine with a four-coordinate cobalt imido ($^{\text{ArL}}$)Co(NAd)(py) ($K_{\text{a}} = 8.04 \text{ M}^{-1}$), as determined by ¹H NMR titration experiments. Kinetic studies revealed that pyridine binding slows down the formation of the tetrazido complex by blocking azide coordination to the Co^{III} imido. Further, ($^{\text{ArL}}$)Co(NR)(py) displays enhanced C–H amination reactivity compared to that of the pyridine-free complex, enabling higher catalytic turnover numbers under milder conditions. The mechanism of C–H amination was probed via kinetic isotope effect experiments [$k_{\text{H}}/k_{\text{D}} = 10.2(9)$] and initial rate analysis with *para*-substituted azides, suggesting a two-step radical pathway. Lastly, the enhanced reactivity of ($^{\text{ArL}}$)Co(NR)(py) can be correlated to a higher spin-state population, resulting in a decreased crystal field due to a geometry change upon pyridine coordination.

Graphical Abstract

*Corresponding Author betley@chemistry.harvard.edu.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.9b01262.

General experimental considerations and procedures, multinuclear NMR data, solid state molecular structures (PDF)

Crystallography data (CIF)

The authors declare no competing financial interest.

1. INTRODUCTION

Direct incorporation of a nitrene functionality into unreactive C–H bonds is a challenging^{1,2} but appealing transformation given the ubiquity of N-functionalities in natural products,³ agrochemicals, and synthetic precursors.⁴ Thus, as an alternative to the traditional approach utilizing preactivated substrates, transition-metal-catalyzed C–H amination has received great attention.^{5–9} More specifically, the synthesis of late, first-row transition-metal-bound nitrene complexes (e.g., imido, $M-(NR^{2-})$,^{10–12} iminyl, $M(^2NR^-)$,^{13–22} nitrene adducts, $M-(^3NR)$ ^{23–25}) is of interest as such species have been invoked as viable and cost-effective nitrene group transfer reagents.

Toward this end, our group has recently demonstrated intermolecular C–H amination mediated by a three-coordinate dipyrinato Fe^{III} imido, ascribing its reactivity to an atypical high-spin electronic configuration.²⁶ Cobalt imidos, specifically those featuring a covalently bound NR^{2-} fragment (i.e., $Co^{III}(NR)$), however, have been relatively less explored in the field of C–H amination as the majority of these species feature closed-shell electronic configurations, resulting in remarkably stable compounds.^{27–34} Although a handful of cobalt imidos display C–H activation or functionalization, such reactivity is still limited to ligand functionalization, precluding productive C–N bond formation. For example, a Co^{III} imido ($Tp^{tBu,Me}$)- $Co^{III}(NAd)$ reported by Theopold displays amination of one of the peripheral *tert*-butyl C–H bonds of the tris(pyrazolyl)borate ligand.³⁵ The amination reactivity is ascribed to a thermally accessible triplet spin state which serves to destabilize the $Co-N_{imido}$ bond, rendering it more reactive. Recently, three-coordinate $(IMes)Co^{IV}(NDipp)_2$ and $[(IMes)Co^V(NDipp)_2]-[BAr^F_4]$ bisimido complexes were reported by Deng's group.³⁶ Interestingly, the Co^{IV} imido ($S = 1/2$) undergoes intra-molecular C–H activation of a benzylic C–H bond from the IMes ligand (with a subsequent C–N forming step), whereas the analogous Co^V imido ($S = 0$) is relatively inert. A leading feature of reactive cobalt imidos is their open-shell spin ground state or thermally accessible low-lying excited states that give rise to radical character along the metal imido bond vector to generate more reactive entities.

Similarly, our group has reported the synthesis of three-coordinate dipyrinato Co^{III} imidos featuring open-shell electronic configurations.³⁷ Studies revealed that $(ArL)-Co^{III}(N^tBu)$

(^{Ar}L = 5-mesityl-1,9-(2,4,6-Ph₃C₆H₂)dipyrrin) exhibits thermal population of spin excited states ($S = 0 \rightarrow 2$), whereas the mesityl analogue (^{Ar}L)Co^{III}(NMe₃) adopts an $S = 1$ electronic ground state. (^{Ar}L)Co^{III}(N^tBu) displays typical two-electron nitrene group transfer reactivity with PMe₂Ph to quantitatively generate phosphinimide PhMe₂P(N^tBu); however, no C–H bond activation was observed. In contrast, (^{Ar}L)Co^{III}(NMe₃) readily undergoes intramolecular benzylic C–H activation to afford a metallacycloindoline (^{Ar}L)Co(κ^2 -NHC₆H₂-2,4-Me₂-6-CH₂). Although neither imido complex is competent for intermolecular C–H activation, we hypothesized that given their propensity to adopt open-shell electronic configurations, such cobalt imidos might manifest intra-molecular C–H amination reactivity instead. We have previously demonstrated the catalytic synthesis of substituted N-heterocycles from linear alkyl azides via intramolecular C–H amination mediated by dipyrinato Fe^{III} iminyl intermediates.^{14,38} Likewise, we envisioned that a cobalt linear alkyl imido complex could generate the substituted pyrrolidine product upon intramolecular C–H amination. Thus, we sought to examine the following questions: (1) can we synthesize dipyrinato Co^{III} imido complexes using linear alkyl azide substrates? (2) If possible, are these imidos competent for intramolecular C–H amination to afford substituted pyrrolidines? Furthermore, (3) is the amination reaction catalytic? To this end, we report herein the intramolecular catalytic C–H amination mediated by dipyrinato Co^{III} imidos to afford N-heterocyclic products. In addition to the structural evidence of a Co^{III} imido as the species responsible for the desired reactivity, kinetic studies were conducted to elaborate mechanistic details of the Co^I/Co^{III} catalytic cycle. Lastly, our findings on ligand-accelerated C–H functionalization reactivity will be discussed.

2. RESULTS AND DISCUSSION

2.1. Synthesis of Cobalt Synthons.

We have previously reported the preparation of dipyrinato Co^{II} and Co^I complexes, (^{Ar}L)CoCl(py) and (^{Ar}L)Co(py), respectively, which possess 1 equiv of pyridine bound to the metal center.³⁷ In an attempt to synthesize a more sterically accessible cobalt synthon, we employed a metalation protocol used for our recently described solvent-free dipyrin Fe congener.²⁶ As such, refluxing (^{Ar}L)Li with anhydrous CoBr₂ in toluene for 15 h produced the three-coordinate Co^{II} complex (^{Ar}L)Co^{II}Br (**1**), as determined by X-ray crystallographic analysis of single crystals grown from a saturated solution of **1** in a hexane/benzene mixture (2:1) stored at –35 °C (Figure 1a). Chemical reduction of **1** with potassium graphite (KC₈) in a thawing benzene solution cleanly afforded the reduced compound (^{Ar}L)Co^I (**2**) (Scheme 1). Crystals suitable for X-ray diffraction were grown from a 2:1 hexane/benzene mixture at –35 °C, and the solid-state structure is shown in Figure 1b.

Interestingly, **2** exhibits ¹H NMR chemical shifts identical to those observed for the pyridine-bound Co^I analogue, (^{Ar}L)Co-(py). However, the solid-state structure of **2** is distinct from that of (^{Ar}L)Co(py),³⁷ displaying an intramolecular η^6 -arene interaction between the cobalt and one of the *ortho*-phenyl substituents from the 2,4,6-Ph₃C₆H₂ aryl groups (Figure 1b).

2.2. Reactivity of (^{Ar}L)Co (**2**) with Alkyl Azides.

With the Co^I starting material **2** in hand, we sought to explore its reactivity with a variety of substituted linear alkyl azides. We envisioned that treatment of **2** with a linear alkyl azide would furnish the corresponding Co^{III} imido, which could subsequently generate the substituted pyrrolidine via intramolecular C–H amination.

To test our hypothesis, we subjected a series of aliphatic azides such as 1-azidopentane, 1-azido-4-methylpentane, and (4-azidobutyl)benzene to a solution of **2** in benzene-*d*₆. In each instance, we observed immediate consumption of the azides and formation of the corresponding linear imine along with regeneration of **2** instead of our anticipated cyclized products (Scheme 2). We postulated that the imines are generated via a rapid α -H-atom abstraction (α -HAA) mediated by a Co^{III} imido intermediate.³⁹ To avoid such undesired reactivity, α -gem-dimethyl-substituted alkyl azides were employed. As a result, slow addition of 1 equiv of 2-azido-2-methylhexane or 2-azido-2,5-dimethylhexane to a solution of **2** cleanly afforded a new paramagnetic species by ¹H NMR spectroscopy. Single crystals of these species suitable for X-ray diffraction were obtained from a benzene/hexane solution at –35 °C to confirm formation of the corresponding Co^{III} imidos **3** and **4**, respectively (Scheme 2). The X-ray crystal structures of both **3** and **4** revealed a trigonal planar geometry around the cobalt center, which is consistent with our previously reported Co imido, (^{Ar}L)-Co^{III}(N^tBu) (Figures 1c and S-47).³⁷ Complexes **3** and **4** display Co–N_{imido} distances (**3**, 1.616(3); **4**, 1.634(5) Å) and Co–N_{imido}–C angles (**3**, 177.5(3)°; **4**, 178.5(5)°) consistent with metrics for (^{Ar}L)Co^{III}(N^tBu) (Co–N_{imido}: 1.609(3) Å, Co–N_{imido}–C angle: 178.8(1)°).

Gratifyingly, heating each of the isolated imidos **3** and **4** to 100 °C in benzene-*d*₆ furnished the cyclized products 2,2,5-trimethylpyrrolidine and 2,2,5,5-tetramethylpyrrolidine in 13 and 9% yield, respectively, as determined by ¹H NMR. Although the yields are low, the formation of N-heterocycles demonstrates the capability of **3** and **4** to perform intramolecular C–H amination. Moreover, regeneration of **2** was observed by ¹H NMR upon consumption of the imidos, which suggests the potential viability of **2** as a catalyst.

To test whether **2** can indeed serve as a catalyst, a 10 mol % benzene-*d*₆ solution of **2** was exposed to 2-azido-2-methylhexane or 2-azido-2,5-dimethylhexane. Surprisingly, in each case, formation of new paramagnetic species distinct from the corresponding imido complex was observed via ¹H NMR, and no desired cyclized product was detected. The same paramagnetic species can be independently prepared either from reaction of **2** with 2 equiv of azide or from reaction of the isolated imido with one additional equivalent of azide in benzene at room temperature (Scheme 2). The X-ray crystal structures of these species revealed the formation of four-coordinate cobalt tetrazido complexes, (^{Ar}L)Co(κ^2 -N₄R₂) (R: CMe₂Bu, **5**; CMe₂(CH₂)₂CHMe₂, **6**) (Figures 1d and S-50). Similar reactivity was previously reported by our group utilizing the same ligand platform with both iron²⁶ and cobalt.³⁷ Such four-coordinate tetrazido species are presumably generated via a rapid trapping of a second equivalent of azide by a three-coordinate imido.^{40–42} Unfortunately, these cobalt tetrazido complexes are remarkably stable. No decay to the corresponding imido or pyrrolidine was observed upon heating **5** or **6** to 100 °C in benzene-*d*₆. Likewise, heating either tetrazido complex in the presence of excess alkyl azides yields no reaction.

The foregoing results demonstrate that, under catalytic conditions, the in situ generated cobalt imido is rapidly converted into catalytically inactive tetrazido species before undergoing intramolecular C–H bond activation, thus preventing potential catalytic turnover. Therefore, to achieve the desired catalytic intramolecular C–H amination, the C–H bond activation step must kinetically outcompete formation of the tetrazido species. We hypothesized that cleavage of relatively weak C–H bonds such as benzylic C–H bonds may lower the activation energy barrier for C–H activation such that intramolecular cyclization occurs faster than bimolecular tetrazido formation.

2.3. Catalytic C–H Amination.

To test this hypothesis, we studied the reactivity of **2** with (4-azido-4-methylpentyl)-benzene (**7**), where a potential benzylic C–H amination could take place to generate 2,2-dimethyl-5-phenylpyrrolidine (**9**) as a product (Scheme 3).

The reaction progression of azide **7** with 10 mol % of **2** in benzene-*d*₆ was monitored by ¹H NMR spectroscopy at temperatures ranging from 25 to 80 °C. A time progression for the reaction performed at 50 °C highlights the paramagnetically shifted resonances for **2** (Figure 2a), the conversion of **2** to the corresponding imido **8** upon addition of azide **7** (10 equiv) initially (Figure 2b and Scheme 3) and after 1 h (Figure 2c), and the final spectrum for the tetrazido complex (Figure 2d). As illustrated in Figure 2a–d, consumption of **8** occurred via two different pathways over time: (1) formation of catalytically inactive tetrazido species and (2) generation of the desired product **9** along with regeneration of **2**, which is rapidly reoxidized to **8** (Figure 2c), indicating that **8** is the catalytic resting state. Once all imido **8** is converted to the tetrazido species, there is no further consumption of azide **7** and the reaction stops (Figure 2d).

As the reaction temperature increases, the yield of pyrrolidine **9** increases, reaching a maximum of 78% at 80 °C. To better understand the amination reaction's temperature dependence, we surveyed the ratio between unreacted azide **7** and pyrrolidine **9** as a function of temperature by ¹H NMR (Figure 3). At low temperature, a considerable amount of unconsumed azide **7** is present as C–H amination is slow. As a result, the imido intermediate **8** is exposed to large concentrations of azide **7**, leading to predominant formation of the corresponding tetrazido complex. As the temperature increases, however, the amination pathway becomes more favorable, promoting faster consumption of **7**, higher yields of **9**, and minimizing formation of the tetrazido complex.

2.4. Kinetic Analysis of Catalytic C–H Amination.

To establish a detailed mechanism for the catalytic amination reaction, we performed various kinetic experiments. As shown in Figure 2b, the oxidation of **2** with excess azide **7** quantitatively affords imido **8** instantaneously, allowing us to estimate the concentration of **8** based on the initial concentration of catalyst **2**. Furthermore, the initial concentration of **8** remains constant during early stages of the catalytic reaction as the concentration of azide **7** is high enough to reoxidize **2** to **8** upon pyrrolidine formation, enabling initial rate kinetic analysis under pseudo-steady-state conditions. Additionally, the initial rate kinetic analysis was necessary to exclude rate diminishing caused by slow formation of the catalytically

inactive tetrazido species. We first examined the reaction order of amination with respect to **8** by varying the catalyst loading. The rates were determined by measuring the concentration of product **9** (<10% yield) over time at 50 °C in benzene-*d*₆ via ¹H NMR spectroscopy. As shown in Figure 4a, a plot of the concentration of the imido (i.e., [**8**]) versus rates of product formation displays a linear correlation, suggesting that C–H amination is first order in imido **8**.

A similar kinetic analysis was conducted holding the catalyst concentration constant and varying azide concentration ([**7**]) to reveal a zeroth-order dependence in azide **7** (Figure 4b), corroborating the unimolecular nature of the amination reaction. Lastly, we performed a kinetic isotope effect (KIE) experiment to investigate the rate-determining step during catalysis. The KIE value was evaluated by comparing the initial rates of 5 mol % of **2** with **7** and its bisdeuterated form at the benzylic site (4-azido-4-methylpentyl-1,1-*d*₂)benzene (**7-*d*₂**). The observed ratio of the rate constants (*k*_H/*k*_D) was determined to be 7.6(2), which is in agreement with the C–H bond activation step being rate-determining (Figure 4c).⁴³ Similarly, the intramolecular KIE value obtained by employing a stoichiometric amount of monodeuterated azide **7-*d*₁** showed selectivity for the C–H bond three times greater than that of the C–D bond (Figure S-10).

Given all the kinetic data, we propose the following catalytic sequence: (1) fast oxidation of **2** by an alkyl azide affords Co^{III} imido **8**; (2) **8** undergoes intramolecular H–atom abstraction to generate a transient Co^{II} amide (**Int**, Scheme 4), followed by (3) rapid radical recombination to afford the cyclized product **9**. Alternatively, a direct C–H insertion pathway from the Co^{III} imido cannot be fully excluded (Figure S-3).

2.5. Mechanism-Driven Improvement of Catalytic Performance.

To the best of our knowledge, this is the first example of catalytic C–H amination mediated by a spectroscopically observable Co^{III} imido, whose identity can be inferred from the structurally characterized analogues **3** and **4**. Furthermore, it is noteworthy that the catalytic C–H amination we describe herein displays a distinct kinetic profile from the similar ring-closing catalysis mediated by a Co^{III} iminyl radical intermediate (in particular, the system reported by Zhang, de Bruin, and co-workers) as opposed to a Co^{III} imido species.^{44–46} Specifically, as suggested by the KIE experiments, we have observed that the C–H activation step as opposed to activation of the azide moiety is rate-determining. Additionally, in contrast to our previously reported Fe-based catalysis, which requires Boc-protection of the pyrrolidine products to regenerate the active catalyst,^{14,38} the Co^I/Co^{III} catalytic cycle showcased herein is capable of yielding free saturated cyclic amines without requiring product sequestration to eliminate product inhibition. Encouraged by these results, we attempted to further optimize the reaction conditions. Lowering the catalyst loading to 5 mol %, however, afforded 31% of pyrrolidine **9** at 80 °C, a significantly diminished yield compared to 78% obtained using 10 mol % of **2**. The diminished yield is consistent with our proposed mechanism that formation of tetrazido complex is favored at relatively high concentrations of azide. Thus, whereas catalysis should be most effective under slow addition of the azide substrate or dilute reaction conditions, reaction time and volumetric throughput would therefore be limited, respectively. An alternative strategy to suppress

tetrazido formation is to employ an ancillary ligand that would competitively bind the imido intermediate, thereby retarding or eliminating tetrazido formation.

To this end, pyridine was a suitable option given its modular steric and electronic profile. To study the binding dynamics between pyridine and a three-coordinate Co^{III} imido without concern for subsequent H-atom abstraction, an adamantyl-substituted imido (**10**) was prepared as a model complex. Stoichiometric reaction of **2** with 1-azidoadamantane (N₃Ad) cleanly produced (^{Ar}L)Co(NAd) **10** in 87% isolated yield. Akin to imidos **3** and **4**, the solid-state structure of **10** displays a trigonal-planar geometry with a Co–N_{imido} distance of 1.620(7) Å and a nearly linear Co–N_{imido}–C angle of 171.4(6)° (Figure 5a).

With **10** in hand, we examined the complexation ratio between the three-coordinate imido and pyridine in solution using a Job plot analysis.⁴⁷ The maximum value in the Job plot (Figure 5b) appeared at 0.5 mole fraction of **10** and pyridine, suggesting a 1:1 binding stoichiometry to afford (**10-py**) in solution. Furthermore, the binding equilibrium constant of pyridine to **10** ($K_a = 8.04 \text{ M}^{-1}$) was determined by tracking the changes in ¹H NMR chemical shifts of **10** as a function of pyridine concentration (Figure 5c).⁴⁸

Although the binding affinity of pyridine is weak, the potential kinetic effect resulting from pyridine coordination on the rate of formation of the tetrazido complex may still be significant. Thus, we measured the conversion rate of isolated imido **10** to the corresponding tetrazido complex (^{Ar}L)Co(κ^2 -N₄Ad₂) (**11**, Figure 6a) as a function of pyridine-*d*₅ concentration. The rate obtained in the absence of pyridine was defined as 1, and the following relative rates (k_{rel}) were determined at 50% formation of **11**. Indeed, a slower formation of **11** was observed at higher concentrations of pyridine-*d*₅, indicating that, as more pyridine-bound imido **10-py** is present in solution, formation of the tetrazido complex is diminished (Figure 6b). These results support our original hypothesis that blocking the fourth coordination site of a three-coordinate imido with an additional ligand offers steric protection against tetrazido formation.

Given these findings, we anticipated that yields of **9** may be improved in the presence of pyridine. Indeed, a 5 mol % catalytic reaction with **7** in the presence of 10 equiv of pyridine-*d*₅ with respect to **2** afforded 90% of **9** at 80 °C, greatly exceeding the amount of pyrrolidine product formed in the absence of pyridine. Furthermore, as the formation of the corresponding tetrazido complex is suppressed, regenerated Co^I catalyst (^{Ar}L)Co(py) is observed when the reaction is completed (Figure S-19). More surprisingly, improved yields of **9** were observed, and increased rates of C–H amination were also detected, implying enhanced amination reactivity of the pyridine-bound imido (**8-py**) relative to the three-coordinate imido **8**. Similar to the kinetic experiments described in section 2.4, initial rates of the C–H amination were measured at varying concentrations of pyridine-*d*₅ under pseudo-steady-state conditions as the oxidation of **2** by azide **7** remained instantaneous and quantitative with pyridine-*d*₅ in solution (Table 1).

As shown in Table 1, in the absence of pyridine, an attempt to measure the initial rate of amination at 25 °C was not successful because the conversion of imido **8** to **9** is too slow to be measured at this temperature (Table 1, entry 1). In contrast, an increased rate is noted at

the same temperature in the presence of 5 mol % of pyridine-*d*₅ (Table 1, entry 2). Likewise, the resulting initial rate increases as the concentration of pyridine-*d*₅ increases, displaying an overall saturation kinetic behavior in pyridine-*d*₅ (Figure 6c). Furthermore, pyridine-*d*₅ can be used as a solvent to produce 92% of **9** with 1 mol % of catalyst **2** at 80 °C. To summarize, pyridine slows down the formation of the undesired tetrazido complex and functions as a cocatalyst, permitting higher catalytic turnover numbers under milder reaction conditions.

2.6. Mechanistic Study of C–H Amination in the Presence of Pyridine.

Excited by the enhanced C–H functionalization reactivity of **8-py**, we investigated whether the mechanism of C–H activation changes in the presence of pyridine. A KIE value of 10.2(9), which is above the classical limit, was observed in the presence of 10 equiv of pyridine-*d*₅ with respect to **2**, suggesting a stepwise radical pathway for C–H amination (Figure S-9).⁴⁹ To further support a radical pathway, initial rates of amination were measured for a series of *para*-substituted azides under the same conditions. As a result, both electron-donating and electron-withdrawing substituted azides displayed slightly increased rates compared to the parent azide **7** (Table 2), suggesting stabilization of benzylic carboradical intermediates (**Int** or **Int-py**) upon H-atom abstraction. Plotting log k_R versus σ^+ parameters (Figure S-12) results in a nonlinear correlation between reaction rate and electron-donating ability of the *para*-substituent on **7-X**. However, plotting log k_R versus the spin delocalization parameters (σ_{mb} , σ_{jj}^*) developed by Jiang and co-workers⁵⁰ yields a more linear relationship (Figure 7). Multiple coefficient linear regression for the dual-parameter equation $\log k_R = \rho_{mb}\sigma_{mb} + \rho_{jj}^*\sigma_{jj}^*$ provided $\rho_{mb} = -0.25$ and $\rho_{jj}^* = 0.72$. The positive value of ρ_{jj}^* indicates that the *para*-substituent (X) delocalizes spin in the transition state. Furthermore, the large $|\rho_{jj}^*/\rho_{mb}|$ value of 2.9 indicates that the spin delocalization effect dominates in the amination reaction, similar to the previously reported iron-catalyzed amination reaction.³⁹

Taking all mechanistic studies together, our proposed catalytic cycle for the intramolecular C–H amination in the presence of pyridine is illustrated in Scheme 4. Several mechanistic features are of note: (1) both **8** and **8-py** are competent species for catalytic C–H amination to afford substituted pyrrolidine **9** in a two-step radical-type pathway; (2) **8-py** displays a rate of amination drastically improved compared to that of **8** along with slower formation of the catalytically inactive tetrazido complex; (3) **2** exhibits an equilibrium with pyridine, existing as either (^{Ar}L)Co(py) or (^{Ar}L)Co(py)₂, both of which are active catalysts to readily furnish **8-py** (Figure S-19).

2.7. Catalysis with Substituted Pyridines.

Given the utility of added pyridine to facilitate the intramolecular amination catalysis, we sought to determine the impact of changing the pyridine donor on catalysis. To this end, the amination reaction was run with azide **7** using 1 mol % of **2** in the presence of 20 equiv of dimethylaminopyridine (4-(Me₂N)py) in benzene-*d*₆. The reaction afforded 91% of **9** at 80 °C, similar to the yield obtained from the reaction conducted in neat pyridine. Motivated by this result, we sought to investigate the effect of the *para*-substituent on pyridine on the reaction rates and yields of C–H amination. Specifically, we attempted to establish differences in values of k_2 among a series of *para*-substituted pyridines, where k_1 and k_2 are

the first-order rate constants of C–H amination mediated by **8** and **8-py'**, respectively (eq 1).
⁵¹ Assuming that k_1 is negligible at 25 °C based on our experimental results (Table 1, entry 1), the rate law can be further simplified (eq 2).

$$\frac{d[\mathbf{9}]}{dt} = k_1[\mathbf{8}] + k_2[\mathbf{8-py}'] \quad (1)$$

$$\frac{d[\mathbf{9}]}{dt} \approx k_2[\mathbf{8-py}'] \text{ at } 25^\circ\text{C} \quad (2)$$

As shown in Table 3, electron-rich pyridines such as 4-(*t*Bu)py and 4-(Me₂N)py exhibit faster conversions as well as higher yields of **9** compared to pyridine. Both [**8-py'**] and k_2 of each of the *para*-substituted pyridines were calculated based on the independently determined equilibrium binding constants using imido **10** (K_a , Table 3). As a result, the values of k_2 for pyridine, 4-(*t*Bu)py, and 4-(Me₂N)py are similar, suggesting that the improved rates are mostly attributable to the higher concentration of the corresponding pyridine-bound imido complex and not to k_2 . Interestingly, we found that the electron-poor 4-(F₃C)py displays a stronger binding affinity than the unsubstituted pyridine yet slows formation of **9**, resulting in a k_2 considerably smaller than that of electron-rich pyridines (Table 3, entry 4). The diminished rate employing 4-(F₃C)py suggests that electronic effects may also influence the reactivity of the pyridine-bound imidos, although additional studies are necessary to evaluate such contributions.

2.8. Spin-State Consideration of the Pyridine-Bound Imido.

We sought to determine if the pyridine-bound imido displays a different electronic structure as compared to the three-coordinate, pyridine-free imido. Our attempts to isolate **10-py** were unsuccessful due to the weak binding affinity of pyridine. Thus, we measured the solution magnetic moment of **10** as a function of concentration of 4-(Me₂N)py, assuming that the total magnetic moment (μ_{eff}) is the summation of two individual components, μ'_{eff} and μ''_{eff} arising from **10** and **10-(4-(Me₂N)py)**, respectively (eq 3). The mole fraction (χ) of each species at a given concentration of 4-(Me₂N)py was calculated based on the K_a of 4-(Me₂N)py.

$$\mu_{\text{eff}} = \mu'_{\text{eff}}(\chi^{\mathbf{10}}) + \mu''_{\text{eff}}(\chi^{\mathbf{10-dmap}}) \quad (3)$$

In the absence of 4-(Me₂N)py, where μ_{eff} is the same as μ'_{eff} , **10** shows a μ_{eff} of 2.5 μ_B , resulting from its $S = 0$ ground state (by analogy to (ArL)Co(N^{*t*}Bu)) and population of thermally accessible excited states (e.g., $S = 2$). As the concentrations of 4-(Me₂N)py increases, the total solution magnetic moment of the mixture increases (Figure 8a), resulting in a μ''_{eff} of 4.0 μ_B . The μ''_{eff} value suggests that the pyridine-bound imido complex could change the spin ground state from the singlet, three-coordinate imidos (e.g., $S = 1, 2$), or pyridine coordination could make the spin excited states thermally more accessible.

The observed changes in solution magnetic moments of **10** as a function of 4-(Me₂N)py concentration can be rationalized by considering the frontier molecular orbitals (FMOs). We have previously proposed a qualitative FMO description of a three-coordinate cobalt imido (^AL)Co(N^tBu) to describe its thermal behavior (Figure 8b).³⁷ Those FMOs would be further perturbed upon coordination of a fourth ligand and thus give rise to a different electronic configuration. Pyridine coordination could result in a trigonal monopyramidal geometry if the pyridine simply associates axially, or the geometry could undergo a tetrahedral distortion as has been observed for pyridine coordination in Holland's iron imido work.^{51–53} To this end, we have performed geometry optimizations for **10-py** considering an *S* = 0, 1, or 2 spin ground state, and the results suggest a trigonal monopyramidal geometry for all cases (Table S-10). Moreover, at the optimized geometries of **10-py**, corresponding *S* = 1 and 2 spin ground states were found to be more stable than that for the *S* = 0 spin state by 6.1 and 7.7 kcal/mol, respectively. As such, we will present our FMO analysis using the *S* = 2 spin ground state as an example, although we cannot fully exclude of the possibility for an *S* = 1 spin ground state given the subtle energy difference between the *S* = 1 and 2 spin states.

A geometry change from trigonal planar to trigonal monopyramidal upon pyridine coordination impacts both the σ - and π -bonding within the molecule. Assuming that pyridine approaches along the *y*-axis, the imide fragment falls below the *xz* plane (Figure 8c). Consequently, the π -overlap between the Co and imido N (Co 3d_{*xz*} + N 2p_{*x*}; Co 3d_{*yz*} + N 2p_{*y*}) is diminished, lowering both d_{*x*} = ($\sigma_{\text{L}}^* + \pi_{\text{N}_x}^*$) and d_{*yz*}($\pi_{\text{N}_y}^*$) in energy. In contrast, 3d_{*z*}² and 3d_{*x*}² - y²($\pi_{\text{N}_y}^*$) are destabilized ($\sigma_{\text{Co-Npy}}^*$), whereas d_{*xy*} remains nominally nonbonding. The net result of the geometric configuration is to decrease the overall 3d energy gap (i.e., (3d_{*xz*} - 3d_{*xy*}) is diminished), leading to the higher spin-state populations observed (Figure 8a). The qualitative orbital interaction analysis agrees with the density functional theory (DFT) results (Figure S-55); however, further studies will be necessary to quantitatively analyze the frontier molecular orbital compositions given that the extent of Co–N_{imido} π -interaction varies as a function of the imido fragment deviation from the *xz* plane.

2.9. Origin of the Enhanced Reactivity of the Pyridine-Bound Imido.

Many transition metal complexes featuring metal–ligand multiple bonds display enhanced group transfer reactivity upon coordination of an exogenous ligand.^{54–57} For example, epoxidation of olefins mediated by either a Cr^V-oxo⁵⁸ or Mn^V-oxo⁵⁹ complex proceeds more efficiently upon ligation of pyridine *N*-oxide to these metal-oxo active species. This effect has also been observed for H-atom abstraction reactivity; however, the reported transition metal complexes are not capable of productive group transfer. The Goldberg group has demonstrated the enhanced H-atom abstraction reactivity of a Mn^V-oxo corrolazine complex in the presence of halide anions.⁶⁰ Similarly, the Holland group showcased that an iron imido complex (^{Me}L)Fe(NAd) undergoes H atom transfer only upon ligation of 4-(^tBu)py.⁵¹ Given these precedents, we propose the following potential explanation to understand how pyridine influences the rate of C–H amination in this study. Based on our FMO analysis and the theoretical studies conducted on **10-py** (section 2.8), we propose that the pyridine-bound, four-coordinate Co imido complex features an attenuated Co–N bond

order by accessing higher spin states as compared to the three-coordinate analogue. Furthermore, as the DFT results suggest, the trigonal monopyramidal geometry obtained upon pyridine ligation may resemble the structure of the transition state for H-atom abstraction more closely. As such, the H atom abstraction step should be more kinetically favored.

3. CONCLUSIONS

The foregoing results demonstrate catalytic C–H amination mediated by dipyrin-supported Co^{III} imidos to afford substituted pyrrolidines from alkyl azides. Although our previously reported cobalt-based alkyl imido (^{Ar}L)Co(N^tBu) does not display intermolecular C–H activation reactivity, our strategy to introduce intramolecular benzylic C–H bonds proximal to the imido moiety promoted C–H bond amination. Kinetic studies support the viability of the structurally characterized three-coordinate dipyrinato Co^{III} imido as a competent nitrene group transfer species. Furthermore, the desired C–H amination reactivity was improved in the presence of pyridine. We found that pyridine plays critical roles during the catalytic reaction: (1) pyridine coordination prevents formation of a catalytically inactive four-coordinate cobalt tetrazido complex and, more importantly, (2) the resulting pyridine-bound Co^{III} imido displays enhanced rate of C–H amination. As a result, we were able to accomplish greater catalytic performance under milder conditions with pyridine as a cocatalyst. We rationalize the improved nitrene group transfer reactivity of the pyridine-bound imido might be resulting from perturbations in its geometric and electronic structures along with potential changes in reaction thermodynamics. We believe this study demonstrates the importance of understanding the salient features of such transition metal ligand multiple bonds to better design new catalysts. Given our findings, we are currently interested in further tailoring our systems to expand the group transfer catalysis protocol.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

This work was supported by a grant from the NIH (GM-115815), the Dreyfus Foundation (Teacher-Scholar Award to T.A.B.), and Harvard University. Y.B. gratefully acknowledges a Kwanjeong Graduate Student Fellowship.

REFERENCES

- (1). Labinger JA; Bercaw JE Understanding and exploiting C–H bond activation. *Nature* 2002, 417, 507–514. [PubMed: 12037558]
- (2). Bergman RG C–H activation. *Nature* 2007, 446, 391–393. [PubMed: 17377575]
- (3). O'Hagan D. Pyrrole, pyrrolidine, pyridine, piperidine and tropane alkaloids. *Nat. Prod. Rep* 2000, 17 (5), 435–446. [PubMed: 11072891]
- (4). Franzén J; Marigo M; Fielenbach D; Wabnitz TC; Kjaersgaard A; Jørgensen KA A general organocatalyst for direct α -functionalization of aldehydes: stereoselective C–C, C–N, C–F, C–Br, and C–S bond-forming reactions. Scope and mechanistic insights. *J. Am. Chem. Soc* 2005, 127 (51), 18296–18304. [PubMed: 16366584]
- (5). Park Y; Kim Y; Chang S. Transition metal-catalyzed C–H amination: scope, mechanism, and applications. *Chem. Rev* 2017, 117 (13), 9247–9301. [PubMed: 28051855]

- (6). Hazelard D; Nocquet P-A; Compain P. Catalytic C–H amination at its limits: challenges and solutions. *Org. Chem. Front* 2017, 4 (12), 2500–2521.
- (7). Zalatan DN; Du Bois J. Metal-catalyzed oxidations of C–H to C–N bonds. *Top. Curr. Chem* 2009, 292, 347–378.
- (8). Ryabov AD Mechanisms of intramolecular activation of C–H bonds in transition metal complexes. *Chem. Rev* 1990, 90 (2), 403–424.
- (9). Dick AR; Sanford MS Transition metal catalyzed oxidative functionalization of carbon-hydrogen bonds. *Tetrahedron* 2006, 62 (11), 2439–2463.
- (10). Nugent WA; Mayer JM Metal–Ligand Multiple Bonds: The Chemistry of Transition Metal Complexes Containing Oxo, Nitrido, Imido, or Alkylidyne Ligands; Wiley-Interscience: Georgetown, 1988.
- (11). Saouma CT; Peters JC $M=E$ and $M=E$ complexes of iron and cobalt that emphasize three-fold symmetry ($E = O, N, NR$). *Coord. Chem. Rev* 2011, 255 (7), 920–937. [PubMed: 21625302]
- (12). Mehn MP; Peters JC Mid-to-high-valent imido and nitrido complexes of iron. *J. Inorg. Biochem* 2006, 100 (4), 634–643. [PubMed: 16529818]
- (13). Iovan DA; Betley TA Characterization of iron–imido species relevant for N–group transfer chemistry. *J. Am. Chem. Soc* 2016, 138 (6), 1983–1993. [PubMed: 26788747]
- (14). Hennessy ET; Betley TA Complex N-heterocycle synthesis via iron-catalyzed, direct C–H bond amination. *Science* 2013, 340 (6132), 591–595. [PubMed: 23641113]
- (15). King ER; Hennessy ET; Betley TA Catalytic C–H bond amination from high-spin iron imide complexes. *J. Am. Chem. Soc* 2011, 133 (13), 4917–4923. [PubMed: 21405138]
- (16). Kuijpers PF; van der Vlugt JI; Schneider S; de Bruin B. Nitrene radical intermediates in catalytic synthesis. *Chem. - Eur. J* 2017, 23 (56), 13819–13829. [PubMed: 28675476]
- (17). Lyaskovskyy V; Suarez AIO; Lu H; Jiang H; Zhang XP; de Bruin B. Mechanism of cobalt(II) porphyrin-catalyzed C–H amination with organic azides: radical nature and H–atom abstraction ability of the key cobalt(III)-Nitrene intermediates. *J. Am. Chem. Soc* 2011, 133 (31), 12264–12273. [PubMed: 21711027]
- (18). Suarez AIO; Lyaskovskyy V; Reek JNH; van der Vlugt JI; de Bruin B. Complexes with nitrogen-centered radical ligands: classification, spectroscopic features, reactivity, and catalytic applications. *Angew. Chem., Int. Ed* 2013, 52 (48), 12510–12529.
- (19). Goswami M; Lyaskovskyy V; Domingos SR; Buma WJ; Woutersen S; Troeppner O; Ivanovi -Burmazovi I; Lu H; Cui X; Zhang XP; Reijerse EJ; DeBeer S; van Schooneveld MM; Pfaff FF; Ray K; de Bruin B. Characterization of porphyrin Co(III)-'nitrene radical' species relevant in catalytic nitrene transfer reactions. *J. Am. Chem. Soc* 2015, 137 (16), 5468–5479. [PubMed: 25844713]
- (20). Lu H; Li C; Jiang H; Lizardi CL; Zhang XP Chemoselective amination of propargylic C(sp³)–H bonds by cobalt(II)-based metalloradical catalysis. *Angew. Chem., Int. Ed* 2014, 53 (27), 7028–7032.
- (21). Lu H; Hu Y; Jiang H; Wojtas L; Zhang XP Stereoselective radical amination of electron-deficient C(sp³)–H bonds by Co(II)-based metalloradical catalysis: direct synthesis of α -amino acid derivatives via α -C–H amination. *Org. Lett* 2012, 14 (19), 5158–5161. [PubMed: 22998175]
- (22). Lu H; Jiang H; Hu Y; Wojtas L; Zhang XP Chemoselective intramolecular allylic C–H amination versus C–C aziridination through Co(II)-based metalloradical catalysis. *Chem. Sci* 2011, 2 (12), 2361–2366.
- (23). Conradie J; Ghosh A. Electronic structure of an iron-porphyrin-nitrene complex. *Inorg. Chem* 2010, 49 (1), 243–248. [PubMed: 19994873]
- (24). Corona T; Ribas L; Rovira M; Farquhar ER; Ribas X; Ray K; Company A. Characterization and reactivity studies of a terminal copper–nitrene species. *Angew. Chem., Int. Ed* 2016, 55 (45), 14005–14008.
- (25). Berry JF Terminal nitrido and imido complexes of the late transition metals. *Comments Inorg. Chem* 2009, 30 (1–2), 28–66.
- (26). Wilding MJT; Iovan DA; Betley TA High-spin iron imido complexes competent for C–H bond amination. *J. Am. Chem. Soc* 2017, 139 (34), 12043–12049. [PubMed: 28777558]

- (27). Jenkins DM; Betley TA; Peters JC Oxidative group transfer to Co(I) affords a terminal Co(III) imido complex. *J. Am. Chem. Soc* 2002, 124 (38), 11238–11239. [PubMed: 12236716]
- (28). Betley TA; Peters JC Dinitrogen chemistry from trigonally coordinated iron and cobalt platforms. *J. Am. Chem. Soc* 2003, 125 (36), 10782–10783. [PubMed: 12952446]
- (29). Hu X; Meyer K. Terminal cobalt(III) imido complexes supported by tris(carbene) ligands: imido insertion into the cobalt-carbene bond. *J. Am. Chem. Soc* 2004, 126 (50), 16322–16323. [PubMed: 15600324]
- (30). Dai X; Kapoor P; Warren TH [Me NN]Co(η^6 -toluene): O=O, N=N, and O=N Bond cleavage provides β -diketiminato cobalt μ -oxo and imido complexes. *J. Am. Chem. Soc* 2004, 126 (15), 4798–4799. [PubMed: 15080682]
- (31). Cowley RE; Bontchev RP; Sorrell J; Sarracino O; Feng Y; Wang H; Smith JM Formation of a cobalt(III) imido from a cobalt(II) amido complex. Evidence for proton-coupled electron transfer. *J. Am. Chem. Soc* 2007, 129 (9), 2424–2425. [PubMed: 17288417]
- (32). Jones C; Schulten C; Rose RP; Stasch A; Aldridge S; Woodul WD; Murray KS; Moubaraki B; Brynda M; La Macchia G; Gagliardi L. Amidinato- and guanidinato-cobalt(I) complexes: characterization of exceptionally short Co–Co interactions. *Angew. Chem., Int. Ed* 2009, 48 (40), 7406–7410.
- (33). Liu Y; Du J; Deng L. Synthesis, structure, and reactivity of low-spin cobalt(II) imido complexes [(Me3P)3Co(NAr)]. *Inorg. Chem* 2017, 56 (14), 8278–8286. [PubMed: 28661656]
- (34). Du J; Wang L; Xie M; Deng L. A two-coordinate cobalt(II) imido complex with NHC ligation: synthesis, structure, and reactivity. *Angew. Chem., Int. Ed* 2015, 54 (43), 12640–12644.
- (35). Shay DT; Yap GPA; Zakharov LN; Rheingold AL; Theopold KH Intramolecular C–H activation by an open-shell cobalt(III) imido complex. *Angew. Chem., Int. Ed* 2005, 44 (10), 1508–1510.
- (36). Zhang L; Liu Y; Deng L. Three-coordinate cobalt(IV) and cobalt(V) imido complexes with N-heterocyclic carbene ligation: synthesis, structure, and their distinct reactivity in C–H bond amination. *J. Am. Chem. Soc* 2014, 136 (44), 15525–15528. [PubMed: 25330361]
- (37). King ER; Sazama GT; Betley TA Co(III) imidos exhibiting spin crossover and C–H bond activation. *J. Am. Chem. Soc* 2012, 134 (43), 17858–17861. [PubMed: 23043624]
- (38). Iovan DA; Wilding MJT; Baek Y; Hennessy ET; Betley TA Diastereoselective C–H bond amination for disubstituted pyrrolidines. *Angew. Chem., Int. Ed* 2017, 56 (49), 15599–15602.
- (39). Hennessy ET; Liu RY; Iovan DA; Duncan RA; Betley TA Iron-mediated intermolecular N-group transfer chemistry with olefinic substrates. *Chem. Sci* 2014, 5 (4), 1526–1532.
- (40). Ceni S; Gallo E; Caselli A; Ragaini F; Fantauzzi S; Piangiolino C. Coordination chemistry of organic azides and amination reactions catalyzed by transition metal complexes. *Coord. Chem. Rev* 2006, 250 (11), 1234–1253.
- (41). Meyer KE; Walsh PJ; Bergman RG A mechanistic study of the cycloaddition-cycloreversion reactions of zirconium-imido complex Cp2Zr(N-t-Bu)(THF) with organic imines and organic azides. *J. Am. Chem. Soc* 1995, 117 (3), 974–985.
- (42). Michelman RI; Bergman RG; Andersen RA Synthesis, exchange reactions, and metallacycle formation in osmium(II) imido systems: formation and cleavage of osmium-nitrogen bonds. *Organo-metallics* 1993, 12 (7), 2741–2751.
- (43). Simmons EM; Hartwig JF On the interpretation of deuterium kinetic isotope effects in C–H bond functionalizations by transition-metal complexes. *Angew. Chem., Int. Ed* 2012, 51 (13), 3066–3072.
- (44). Kuijpers PF; Tiekink MJ; Breukelaar WB; Broere DLJ; van Leest NP; van der Vlugt JI; Reek JNH; de Bruin B. Cobalt-porphyrin-catalysed intramolecular ring-closing C–H amination of aliphatic azides: a nitrene-radical approach to saturated heterocycles. *Chem. - Eur. J* 2017, 23 (33), 7945–7952. [PubMed: 28332743]
- (45). Li C; Lang K; Lu H; Hu Y; Cui X; Wojtas L; Zhang XP Catalytic radical process for enantioselective amination of C(sp3)–H bonds. *Angew. Chem., Int. Ed* 2018, 57 (51), 16837–16841.
- (46). Lu H; Lang K; Jiang H; Wojtas L; Zhang XP Intramolecular 1,5-C(sp3)–H radical amination via Co(II)-based metalloradical catalysis for five-membered cyclic sulfamides. *Chem. Sci* 2016, 7 (12), 6934–6939. [PubMed: 28138382]

- (47). Renny JS; Tomasevich LL; Tallmadge EH; Collum DB Method of continuous variations: applications of Job plots to the study of molecular associations in organometallic chemistry. *Angew. Chem., Int. Ed* 2013, 52 (46), 11998–12013.
- (48). Fielding L. Determination of association constants (K_a) from solution NMR data. *Tetrahedron* 2000, 56 (34), 6151–6170.
- (49). Hammes-Schiffer S; Stuchebrukhov AA Theory of coupled electron and proton transfer reactions. *Chem. Rev* 2010, 110 (12), 6939–6960. [PubMed: 21049940]
- (50). Jiang X-K Establishment and successful application of the $\sigma_{jj}^{\bullet}$ scale of spin-delocalization substituent constants. *Acc. Chem. Res* 1997, 30 (7), 283–289.
- (51). Cowley RE; Eckert NA; Vaddadi S; Figg TM; Cundari TR; Holland PL Selectivity and mechanism of hydrogen atom transfer by an isolable imidoiron(III) complex. *J. Am. Chem. Soc* 2011, 133 (25), 9796–9811. [PubMed: 21563763]
- (52). Cowley RE; Holland PL Ligand effects on hydrogen atom transfer from hydrocarbons to three-coordinate iron imides. *Inorg. Chem* 2012, 51 (15), 8352–8361. [PubMed: 22800175]
- (53). Eckert NA; Vaddadi S; Stoian SA; Lachicotte RJ; Cundari TR; Holland PL Coordination number dependence of reactivity in an imidoiron(III) complex. *Angew. Chem., Int. Ed* 2006, 45 (41), 6868–6871.
- (54). Berrisford DJ; Bolm C; Sharpless KB Ligand-accelerated catalysis. *Angew. Chem., Int. Ed. Engl* 1995, 34 (10), 1059–1070.
- (55). Man W-L; Lam WWY; Yiu S-M; Lau T-C; Peng S-M Direct aziridination of alkenes by a cationic (salen)ruthenium(VI) nitrido complex. *J. Am. Chem. Soc* 2004, 126 (47), 15336–15337. [PubMed: 15563133]
- (56). Jacobsen EN; Marko I; Mungall WS; Schroeder G; Sharpless KB Asymmetric dihydroxylation via ligand-accelerated catalysis. *J. Am. Chem. Soc* 1988, 110 (6), 1968–1970.
- (57). Jacobsen EN; Marko I; France MB; Svendsen JS; Sharpless KB Kinetic role of the alkaloid ligands in asymmetric catalytic dihydroxylation. *J. Am. Chem. Soc* 1989, 111 (2), 737–739.
- (58). Samsel EG; Srinivasan K; Kochi JK Mechanism of the chromium-catalyzed epoxidation of olefins. Role of oxochromium(V) cations. *J. Am. Chem. Soc* 1985, 107 (25), 7606–7617.
- (59). Finney NS; Pospisil PJ; Chang S; Palucki M; Konsler RG; Hansen KB; Jacobsen EN On the viability of oxametallacyclic intermediates in the (salen)Mn-catalyzed asymmetric epoxidation. *Angew. Chem., Int. Ed. Engl* 1997, 36 (16), 1720–1723.
- (60). Prokop KA; de Visser SP; Goldberg DP Unprecedented rate enhancements of hydrogen-atom transfer to a manganese(V)–oxo corrolazine complex. *Angew. Chem., Int. Ed* 2010, 49 (30), 5091–5095.

Figure 1.

Solid-state structures for (a) $(\text{ArL})\text{Co}^{\text{II}}\text{Br}$ (**1**), (b) $(\text{ArL})\text{Co}^{\text{I}}$ (**2**), (c) $(\text{ArL})\text{Co}(\text{NR})$ (**3**), $\text{R} = \text{C}(\text{CH}_3)_2(\text{CH}_2)_3\text{CH}_3$, and (d) $(\text{ArL})\text{Co}(\kappa^2\text{-N}_4\text{R}_2)$ (**5**), $\text{R} = \text{C}(\text{CH}_3)_2(\text{CH}_2)_3\text{CH}_3$ at 100 K, with the thermal ellipsoids set at the 50% probability level (hydrogen atoms and solvent molecules are omitted for clarity; Co, aquamarine; C, gray; N, blue). Selected bond lengths ($\text{\AA}$) and angles ($^\circ$) for (**3**): Co–N1, 1.931(3); Co–N2, 1.926(3); Co–N_{imido}, 1.616(3); Co–N_{imido}–C, 177.5(3).

Scheme 1.
Synthesis of 2

Scheme 2.
Reactivity between 2 and Various Linear Alkyl Azides

Scheme 3.
Intramolecular C–H Amination of Benzylic C–H Bonds

Figure 2.

(a) ^1H NMR spectrum of **2** in benzene- d_6 . (b) ^1H NMR spectrum taken following addition of 10 equiv of azide **7** in benzene- d_6 . Paramagnetic chemical shifts corresponding to imido **8** are colored in blue. (c) ^1H NMR spectrum taken after 1 h and the chemical shifts corresponding to tetrazen $(\text{ArL})\text{Co}(\kappa^2\text{-N}_4\text{R}'_2)$ are colored in red. (d) Reaction completed.

Figure 3. Diamagnetic region in ^1H NMR spectra of the reactions conducted at various temperatures. The singlet at 3.3 ppm corresponds to 1,3,5-trimethoxybenzene, which was used as an internal standard for integration.

Figure 4.

(a) First-order dependence on the concentration of **8** at 50 °C in C_6D_6 . (b) Zeroth-order dependence on the concentration of **7** at 50 °C in C_6D_6 . (c) Kinetic isotope effect study employing **7**/**7**- d_2 at 50 °C in benzene- d_6 .

Scheme 4.
Proposed Catalytic Cycle

Figure 5.

(a) Solid-state structure of $(^{Ar}L)Co(NAd)$ (**10**) at 100 K with the thermal ellipsoids set at the 50% probability level (hydrogen atoms and solvent molecules are omitted for clarity; Co, aquamarine; C, gray; N, blue). (b) Job plot of **10** and pyridine at 25 °C. In all cases, $[10] + [py] = 4.25$ mM in benzene- d_6 . (c) 1H NMR titration curve of **10** at 25 °C in benzene- d_6 . Black dashed line represents a result of fitting based on 1:1 binding isotherms.

Figure 6.

(a) Solid-state structure of $(\Lambda^r\text{L})\text{Co}(\kappa^2\text{-N}_4\text{Ad}_2)$ (**11**) at 100 K with the thermal ellipsoids set at the 50% probability level (hydrogen atoms and solvent molecules are omitted for clarity; Co, aquamarine; C, gray; N, blue). (b) Observed rates of formation of **11** as a function of $[\text{py-d}_5]$ at 25 °C. In all cases, $[\mathbf{10}] = 8.50$ mM and $[\text{N}_3\text{Ad}] = 85.0$ mM in benzene- d_6 . (c) Observed rates of formation of **9** as a function of $[\text{py-d}_5]$ at 25 °C. In all cases, $[\mathbf{2}] = 4.25$ mM and $[\mathbf{7}] = 85.0$ mM in benzene- d_6 .

Figure 7. Linear free energy correlation of $\log k_R$ vs $(\sigma_{mb}, \sigma_{JJ})$ for the amination reaction of *p*-substituted aryl azide substrates listed in Table 2, where $\rho_{mb} = -0.25$, $\rho_{JJ} = 0.72$.

Figure 8.

(a) Solution magnetic moments of mixture of **10** and **10**-(4-Me₂Npy) as a function of [4-Me₂Npy] at 25 °C in benzene-*d*₆. (b) Frontier MO description of (^{Ar}L)Co(NR'). (c) Frontier MO description of (^{Ar}L)Co(NR')(py).

Table 1.Effect of Pyridine Concentration on C–H Amination Reactivity^a

entry	[py- <i>d</i> ₅] (mM)	rate (mM/min)	yield (%) ^b
1	0	N/A ^c	<10
2	4.3	0.52(3)	21
3	8.5	0.70(5)	22
4	26	0.813(5)	35
5	89	1.12(2)	89
6	174	1.45(4)	91
7	300	1.63(4)	93
8	430	1.77(3)	93

^aReactions were conducted with pyridine-*d*₅ to manipulate kinetic experiments with ¹H NMR spectroscopy. In all cases, [2] = 4.25 mM and [7] = 85.0 mM in benzene-*d*₆ at 25 °C.

^b¹H NMR yields using 1,3,5-trimethoxybenzene as an internal standard.

^cInitial rate is not available due to a slower formation of **9** at 25 °C.

Table 2.

Initial Rates and Final Yields of 9 with *para*-Substituted Azide Substrates

<p>(7-X) $\xrightarrow[25\text{ }^{\circ}\text{C, C}_6\text{D}_6]{5\text{ mol\% (2) [py-}d_5] = 42.5\text{ mM}}$ (9-X)</p>		
entry	<i>p</i> -substituent (X)	relative rate (k_R/k_H)
1	-H	1.00
2	-CH ₃	1.11
3	-OCH ₃	1.39
4	-CF ₃	1.41

Table 3.Catalytic Activity of **2** with *para*-Substituted Pyridines

entry	<i>p</i> -RC ₅ H ₄ N	<i>K</i> _a (M ⁻¹) ^{<i>a</i>}	<i>k</i> _R / <i>k</i> _H	NMR yield (%) ^{<i>b</i>}	<i>k</i> ₂ (min ⁻¹)
1	H	8.04	1.00	36.4	1.25
2	^t Bu	12.9	1.7(4)	89.4	1.42
3	NMe ₂	41.8	3.1(7)	90.8	1.24
4	CF ₃	12.5	0.36(2)	20.4	0.31

^{*a*}Binding constants at 25 °C in benzene-*d*₆.^{*b*}¹H NMR yields using 1,3,5-trimethoxybenzene as an internal standard.