Published in final edited form as:

Childs Nerv Syst. 2019 February ; 35(2): 373–378. doi:10.1007/s00381-018-3968-6.

Onyx embolization of a ruptured anterior inferior cerebellar artery in a neonate

Travis S CreveCoeur¹, Afshin Salehi¹, Brandon A Miller², Matthew J Austin³, Dewitte T Cross III⁴, Matthew D Smyth¹

¹Department of Neurosurgery, St. Louis Children's Hospital, Washington University School of Medicine, St. Louis, MO, USA

²Department of Neurosurgery, University of Kentucky, Lexington, KY, USA

³Department of Interventional Neuroradiology, Lahey Hospital & Medical Center, Tufts University School of Medicine, Burlington, MA, USA

⁴Department of Endovascular Surgical Neuroradiology, Mallinckrodt Institute of Radiology, Washington University School of Medicine, St. Louis, MO, USA

Abstract

Aneurysmal subarachnoid hemorrhage (SAH) is rare in neonates. The authors present a unique report of a neonate with SAH from anterior inferior cerebellar artery (AICA) aneurysm rupture that was successfully treated with Onyx embolization. This case report demonstrates the utility of Onyx embolization for posterior circulation aneurysms in neonates and the successful management of SAH in this population.

Keywords

Subarachnoid hemorrhage; Anterior inferior cerebellar artery; Onyx embolization; Liquid embolic agent; Aneurysm; Posterior circulation

Subarachnoid hemorrhage (SAH) in neonates is rare. While endovascular and microsurgery are the most common treatments for ruptured aneurysms, liquid embolization presents another option for previously inaccessible or difficult-to-treat lesions. We present a unique report of a neonate who suffered SAH secondary to rupture of an anterior inferior cerebellar artery (AICA) aneurysm that was successfully treated with Onyx embolization.

Case report

History and physical

An otherwise healthy four-day old Hispanic male with an unremarkable birth history was noted to have nausea and diarrhea. The following morning, his mother noticed decreased oral intake and lethargy. Later that day, he developed disconjugate gaze and was admitted to

Travis S. CreveCoeur, Crevecoeur.travis@wustl.edu.

Conflict of interest The authors declare that they have no conflict of interest.

St. Louis Children's Hospital. Upon arrival, the patient was lethargic with sluggish pupils and medial right eye deviation. An initial head CT without contrast (Fig. 1a) demonstrated acute SAH with mass effect on the pons as well as diffuse intraventricular hemorrhage (IVH) and hydrocephalus. CT angiography (CTA) (Fig. 1b) revealed a small right-sided posterior circulation aneurysm. He underwent conventional cerebral angiography after being intubated due to apneic episodes.

Operation

The right femoral artery was accessed using a 4-French sheath. The right vertebral artery was selected with a 4-French JB2 catheter. Initial angiography (Fig. 2a) demonstrated a 1.6-mm fusiform aneurysm arising from a right AICA inferior branch proximally. The diameter of the parent AICA at its origin was 0.8 mm before bifurcating into its superior and inferior branches just proximal to basilar artery. A multidisciplinary discussion resulted in a recommendation to pursue endovascular treatment.

A Headway Duo micro-catheter (MicroVention, Aliso Viejo CA) and Transend platinum EX micro-wire (Stryker, Kalamazoo MI) were initially used but we were unable to access the AICA. A second attempt was made with a Synchro-14 soft pre-shaped micro-wire (Stryker, Kalamazoo MI). With a JB2 catheter in the proximal right vertebral artery, the micro-catheter and micro-wire were used to select the inferior branch of the right anterior inferior cerebellar artery.

Immediate selective right AICA angiography through the micro-catheter demonstrated contrast extravasation (Fig. 2b). Embolization with Onyx-34 (Medtronic, Irvine CA) resulted in occlusion of the inferior right AICA branch as well as the small fusiform aneurysm arising from that branch (Fig. 2c). There was Onyx extravasation into the subarachnoid space surrounding the inferior right AICA branch and aneurysm. No Onyx refluxed into the basilar artery. At the completion of embolization, the micro-catheter was withdrawn from the AICA without complication.

Angiography (Fig. 2d) immediately after treatment demonstrated complete obliteration of the inferior branch of the AICA and aneurysm. Anterograde flow in the superior branch of the AICA was maintained. There was no evidence of thrombus formation or distal vessel occlusion. The JB2 catheter was removed, and no further angiography was done due to weight-based contrast dose limitations. Hemostasis was achieved at the femoral artery puncture site after sheath removal using manual compression. The patient was transferred back to the pediatric ICU in stable neurological condition. In total, the infant was exposed to 762 mGy of radiation along with 18 mL of Optiray-320 (Mallinckrodt Pharmaceuticals, St. Louis, MO) contrast administered.

Post operative course

Immediately after surgery, a head CT (Fig. 3a) revealed diffuse intraventricular hemorrhage with worsening ventriculomegaly; thus, a right-sided extraventricular drain (EVD) was placed. On post-operative day 3, video EEG showed clinical seizures, characterized by left arm posturing and rhythmic arm movement, and subclinical seizures arising from the left frontal and right central head regions. Each seizure lasted 30 s to 5 min and resolved with a

single-dose Ativan and increased Keppra dose. The following day (post-operative day 4), the patient experienced five brief (< 30 s) subclinical seizures from the right central regions, which resolved by increasing Keppra to maximum dose. No further seizures were observed in the subsequent 48 h and video EEG was discontinued. Genetic workup during hospitalization was unremarkable.

Surveillance MR angiography (MRA) was performed post-operative day 7 which demonstrated stable aneurysm obliteration and vasospasm in the basilar, and bilateral anterior and middle cerebellar arteries. Permissive hypertension (SBP > 95, or 110 above previous baseline), aggressive hydration, and nimodipine were initiated. Repeat MRA and CT were performed on post-operative day 14 and 18, respectively, and both showed improved clearance of the hemorrhage and attenuation of the vasospasm. Nimodipine was continued for a total of 21 days. The EVD was removed on post-operative day 20 when the CSF had cleared. On post-operative day 23, he developed a fixed medial right eye deviation with inability to close the eye. Head CT (Fig. 3b) showed interval appearance of new intraventricular hemorrhage with blood around the site of the EVD catheter, as well as increased size of the lateral and third ventricles. A 4-cm McComb reservoir was placed the following day and returned xanthochromic CSF. The reservoir was tapped a total of 4 times over 14 days. He was ultimately discharged 3 weeks later after one other episode of seizure that resolved with a single dose of Ativan for a total hospital stay of 46 days. At discharge, our patient had still displayed signs of lateral pontine syndrome. He displayed a right-sided seventh nerve palsy, with a fixed right-eye dilation at 2 mm and his left pupil sluggish. His eyes deviated inward and displayed intermittent horizontal nystagmus. Per audiology, he presented with inadequate auditory brain response (ABR) test under natural sleep.

Post hospital course

Fifteen month after his initial intervention, our patient continues to improve developmentally through stretching with physical therapy. Although he has slight increase in the third and fourth ventricular size with sulcal effacement (Fig. 4), he has been weaned off his ventricular taps. His mother has reported no acute events, or neurologic deterioration. He can now lift his head with his own weight despite still needing minor assistance for head and trunk control. There is noticeable improvement to his lateral pontine syndrome symptoms with increased movements on the right side of his face. However, it still remains flaccid in comparison with his left. Concerned for exposure keratopathy, he underwent a tarsorrhaphy which left him with an eye patch on his eyelid. On neurologic exam, he has symmetric and normal tone in all extremities with his anterior fontanelle noted to be full and bulging. His head circumference has increased to the 80th percentile (previous 55th percentile). Given these findings, the next course of action is placement of a right occipital ventriculoperitoneal shunt to relieve his hydrocephalus.

Discussion

This is the first reported case of SAH caused by posterior circulation aneurysm rupture repaired with Onyx embolization as well as the first case report of a posterior circulation aneurysm in an infant treated with this technique. Pediatric aneurysms account for less than

5 of all aneurysms and are more likely to occur in males, have fusiform character, and tend to be giant in presentation (>2.5 cm) [1–4]. Posterior circulation aneurysms account for 17–21 of pediatric aneurysms [1, 3], and within this group 24 are less than 12 months of age [5]. A recent review of ruptured aneurysms in infants found only two reports of posterior circulation aneurysms [6].

AICA aneurysms account for 1 of all pediatric aneurysms [4]. Mohammad and colleagues reported a partially thrombosed giant basilar artery at the level of the AICA [7] in a 15-year-old boy with sensorineural hearing loss, while another group reported a traumatic intracranial aneurysm in an 8-year-old boy who suffered concurrent arteriovenous malformation after a traffic accident [8]. Thus, to our knowledge, this report serves as a novel presentation of a primary AICA aneurysm rupture in the neonatal population and the only report of a posterior circulation aneurysm in this population treated with Onyx embolization.

Interventions for pediatric aneurysms have mirrored the adult population with the shifting of the current treatment paradigm to favor of endovascular treatment [9]. Endovascular treatment is less invasive and allows for access to difficult-to-expose aneurysms [10]. While some institutions have reported a higher complete obliteration rate with microsurgery versus endovascular treatment [11], analysis of the Kids Inpatient Database (KID) demonstrated a higher hospital mortality, complication rate, and overall costs of surgical clipping when compared to endovascular modality [12].

Within the realm of endovascular interventions traditional methods (coil) stenting) flow diversion) have required larger and less flexible catheters that are inaccessible to smaller and distal aneurysms, which may result in rupture of nearby vasculature [13]. Liquid embolization offers a unique solution to this problem, as well as the potential to fill the aneurysm completely [14]. n-butyl cyanoacrylate (nBCA) is preferred by some operators but it can have issues with the delivery precision due to its adhesiveness to ionic compounds, such as blood [15]. Furthermore, potential mistiming of embolization with an nBCA could result in reflux into the basilar artery lumen, which would highly complicate the case. The Onyx liquid embolic system was developed as a nonadhesive liquid composed of ethylene vinyl alcohol copolymer (EVOH) dissolved in dimethyl sulfoxide (DMSO) with tantalum precipitate. As the Onyx solution is injected, the DMSO will wash away in the vasculature while the EVOH and tantalum form a polymeric embolus [16]. Significant disadvantages of using this intervention are operator dependence, learning curve, costs, and inability to be redacted [15]. The pediatric population presents a particular issue with a higher susceptibility to toxicity due to relatively low weight. Within the literature, Onyx has been used with success in the pediatric population with infectious intracranial aneurysms, traumatic intracranial aneurysms, pseudoaneurysms, and arteriovenous malformations (AVMs) [14, 17–19]. In the adult population, Onyx has been used to treat a variety of pathologies, including an AICA flow-dependent aneurysm [20].

The AICA is small in diameter and highly variable in its course, which made the technical aspect of endovascular treatment rather difficult. The AICA may branch from the vertebral, basilar, or posterior inferior cerebellar artery (PICA) [21]. An AICA that is strongly

developed has been defined as > 1.2 mm in an adult brain [22]; thus, in a pre-mature neonate, the vessel would be expected to be far less developed and more sensitive to endovascular shear forces. The risk of hemorrhage related to catheterization of very small vessels of the cerebral vasculature is ever present. The extravasation of contrast seen in our patient could have been from the dissection of a fragile AICA or re-rupture of the AICA aneurysm and likely not from wire perforation, as the wire was seen to follow the lumen of the vessel on roadmapping. The decision to use a micro-catheter with a smaller distal tip OD may have been considered but would not necessarily have altered outcomes. Selecting the 0.53-mm distal tip OD Headway Duo (MicroVention, Aliso Viejo CA) as the micro-catheter of choice preserved the option of coil embolization in addition to liquid embolic agent embolization. The use of other catheters, which may have a smaller distal tip OD, would have been a viable technique for an nBCA embolization only. However, the operators decided that having the flexibility of treatment options using the Headway Duo (MicroVention, Aliso Viejo CA) micro-catheter provided outweighed the value in treating this particular patient than the slight decrease in distal tip OD achieved by other less flexible catheters and nBCA.

References

1. Huang J, McGirt MJ, Gailloud P, Tamargo RJ (2005) Intracranial aneurysms in the pediatric population: case series and literature review. *Surg Neurol* 63:424–432 [PubMed: 15883063]
2. Buis DR, Ouwerkerk WJR, Takahata H, Vandertop WP (2006) Intracranial aneurysms in children under 1 year of age: a systematic review of the literature. *Childs Nerv Syst* 22:1395–1409. 10.1007/s00381-006-0142-3 [PubMed: 16807726]
3. Aryan HE, Giannotta SL, Fukushima T, Park MS, Ozgur BM, Levy ML (2006) Aneurysms in children: review of 15 years experience. *J Clin Neurosci* 13:188–192 [PubMed: 16446095]
4. Beez T, Steiger H-J, Hänggi D (2016) Evolution of management of intracranial aneurysms in children. *J Child Neurol* 31:773–783. 10.1177/0883073815609153 [PubMed: 26516106]
5. Liang J, Bao Y, Zhang H, Wrede KH, Zhi X, Li M, Ling F (2009) The clinical features and treatment of pediatric intracranial aneurysm. *Childs Nerv Syst* 25:317–324. 10.1007/s00381-008-0725-2 [PubMed: 18839183]
6. Lyon KA, Arrey EN, Haider AS, Jeevan DS, Benardete EA (2017) Endovascular treatment of a large ruptured middle cerebral artery bifurcation aneurysm in a 5-week-old infant: case report. *J Neurosurg Pediatr* 20:1–7. 10.3171/2017-5.PEDS17116. [PubMed: 28438070]
7. Mohammad LM, Coon AL, Carlson AP (2017) Resolution of giant basilar artery aneurysm compression and reversal of sensorineural hearing loss with use of a flow diverter: case report. *J Neurosurg Pediatr* 20:1–5. 10.3171/2016.9.PEDS16428 [PubMed: 28438070]
8. Anami H, Aihara Y, Kawashima A (2014) A case report of a pediatric traumatic aneurysm with arteriovenous (A-V) fistula CASE-BASED UPDATE. *Childs Nerv Syst* 30:2003–2008. 10.1007/s00381-014-2551-z [PubMed: 25227169]
9. Vasan R, Patel J, Sweeney JM, Carpenter AM, Downes K, Youssef AS, van Loveren H, Agazzi S (2013) Pediatric intracranial aneurysms: current national trends in patient management and treatment. *Childs Nerv Syst* 29:451–456. 10.1007/s00381-012-1945-z [PubMed: 23288368]
10. Takemoto K, Tateshima S, Golshan A, Gonzalez N, Jahan R, Duckwiler G, Vinuela F (2014) Endovascular treatment of pediatric intracranial aneurysms: a retrospective study of 35 aneurysms. *J Neurointerv Surg* 6:432–438. 10.1136/neurintsurg-2013-010852 [PubMed: 23986132]
11. Sanai N, Auguste KI, Lawton MT (2010) Microsurgical management of pediatric intracranial aneurysms. *Childs Nerv Syst* 26: 1319–1327 [PubMed: 20625742]
12. Alawi A, Edgell RC, Elbabaa SK, Callison RC, Khalili YA, Allam H, Alshekhlee A (2014) Treatment of cerebral aneurysms in children: analysis of the Kids Inpatient Database. *J Neurosurg Pediatr* 14:23–30. 10.3171/2014.4.PEDS13464 [PubMed: 24835049]

13. Gross BA, Smith ER, Scott RM, Orbach DB (2015) Intracranial aneurysms in the youngest patients: characteristics and treatment challenges. *Pediatr Neurosurg* 50:18–25. 10.1159/000370161 [PubMed: 25721683]
14. Reig AS, Simon S, Mericle RA (2009) Embolization of a giant pediatric, posttraumatic, skull base internal carotid artery aneurysm with a liquid embolic agent. *J Neurosurg Pediatr* 4:449–452. 10.3171/2009.6.PEDS0953 [PubMed: 19877778]
15. Leyon JJ, Littlehales T, Rangarajan B, Hoey ET, Ganeshan A (2014) Endovascular embolization: review of currently available embolization agents. *Curr Probl Diagn Radiol* 43:35–53. 10.1067/j.cpradiol.2013.10.003 [PubMed: 24290201]
16. Saeed Kilani M, Izaaryene J, Cohen F, Varoquaux A, Gaubert JY, Louis G, Jacquier A, Bartoli JM, Moulin G, Vidal V (2015) Ethylene vinyl alcohol copolymer (Onyx®) in peripheral interventional radiology: indications, advantages and limitations. *Diagn Interv Imaging* 96:319–326 [PubMed: 25704146]
17. Flores BC, Patel AR, Braga BP, et al. (2016) Management of infectious intracranial aneurysms in the pediatric population. 1205–1217. 10.1007/s00381-016-3101-7
18. Medel R, Crowley RW, Hamilton DK, Dumont AS (2009) Endovascular obliteration of an intracranial pseudoaneurysm: the utility of Onyx. *J Neurosurg Pediatr* 4:445–448. 10.3171/2009.6.PEDS09104 [PubMed: 19877777]
19. de Castro-Afonso LH, Nakiri GS, Oliveira RS, Santos MV, Santos ACD, Machado HR, Abud DG (2016) Curative embolization of pediatric intracranial arteriovenous malformations using Onyx: the role of new embolization techniques on patient outcomes. *Neuroradiology* 58:585–594. 10.1007/s00234-016-1666-1 [PubMed: 26909692]
20. Mahmoud M, El Serwi A, Alaa Habib M, Abou Gamrah S (2012) Endovascular treatment of AICA flow dependent aneurysms: a report of three cases and review of the literature. *Interv Neuroradiol* 18:449–457 [PubMed: 23217640]
21. Naidich TP, Kricheff II, George AE, Lin JP (1976) The normal anterior inferior cerebellar artery. Anatomic-radiographic correlation with emphasis on the lateral projection. *Radiology* 119:355–373. 10.1148/119.2.355 [PubMed: 1265265]
22. Woischneck D, Hussein S (1991) The anterior inferior cerebellar artery (AICA): clinical and radiological significance. *Neurosurg Rev* 14:293–295. 10.1007/BF00383265 [PubMed: 1791944]

Fig 1.

CT neuroimaging on presentation. Head CT without contrast demonstrating a large amount of subarachnoid hemorrhage within the basal cisterns with diffuse intraventricular hemorrhage and ventriculomegaly. **b** CT demonstrating $4 \times 6 \times 7$ -mm hematoma in the right perimesencephalic cistern with mass effect on the pons (arrow). There is blood within the basal cisterns, fourth ventricle, and cisterna magna. **c** CT angiogram windowed for arterial conspicuity demonstrating a small aneurysm in the right perimesencephalic region

Fig 2.

Cerebral angiography. Right vertebral artery angiogram with magnified frontal view of the posterior fossa demonstrating a 1,6-mm fusiform aneurysm of the right AICA in an inferior branch proximally (arrow), plus decreased cerebellar staining in the right medial cerebellum. **b** Right AICA angiogram with magnified frontal view of the posterior fossa demonstrating contrast extravasation from the AICA aneurysm. **c** Right vertebral artery angiogram with magnified frontal view of the posterior fossa after endovascular sacrifice of the inferior branch of the right AICA by Onyx-34 (Micro Therapeutics, Irvine CA) embolization showing complete opacification of the aneurysm and no evidence of residual extravasation. **d** Frontal view of Onyx-34 (Micro Therapeutics, Irvine CA) cast

Fig 3.

Post-operative CT neuroimaging. Diffuse intraventricular hemorrhage with subtle slightly worsened ventriculomegaly status post Onyx-34 (Micro Therapeutics, Irvine CA) embolization. There is midline shift to the right thalamus and brainstem due to hematoma. **b** Post-operative day 23, interval removal of right frontal approach EVD with increase in size of both lateral ventricles despite improvement of intraventricular hemorrhage. **c** Post-operative day 27, stable ventriculomegaly following placement of a new ventricular reservoir

Fig 4.

Follow-up MRI neuroimaging. Fifteen-month follow-up visit with T1- and T2-weighted images (left and right respectively) show interval increase in ventricular dimensions at the level of the medullary-pontine junction. The ventricular catheter was intact with tip in the left lateral ventricle(not shown)