Published in final edited form as:

J Am Chem Soc. 2019 October 23; 141(42): 16944–16953. doi:10.1021/jacs.9b09015.

Direct Manipulation of Metal Imido Geometry: Key Principles to Enhance C–H Amination Efficacy

Yunjung Baek, Elisabeth T. Hennessy, Theodore A. Betley*

Department of Chemistry and Chemical Biology, Harvard University, 12 Oxford Street, Cambridge, Massachusetts 02138, United States

Abstract

We report the catalytic C–H amination mediated by an isolable Co^{III} imido complex $(^{\text{Tr}}\text{L})\text{Co}(\text{NR})$ supported by a sterically demanding dipyrromethene ligand ($^{\text{Tr}}\text{L}$ = 5-mesityl-1,9-(trityl)dipyrin). Metalation of $(^{\text{Tr}}\text{L})\text{Li}$ with CoCl_2 in THF afforded a high-spin ($S = 3/2$) three-coordinate complex $(^{\text{Tr}}\text{L})\text{CoCl}$. Chemical reduction of $(^{\text{Tr}}\text{L})\text{CoCl}$ with potassium graphite yielded the high-spin ($S = 1$) Co^{I} synthon $(^{\text{Tr}}\text{L})\text{Co}$ which is stabilized through an intramolecular η^6 -arene interaction. Treatment of $(^{\text{Tr}}\text{L})\text{Co}$ with a stoichiometric amount of 1-azidoadamantane (AdN_3) furnished a three-coordinate, diamagnetic Co^{III} imide $(^{\text{Tr}}\text{L})\text{Co}(\text{NAd})$ as confirmed by single-crystal X-ray diffraction, revealing a rare trigonal pyramidal geometry with an acute $\text{Co}-\text{N}_{\text{imido}}-\text{C}$ angle $145.0(3)^\circ$. Exposure of 1–10 mol % of $(^{\text{Tr}}\text{L})\text{Co}$ to linear alkyl azides (RN_3) resulted in catalytic formation of substituted *N*-heterocycles via intramolecular C–H amination of a range of C–H bonds, including primary C–H bonds. The mechanism of the C–N bond formation was probed via initial rate kinetic analysis and kinetic isotope effect experiments [$k_{\text{H}}/k_{\text{D}} = 38.4(1)$], suggesting a stepwise H–atom abstraction followed by radical recombination. In contrast to the previously reported C–H amination mediated by $(^{\text{Ar}}\text{L})\text{Co}(\text{NR})$ ($^{\text{Ar}}\text{L}$ = 5-mesityl-1,9-(2,4,6- $\text{Ph}_3\text{C}_6\text{H}_2$)dipyrin), $(^{\text{Tr}}\text{L})\text{Co}(\text{NR})$ displays enhanced yields and rates of C–H amination without the aid of a cocatalyst, and no catalyst degradation to a tetrazene species was observed, as further supported by the pyridine inhibition effect on the rate of C–H amination. Furthermore, $(^{\text{Tr}}\text{L})\text{Co}(\text{NAd})$ exhibits an extremely low one-electron reduction potential ($E_{\text{red}}^\circ = -1.98 \text{ V}$ vs $[\text{Cp}_2\text{Fe}]^{+/0}$) indicating that the highly basic terminal imido unit contributes to the driving force for H–atom abstraction.

Graphical Abstract

*Corresponding Author betley@chemistry.harvard.edu.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.9b09015.

General experimental considerations and procedures, multinuclear NMR data, electrochemistry data, kinetic analysis, magnetometry data, solid state molecular structures, computational details, and crystallography data (PDF)

Crystallographic CIF files for complexes 1–4 (CIF)

The authors declare no competing financial interest.

1. INTRODUCTION

The development of a cost-effective and efficient transition-metal catalyst for direct C–H functionalization has been of interest to synthesize value-added commodity chemicals from simple hydrocarbon feedstocks.^{1,2} To this end, the use of bioinspired transition metal complexes featuring metal–ligand multiple bonds (MLMBs) has emerged as an appealing C–H functionalization methodology.^{3–5} In this regard, the iron-oxo complex (compound I) of Cytochrome P450 demonstrates nature’s key strategy to render the iron-oxo multiple bond highly reactive, thereby enhancing its O–atom transfer efficacy.⁶ Specifically, the open-shell electronic configuration, high oxidation state of compound I, and highly basic oxo unit of its one-electron reduced form have been proposed as leading features enabling facile C–H hydroxylation at ambient conditions.^{7,8}

As such, the isoelectronic synthetic analogues, metal nitrenoid complexes (e.g., imido, $M(NR^{2-})$;^{4,9–11} iminyl, $M(2NR^-)$;^{12–19} nitrene adducts, $M(3NR)^{20–23}$) have received great attention as viable N–group transfer reagents into unactivated C–H bonds. In particular, late, first-row transition metal-based nitrenoid complexes have been extensively studied for direct C–N bond formation (Mn,²⁴ Fe,^{18,19,25–34} Co,^{35–43} Ni,^{10,11,44,45} and Cu^{23,46,47}). Among the structurally characterized metal nitrenoid complexes, however, only a few manifest productive C–H amination due to the typically strong MLMBs between the metal ion and nitrene unit.^{33,34,38,39,44,46} Thus, in order to improve the efficacy of C–H amination, understanding the key factors governing the lability of the terminal nitrenoid moiety is necessary. Akin to the enzymatic C–H hydroxylation, assessing redox potentials and basicity of relevant species (e.g., imides or amides) involved in C–H amination processes operating via H–atom abstraction (HAA) is essential to comprehend the thermodynamic and kinetic properties of the desired transformations.^{48–54}

From a synthetic perspective, these properties (E_{red}° and pK_a) can be tuned by various elements such as identity or oxidation state of a metal center(s),⁵⁵ electronic configuration of metal nitrenoid complexes,^{19,33,56} and the local geometry of the terminal nitrenoid moiety.⁵⁷ Previously, we have demonstrated that manipulation of the spin state of the metal imide³³ and the spin density at the nitrogen atom of the nitrenoid unit¹⁹ can serve as a synthetic approach to weaken the MLMBs and enable facile C–H amination.

In contrast, the direct manipulation of the geometry of a metal nitrenoid and its effect on C–H amination efficacy is relatively less explored. Nonetheless, Holland and co-workers have proposed that the geometry of the terminally bound imido unit can alter the basicity of the metal imido complex. They found enhanced C–H activation reactivity from the β -diketiminate Fe^{III} imido complex upon pyridine coordination.^{57–59} The pyridine-bound iron imide was proposed as more bent and basic compared to the pyridine-free iron imide. However, structural characterization of the bent imide is absent, and thus, a direct correlation between the bent imido geometry and its basicity is lacking.

Recently, our group showcased that the geometry of the imido unit can also affect the rate of C–H amination.³⁸ We have demonstrated the intramolecular C–H amination mediated by (^{Ar}L)Co(NR) (^{Ar}L = 5-mesityl-1,9-(2,4,6-Ph₃C₆H₂)dipyrrin) and found that addition of pyridine improves both yields and rates of C–H amination. Exogenous pyridine coordinates to the trigonal planar (^{Ar}L)Co(NR) to generate the pyridine-bound imide (^{Ar}L)Co(NR)(py) providing steric protection to circumvent formation of a catalytically inactive cobalt tetrazido complex. More importantly, the rate of C–H amination was accelerated in the presence of pyridine as cocatalyst. We have ascribed the enhanced rate of C–H amination from (^{Ar}L)Co(NR)(py) to the bent and attenuated MLMB (Co–NR) as suggested by density functional theory (DFT) calculations, yet isolation of the bent (^{Ar}L)Co(NR)–(py) was hampered due to the weak binding affinity of pyridine.

Motivated by these precedents, we postulated that generating a metal-bound nitrenoid unit in a bent geometry can be an alternative strategy to engender a more reactive MLMB and thus improve C–H amination efficacy both kinetically and thermodynamically.⁵² Toward this, we sought to design a ligand that would enable synthesis of a cobalt imido complex featuring a significantly bent nitrenoid unit in the absence of exogenous ligands (e.g., pyridine). Specifically, we sought to investigate the following questions: (1) can we synthesize a dipyrrinato Co^{III} imido complex featuring a bent nitrenoid unit by modifying the substituents on the dipyrrin scaffold? If possible, (2) is the imide competent for catalytic C–H amination to afford substituted N-heterocycles? And (3) how does the geometry of the metal-bound nitrenoid moiety alter the efficacy and reactivity profile for C–H amination with respect to yields, rates, and reaction scope?

Toward this goal, we report herein the synthesis of a trityl-substituted dipyrromethene platform (^{Tr}L) (^{Tr}L = 5-mesityl-1,9-(trityl)dipyrrin) that provides access to a robust Co^I catalyst (^{Tr}L)Co and the corresponding bent Co^{III} imido complex (^{Tr}L)Co(NR), which enables a facile Co^{I/III} catalytic C–H amination. Detailed mechanistic studies on the C–H amination processes establish the improved C–H amination efficacy of (^{Tr}L)Co(NR) as compared to the previously reported linear (^{Ar}L)Co(NR), demonstrating the importance of modulating the primary coordination sphere to effect more reactive and synthetically applicable C–H amination catalysis.

2. RESULTS AND DISCUSSION

2.1. Synthesis and Characterization of Cobalt Synthons.

As discussed in the Introduction, the ligand design principle to synthesize a bent Co^{III} imide was informed by the previously studied ligand-accelerated C–H amination processes mediated by (ArL)Co(NR) (ArL = 5-mesityl-1,9-(2,4,6-Ph₃ C₆ H₂)dipyrin).³⁸ Specifically, we sought to emulate the primary coordination environment of (ArL)Co(NR)(py) by modifying the flanking groups of the dipyrin scaffold. To this end, trityl moieties were considered to be more voluminous ligand flanking units than the 2,4,6-triphenylphenyl substituents of (ArL). Thus, we envisioned that the steric profile of the trityl groups may enforce a bent geometry on the nitrenoid unit and eliminate formation of cobalt tetrazido complexes even in the absence of exogenous pyridine. Additionally, we proposed that one of the phenyl rings of the trityl substituents may stabilize the low-coordinate cobalt center via an intramolecular η^6 -arene interaction affording a robust Co^I synthon akin to the previously reported iron and cobalt synthons such as (ArL)Fe³³ and (ArL)Co.³⁸

With these design principles in mind, we first prepared 2-trityl-1*H*-pyrrole via Friedel–Crafts alkylation of excess pyrrole with trityl chloride to avoid bis-alkylation. Following literature procedures,¹⁸ subsequent condensation with mesitylaldehyde dimethyl acetal followed by oxidation with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) afforded (TrL)H (Scheme 1a). Subsequent deprotonation of (TrL)H with PhLi yielded the lithiated scaffold (TrL)Li. Metalation of (TrL)Li with CoCl₂ in THF for 20 h at room temperature afforded a three-coordinate complex (TrL)CoCl (**1**) (Scheme 1b). Single crystals of **1** suitable for X-ray diffraction were grown from a layered 1:2 benzene/hexanes solution at room temperature. In contrast to the previously reported trigonal planar Co^{II} complex (ArL)CoBr, **1** adopts a trigonal pyramidal geometry, displaying an η^1 -arene interaction between the cobalt and one of the *ortho* carbons from the trityl substituents (Figure 1a). The discrete η^1 -arene interaction [Co–C_{ortho} = 2.345(7) Å] is of note as it occupies the equatorial position of the dipyrin plane positioning the chloride on the axial site. Solid-state magnetic susceptibility of **1** was measured with super-conducting quantum interference device magnetometry (SQUID) and is consistent with a high-spin ground state (*S* = 3/2) (Figure 2a).

Chemical reduction of **1** with KC₈ in a thawing benzene solution cleanly furnished the reduced species (TrL)Co (**2**), which exhibits an intramolecular η^6 -arene interaction between the cobalt and one of the phenyl groups of the trityl substituents (Scheme 1b, Figure 1b). The solid-state magnetic susceptibility of **2** is consistent with an *S* = 1 high-spin ground state (Figure 2b). The cyclic voltammogram of **2** measured in a non-coordinating solvent 1,2-difluorobenzene reveals two sequential one-electron oxidations (*E*_{1/2} = –0.80, 0.75 V vs [Cp₂Fe]⁺⁰) and a single one-electron reduction (–2.1 V, Figure 3a). The observed redox potentials are almost identical with the ones observed for its congener (ArL)Co, suggesting that **2** is a suitable alternative Co^I synthon to further explore two-electron oxidative group transfer reactivity (Figure 3b).

2.2. Synthesis and Characterization of (TrL)Co(NAd).

With the (TrL)Co^I synthon in hand, we sought to investigate its reactivity toward organoazides. Treating a benzene-*d*₆ solution of **2** with a stoichiometric amount of 1-azidoada-mantane (AdN₃) immediately furnished a diamagnetic species as verified by ¹H and ¹³C NMR spectroscopies. Single crystals of the product obtained from a 1:1 benzene/hexanes mixture stored at room temperature were analyzed by single crystal X-ray diffraction to reveal formation of a three-coordinate Co^{III} imido complex (TrL)Co(NAd) (**3**) (Scheme 2 and Figure 1c–d).

Notably, **3** exhibits a unique distorted trigonal pyramidal geometry with a significantly bent imido moiety [$\angle\text{Co-N-C} = 145.0(3)^\circ$]. Akin to **1**, a relatively long η^1 -arene interaction between the cobalt and one of the *ortho* carbons from the trityl groups is present ($\text{Co-C}_{ortho} = 2.756 \text{ \AA}$) orienting the nitrenoid unit orthogonal to the dipyrin plane. Furthermore, the sterically encumbered trityl substituents shield the trans position of the nitrenoid moiety, enforcing an acute $\angle\text{L-Co-NAd}$ (119.7°) as well as blocking the fourth coordination site of **3** as further evidenced by the space-filling model (Figure S–28). Consequently, no bimolecular reactivity affording the four-coordinate tetrazido complex was detected upon heating a benzene-*d*₆ solution of **3** with excess amount of AdN₃ (Scheme 2).

We note that only a handful of structurally characterized metal imides display a substantially bent nitrenoid unit. For late, first-row transition metals, a four-coordinate bis(pyrrolyl)-pyridine Fe^{IV} imide⁶¹ and five-coordinate Fe^{IV} imide⁶² supported by a tetradentate N-heterocyclic (NHC) ligand display an Fe–N–C angle of $140.5(3)^\circ$ and $150.07(10)^\circ$, respectively. Hillhouse and co-workers have shown the synthesis of a three-coordinate nickel imide featuring a bent nitrenoid moiety [$\angle\text{Ni-N-C} = 127.3(3)^\circ$] using a sterically encumbered aryl azide (e.g., 2,5-dimesitylphenylazide, 2,6-bis(2,6-diisopropylphenyl)phenylazide).⁴⁵ Given that those bent M–N–C linkages were induced by either multidentate chelating ligands or steric properties of the R substituents on the metal-bound imido (NR) group, furnishing the bent imido complex **3** via direct manipulation of the primary coordination environment with the low-coordinate bidentate dipyrin ligand is significant.

Having isolated the bent imido **3**, we sought to understand its electronic structure. As discussed, the diamagnetic spin ground state of **3** is further supported by the temperature-independent susceptibility data (Figure 2c), which is consistent with the vast majority of Co^{III} imides including the analogous dipyrin alkyl Co^{III} imide (ArL)Co(NR) reported by our group.^{37–43} However, unlike (ArL)Co(NR), no thermal population of spin excited states (e.g., $S = 0 \rightarrow 2$) was observed (Figure 2c inset).³⁷

To further elucidate the frontier molecular orbital interactions in **3**, DFT calculations were conducted (SI–59).^{63,64} Interestingly, owing to the bent geometry, HOMO, HOMO – 1, and HOMO – 2 of **3** display almost non-bonding or significantly disrupted π^* interactions with respect to the imido unit. Instead, σ^* or π^* contributions from the dipyrin become dominant. The non-bonding interaction with $3d_{x^2-y^2}$ and the slight π -overlap between the 2p orbitals of the N_{imido} and Co $3d_z^2$, $3d_{yz}$, and $3d_{xy}$ are illustrated in Figure 4.

2.3. Catalytic C–H Amination.

2.3.1. Steric Protection from (^{Tr}L) Leads to Higher Yields of C–H Amination.

—Encouraged by the distorted MLMB of **3** and its steric shielding to prevent the catalyst degradation, we speculated that analogous (^{Tr}L)Co(NR) species featuring a linear alkyl moiety would greatly benefit the intramolecular C–H amination to afford *N*-hetetrocycles. To test this hypothesis, (4-azido-4-methylpentyl) benzene (**5**) was exposed to 10 mol % of **2** in benzene-*d*₆. Gratifyingly, the desired product 2,2-dimethyl-5-phenylpyrrolidine (**6**) was quantitatively obtained at room temperature along with regenerated catalyst **2** (Figure S–2). We note that under the same conditions ([**5**] = 85.0 mM at 25 °C), 10 mol % of (^{Ar}L)Co yielded only 13% of **6** due to concomitant formation of a catalytically inactive tetrazido complex (^{Ar}L)Co(κ^2 -N₄R₂, R = C(CH₃)₂(CH₂)₃Ph),³⁸ highlighting that absence of the off-cycle pathway is critical to improve the yields of C–H amination. As a result, even with 1 mol % of **2** in benzene-*d*₆, **6** was obtained in 91% yield at 80 °C.

The steric protection of the trityl substituents is further supported by the effect of pyridine on the rates of catalytic C–H amination. Given that the fourth-coordination site of **3** is blocked by the trityl substituents, we hypothesized that pyridine would behave as an inhibitor as opposed to an accelerant in contrast with the C–H amination mediated by (^{Ar}L)Co(NR). Indeed, unlike (^{Ar}L)Co(NR), which binds 1 equiv of pyridine in solution,³⁸ **3** showed no binding affinity toward pyridine. Moreover, at 50 °C, slower rates of C–H amination were observed when exposing **5** to 10 mol % of **2** at higher concentrations of pyridine-*d*₅ (Figure S-7). A double-reciprocal plot suggests that pyridine-*d*₅ and **5** compete for the same coordination site of **2**,⁶⁵ and thus, the desired C–H amination is hampered in the presence of pyridine-*d*₅ (Figures 5a and S-9). Job plot analysis further supports that **2** readily binds pyridine in benzene-*d*₆ to furnish the (^{Tr}L)Co(py) (**4**) adduct (Scheme 2 and Figure 5b).⁶⁶ Furthermore, single-crystal X-ray diffraction analysis of **4** verifies that pyridine occupies the same axial position as the imido moiety of **3** (Figure 5c). The solid-state structure of **4** displays a η^2 -arene interaction with one of the *ortho* and *meta* carbons from the trityl groups [Co–C_{ortho} = 2.0520(8) Å and Co–C_{meta} = 2.239(8) Å]. These results are in line with the aforementioned steric environment of **3** suggesting that the fourth coordination site of (^{Tr}L)Co(NR) is no longer available.

Thus, by simply switching the flanking groups of the Co^I catalyst from the 2,4,6-triphenylphenyl to trityl, pyridine becomes a competitive inhibitor as opposed to a cocatalyst for the desired C–H amination. These results demonstrate that the pyridine ligation effect precluding tetrazene formation, which is observed for the previously studied amination catalysis with the (^{Ar}L) scaffold is successfully incorporated within the (^{Tr}L) ligand. Furthermore, considering that many other attempted N-group transfer reactions utilizing organoazides as a nitrene source are hampered by the reductive coupling of organoazides furnishing tetrazene^{37,38,67} or hexazene⁶⁸ complexes, we believe that the study described herein provides a ligand design to circumvent loss of reactive metal nitrenoid intermediates.

2.3.2. Mechanism of the C–H Amination.—Next, we sought to investigate the mechanism of C–H amination mediated by (^{Tr}L)Co(NR) and explore how the unusual

geometry of the cobalt imide affects its C–H amination reactivity as compared to that of (^{Ar}L)Co(NR).

To this end, the reaction progression was monitored by ¹H NMR spectroscopy (Figure 6a). In the presence of a 10-fold excess amount of azide substrate **5**, (^{Tr}L)Co (**2**) immediately converted to the corresponding Co^{III} imide (**7**) at 25 °C as verified by the diagnostic ¹H NMR chemical shifts (Figure S-2). At early stages of the reaction, imido **7** is the resting species as the concentration of azide **5** is sufficient to continuously oxidize **2** as soon as **7** generates pyrrolidine **6** (Figure 6b inset). Indeed, the rate of the azide consumption ($-d[\mathbf{5}]/dt$) is almost identical to the rate of product formation ($d[\mathbf{6}]/dt$) at initial stages of the catalytic reaction (SI-17). Thus, while the concentration of **7** is constant at early stages of the reaction (**6** < 10%), the rate of formation of **6** is zeroth order in [**5**] and first order in [**7**], corroborating that **7** is competent for C–H amination in a unimolecular fashion (Figure 6b–c).

Furthermore, a kinetic isotope effect (KIE) value was evaluated by comparing the initial rates of 10 mol % of **2** with **5** and (4-azido-4-methylpentyl-1,1-*d*₂) benzene (**5-d**₂). The observed ratio of the rate constants (k_H/k_D) was determined to be 38.4(1) at 25 °C (Figure 7), consistent with a rate-determining homolytic C–H bond cleavage.⁶⁹ Given all the kinetic data, we propose the following Co^{I/III} catalytic cycle (Scheme 3): fast oxidation of **2** by an alkyl azide affords a Co^{III} imide which undergoes intramolecular HAA; subsequent radical recombination of the resulting Co^{II} amide affords the cyclized products. However, we note that a direct C–H insertion pathway from the Co imido complex cannot be fully excluded.

2.3.3. Enhancement in Rates of C–H Amination.—Surprisingly, the rate of C–H amination mediated by (^{Tr}L)Co(NR) is significantly faster compared to (^{Ar}L)Co(NR). Specifically, the rate constant (*k*) for the benzylic C–H amination mediated by **7** to afford **6** is 0.063 min^{−1} at 25 °C, whereas the same magnitude of *k* = 0.062 min^{−1} was observed for the analogous (^{Ar}L)Co(NR) at 50 °C.³⁸ We propose the faster conversion is a result of the bent imido moiety of **7** (as observed for **3**) that likely resembles the transition state geometry for HAA more closely than the linear (^{Ar}L)Co(NR). Therefore, the enhanced C–H amination rate of **7** supports our original hypothesis, where a bent nitrenoid unit may enhance the rate of C–H amination as less structural rearrangement is required for HAA. Overall, these results highlight that a bent MLMB linkage, which is enforced by the (^{Tr}L) scaffold, is critical to enhance the rate of C–H amination, thus kinetically improving the overall C–H amination process.

2.3.4. Scope of C–H Amination.—In addition to the enhanced rates, (^{Tr}L)Co(NR) is capable of functionalizing a range of C–H bonds, even primary C–H bonds in a catalytic fashion (Table 1). Previously, we were only able to functionalize benzylic C–H bonds from the analogous three-coordinate metal imides M(NR^{2−}): (^{Ar}L)Fe(NAd)³³ and (^{Ar}L)Co(NR).³⁸ Thus, the ability of (^{Tr}L)Co(NR) to functionalize C–H bonds stronger than a benzylic C–H bond catalytically is noteworthy. As such, we sought to address the elements rendering (^{Tr}L)Co(NR) a more potent oxidant relative to the analogous (^{Ar}L)Co(NR). Given that both imides (^{Tr}L)Co(NR) and (^{Ar}L)Co(NR) accomplish intramolecular C–H amination via an initial rate-limiting HAA step, as suggested by the measured KIE values,³⁸ we attempted to

evaluate thermodynamic parameters such as (E_{red}° and $\text{p}K_{\text{a}}$ as described in eq 1, where the constant C represents the standard potential for proton reduction in aprotic solvent.⁴⁹ We hypothesized that the difference in driving force for HAA from the respective imides is proportional to the difference in bond dissociation energies (BDEs) of the N–H bonds of the corresponding Co^{II} amide complexes (Scheme 4).

$$\text{BDE of N-H (kcal/mol)} = 23.06E_{\text{red}}^{\circ} + 1.37\text{p}K_{\text{a}} + C \quad (1)$$

To this end, we used **3** and its isolable congener ($^{\text{Ar}}\text{L}$)Co(NAd) (**8**) as model complexes to probe their one-electron reduction potentials (E_{red}° values for $\text{Co}^{\text{III/II}}$ couples) via electrochemical analysis. Cyclic voltammetry experiments of **3** and **8** were conducted using 0.1 M [$^{\text{t}}\text{Bu}_4\text{N}$][PF₆] electrolyte in the non-coordinating solvent 1,2-difluorobenzene to obviate exogenous ligand coordination that may affect the E_{red}° . Interestingly, both **3** and **8** display reversible one electron reductions at $E_{1/2}$ of -1.98 and -1.93 V (vs [Cp₂Fe]^{+/0}), respectively (Figure 8).

The observed low reduction potentials suggest that the driving force for HAA can be ascribed to the high basicity of the one-electron reduced anionic imides [$\text{LCo}(\text{NR})$][−], which are equivalent to the conjugate base of amides [$\text{LCo}(\text{NHR})$] **3a** and **8a**, respectively (Scheme 4).⁵⁵ Furthermore, given the almost identical E_{red}° values of **3** and **8**, we believe that the greater oxidizing power of ($^{\text{Tr}}\text{L}$)Co(NR) compared to ($^{\text{Ar}}\text{L}$)Co(NR) arises from a more basic imido moiety of its one-electron reduced form [$(^{\text{Tr}}\text{L})\text{Co}(\text{NR})$][−].

Unfortunately, our attempts to directly probe the $\text{p}K_{\text{a}}$ of the respective amide complexes ($^{\text{Tr}}\text{L}$)Co(NHAd) (**3a**) and ($^{\text{Ar}}\text{L}$)Co(NHAd) (**8a**) were unsuccessful due to their high propensity for subsequent HAA resulting in formation of 1-adamantylamine.⁵² Instead, the difference in $\text{p}K_{\text{a}}$ between **3a** and **8a** was estimated by accounting for the difference in BDEs of the C–H bonds (6 kcal/mol) that can be activated by the respective imides intramolecularly at 25 °C. Specifically at 25 °C, ($^{\text{Tr}}\text{L}$)Co(NR) accomplishes amination of secondary C–H bonds (96 kcal/mol, R = C(CH₃)₂(CH₂)₃CH₃) affording 2,2,5-trimethylpyrrolidine, whereas ($^{\text{Ar}}\text{L}$)Co(NR) can only functionalize benzylic C–H bonds (90 kcal/mol, R = C(CH₃)₂(CH₂)₃Ph) to afford **6**. Accordingly, the conjugate base of **3a** [$(^{\text{Tr}}\text{L})\text{Co}(\text{NR})$][−] is estimated to be about 10⁵ times more basic than the conjugate base of **8a** [$(^{\text{Ar}}\text{L})\text{Co}(\text{NR})$][−]. The observed difference in $\text{p}K_{\text{a}}$ is further supported by DFT calculations,⁵² yet evaluation of the absolute $\text{p}K_{\text{a}}$ of each amide complex needs to be further examined (SI-41).

This observed trend in basicity is in agreement with literature reports proposing that a bent imido moiety is more basic compared to a linear one.⁵⁷ Thus, these results demonstrate that altering the local geometry of the terminally bound metal imido unit subsequently affects its basicity as well and, thereby, enhances its oxidizing power and broadens its C–H amination reaction scope.

3. CONCLUSIONS

The foregoing results illustrate how rational ligand design allowed the synthesis of a robust Co^{I} catalyst and isolation of a bent Co^{III} imide that promotes facile catalytic C–H amination. The previously studied pyridine-accelerated C–H amination process mediated by $(^{\text{Ar}}\text{L})\text{Co}(\text{NR})$ informed the design principles for the synthesis of a dipyrromethene ligand ($^{\text{Tr}}\text{L}$) featuring sterically encumbered trityl substituents. The subtle modification of the supporting ligand drastically altered the primary coordination environment of a Co^{III} imide to force the terminal imido unit perpendicular to the dipyrin ligand plane, imposing a bent geometry. Kinetic and thermodynamic analyses of the C–H amination mediated by either $(^{\text{Tr}}\text{L})\text{Co}(\text{NR})$ or $(^{\text{Ar}}\text{L})\text{Co}(\text{NR})$ highlight that the unique structure of $(^{\text{Tr}}\text{L})\text{Co}(\text{NR})$ is key for the significantly enhanced C–H amination efficacy as manifested in yields, rates, and oxidizing power. In particular, the sterically demanding trityl substituents prevent catalyst degradation (via tetrazene formation), contributing to higher yields of C–H amination. More importantly, the atypical trigonal pyramidal geometry of the Co^{III} imide with a significantly bent imide resembles the transition state for HAA more closely, and thereby, $(^{\text{Tr}}\text{L})\text{Co}(\text{NR})$ displays enhanced rates of C–H amination compared to the trigonal planar $(^{\text{Ar}}\text{L})\text{Co}(\text{NR})$ possessing a linear imido geometry. Lastly, the more bent, and thus more basic, one-electron reduced species of $(^{\text{Tr}}\text{L})\text{Co}(\text{NR})$ provides a greater thermodynamic driving force for HAA enabling C–H amination with a broader substrate scope at milder conditions. We believe that this study provides an alternative synthetic strategy to generate highly reactive metal imido complexes via direct manipulation of the geometry of MLMBs to accomplish more applicable catalytic transformations.

4. EXPERIMENTAL SECTION

General experimental considerations, synthesis and spectroscopic characterizations, X-ray diffraction techniques, kinetic analysis, electrochemistry measurements, and DFT results are all provided in the Supporting Information.

4.1. Preparation of $(^{\text{Tr}}\text{L})\text{CoCl}$ (**1**).

In an oven-dried 20 mL vial, anhydrous CoCl_2 (0.105 g, 0.81 mmol, 1.2 equiv) was dissolved in THF and frozen in a liquid nitrogen cooled cold-well. To a frozen solution, a solution of $(^{\text{Tr}}\text{L})\text{Li}$ (0.52 g, 0.68 mmol, 1 equiv) in THF was added and stirred for 20 h at room temperature. The resulting dark red reaction mixture was filtered through a coarse glass frit with Celite to remove LiCl and the excess CoCl_2 . The Celite was washed with an additional 20 mL benzene. The collected filtrate was frozen and benzene was removed by sublimation *in vacuo* to afford **1** as a dark red powder (0.52 g, 90%). Crystals suitable for X-ray diffraction were grown from a 2:1 *n*-hexane/benzene layered solution at room temperature. ^1H NMR (600 MHz, 295 K, C_6D_6): δ /ppm 68.14, 23.15, 20.56, 4.70, 0.64, –1.16, –10.13, –37.86. Anal. calc. for $\text{C}_{56}\text{H}_{45}\text{ClCoN}_2$: C 80.04, H 5.40, N 3.33; Found: C 80.00, H 5.29, N 3.45.

4.2. Preparation of (^{Tr}L)Co (2).

In an oven-dried 20 mL vial, **1** (0.4 g, 0.476 mmol, 1 equiv) was dissolved in 10 mL of benzene and added to a frozen solution of potassium graphite (KC₈, 0.071 g, 0.524 mmol, 1.1 equiv). The reaction mixture was slowly warmed up to room temperature and stirred for 3 h at room temperature. The dark red-orange mixture was filtered through a coarse glass frit with Celite to remove graphite. The Celite was washed with an additional 20 mL benzene. The collected filtrate was frozen and benzene was removed by sublimation *in vacuo* to afford **2** as a dark orange powder. (0.3 g, 78%). Crystals suitable for X-ray diffraction were grown from a 1:1 *n*-hexane/benzene mixture at room temperature. ¹H NMR (600 MHz, 295 K, C₆D₆): δ/ppm 49.82, 21.04, 14.90, 5.84, 5.16, 4.54, 0.96, -2.51. Anal. calc. for C₅₆H₄₅CoN₂: C 83.56, H 5.64, N 3.48; Found: C 83.45, H 5.50, N 3.39.

4.3. Preparation of (^{Tr}L)Co(NAd) (3).

In an oven-dried 20 mL vial, a solution of 1-azidoadamantane (22 mg, 0.12 mmol, 1.0 equiv) in benzene was added to a stirring solution of **2** (100 mg, 0.12 mmol, 1.0 equiv) in benzene to observe an immediate color change from dark red to dark purple. After stirring for 30 min at room temperature, the reaction mixture was frozen, and benzene was removed by sublimation *in vacuo* to afford **3** as a purple powder (109 mg, 92%). Crystals suitable for X-ray diffraction were grown from a 1:1 *n*-hexane/benzene mixture at room temperature. ¹H NMR (600 MHz, 295 K, C₆D₆): δ/ppm 7.94 (d, 12 H), 7.23 (d, 2 H), 6.97 (t, 6 H), 6.84 (s, 2 H), 6.67 (t, 11 H), 6.03 (d, 2 H), 2.26 (s, 5 H), 2.20 (s, 3 H), 1.80 (s, 3 H), 1.63 (t, 6 H), 1.08 (s, 6 H). ¹³C{¹H} NMR (125MHz, C₆D₆): δ/ppm 192.23, 153.36, 151.26, 146.88, 137.28, 135.37, 134.93, 132.14, 129.57, 128.59, 125.51, 124.41, 69.27, 61.65, 61.12, 37.54, 33.30, 31.03, 21.25, 20.69. Anal. calc. for C₆₆H₆₀CoN₃: C 83.08, H 6.34, N 4.40; Found: C 82.72, H 6.52, N 4.71.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

T.A.B. acknowledges this work was supported by a grant from the NIH (GM-115815), the Dreyfus Foundation (Teacher-Scholar Award), and Harvard University. Y.B. is thankful for a Kwanjeong Graduate Student Fellowship.

REFERENCES

- (1). Labinger JA; Bercaw JE Understanding and exploiting C–H bond activation. *Nature* 2002, 417, 507. [PubMed: 12037558]
- (2). Bergman RG C–H activation. *Nature* 2007, 446, 391. [PubMed: 17377575]
- (3). Davies HML; Manning JR Catalytic C–H functionalization by metal carbenoid and nitrenoid insertion. *Nature* 2008, 451, 417. [PubMed: 18216847]
- (4). Saouma CT; Peters JC M≡E and M=E complexes of iron and cobalt that emphasize three-fold symmetry (E = O, N, NR). *Coord. Chem. Rev* 2011, 255 (7), 920–937. [PubMed: 21625302]
- (5). Nugent WA; Mayer JM Metal-ligand multiple bonds: the chemistry of transition metal complexes containing oxo, nitrido, imido, or alkylidyne ligands; Wiley-Interscience: New York, 1988.
- (6). Ortiz de Montellano PR Cytochrome P450: structure, mechanism, and biochemistry, 4th ed.; Kluwer Academic/Plenum Publishers: New York, 2005.

- (7). Hill EA; Weitz AC; Onderko E; Romero-Rivera A; Guo Y; Swart M; Bominaar EL; Green MT; Hendrich MP; Lacy DC; Borovik AS Reactivity of an Fe^{IV}-oxo complex with protons and oxidants. *J. Am. Chem. Soc* 2016, 138 (40), 13143–13146. [PubMed: 27647293]
- (8). Green MT; Dawson JH; Gray HB Oxoiron(IV) in chloroperoxidase compound II is basic: implications for P450 chemistry. *Science* 2004, 304 (5677), 1653. [PubMed: 15192224]
- (9). Zhang L; Deng L. C–H bond amination by iron-imido/nitrene species. *Chin. Sci. Bull* 2012, 57 (19), 2352–2360.
- (10). Waterman R; Hillhouse GL η^2 -Organoazide complexes of nickel and their conversion to terminal imido complexes via dinitrogen extrusion. *J. Am. Chem. Soc* 2008, 130 (38), 12628–12629. [PubMed: 18729364]
- (11). Mindiola DJ; Hillhouse GL Terminal amido and imido complexes of three-coordinate nickel. *J. Am. Chem. Soc* 2001, 123 (19), 4623–4624. [PubMed: 11457258]
- (12). Olivos Suarez AI; Jiang H; Zhang XP; de Bruin B. The radical mechanism of cobalt(II) porphyrin-catalyzed olefin aziridination and the importance of cooperative H-bonding. *Dalton Trans.* 2011, 40 (21), 5697–5705. [PubMed: 21483935]
- (13). Kuijpers PF; van der Vlugt JI; Schneider S; de Bruin B. Nitrene radical intermediates in catalytic synthesis. *Chem. - Eur. J* 2017, 23 (56), 13819–13829. [PubMed: 28675476]
- (14). Lyaskovskyy V; Suarez AIO; Lu H; Jiang H; Zhang XP; de Bruin B. Mechanism of cobalt(II) porphyrin-catalyzed C–H amination with organic azides: radical nature and H–atom abstraction ability of the key cobalt(III)–nitrene intermediates. *J. Am. Chem. Soc* 2011, 133 (31), 12264–12273. [PubMed: 21711027]
- (15). Suarez AIO; Lyaskovskyy V; Reek JNH; van der Vlugt JI; de Bruin B. Complexes with nitrogen-centered radical ligands: classification, spectroscopic features, reactivity, and catalytic applications. *Angew. Chem., Int. Ed* 2013, 52 (48), 12510–12529.
- (16). Kuijpers PF; Tiekink MJ; Breukelaar WB; Broere DLJ; van Leest NP; van der Vlugt JI; Reek JNH; de Bruin B. Cobalt-porphyrin-catalysed intramolecular ring-closing C–H amination of aliphatic azides: a nitrene-radical approach to saturated heterocycles. *Chem. - Eur. J* 2017, 23 (33), 7945–7952. [PubMed: 28332743]
- (17). Goswami M; Lyaskovskyy V; Domingos SR; Buma WJ; Woutersen S; Troeppner O; Ivanovi - Burmazovi I; Lu H; Cui X; Zhang XP; Reijerse EJ; DeBeer S; van Schooneveld MM; Pfaff FF; Ray K; de Bruin B. Characterization of porphyrin-Co(III)-'nitrene radical' species relevant in catalytic nitrene transfer reactions. *J. Am. Chem. Soc* 2015, 137 (16), 5468–5479. [PubMed: 25844713]
- (18). King ER; Hennessy ET; Betley TA Catalytic C–H bond amination from high-spin iron imido complexes. *J. Am. Chem. Soc* 2011, 133 (13), 4917–4923. [PubMed: 21405138]
- (19). Wilding MJT; Iovan DA; Wrobel AT; Lukens JT; MacMillan SN; Lancaster KM; Betley TA Direct comparison of C–H bond amination efficacy through manipulation of nitrogen-valence centered redox: imido versus iminyl. *J. Am. Chem. Soc* 2017, 139 (41), 14757–14766. [PubMed: 28937756]
- (20). Conradie J; Ghosh A. Electronic structure of an iron-porphyrin–nitrene complex. *Inorg. Chem* 2010, 49 (1), 243–248. [PubMed: 19994873]
- (21). Corona T; Ribas L; Rovira M; Farquhar ER; Ribas X; Ray K; Company A. Characterization and reactivity studies of a terminal copper–nitrene species. *Angew. Chem., Int. Ed* 2016, 55 (45), 14005–14008.
- (22). Berry JF Terminal nitrido and imido complexes for the late transition metals. *Comments Inorg. Chem* 2009, 30 (1–2), 28–66.
- (23). Carsch KM; DiMucci IM; Iovan DA; Li A; Zheng S-L; Titus CJ; Lee SJ; Irwin KD; Nordlund D; Lancaster KM; Betley TA Synthesis of a copper-supported triplet nitrene complex pertinent to copper-catalyzed amination. *Science* 2019, 365 (6458), 1138–1143. [PubMed: 31515388]
- (24). Zdilla MJ; Dexheimer JL; Abu-Omar MM Hydrogen atom transfer reactions of imido manganese(V) corrole: one reaction with two mechanistic pathways. *J. Am. Chem. Soc* 2007, 129 (37), 11505–11511. [PubMed: 17718564]

- (25). Brown SD; Betley TA; Peters JC A Low-spin d5 iron imide: nitrene capture by low-coordinate iron(I) provides the 4-coordinate Fe(III) complex [PhB(CH2PPh2)3]Fe≡N-p-tolyl. *J. Am. Chem. Soc* 2003, 125 (2), 322–323. [PubMed: 12517130]
- (26). Brown SD; Peters JC Ground-state singlet L3Fe-(μ-N)-FeL3 and L3Fe(NR) complexes featuring pseudotetrahedral Fe(II) centers. *J. Am. Chem. Soc* 2005, 127 (6), 1913–1923. [PubMed: 15701026]
- (27). Betley TA; Peters JC Dinitrogen chemistry from trigonally coordinated ion and cobalt platforms. *J. Am. Chem. Soc* 2003, 125 (36), 10782–10783. [PubMed: 12952446]
- (28). Thomas CM; Mankad NP; Peters JC Characterization of the terminal iron(IV) imides {[PhBPtBu2(pz')][FeIV:NAd]}+. *J. Am. Chem. Soc* 2006, 128 (15), 4956–4957. [PubMed: 16608321]
- (29). Bart SC; Lobkovsky E; Bill E; Chirik PJ Synthesis and hydrogenation of bis(imino)pyridine iron imides. *J. Am. Chem. Soc* 2006, 128 (16), 5302–5303. [PubMed: 16620076]
- (30). Bowman AC; Milsmann C; Bill E; Turner ZR; Lobkovsky E; DeBeer S; Wieghardt K; Chirik PJ Synthesis and electronic structure determination of N-alkyl-substituted bis(imino)pyridine iron imides exhibiting spin crossover behavior. *J. Am. Chem. Soc* 2011, 133 (43), 17353–17369. [PubMed: 21985461]
- (31). Cowley RE; DeYonker NJ; Eckert NA; Cundari TR; DeBeer S; Bill E; Ottenwaelder X; Flaschenriem C; Holland PL Three-coordinate terminal imidoiron(III) complexes: structure, spectroscopy and mechanism of formation. *Inorg. Chem* 2010, 49 (13), 6172–6187. [PubMed: 20524625]
- (32). Wang L; Hu L; Zhang H; Chen H; Deng L. Three-coordinate iron(IV) bisimido complexes with aminocarbene ligation: synthesis, structure, and reactivity. *J. Am. Chem. Soc* 2015, 137 (44), 14196–14207. [PubMed: 26505122]
- (33). Wilding MJT; Iovan DA; Betley TA High-spin iron imido complexes competent for C–H bond amination. *J. Am. Chem. Soc* 2017, 139 (34), 12043–12049. [PubMed: 28777558]
- (34). Iovan DA; Betley TA Characterization of iron-imido species relevant for N-group transfer chemistry. *J. Am. Chem. Soc* 2016, 138 (6), 1983–1993. [PubMed: 26788747]
- (35). Du J; Wang L; Xie M; Deng L. A two-coordinate cobalt(II) imido complex with NHC ligation: synthesis, structure, and reactivity. *Angew. Chem., Int. Ed* 2015, 54 (43), 12640–12644.
- (36). Liu Y; Du J; Deng L. Synthesis, structure, and reactivity of low-spin cobalt(II) imido complexes [(Me3P)3Co(NAr)]. *Inorg. Chem* 2017, 56 (14), 8278–8286. [PubMed: 28661656]
- (37). King ER; Sazama GT; Betley TA Co(III) imidos exhibiting spin crossover and C–H bond activation. *J. Am. Chem. Soc* 2012, 134 (43), 17858–17861. [PubMed: 23043624]
- (38). Baek Y; Betley TA Catalytic C–H amination mediated by dipyrin cobalt imidos. *J. Am. Chem. Soc* 2019, 141 (19), 7797–7806. [PubMed: 31016975]
- (39). Shay DT; Yap GPA; Zakharov LN; Rheingold AL; Theopold KH Intramolecular C–H activation by an open-shell cobalt(III) imido complex. *Angew. Chem., Int. Ed* 2005, 44 (10), 1508–1510.
- (40). Hu X; Meyer K. Terminal cobalt(III) imido complexes supported by tris(carbene) ligands: imido insertion into the cobalt–carbene bond. *J. Am. Chem. Soc* 2004, 126 (50), 16322–16323. [PubMed: 15600324]
- (41). Jones C; Schulten C; Rose RP; Stasch A; Aldridge S; Woodul WD; Murray KS; Moubaraki B; Brynda M; La Macchia G; Gagliardi L. Amidinato- and guanidinato-cobalt(I) complexes: characterization of exceptionally short Co–Co interactions. *Angew. Chem., Int. Ed* 2009, 48 (40), 7406–7410.
- (42). Dai X; Kapoor P; Warren TH [Me2NN]Co(η6-toluene): O=O, N=N, and O=N bond cleavage provides β-diketiminato cobalt μ-oxo and imido complexes. *J. Am. Chem. Soc* 2004, 126 (15), 4798–4799. [PubMed: 15080682]
- (43). Jenkins DM; Betley TA; Peters JC Oxidative group transfer to Co(I) affords a terminal Co(III) imido complex. *J. Am. Chem. Soc* 2002, 124 (38), 11238–11239. [PubMed: 12236716]
- (44). Laskowski CA; Miller AJM; Hillhouse GL; Cundari TR A Two-coordinate nickel imido complex that effects C–H amination. *J. Am. Chem. Soc* 2011, 133 (4), 771–773. [PubMed: 21175213]
- (45). Harrold ND; Hillhouse GL Strongly bent nickel imides supported by a chelating bis(N-heterocyclic carbene) ligand. *Chem. Sci* 2013, 4 (10), 4011–4015.

- (46). Aguila MJB; Badiei YM; Warren TH Mechanistic insights into C–H amination via dicopper nitrenes. *J. Am. Chem. Soc* 2013, 135 (25), 9399–9406. [PubMed: 23656170]
- (47). Bakhoda A; Jiang Q; Bertke JA; Cundari TR; Warren TH Elusive terminal copper aryl nitrene intermediates. *Angew. Chem., Int. Ed* 2017, 56 (23), 6426–6430.
- (48). Stone KL; Borovik AS Lessons from nature: unraveling biological C–H bond activation. *Curr. Opin. Chem. Biol* 2009, 13 (1), 114–118. [PubMed: 19297238]
- (49). Warren JJ; Tronic TA; Mayer JM Thermochemistry of proton-coupled electron transfer reagents and its implications. *Chem. Rev* 2010, 110 (12), 6961–7001. [PubMed: 20925411]
- (50). Bordwell FG; Cheng JP; Harrelson JA Homolytic bond dissociation energies in solution from equilibrium acidity and electrochemical data. *J. Am. Chem. Soc* 1988, 110 (4), 1229–1231.
- (51). Dhar D; Tolman WB Hydrogen atom abstraction from hydrocarbons by a copper(III)-hydroxide complex. *J. Am. Chem. Soc* 2015, 137 (3), 1322–1329. [PubMed: 25581555]
- (52). Nieto I; Ding F; Bontchev RP; Wang H; Smith JM Thermodynamics of hydrogen atom transfer to a high-valent iron imido complex. *J. Am. Chem. Soc* 2008, 130 (9), 2716–2717. [PubMed: 18266366]
- (53). Cowley RE; Bontchev RP; Sorrell J; Sarracino O; Feng Y; Wang H; Smith JM Formation of a cobalt(III) imido from a cobalt(II) amido complex. Evidence for proton-coupled electron transfer. *J. Am. Chem. Soc* 2007, 129 (9), 2424–2425. [PubMed: 17288417]
- (54). Scepaniak JJ; Young JA; Bontchev RP; Smith JM Formation of ammonia from an iron nitrido complex. *Angew. Chem., Int. Ed* 2009, 48 (17), 3158–3160.
- (55). Parsell TH; Yang M-Y; Borovik AS C–H bond cleavage with reductants: re-investigating the reactivity of monomeric Mn^{III/IV}-oxo complexes and the role of oxo ligand basicity. *J. Am. Chem. Soc* 2009, 131 (8), 2762–2763. [PubMed: 19196005]
- (56). Zhang L; Liu Y; Deng L. Three-coordinate cobalt(IV) and cobalt(V) imido complexes with N-heterocyclic carbene ligation: synthesis, structure, and their distinct reactivity in C–H bond amination. *J. Am. Chem. Soc* 2014, 136 (44), 15525–15528. [PubMed: 25330361]
- (57). Cowley RE; Eckert NA; Vaddadi S; Figg TM; Cundari TR; Holland PL Selectivity and mechanism of hydrogen atom transfer by an isolable imidoiron(III) complex. *J. Am. Chem. Soc* 2011, 133 (25), 9796–9811. [PubMed: 21563763]
- (58). Cowley RE; Holland PL Ligand effects on hydrogen atom transfer from hydrocarbons to three-coordinate iron imides. *Inorg. Chem* 2012, 51 (15), 8352–8361. [PubMed: 22800175]
- (59). Eckert NA; Vaddadi S; Stoian S; Lachicotte RJ; Cundari TR; Holland PL Coordination-number dependence of reactivity in an imidoiron(III) complex. *Angew. Chem., Int. Ed* 2006, 45 (41), 6868–6871.
- (60). Chilton NF; Anderson RP; Turner LD; Soncini A; Murray KS PHI: A powerful new program for the analysis of anisotropic monomeric and exchange-coupled polynuclear d- and f- block complexes. *J. Comput. Chem* 2013, 34 (13), 1164–1175. [PubMed: 23386394]
- (61). Searles K; Fortier S; Khusniyarov MM; Carroll PJ; Sutter J; Meyer K; Mindiola DJ; Caulton KG A cis-divacant octahedral and mononuclear iron(IV) imide. *Angew. Chem., Int. Ed* 2014, 53 (51), 14139–14143.
- (62). Anneser MR; Elpitiya GR; Townsend J; Johnson EJ; Powers XB; DeJesus JF; Vogiatzis KD; Jenkins DM Unprecedented five-coordinate iron(IV) imides generate divergent spin states based on the imide R-groups. *Angew. Chem., Int. Ed* 2019, 58 (24), 8115–8118.
- (63). Perdew JP Density-functional approximation for the correlation energy of the inhomogeneous electron gas. *Phys. Rev. B: Condens. Matter Mater. Phys* 1986, 33 (12), 8822–8824.
- (64). Becke AD Density-functional exchange-energy approximation with correct asymptotic behavior. *Phys. Rev. A: At., Mol., Opt. Phys* 1988, 38 (6), 3098–3100.
- (65). Lineweaver H; Burk D. The determination of enzyme dissociation constants. *J. Am. Chem. Soc* 1934, 56 (3), 658–666.
- (66). Renny JS; Tomasevich LL; Tallmadge EH; Collum DB Method of continuous variations: applications of Job plots to the study of molecular associations in organometallic chemistry. *Angew. Chem., Int. Ed* 2013, 52 (46), 11998–12013.

- (67). Bowman AC; Tondreau AM; Lobkovsky E; Margulieux GW; Chirik PJ Synthesis and electronic structure diversity of pyridine(diimine)iron tetrazene complexes. *Inorg. Chem* 2018, 57 (16), 9634–9643. [PubMed: 29620870]
- (68). Cowley RE; Elhaik J; Eckert NA; Brennessel WW; Bill E; Holland PL A bridging hexazene (RNNNNNR) ligand from reductive coupling of azides. *J. Am. Chem. Soc* 2008, 130 (19), 6074–6075. [PubMed: 18419120]
- (69). Hammes-Schiffer S; Stuchebrukhov AA Theory of coupled electron and proton transfer reactions. *Chem. Rev* 2010, 110 (12), 6939–6960. [PubMed: 21049940]

Figure 1. Solid-state structures of (a) $(\text{TrL})\text{CoCl}$ (**1**), (b) $(\text{TrL})\text{Co}$ (**2**), and (c, d) $(\text{TrL})\text{Co(NAd)}$ (**3**) at 100 K with thermal ellipsoids set at 50% probability (hydrogen atoms and solvent molecules are omitted for clarity; Co aquamarine, C gray, N blue, Cl green). Selected bond lengths (Å) and angles (deg) for **1**: Co–C_{ortho}, 2.345(7). **3**: Co–C_{ortho}, 2.756; Co–N, 1.636(3); $\angle\text{Co–N–C} = 145.0(3)$.

Figure 2.

Variable-temperature susceptibility data of (a) (TrL)CoCl (**1**) and (b) (TrL)Co (**2**) collected at 1.0 T from 5 to 300 K. (inset) Reduced magnetization data collected at three fields (1, 4, 7 T) over the temperature range 1.8–10 K. Magnetization fit parameters obtained with PHI:⁶⁰ for **1**, $S = 3/2$, $g = 2.4$, $D = -48.1 \text{ cm}^{-1}$, $|E/D| = 0.11$ and, for **2**, $S = 1$, $g = 2.03$, $D = 25.98 \text{ cm}^{-1}$, $|E/D| = 0.085$. (c) Variable-temperature susceptibility data of (TrL)Co(NAd) (**3**) collected at 0.5 and 1.0 T upon heating and cooling (5–300 K and 300–5 K).

Figure 3. Cyclic voltammograms of (a) **2** and (b) (^{Ar}L)Co obtained in 1,2-difluorobenzene at room temperature using 0.1 M [ⁿBu₄N]-[PF₆] electrolyte. Potentials were referenced to the [Cp₂Fe]^{+/0} couple. Scan rate = 100 mV/s.

Figure 4.
Frontier molecular orbital description of **3**.

Figure 5.

(a) Double-reciprocal plot with 10 mol % of **2** at varying concentrations of pyridine- d_5 at 50 °C. (b) Job plot analysis between **2** and pyridine in benzene- d_6 at 25 °C. (c) Solid-state structures of $(^{\text{Tr}}\text{L})\text{Co(py)} (4)$ at 100 K with thermal ellipsoids set at 50% probability (hydrogen atoms and solvent molecules are omitted for clarity; Co aquamarine, C gray, N blue). Selected bond lengths (Å): $\text{Co}-\text{C}_{\text{ortho}}$ 2.050(8); $\text{Co}-\text{C}_{\text{meta}}$ 2.239(8).

Figure 6.

(a) Full reaction progression with 10 mol % of **2** at 25 °C in benzene- d_6 . (b) Zeroth-order dependence on the concentration of **5** at 25 °C in benzene- d_6 . (inset) Initial stages of reaction showing the concentration of **7** is constant. A dashed line indicates the region less than 10% of **6**. (c) First-order dependence on the concentration of **7** at 25 °C benzene- d_6 .

Figure 7.
Kinetic isotope effect experiments with 10 mol % of **2** at 25 °C in benzene-*d*₆.

Figure 8.

Cyclic voltammograms of **3** and **8** obtained in 1,2-difluorobenzene at room temperature using 0.1 M $[\text{nBu}_4\text{N}][\text{PF}_6]$ electrolyte. Potentials were referenced to the $[\text{Cp}_2\text{Fe}]^{+/0}$ couple. Scan rate = 100 mV/s.

Scheme 1.
Ligand and Metal Complex Syntheses

Scheme 2.
Reactivity Assessment of 2

Scheme 3.
Proposed Catalytic Cycle

Scheme 4. Thermodynamic Square Scheme for HAA from Co^{III} Imides 3 and 8

Table 1.

Catalytic C–H Amination Using **2** and Alkyl Azides^a ^aIsolated yields from reactions conducted with 85.0 mM of azide substrate.^b1 mol % of **2**.^c5 mol % of **2**.^d10 mol % of **2**.