Published in final edited form as:

J Am Chem Soc. 2018 September 26; 140(38): 11906–11910. doi:10.1021/jacs.8b07418.

Oxidative Coupling with Zr(IV) Supported by a Non-Innocent Anthracene-Based Ligand: Application to the Catalytic Cotrimerization of Alkynes and Nitriles to Pyrimidines

Choon Heng Low, Jeffrey N. Rosenberg, Marco A. Lopez, Theodor Agapie*

Division of Chemistry and Chemical Engineering, California Institute of Technology, 1200 East California Boulevard MC 127-72, Pasadena, California 91125, United States

Abstract

We report the synthesis and reactivity of Zr complexes supported by a 9,10-anthracenediyl-linked bisphenoxide ligand, **L**. $\text{Zr}^{\text{IV}}\text{LBn}_2$ (**1**) undergoes facile photolytic reduction with concomitant formation of bibenzyl and $\text{Zr}^{\text{IV}}\text{L}(\text{THF})_3$ (**2**), which displays a two-electron reduced anthracene moiety. Leveraging ligand-stored reducing equivalents, **2** promotes the oxidative coupling of internal and terminal alkynes to isolable zirconacyclopentadiene complexes, demonstrating the reversible utilization of anthracene as a redox reservoir. With diphenylacetylene under CO, cyclopentadienone is formed stoichiometrically. **2** is competent for the catalytic formation of pyrimidines from alkynes and nitriles. Mechanistic studies suggest that selectivity for pyrimidine originates from preferred formation of an azazirconacyclopentadiene intermediate, which reacts preferentially with nitriles over alkynes.

Graphical Abstract

*Corresponding Author: agapie@caltech.edu.

Supporting Information

Experimental procedures for the synthesis of **LH2**, complexes **1-5**, detailed characterization data, supplementary figures (PDF) and crystallographic details (CIF) for complexes **1**, **2**, **3a**, and **5**. The Supporting Information is available free of charge on the ACS Publications website.

The authors declare no competing financial interests.

The use of redox-active ligands has expanded the reaction toolkit for chemists, mediating challenging multi-electron chemical transformations not usually observed in their absence.¹ At early metals such as Zr, for which redox-neutral processes like olefin polymerization are prevalent, redox-active ligands can facilitate new reactivity.² Typically, transformations involving redox non-innocent ligands result in formal changes of the ligand redox state without substantial changes in ligand coordination mode. Ligands that may change coordination environment around the metal are expected to result in more facile reactivity if adapted to the electronic and steric demands of the metal center. Our group and others have demonstrated the utility of labile and redox non-innocent pendant arene ligands in the development of new chemical reactivity, including CO cleavage and coupling, metal phosphide formation and coupling, metal nitride CO coupling, cross-coupling chemistry, and CO₂ activation.³ In most of these systems, the arene ligands are based on substituted benzene, requiring very negative potentials for formal reduction. Due to a smaller loss of aromaticity in its π -system,⁴ anthracene displays a more accessible reduction potential,⁵ allowing for formation of a dianionic state that coordinates metal ions at the bridgehead positions,⁶ though such transition metal complexes are very rare.⁷ We envisioned that the anthracene motif will promote novel reactivity by functioning as a reductively non-innocent ligand at milder potentials than benzene, facilitating redox chemistry at the pendant arene. Additionally, the formation of new bonds to the metal filling coordination sites opened during reactions such as reductive elimination is expected to facilitate reaction turnover and catalysis (Figure 1).

Syn-9,10-anthracenediyl-linked bisphenol (**LH₂**) was synthesized from commercially available starting materials in six steps (see SI). Protonolysis of tetrabenzyl zirconium (ZrBn₄) with **LH₂** provides bisbenzyl complex **1**. Single-crystal X-ray diffraction (XRD) studies (Figure 2) reveal long distances between Zr and the anthracene group which correlate with the C-C distances in the arene to suggest no significant interaction of the metal with the anthracene π -system.

Toward promoting reductive elimination from **1**, photolysis was investigated. After 18 h irradiation, ¹H NMR spectroscopy shows the loss of the benzylic signal in **1**, and the formation of bibenzyl and a new species, **2** (Scheme 1). The solid-state structure of **2** (Figure 2) shows coordination to Zr of the two phenoxide donors, three THF molecules and the anthracene linker. The anthracene motif binds in an η^4 fashion via C1, C2, C7, and C8, with Zr–C distances in the range of 2.432(1)–2.525(1) Å indicative of strong interactions. These are slightly longer than the Zr–benzyl distances in **1** (2.257(2)–2.272(2) Å). Significant lengthening of the C1–C2, C1–14, C7–C8 and C8–C9 bonds suggests disruption of aromaticity in the arene. Overall, the structural parameters observed are consistent with a two-electron reduced anthracene motif.^{6a–e} Although oxidatively or photolytically induced C–C reductive eliminations have been reported at Zr(IV),^{2d,8} such transformations are rare, and the reduced metal species can undergo undesired side reactivity such as CH activation.^{8e}

Complex **2**, although formally displaying a Zr(IV) center, stores two reducing equivalents in the anthracene motif. The possibility of **2** performing anthracene-based redox chemistry was investigated towards the oxidative cyclometallation of alkynes, a reaction with precedent for Zr(II).⁹ Heating a solution of **2** with diphenylacetylene (two equiv.) at 90 °C led to a red-to-

yellow color change and formation of a 1:2 Zr:alkyne species (**3a**) by ^1H NMR spectroscopy. The solid-state structure of **3a** shows the formation of a zirconacyclopentadiene by the oxidative coupling of two alkynes with two reducing equivalents originating from the ligand (Figure 2). Within the zirconacyclopentadiene ring, localized double bonds at C58–C59 (1.357(2) Å) and C60–C61 (1.354(3) Å) are observed.

With examples of both reductive and oxidative C–C bond formation involving redox at the pendant anthracene, we probed substrates with potential for regeneration of a masked Zr(II) complex. Extended heating of **3a** in the presence of excess diphenylacetylene did not effect further reactivity, likely due to steric constraints. Treatment of *in situ*-generated **3a** with CO at 90 °C results in the formation of a new major species assigned as **4**, a 1:1 adduct between zirconium and a tetraphenylcyclopentadienone (CPD) molecule (Scheme 1). Independent synthesis by mixing one equiv. CPD with **2**, resulting in a matching ^1H NMR spectrum, support this assignment. The synthesis of cyclopentadienones via [2+2+1] coupling of two alkynes and CO has been reported but is limited to mid-to-late transition metal complexes.¹⁰ CO insertion into zirconacyclopentadiene has not been reported, to our knowledge, though related systems involving CO-alkyne chemistry are known for Zr.¹¹ Conceptually related, the catalytic aza-Pauson-Khand reaction involving Ti-imido complexes and alkynes to afford pyrroles, with diazenes acting as a nitrene source has been reported.¹² Although turnover was not achieved for CPD formation, this reaction demonstrates that coupling of three substrates is possible, with an overall process that requires both oxidation and reduction of the anthracene moiety.

To examine if a sterically more open zirconacyclopentadiene promotes further reactivity, terminal alkyne phenylacetylene was tested. Addition of phenylacetylene (two equiv.) to **2** in benzene at room temperature led to a rapid red-to-yellow color change. The new species (**3b**) was assigned as a zirconacyclopentadiene complex structurally analogous to **3a** (^1H NMR spectroscopy). Coupling of terminal alkynes by zirconium is rare,¹³ primarily attributed to incompatibility of the acidic acetylenic proton with low-valent Zr(II) species and/or precursors.¹⁴

Heating **3b** in benzene to 90 °C in the presence of excess phenylacetylene did not result in further insertion. However, heating of **3b** in the presence of one equiv. of *p*-tolunitrile (TolCN) and two equiv. of phenylacetylene led to the almost complete consumption of TolCN after one hour, with complex **3b** still present (^1H NMR spectroscopy). GC-MS analysis showed the formation of a mixture of homo- and heterotrimerized products, suggesting that **2** may be a competent precatalyst for the cotrimerization of alkynes and nitriles.

To further investigate this reactivity, benzonitrile and phenylacetylene were chosen as model substrates. A mixture of alkyne was heated at 90 °C with excess nitrile (7.5 equiv.), to disfavor alkyne trimerization,¹⁵ in the presence of 5 mol% of **2**. After two hours, highly selective formation of 2,4,6-triphenylpyrimidine (**6a**) was observed, in 53% yield (Table 1, entry 1). GC-MS and NMR analysis showed no detectable formation of triphenylbenzene or triphenyltriazine, and less than 2% of triphenylpyridine. Pyrimidines feature in a wide variety of active pharmaceuticals and natural products, and as ligands in coordination

complexes.¹⁶ Syntheses of pyrimidines via the highly efficient and atom-economical [2+2+2] cycloaddition route¹⁷ have been scarce.¹⁸ This has been attributed to the difficulty in incorporation of multiple nitrogen atoms via cycloaddition due to the more reactive nature of alkynes compared to nitriles,¹⁹ even when employing the nitrile as solvent.²⁰ Of those, many are either stoichiometric or require high catalyst loadings (20%),^{18d–g} the use of nitriles as a solvent,^{18h,18i} or tethered alkyne-nitrile substrates.^{18j} The observation of catalytic cycloaddition involving a nitrile is particularly notable for an early metal. Although there are several systems for pyridine synthesis based on Zr in combination with Ni²¹ or Cu,²² and Ti,²³ these reactions are not catalytic, likely due to the strong binding of the nitrogen moiety to the highly Lewis acidic metal center. For the present catalyst, the propensity of the anthracene moiety to accept reducing equivalents is proposed to facilitate reductive cyclization, promoting turnover.

Increasing the reaction temperature to 105 °C in toluene led to quantitative yields of **6a**, with longer reaction times leading to triazine formation after alkyne substrate has been consumed (Table 1, entries 2 and 3). The nitrile excess can be reduced without appreciable loss in selectivity, but results in slower conversions (entries 4 and 5). At 2:1 PhCN:PhCCH, high selectivity for pyrimidine (94%) is retained, albeit with a lower yield of ~80% (entry 6). Notably, even excess alkyne (1:5 PhCN:PhCCH), the pyrimidine is favored over pyridine. Catalyst loading can be lowered to 3 mol%, while providing similarly high yields and selectivities after five hours (entry 8). Control experiments ran with simple Zr complexes (ZrBn₄, ZrCl₄ and ZrBn₂Cl₂) reduced by Mg(THF)₃(anthracenide), or with KC₈ or photolytically in the presence of anthracene, did not result in catalysis (see SI).

Using the optimized conditions (Table 1, entry 4), the scope of pyrimidine synthesis was explored (Table 2). Internal dialkyl alkyne 5-decyne provides **6b** quantitatively, while diphenylacetylene is not competent. Terminal alkyl alkynes (Table S1) did not produce any cotrimerized products detectable by GC-MS, with the exception of trimethylsilylacetylene, which quantitatively afforded **6c**. A variety of nitriles were also found to be competent for catalysis (Table 2, **6d**, **6e** and **6f**) though aryl nitriles with O or N-containing substituents and alkyl nitriles (Table S2) did not lead to cyclotrimerized products. Addition of an equimolar amount of acetonitrile to PhCN under optimized conditions completely shuts down generation of **6a**, suggesting that it acts as a strong inhibitor, likely through competitive binding to the metal.

To gain insight into the mechanistic basis of product selectivity, the synthesis of azazirconacycle **5**, a potential catalytic intermediate, was targeted. Addition of two equiv. TolCN followed by one equiv. PhCCH to **2** results in the formation of azazirconacycle **5**. XRD studies took advantage of the distinct aryl groups to unambiguously assign N-coordination to Zr and 2,4-diaryl substitution (Figure S37). Although we could not rule out the initial reversible formation of a nitrile-nitrile coupled complex²⁴ of the type previously reported for group IV metals,²⁵ in the presence of both nitrile and alkyne, complex **5** is the major Zr product. Stoichiometric reactions carried out from the isolated (aza)zirconacycles, **3b** and **5**, provide insight into catalytically relevant pathways (Scheme 2). Reaction of **3b** or **5** at 90 °C with PhCN results in pyridine or pyrimidine formation, respectively, while neither complex reacts in the presence of additional PhCCH (*vide supra*). GC-MS analysis of the

reaction of **5** with PhCN showed predominantly the formation of pyrimidine with mixed aryl substituents, with minor formation of 2,4,6-triphenylpyrimidine and 6-phenyl-2,4-di(tolyl)pyrimidine, indicating that the coupling of alkyne and nitrile is reversible, but is slow relative to pyrimidine formation. Based on the selectivity in catalysis, pathway B is strongly favored over pathway A, leading to the high selectivities observed. The preference for the formation of **5** rather than **3b** from a mixture of nitrile and alkyne is likely due to the precoordination of nitrile. The small amount of pyridine byproduct results from the formation of **3b** as a minor component through Pathway A.

In summary, we have synthesized and characterized a series of zirconium complexes supported by a bisphenoxide ligand displaying a 9,10-anthracenediyl motif. The ligand exhibits both redox non-innocence and hemilability, facilitating two-electron chemistry at a Zr(IV) center such as the photolytic reductive elimination of bibenzyl, oxidative coupling of unsaturated organic substrates such as alkynes and nitriles and their subsequent reductive elimination to CPD or N-containing heterocycles. We have demonstrated an efficient Zr-catalyzed three-component [2+2+2] cycloaddition of alkynes and aryl nitriles to selectively afford tri- and tetra-substituted pyrimidines. The catalyst displays excellent selectivity even without the use of excess nitrile. The ability of this Zr catalyst to turnover, in contrast to other early metal systems is proposed to stem from the ability of the anthracene motif to promote redox chemistry and product dissociation. Given the established efficient stoichiometric oxidative coupling chemistry characteristic of Zr,^{9b,26} the development of new types of catalysis based on it is of interest for a variety of applications.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

We are grateful to the Dreyfus Foundation (T.A.), the Agency of Science, Technology and Research (A*STaR), Singapore (C.H.L.), and the Caltech's SURF (J.N.R.) and WAVE (M.A.L.) programs for funding. We thank Lawrence M. Henling, and Michael K. Takase for crystallographic assistance, and David VanderVelde for advice on NMR. The NSF (CRIF:MU CHE0639094) and NIH (RR027690) are thanked for instrument grants to Caltech.

REFERENCES

- (1). (a)Chirik PJ; Wieghardt K, "Radical Ligands Confer Nobility on Base-Metal Catalysts." *Science* 2010, 327, 794–795; [PubMed: 20150476] (b)Luca OR; Crabtree RH, "Redox-active ligands in catalysis." *Chem. Soc. Rev* 2013, 42, 1440–1459; [PubMed: 22975722] (c)Lyaskovskyy V; de Bruin B, "Redox Non-Innocent Ligands: Versatile New Tools to Control Catalytic Reactions." *ACS Catalysis* 2012, 2, 270–279.
- (2). (a)Blackmore KJ; Lal N; Ziller JW; Heyduk AF, "Catalytic Reactivity of a Zirconium(IV) Redox-Active Ligand Complex with 1,2-Diphenylhydrazine." *J. Am. Chem. Soc* 2008, 130, 2728–2729; [PubMed: 18266370] (b)Blackmore KJ; Ziller JW; Heyduk AF, "Oxidative Addition" to a Zirconium(IV) Redox-Active Ligand Complex." *Inorg. Chem* 2005, 44, 5559–5561; [PubMed: 16060604] (c)Nguyen AI; Zarkesh RA; Lacy DC; Thorson MK; Heyduk AF, "Catalytic nitrene transfer by a zirconium(iv) redox-active ligand complex." *Chemical Science* 2011, 2, 166–169; (d)Haneline MR; Heyduk AF, "C–C Bond-Forming Reductive Elimination from a Zirconium(IV) Redox-Active Ligand Complex." *J. Am. Chem. Soc* 2006, 128, 8410–8411; [PubMed: 16802801] (e)Heins SP; Wolczanski PT; Cundari TR; MacMillan SN, "Redox non-innocence

permits catalytic nitrene carbonylation by (dadi)Ti=NAd (Ad = adamantyl).” *Chemical Science* 2017, 8, 3410–3418. [PubMed: 28507712]

- (3). (a) Buss JA; Agapie T, “Mechanism of Molybdenum-Mediated Carbon Monoxide Deoxygenation and Coupling: Mono- and Dicarbonyl Complexes Precede C–O Bond Cleavage and C–C Bond Formation.” *J. Am. Chem. Soc* 2016, 138, 16466–16477; [PubMed: 27936655] (b) Buss JA; Agapie T, “Four-electron deoxygenative reductive coupling of carbon monoxide at a single metal site.” *Nature* 2016, 529, 72–75; [PubMed: 26689364] (c) Horak KT; Velian A; Day MW; Agapie T, “Arene non-innocence in dinuclear complexes of Fe, Co, and Ni supported by a para-terphenyl diphosphine.” *Chem. Commun* 2014, 50, 4427–4429; (d) Henthorn JT; Lin S; Agapie T, “Combination of Redox-Active Ligand and Lewis Acid for Dioxygen Reduction with π -Bound Molybdenum–Quinonoid Complexes.” *J. Am. Chem. Soc* 2015, 137, 1458–1464; [PubMed: 25577950] (e) Velian A; Lin S; Miller AJM; Day MW; Agapie T, “Synthesis and C–C Coupling Reactivity of a Dinuclear NiI–NiI Complex Supported by a Terphenyl Diphosphine.” *J. Am. Chem. Soc* 2010, 132, 6296–6297; [PubMed: 20397653] (f) Kurogi T; Ishida Y; Kawaguchi H, “Synthesis of titanium and zirconium complexes supported by a p-terphenoxide ligand and their reactions with N₂, CO₂ and CS₂.” *Chem. Commun* 2013, 49, 11755–11757; (g) Buss J. V., David; Agapie Theodor, *J. Am. Chem. Soc* 2018, Submitted; (h) Buss JA; VanderVelde DG; Agapie T, “Lewis Acid Enhancement of Proton Induced CO₂ Cleavage: Bond Weakening and Ligand Residence Time Effects.” *J. Am. Chem. Soc* 2018, 140, 10121–10125; [PubMed: 30032606] (i) Plundrich GT; Wadepohl H; Clot E; Gade LH, “ η^6 -Arene-Zirconium-PNP-Pincer Complexes: Mechanism of Their Hydrogenolytic Formation and Their Reactivity as Zirconium(II) Synthons.” *Chem. Eur. J* 2016, 22, 9283–9292; [PubMed: 27258989] (j) Jilek RE; Jang M; Smolensky ED; Britton JD; Ellis JE, “Structurally Distinct Homoleptic Anthracene Complexes, [M(C₁₄H₁₀)₃]₂, M = Titanium, Zirconium, Hafnium: Tris(arene) Complexes for a Triad of Transition Metals.” *Angew. Chem. Int. Ed* 2008, 47, 8692–8695.
- (4). Schleyer P. v. R.; Manoharan M; Jiao H; Stahl F, “The Acenes: Is There a Relationship between Aromatic Stabilization and Reactivity?” *Org. Lett* 2001, 3, 3643–3646. [PubMed: 11700102]
- (5). Connelly NG; Geiger WE, “Chemical Redox Agents for Organometallic Chemistry.” *Chem. Rev* 1996, 96, 877–910. [PubMed: 11848774]
- (6). (a) Roitershtein DM; Rybakova LF; Petrov ES; Ellern AM; Antipin MY; Struchkov YT, “Lutetium complexes with an anthracene dianion ligand. Molecular structure of [(C₅H₅)₂Lu(C₁₄H₁₀)] [Na⁺(diglyme)₂].” *J. Organomet. Chem* 1993, 460, 39–45; (b) Bogdanovic B, “Magnesium anthracene systems and their application in synthesis and catalysis.” *Acc. Chem. Res* 1988, 21, 261–267; (c) Fedushkin IL; Bochkarev MN; Dechert S; Schumann H, “A Chemical Definition of the Effective Reducing Power of Thulium(II) Diiodide by Its Reactions with Cyclic Unsaturated Hydrocarbons.” *Chem. Eur. J* 2001, 7, 3558–3563; [PubMed: 11560327] (d) Lehmkuhl H; Mehler K; Shakoor A; Krüger C; Tsay Y-H; Benn R; Rufi ska A; Schroth G, “9,10-Dihydro-9,10-anthrylen- Verbindungen des Aluminiums.” *Chem. Ber* 1985, 118, 4248–4258; (e) Lehmkuhl H; Shakoor A; Mehler K; Krüger C; Angermund K; Tsay Y-H, “[9,10-Dihydro-9,10-bis(trimethylsilyl)-9,10-anthrylen]magnesium: Molekülstruktur eines „Anthracenmagnesiums”. ” *Chem. Ber* 1985, 118, 4239–4247; (f) Bogdanovi B; Janke N; Krüger C; Mynott R; Schlichte K; Westeppe U, “Synthesis and Structure of 1,4-Dimethylantracenemagnesium-3thf and μ -Trichlorodimagnesium-6thf(1+) Anthracenide.” *Angew. Chem. Int. Ed* 1985, 24, 960–961.
- (7). (a) Fryzuk Michael D; Jafarpour L; Kerton Francesca M; Love Jason B; Rettig Steven J, “Dinuclear π Complexes of Yttrium and Lutetium with Sandwiched Naphthalene and Anthracene Ligands: Evidence for Rapid Intramolecular Inter-Ring Rearrangements.” *Angew. Chem. Int. Ed* 2000, 39, 767–770; (b) Huang W; Khan SI; Diaconescu PL, “Scandium Arene Inverted-Sandwich Complexes Supported by a Ferrocene Diamide Ligand.” *J. Am. Chem. Soc* 2011, 133, 10410–10413; [PubMed: 21657792] (c) Huang W; Abukhalil PM; Khan SI; Diaconescu PL, “Group 3 metal stilbene complexes: synthesis, reactivity, and electronic structure studies.” *Chem. Commun* 2014, 50, 5221–5223.
- (8). (a) Erker G, “The reaction of intermediate zirconocene-aryne complexes with C–H bonds in the thermolysis of diarylzirconocenes.” *J. Organomet. Chem* 1977, 134, 189–202; (b) Burk MJ; Tumas W; Ward MD; Wheeler DR, “Oxidation chemistry of d⁰ organometallic complexes.” *J. Am. Chem. Soc* 1990, 112, 6133–6135; (c) Ishikawa T; Ogawa A; Hirao T, “Oxovanadium(V)-induced oxidation of alkenylzirconocenes for facile inter- and intramolecular coupling.” *J.*

Organomet. Chem 1999, 575, 76–79;(d)Margulieux GW; Semproni SP; Chirik PJ, “Photochemically Induced Reductive Elimination as a Route to a Zirconocene Complex with a Strongly Activated N2 Ligand.” Angew. Chem. Int. Ed 2014, 53, 9189–9192;(e)Covert KJ; Mayol A-R; Wolczanski PT, “Carbon-oxygen and related R-X bond cleavages mediated by (silox)3Ti and other Group 4 derivatives (silox = tBu3SiO).” Inorg. Chim. Acta 1997, 263, 263–278.

- (9). (a)Negishi E.-i.; Cederbaum FE; Takahashi T, “Reaction of zirconocene dichloride with alkylolithiums or alkyl grignard reagents as a convenient method for generating a “zirconocene” equivalent and its use in zirconium-promoted cyclization of alkenes, alkynes, dienes, enynes, and diynes.” Tetrahedron Lett. 1986, 27, 2829–2832;(b)Nitschke JR; Zürcher S; Tilley TD, “New Zirconocene-Coupling Route to Large, Functionalized Macrocycles.” J. Am. Chem. Soc 2000, 122, 10345–10352;(c)Nugent WA; Thorn DL; Harlow RL, “Cyclization of diacetylenes to E,E exocyclic dienes. Complementary procedures based on titanium and zirconium reagents.” J. Am. Chem. Soc 1987, 109, 2788–2796;(d)Wielstra Y; Gambarotta S; Meetsma A; De Boer JL; Chiang MY, “Carbon-carbon bond formation versus hydrogen transfer in the reaction of alkynes with mono(cyclopentadienyl)zirconium(II) complexes: regioselective dimerization, cocyclotrimerization, and catalytic cyclotrimerization. The preparation and x-ray structures of Cp(dmpe)XZr[(R)C:C(R')C(R):C(R')] [R = CH3; R' = H, CH3; X = Cl, CH3; dmpe = 1,2-bis(dimethylphosphino)ethane] and of CpZr(dmpe)Cl[C.tplbond.C(t-Bu)][CH:CH(t-Bu)].” Organometallics 1989, 8, 2696–2702.
- (10). (a)Schaefer C; Werz DB; Staeb TH; Gleiter R; Rominger F, “CpCo-Stabilized Cyclopentadienones from Cyclobutadiene Complexes: Experimental and Theoretical Investigations.” Organometallics 2005, 24, 2106–2113;(b)Schoe NE, “Transition metal-mediated cycloaddition reactions of alkynes in organic synthesis.” Chem. Rev 1988, 88, 1081–1119;(c)Gleiter R; Werz DB, “Reactions of Metal-Complexed Carbocyclic 4π Systems.” Organometallics 2005, 24, 4316–4329;(d)Shibata T; Yamashita K; Ishida H; Takagi K, “Iridium Complex Catalyzed Carbonylative Alkyne–Alkyne Coupling for the Synthesis of Cyclopentadienones.” Org. Lett 2001, 3, 1217–1219. [PubMed: 11348198]
- (11). (a)Sikora DJ; Rausch MD, “Reactions of Cp2M(CO)2 and Cp2M(CO)(PPh3) (M = Zr, Hf) with acetylenes: Formation of some metallacyclopentadiene complexes of zirconocene and hafnocene.” J. Organomet. Chem 1984, 276, 21–37;(b)Guram AS; Guo Z; Jordan RF, “Zirconium-mediated reactions of carbon monoxide and alkynes. Insertion chemistry of cationic Zr(IV).eta.2-acyl and alkenyl complexes.” J. Am. Chem. Soc 1993, 115, 4902–4903;(c)Takahashi T; Huo S; Hara R; Noguchi Y; Nakajima K; Sun W-H, “Reaction of Zirconacyclopentadienes with CO in the Presence of n-BuLi. Selective Formation of Cyclopentenone Derivatives from Two Alkynes and CO.” J. Am. Chem. Soc 1999, 121, 1094–1095;(d)Li H; Liu L; Zhao F; Wang C; Wang C; Song Q; Zhang W-X; Xi Z, “Carbonylation of 1-Lithiobutadiene with Carbon Monoxide Followed by Intramolecular Acyllithiation of C=C Double Bond and Intermolecular Acylation with Acid Chloride: Scope, Applications, and Mechanistic Aspects.” J. Org. Chem 2012, 77, 4793–4800. [PubMed: 22524473]
- (12). Gilbert ZW; Hue RJ; Tonks IA, “Catalytic formal [2+2+1] synthesis of pyrroles from alkynes and diazenes via TiII/TiIV redox catalysis.” Nature Chemistry 2015, 8, 63.
- (13). (a)Denhez C; Médégan S; Hélicon F; Namy J-L; Vasse J-L; Szymoniak J, “Reduction of Cp2ZrCl2 with Mischmetall: A New Method for Generating an Efficient “Cp2Zr” Equivalent.” Org. Lett 2006, 8, 2945–2947; [PubMed: 16805523] (b)Zirngast M; Marschner C; Baumgartner J, “Group 4 Metallocene Complexes of Tris(trimethylsilyl)silylacetylene and Related Alkynes.” Organometallics 2008, 27, 2570–2583;(c)Shenyong R; Takashi S; David N; Hiroyuki S; Kiyohiko N; Ken-ichiro K; Zhiying S; Tamotsu T, “Selective Dimerization of Aryl-substituted Terminal Alkynes on Bis(indenyl)zirconocene Derivatives.” Chem. Lett 2011, 40, 1443–1444;(d)Bousrez G; Jaroschik F; Martinez A; Harakat D; Nicolas E; Le Goff XF; Szymoniak J, “Reactivity differences between 2,4- and 2,5-disubstituted zirconacyclopentadienes: a highly selective and general approach to 2,4-disubstituted phospholes.” Dalton Trans. 2013, 42, 10997–11004. [PubMed: 23797680]
- (14). Ma W; Yu C; Chen T; Xu L; Zhang W-X; Xi Z, “Metallacyclopentadienes: synthesis, structure and reactivity.” Chem. Soc. Rev 2017, 46, 1160–1192. [PubMed: 28119972]

- (15). Although alkyne homotrimerization products were not produced when 3b was treated with excess phenylacetylene, in the presence of certain additives (nitriles, CO), their formation was observed.
- (16). (a)Gómez-de la Torre F; de la Hoz A; Jalón FA; Manzano BR; Rodríguez AM; Elguero J; Martínez-Ripoll M, "Pd(II) Complexes with Polydentate Nitrogen Ligands. Molecular Recognition and Dynamic Behavior Involving Pd–N Bond Rupture. X-ray Molecular Structures of $[\{Pd(C_6HF_4)_2\}(bpzpm)]$ and $[\{Pd(\eta^3-C_4H_7)\}_2(bpzpm)](CF_3SO_3)_2$ [$bpzpm = 4,6$ -Bis(pyrazol-1-yl)pyrimidine]." *Inorg. Chem* 2000, 39, 1152–1162; [PubMed: 12526405] (b)Kua J; Xu X; Periana RA; Goddard WA, "Stability and Thermodynamics of the $PtCl_2$ Type Catalyst for Activating Methane to Methanol: A Computational Study." *Organometallics* 2002, 21, 511–525;(c)Warsink S; Chang IH; Weigand JJ; Hauwert P; Chen J-T; Elsevier CJ, "NHC Ligands with a Secondary Pyrimidyl Donor for Electron-Rich Palladium(0) Complexes." *Organometallics* 2010, 29, 4555–4561;(d)Walker SR; Carter EJ; Huff BC; Morris JC, "Variolins and Related Alkaloids." *Chem. Rev* 2009, 109, 3080–3098; [PubMed: 19489543] (e)Undheim K; Benneche T, 6.02 - Pyrimidines and their Benzo Derivatives A2 - Katritzky, Alan R In *Comprehensive Heterocyclic Chemistry II*, Rees CW; Scriven EFV, Eds. Pergamon: Oxford, 1996; pp 93–231.
- (17). (a)Perreault S; Rovis T, "Multi-component cycloaddition approaches in the catalytic asymmetric synthesis of alkaloid targets." *Chem. Soc. Rev* 2009, 38, 3149–3159; [PubMed: 19847348] (b)Gulevich AV; Dudnik AS; Chernyak N; Gevorgyan V, "Transition Metal-Mediated Synthesis of Monocyclic Aromatic Heterocycles." *Chem. Rev* 2013, 113, 3084–3213; [PubMed: 23305185] (c)Chopade PR; Louie J, "[2+2+2] Cycloaddition Reactions Catalyzed by Transition Metal Complexes." *Adv. Synth. Catal* 2006, 348, 2307–2327.
- (18). (a)Hill MD; Movassaghi M, "New Strategies for the Synthesis of Pyrimidine Derivatives." *Chem. Eur. J* 2008, 14, 6836–6844; [PubMed: 18384023] (b)Mahfoudh M; Abderrahim R; Leclerc E; Campagne J-M, "Recent Approaches to the Synthesis of Pyrimidine Derivatives." *Eur. J. Org. Chem* 2017, 2017, 2856–2865;(c)Karad Somnath N; Liu RS, "Regiocontrolled Gold-Catalyzed [2+2+2] Cycloadditions of Ynamides with Two Discrete Nitriles to Construct 4-Aminopyrimidine Cores." *Angew. Chem. Int. Ed* 2014, 53, 9072–9076;(d)You X; Yu S; Liu Y, "Reactions of Zirconocene Butadiyne or Monoyne Complexes with Nitriles: Straightforward Synthesis of Functionalized Pyrimidines." *Organometallics* 2013, 32, 5273–5276;(e)Chen P; Song C.-x.; Wang W.-s.; Yu X.-l.; Tang Y, "TfOH-mediated [2 + 2 + 2] cycloadditions of ynamides with two discrete nitriles: synthesis of 4-aminopyrimidine derivatives." *RSC Advances* 2016, 6, 80055–80058;(f)Burlakov VV; Becker L; Bogdanov VS; Andreev MV; Arndt P; Spannenberg A; Baumann W; Rosenthal U, "Reactivity of a Seven-Membered Zirconacyclocumulene towards CN Multiple Bonds – Formation of Metallaheterocycles by Insertion of $C\equiv N$ and $C=N$ Groups." *Eur. J. Inorg. Chem* 2014, 2014, 5304–5310;(g)Fuji M; Obora Y, "FeCl₃-Assisted Niobium-Catalyzed Cycloaddition of Nitriles and Alkynes: Synthesis of Alkyl- and Arylpyrimidines Based on Independent Functions of NbCl₅ and FeCl₃ Lewis Acids." *Org. Lett* 2017, 19, 5569–5572; [PubMed: 28968133] (h)Sato Y; Yasuda K; Obora Y, "Strategy for the Synthesis of Pyrimidine Derivatives: NbCl₅-Mediated Cycloaddition of Alkynes and Nitriles." *Organometallics* 2012, 31, 5235–5238;(i)Xie L-G; Niyomchon S; Mota AJ; González L; Maulide N, "Metal-free intermolecular formal cycloadditions enable an orthogonal access to nitrogen heterocycles." *Nat. Commun* 2016, 7, 10914; [PubMed: 26975182] (j)Lane TK; Nguyen MH; D'Souza BR; Spahn NA; Louie J, "The iron-catalyzed construction of 2-aminopyrimidines from alkynenitriles and cyanamides." *Chem. Commun* 2013, 49, 7735–7737.
- (19). Liu Y; Yan X; Yang N; Xi C, "Highly regioselective cyclotrimerization of terminal alkynes catalyzed by Fe(II) complexes bearing 2-(benzimidazolyl)-6-(1-(arylimino)ethyl)pyridines." *Catal. Commun* 2011, 12, 489–492.
- (20). (a)Hilt G; Vogler T; Hess W; Galbiati F, "A simple cobalt catalyst system for the efficient and regioselective cyclotrimerisation of alkynes." *Chem. Commun* 2005, 1474–1475;(b)Hsieh J-C; Cheng C-H, "O-Dihaloarenes as aryne precursors for nickel-catalyzed [2 + 2 + 2] cycloaddition with alkynes and nitriles." *Chem. Commun* 2008, 2992–2994.
- (21). Takahashi T; Tsai F-Y; Kotora M, "Selective Formation of Substituted Pyridines from Two Different Alkynes and a Nitrile: Novel Coupling Reaction of Azazirconacyclopentadienes with Alkynes." *J. Am. Chem. Soc* 2000, 122, 4994–4995.

- (22). Takahashi T; Tsai F-Y; Li Y; Wang H; Kondo Y; Yamanaka M; Nakajima K; Kotora M, "Selective Preparation of Pyridines, Pyridones, and Iminopyridines from Two Different Alkynes via Azazirconacycles." *J. Am. Chem. Soc* 2002, 124, 5059–5067. [PubMed: 11982370]
- (23). (a)Hill JE; Balaich G; Fanwick PE; Rothwell IP, "The chemistry of titanacyclopentadiene rings supported by 2,6-diphenylphenoxide ligation: stoichiometric and catalytic reactivity." *Organometallics* 1993, 12, 2911–2924;(b)Tanaka R; Yuza A; Watai Y; Suzuki D; Takayama Y; Sato F; Urabe H, "One-Pot Synthesis of Metalated Pyridines from Two Acetylenes, a Nitrile, and a Titanium(II) Alkoxide." *J. Am. Chem. Soc* 2005, 127, 7774–7780; [PubMed: 15913367] (c)Varela JA; Saá C, "Construction of Pyridine Rings by Metal-Mediated [2 + 2 + 2] Cycloaddition." *Chem. Rev* 2003, 103, 3787–3802. [PubMed: 12964884]
- (24). A new species was formed upon addition of nitrile (2 equiv.) to 2, with displacement of THF. Potential assignments include a bis(nitrile) adduct or a diazazirconacyclopentadiene complex.
- (25). (a)Becker L; Arndt P; Jiao H; Spannenberg A; Rosenthal U, "Nitrile–Nitrile C C Coupling at Group 4 Metallocenes to Form 1-Metalla-2,5-diaza-cyclopenta-2,4-dienes: Synthesis and Reactivity." *Angew. Chem. Int. Ed* 2013, 52, 11396–11400;(b)Becker L; Strehler F; Korb M; Arndt P; Spannenberg A; Baumann W; Lang H; Rosenthal U, "Unusual Nitrile–Nitrile and Nitrile–Alkyne Coupling of Fc C N and Fc C C C N." *Chem. Eur. J* 2014, 20, 3061–3068; [PubMed: 24615841] (c)Kiel GR; Samkian AE; Nicolay A; Witzke RJ; Tilley TD, "Titanocene-Mediated Dinitrile Coupling: A Divergent Route to Nitrogen-Containing Polycyclic Aromatic Hydrocarbons." *J. Am. Chem. Soc* 2018, 140, 2450–2454. [PubMed: 29383934]
- (26). (a)Kiel GR; Patel SC; Smith PW; Levine DS; Tilley TD, "Expanded Helicenes: A General Synthetic Strategy and Remarkable Supramolecular and Solid-State Behavior." *J. Am. Chem. Soc* 2017, 139, 18456–18459; [PubMed: 29215272] (b)Kiel Gavin R; Ziegler Micah S; Tilley TD, "Zirconacyclopentadiene-Annulated Polycyclic Aromatic Hydrocarbons." *Angew. Chem. Int. Ed* 2017, 56, 4839–4844.

Figure 1.
Non-innocent anthracene moiety facilitating reductive elimination.

Figure 2.

Solid-state structures of **1**, **2**, and **3a** and central ring bond metrics of **1** and **2**. Bond distances in Å. Thermal ellipsoids shown at 50% probability. Solvent molecules and hydrogen atoms omitted for clarity.

Scheme 1.

Synthesis and reactivity of bisphenoxide zirconium complexes with pendant anthracene

Scheme 2. Stoichiometric reactions involving catalytically relevant species.

Table 1.

Cycloaddition of phenyl acetylene with benzonitrile under various conditions

$\text{PhCCH (1 equiv.)} + \text{PhCN} \xrightarrow[\text{toluene, 105 } ^\circ\text{C}]{\text{2}}$ <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;"> <p>6a</p> </div> <div style="text-align: center;"> <p>6a'</p> </div> <div style="text-align: center;"> <p>6a''</p> </div> </div>								
entry	[2]/mol %	nitrile (equiv.)	time/h	^a PhCCH consumed/ %	^{b,c} selectivity/%			
					6a	6a'	6a''	
1 ^d	5	7.5	2	53	98	2	nd	
2	5	7.5	1	>99	59	1	40	
3	5	7.5	0.5	>99	99	1	nd	
4	5	6	1	>99	99	1	nd	
5	5	3	5	98	97	3	nd	
6	5	2	8	79	95	5	nd	
7	5	0.2	8	7	55	45	nd	
8	3	6	5	95	98	2	nd	

^a based on GC-MS analysis, averaged over 2 runs;^b based on ¹H NMR integration, averaged over 2 runs;^c nd: not detected by GC-MS;^d run at 90 °C in benzene

Table 2.

Substrate scope of cycloaddition of alkynes with nitriles to pyrimidines

^aProducts (**6**) and yields/% ^aisolated yields averaged over 2 runs