CORRESPONDENCE

Smoking Cessation Attempts and Common Strategies Employed

A Germany-Wide Representative Survey Conducted in 19 Waves From 2016 to 2019 (The Debra Study) and Analyzed by Socioeconomic Status)

by Prof. Dr. Daniel Kotz, PhD MSc MPH, Prof. Dr. med. Anil Batra, and Dr. rer. nat. Sabrina Kastaun in issue 1–2/2020

Not a Smoking Cessation Aid

The authors did not mention how much nicotine is inhaled when e-cigarettes are used purportedly in order to stop smoking (1).

With each cigarette, smokers take in 1 mg of nicotine. According to the tobacco guideline, the components/ingredients of smoking one cigarette are defined as follows: nicotine content 1.0 mg, carbon monoxide 10 mg, tar 10 mg per cigarette. The nicotine content of the cigarette itself is much higher. Swallowing tobacco or drinking an infusion of a few cigarettes is fatal. Tar has an unpleasant effect and prevents smokers from increasing the amount of cigarettes smoked ad infinitum. “Chain smokers” admit to smoking between 35 and 40 cigarettes or two packs per day; their intake is therefore 35–40 mg nicotine; more than that is mostly not tolerated.

Since nicotine does not burn up in e-cigarettes but is vaporized, the nicotine contained in the vaping liquids is almost completely inhaled. In the EU, e-cigarette liquids usually contain up to 20 mg/mL, diluted with flavorings to the ready solution of 10 mg/mL: mango or cucumber flavor are tolerated better than tar.

Lack of Authoritative Measures to Support Abstinence From Nicotine

The authors clearly pointed out that currently in Germany, fewer and fewer smokers stop smoking, and that several factors can be outlined that further stabilize the social gradient in smoking and health behaviors (1). Chronic obstructive pulmonary disease, which accounts for about one-third of all lung transplantations in Germany, can be explained in the vast majority of cases with tobacco inhalation. It is of note that in the current guideline of the German Medical Association according to paragraph 16 section 1 S.1 Nos 2 and 5 of the German Transplant Act, the only formal requirement is six months’ abstinence before admission to the waiting list. Recent studies have found a smoking relapse rate after lung transplantation of up to 15% (2). In my opinion, comparable authoritative measures from transplantation centers are lacking, which before and especially after a transplantation help patients to maintain abstinence—and which should therefore be funded in advance for each patient and, in particular, indepen-

I ask my “vaping” patients regularly for concrete amounts. On average they report that 10 ml vape juice at 10 mg/mL—that is, 100 mg nicotine—lasts them about a day and a half. The nicotine intake is almost twice that of chain smokers, but all of them subjectively expressed the opinion that they smoked less. Fruit esters do not scratch in the throat in the way tar does.

In conclusion: the key statement of the article, that e-cigarettes constitute a form of smoking cessation support, is untenable. DOI: 10.3238/arztebl.2020.0298a

References

1. Kotz D, Batra A, Kastaun S: Smoking cessation attempts and common strategies employed—a Germany-wide representative survey conducted in 19 waves from 2016 to 2019 (The DEBRA Study) and analyzed by socioeconomic status. *Dtsch Arztebl Int* 2020; 117: 7–13.

Dr. med. Ralf Cüppers

Medical psychotherapist, addiction psychiatrist
Flensburg, Germany
ralf@psychotherapeutische-medizin.net

Conflict of interest statement

The author declares that no conflict of interest exists

dently of the patient’s income, not least in order to meet the legislator’s ambiguous claim in § 1 of the transplant act in each and every possible way: “The aim of the law is to promote willingness to donate organs in Germany.” DOI: 10.3238/arztebl.2020.0298b

References

1. Kotz D, Batra A, Kastaun S: Smoking cessation attempts and common strategies employed—a Germany-wide representative survey conducted in 19 waves from 2016 to 2019 (The DEBRA Study) and analyzed by socioeconomic status. *Dtsch Arztebl Int* 2020; 117: 7–13.
2. Hofmann P, Benden C, Kohler M, Schuurmans MM: Smoking resumption after heart or lung transplantation: a systematic review and suggestions for screening and management. *J Thorac Dis* 2018; 10: 4609–18.

Michael Noll-Hussong, MD

Psychosomatic Medicine and Psychotherapy
Saarland University Medical Center
Kirrberger Str. 100
66421 Homburg
Germany
Michael.Noll-Hussong@uks.eu

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

In our current study using data from the German Study on Tobacco Use (DEBRA), we reported that in Germany, e-cigarettes are currently the most common smoking cessation aid, initiated by smokers themselves (1). In order to weight up the opportunities and risks of e-cigarettes for tobacco cessation, it is important to distinguish tobacco smoke from nicotine in traditional cigarettes.

Tobacco smoke contains some 5000 chemicals, of which at least 98% have carcinogenic, cardiovascular, or respiratory effects. Tobacco smoke harms almost every organ in the human body and is the biggest avoidable risk factor for a multitude of disorders, which may result in reduced quality of life and premature death. The nicotine in tobacco is primarily responsible for the development of dependency and withdrawal symptoms, but compared with tobacco smoke it has a relatively low potential for causing harm. A clinical study of more than 3000 patients with chronic obstructive pulmonary disease, for example, did not find any association between the use of nicotine replacement therapy over a time period of up to five years and cardiovascular disorders or other serious events (2).

The aim of smoking cessation is primarily to stop tobacco consumption and the associated inhalation of tobacco smoke completely and permanently. Unfortunately, tobacco abstinence is extremely difficult because nicotine is so addictive. Pharmacological treatments aim to alleviate symptoms that arise primarily as a result of nicotine withdrawal and thus help to prevent a relapse. This is also the approach of nicotine replacement therapy, an established support method whose efficacy has been confirmed in more than 130 clinical studies in more than 64 000 smokers.

In principle, e-cigarettes resemble nicotine replacement therapy: the nicotine in the tobacco cigarette is substituted without the harmful adverse effects of tobacco smoke. The difference to nicotine replacement therapy—which is based mainly on patches, chewing gum, or nicotine mouth spray—is that the amplifier effect of nicotine intake is largely retained and that the method of nicotine administration is not uncoupled from the effect of the nicotine. Furthermore, e-cigarettes are by no means risk-free, but several orders of magnitude less harmful than tobacco smoke (3). Many studies from different countries have shown—in the same way as our recent study from Germany (1)—that e-cigarettes are popular among

smokers as a smoking cessation aid. Our data also show that half of the smokers in Germany who use e-cigarettes as cessation aids use liquids not containing nicotine (1).

The first randomized controlled trials of the efficacy and safety of e-cigarettes in tobacco cessation showed that e-cigarettes compared with placebo can increase tobacco abstinence in the long term without causing serious adverse effects were seen in the study period. In the latest and so far largest study, e-cigarettes were even superior to nicotine replacement therapy in terms of abstinence from tobacco products. But this study also showed that a large proportion of consumers still used these e-cigarettes one year on.

DOI: 10.3238/arztebl.2020.0299

References

1. Kotz D, Batra A, Kastaun S: Smoking cessation attempts and common strategies employed—a Germany-wide representative survey conducted in 19 waves from 2016 to 2019 (The DEBRA Study) and analyzed by socioeconomic status. *Dtsch Arztebl Int* 2020; 117: 7–13.
2. Murray RP, Bailey WC, Daniels K, et al.: Safety of nicotine polacrilex gum used by 3,094 participants in the lung health study. *Chest* 1996; 109: 438–45.
3. National Academies of Sciences Engineering and Medicine: Public health consequences of e-cigarettes. www.nationalacademies.org/hmd/Reports/2018/public-health-consequences-of-e-cigarettes.aspx. (last accessed on 11 March 2020).
4. Hajek P, Phillips-Waller A, Przulj D, et al.: A randomized trial of E-cigarettes versus nicotine-replacement therapy. *N Engl J Med* 2019; 380: 629–37.

Prof. Dr. Daniel Kotz, PhD MSc MPH

Institut für Allgemeinmedizin, Schwerpunkt Suchtforschung und klinische Epidemiologie, Medizinische Fakultät der Heinrich-Heine-Universität Düsseldorf
Research Department of Behavioural Science and Health, Institute of Epidemiology and Health Care, University College London, London, UK
Daniel.Kotz@med.uni-duesseldorf.de

Prof. Dr. med. Anil Batra

Sektion für Suchtmedizin und Suchtforschung,
Universitätsklinik für Psychiatrie und Psychotherapie,
Tübingen

Dr. rer. nat. Sabrina Kastaun

Institut für Allgemeinmedizin, Schwerpunkt Suchtforschung und klinische Epidemiologie, Medizinische Fakultät der Heinrich-Heine-Universität Düsseldorf

Conflict of interest statement

Prof Kotz is the elected representative of the German College of General Practitioners and Family Physicians involved in updating the S3 clinical practice guideline "Screening, Diagnostics, and Treatment of Harmful and Addictive Tobacco Use"

Prof Batra has received funding from Pfizer into a third-party account. He is a behavioral therapist and has developed his own tobacco withdrawal program ("non-smoker in 6 weeks"). He is the coordinator and director of the named S3 guideline "Screening, Diagnostics, and Treatment of Harmful and Addictive Tobacco Use" (AWMF registry No 076–006).

Dr. Kastaun declares that no conflict of interest exists.