Published in final edited form as:

Methods Mol Biol. 2015 ; 1312: 77–86. doi:10.1007/978-1-4939-2694-7_11.

Multiple immunoblots by passive diffusion of proteins from a single SDS-PAGE gel

Biji T. Kurien, PhD[†], R. Hal Scofield, MD^{†,§,‡}

[†] Arthritis and Clinical Immunology Program, Oklahoma Medical Research Foundation, Oklahoma City, Oklahoma, OK 73104, USA

[§] Department of Medicine, University of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma, OK 73104, USA

[‡] Department of Veterans Affairs Medical Center, Oklahoma City, Oklahoma, OK 73104, USA

Summary

Western blotting enables the detection and characterization of proteins of low abundance. Sodium dodecyl sulfate (SDS) polyacrylamide gel separated proteins are normally transferred electrophoretically to nitrocellulose or polyvinylidene difluoride membranes. Here we describe the transfer proteins [Ro 60 (or SSA) autoantigen, 220 and 240 kD spectrin antigens and prestained molecular weight standards] by diffusion from SDS polyacrylamide gels at 37°C. Up to twelve immunoblots can be obtained from a single gel by this method.

Keywords

Non-electrophoretic transfer; Immunoblots; Bi-directional transfer; Nitrocellulose membrane; Autoantibodies; Autoantigens

1. Introduction

Western blotting, involving electrophoretic transfer of proteins to a microporous membrane support with subsequent immunodetection (1), has made a tremendous impact in the field of immunology. Proteins separated on sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS PAGE) are transferred electrophoretically to mainly nitrocellulose or polyvinylidene difluoride (1–4). Protein transfer from gels to membranes has been achieved in three ways: (a) simple diffusion (3) (b) vacuum-assisted solvent flow (5,6) and (c) electrophoretic elution (1,7,8). Only electrophoretic elution has been used widely owing to a variety of reasons, including efficiency, simplicity and length of transfer. Bi-directional transfer procedure was demonstrated in 1980, when Smith and Summers (9) transferred DNA and RNA from polyacrylamide gels to nitrocellulose in 36 h to obtain two blots. Here we demonstrate that proteins can be efficiently transferred non-electrophoretically from SDS-PAGE gels to nitrocellulose membranes. A similar procedure has been used in 1982

(10) to transfer proteins from thin (0.5 mm) native isoelectric focusing gels to nitrocellulose membranes to obtain two blots in 1 h. Diffusion mediated transfer of immunoglobulins from one side of a native gel after isoelectric focusing to an antigen-coated nitrocellulose sheet has been achieved in a similar fashion (11,12) We were able to transfer and immunologically detect a 60,000 molecular weight autoantigen (Ro 60) (Fig. 2) as well as the spectrin antigens (molecular weight >200,000) (Fig. 3) using this method. When prestained molecular weight standards were transferred, only markers up to 118,000 molecular weight could be visualized (these proteins were not visualized immunologically) (Fig. 2 and 3).

Thus, we have obtained up to 12 immunoblots from a single gel using multiple antigens (high, intermediate and low molecular weight proteins) and multiple sera. Subsequently several investigators have also obtained similar results and also in a quantitative manner (*see Chapters 9 and 10*).

2. Materials

Prepare all solutions using ultrapure water (prepared by purifying deionized water, to attain a sensitivity of $18 \text{ M } \Omega \text{ cm}^2$ at 25°C) and analytical grade reagents. Prepare and store all reagents at room temperature (unless indicated otherwise). Diligently follow all waste disposal regulations when disposing waste materials. We do not add sodium azide to the reagents.

2.1. SDS Polyacrylamide Gel

1. Resolving gel buffer: 1.5 M Tris-HCl, pH 8.8. Add about 100 mL water to a 1 L graduated cylinder or a glass beaker (*see Note 1*). Weigh 181.7 g Tris and transfer to the cylinder. Add water to a volume of 900 mL. Mix and adjust pH with HCl (*see Note 2*). Make up to 1 L with water. Store at 4°C .
2. Stacking gel buffer: 0.5 M Tris-HCl, pH 6.8. Weigh 60.6 g Tris and prepare a 1 L solution as in previous step. Store at 4°C .
3. Thirty percent acrylamide/Bis solution (29.2:0.8) acrylamide:bis: Weigh 29.2 g of acrylamide monomer and 0.8 g Bis (cross-linker) and transfer to a 100 mL graduated cylinder containing about 40 mL of water. Add a spatula of AG 501-X8 (D) mixed-resin beads and mix for about 30 min. Make up to 100 mL with water and filter through a $0.45 \mu\text{m}$ Corning filter (*see Note 3*).
4. Store at 4°C , in a bottle wrapped with aluminum foil (*see Note 4*).
5. Ammonium persulfate: 10% solution in water (*see Note 5*).
6. N.N.N. -Tetramethyl-ethylenediamine. Store at 4°C (*see Note 6*).
7. SDS-PAGE running buffer: 0.025 M Tris, pH 8.3, 0.192 M glycine, 0.1% SDS (*see Note 7*).
8. SDS lysis buffer (5x): 0.3 M Tris-HCl (pH 6.8), 10% SDS, 25% β -mercaptoethanol, 0.1% bromophenol blue, 45% glycerol. Leave one aliquot at 4°C for current use and store remaining aliquots at -20°C (*see Note 8*).

9. Bromophenol blue (BPB) solution: Dissolve 0.1 g BPB in 100 mL water.

2.2. Immunoblotting

1. Nitrocellulose membranes.
2. Western blot transfer buffer: 0.025 M Tris, 0.192 M glycine, 20% methanol (*see* Note 9).
3. Tris buffered saline (TBS; 10x): 1.5 M NaCl, 0.1 M Tris-HCl, pH 7.4.
4. TBS containing 0.05% Tween-20 (TBST).
5. Blocking solution: 5% milk in TBS (*see* Note 10). Store at 4° C.
6. Diluent solution: 5% milk in TBST (*see* Note 10). Store at 4° C.
7. Mini PROTEAN® 3 System Glass plates.
8. Medium binder clips (1¼ inch).
9. Plastic container.
10. Wypall X-60-reinforced paper.
11. Nitro blue tetrazolium (NBT)/5-bromo-4-chloro-3-indolyl phosphate (BCIP): Dissolve 1 g NBT in 20 mL of 70% dimethylformamide (DMF). Dissolve 1 g BCIP in 20 mL of 100% DMF. Add 33 µL of BCIP and 66 µL of NBT to 10 mL of alkaline phosphatase buffer just before adding to membrane. Alternatively, use 1-Step™ NBT/BCIP readymade mix.

2.3. Antigens and conjugates

1. Purified red blood cell spectrin, anti-spectrin polyclonal antibody, anti-hemoglobin antibody.
2. BenchMark pre-stained molecular weight standards.
3. Purified bovine Ro 60: Purify Ro 60 as reported (13,14) or purchase from Immunovision, Springdale, AK, USA.
4. Prepare human erythrocyte membranes according to Dodge et al (15).

3. Methods

Carry out all procedures at room temperature unless otherwise specified.

3.1. 10 % Sodium Dodecyl Sulfate Polyacrylamide Gel Electrophoresis (SDS PAGE)

1. Mix 2.5 mL of resolving buffer, 3.33 mL of acrylamide mixture and 4 mL water in a 50 mL conical flask. Degas with helium for 10 min. Add 100 µL of SDS, 80 µL of ammonium persulfate and 10 µL of TEMED and cast gel within a 7.25 cm × 10 cm × 1.5 mm gel cassette. Allow space for stacking gel and gently overlay with isobutanol or water (*see* Note 11).

2. Prepare the stacking gel by mixing 1.25 mL of resolving buffer, 0.67 mL of acrylamide mixture and 3 mL water in a 50 mL conical flask. Degas with helium for 10 min. Add 100 μ L of SDS, 40 μ L of ammonium persulfate and 5 μ L of TEMED. Insert a 10-well gel comb immediately without introducing air bubbles.
3. Heat aliquots of bovine Ro 60, RBC membranes and human spectrin antigens at 95°C for 5 min. Do not add lysis buffer to the prestained protein standard or subject it to heat. Centrifuge the heated samples at $3000 \times g$ for 30 s to bring down the condensate. Load increasing amounts of Ro antigen (1–4 μ g) on one gel and same amounts of spectrin (3 μ g/lane) or RBC membrane antigens on two other gels along with protein standards (10 μ L/well-2 μ g/marker/lane). Add protein standards in every other lane (alternating with spectrin) in the gel with spectrin. Electrophorese at 15 mA until the sample has entered the gel and then continue at 25 mA till the dye front (from the BPB dye in the samples) reached the bottom of the gel (*see* Note 12).
4. Following electrophoresis, pry the gels plates open with the use of a spatula. The gel remains on one of the glass plates. Rinse the gel with water and transfer carefully to a container with western blot transfer buffer.
5. Cut a nitrocellulose membrane to the size of the gel and immerse in methanol. Rinse twice in distilled water and once with transfer buffer.

3.2. Non-electrophoretic Transfer

1. Immediately following SDS-PAGE, when the dye front reaches the end of the gel, turned off the power supply. Separate the gel plates with the help of a spatula or similar tool. Remove the stacking gel.
2. Rinse the gel (still supported by the bottom glass plate) carefully with deionized water to remove traces of SDS-PAGE running buffer.
3. Excise the gel with spectrin antigen such that there is one lane with the protein markers and one with the spectrin antigen.
4. Leave the gels to air dry for 5–10 min (*see* Note 13).
5. Gently lay one nitrocellulose membrane, cut to the shape of the gel, on top of the gel (*see* Note 14).
6. Gently lift the gel-membrane sandwich from the glass plate and place it on a Whatman No. 3 filter (place membrane side directly on the filter paper and the exposed gel side on top) cut to the size of the gel.
7. Place a second nitrocellulose membrane, cut to the shape of the gel, on top of the gel, followed by a Whatman No. 3 filter paper cut similarly (*see* Note 14).
8. Place the nitrocellulose-gel-filter paper sandwich between two mini-PROTEAN^R 3 System glass plates and secure with clamps.

9. Place this assembly in a pre-warmed humidified plastic container (Fig. 1) and incubate at 37°C for 30 min (*see* Note 15). Remove the membranes for immunoblotting (*see* Note 16) (Fig. 2).
10. Repeat this procedure with another set of nitrocellulose membranes and incubate the assembly at 37°C for 2 h to obtain two more blots from the same gel (*see* Note 17).
11. Use the same gel further to obtain blots by repeating this procedure to obtain a total of 12 blots. Incubate gel with the respective membranes for a period of 3 or 4 h to obtain the third and fourth sets of blots. Obtain the fifth and sixth sets of blots (Fig. 3) by incubating with the respective membranes for a period of 9 or 36 h respectively (*see* Note 18).
12. Cut excess membrane to smoothen edges and also cut the spectrin containing membrane into individual lanes (*see* Note 19).
13. Block the membranes with blocking solution for one h.
14. Add appropriate anti-sera to the membranes (anti-spectrin, anti-Ro 60 or control sera) and incubate for 2 h.
15. Rinse membrane strips with deionized water 2–3 three times (*see* Note 20).
16. Wash 5x with TBST, five min each time.
17. Add anti-human IgG or anti-rabbit IgG alkaline phosphatase conjugate (1:5000 dilution, diluted in diluent) and incubate for one h.
18. Wash as in step 14.
19. Add 500 µL of nitroblue tetrazolium/5-bromo-4-chloro-3-indolyl phosphate substrate and let bands develop.
20. Rinse 2–3 three times with deionized water (*see* Note 20).
21. Wash with TBST and arrange strips on paper board inserts (*see* Note 21).

4. Notes

1. Having water at the bottom of the cylinder helps to dissolve the Tris relatively easily, allowing the magnetic stir bar to go to work immediately. If using a glass beaker, the Tris can be dissolved faster if the water is warmed to about 37°C. However, the downside is that care should be taken to bring the solution to room temperature before adjusting pH.
2. Concentrated HCl (12 N) can be used at first to narrow the gap from the starting pH to the required pH. From then on it would be better to use a series of HCl (e.g. 6 N and 1 N) with lower ionic strengths to avoid a sudden drop in pH below the required pH.
3. Wear a mask when weighing acrylamide. To avoid exposing acrylamide to co-workers, cover the weigh boat containing the weighed acrylamide with another

weigh boat (similar size to the original weigh boat containing the weighed acrylamide) when transporting it to the fume hood. Transfer the weighed acrylamide to the cylinder inside the fume hood and mix on a stirrer placed inside the hood. Unpolymerized acrylamide is a neurotoxin and care should be exercised to avoid skin contact. Mixed resin AG 501 –X8 (D) (anion and cation exchange resin) is used when acrylamide solution is made, since it removes charged ions (e.g. free radicals) and allows longer storage. Some investigators store the prepared acrylamide along with this resin in the refrigerator. However, we filter them out before storage. The used mixed resin should be disposed of as hazardous waste. Manufacturer's warning states that this resin is explosive when mixed with oxidizing substances. The resin contains a dye that changes from blue-green to gold when the exchange capacity is exhausted.

4. The acrylamide solution can be stored at 4°C for one month. Acrylamide hydrolyzes to acrylic acid and ammonia. The acrylamide mixture, buffer and water can be prepared in large batches, frozen in aliquots (for greater day-to-day reproducibility) and used indefinitely (*see* Ref. 16). Remove the required amount, bring to room temperature and add the other ingredients for polymerization. However, in our laboratory we make the acrylamide solution fresh about every month when we cast our own gels.
5. We find it is best to prepare this fresh each time.
6. We find that storing at 4°C reduces its pungent smell.
7. Simple method of preparing running buffer: Prepare 10x native buffer (0.25 M Tris, 1.92 M glycine). Weigh 30.3 g Tris and 144 g glycine, mix and make it to 1L with water. Dilute 100 mL of 10X native buffer to 990 mL with water and add 10 mL of 10% SDS. Care should be taken to add SDS solution last, since it makes bubbles.
8. SDS precipitates at 4°C. Therefore, the lysis buffer needs to be warmed prior to use.
9. Dilute 100 mL of 10x native buffer to 800 mL with water and add 200 mL of methanol. Avoid adding methanol directly to the 10x buffer, since it precipitates its ingredients. Even in such a scenario the precipitate can be redissolved by the addition of 800 mL water.
10. Add 100 mL of 10x TBS to a 1L graduated cylinder mL and make it to about 800 mL with water. Transfer 50 g skim milk powder into the cylinder and mix stir until dissolved. Make to 1L with water. Separate 500 mL as the blocking solution. To the remaining 500 mL add 250 µL of Tween-20 (cut end of blue tip to aspirate Tween-20 easily), dissolve and use it as the diluent.
11. The gel cassette was sealed at the base using 1% agarose. Overlay the resolving gel with water for gels having acrylamide concentration lower than 8% and use isobutanol (or isobutanol saturated with water) for gels of 10% or greater (*see* Ref. 17). This overlay prevents contact with atmospheric oxygen, (which inhibits

acrylamide polymerization) in addition to helping to level the resolving gel solution.

12. Centrifuging the samples prior to the run helps remove insoluble debris, which could produce streaks in the protein lanes (revealed when stained with Coomassie blue). Add a drop of 0.1% BPB to the upper chamber buffer. This helps to form a much stronger dye front during the electrophoretic run.
13. Membrane contact with the gel is much better when the gel is not moist. Therefore it is important to dry the gel for 5 to 10 min. The membrane will now stick well to the gel and gel will peel off the bottom glass plate by just lifting the membrane.
14. Hold the two top corners of the membranes with each hand. Lower the bottom part of the membrane first on the lower part of the gel and gently release the membrane little by little to lay the complete membrane on the gel. This will prevent trapping of bubbles in between the gel and the membrane. A 10 mL pipette was used to roll out the air bubbles from the gel membrane sandwich prior to placing in transfer cassette. In the case of the gel with spectrin, cut the membrane to fit the two lanes of the gel.
15. The humid chamber consisted of a closed plastic container with a moist Terri Wipes paper towel at the bottom. The container must be big enough to contain the nitrocellulose-gel-filter paper assembly encased within the glass plates.
16. The second set of two blots was also obtained following incubation with the gel for one hour (*see* Fig. 3B, 3C).
17. While removing the nitrocellulose membranes from the gel for immunoblotting, it would be common to find that the gel comes up stuck to one of the two membranes. To remove this membrane from the gel, place a fresh, dry nitrocellulose membrane on top of the gel and gently lift the gel. The gel becomes stuck to this fresh membrane, thus releasing the other membrane.
18. Gel dries, in spite of placing in humid chamber, when incubated for longer time periods (36 h). Therefore it is best to use the blots obtained after 12 hour incubation with the membrane.
19. Cut a tiny wedge from the bottom left side of the marker lane and the main membrane sheet for orientation purposes. Also, in the case of the membranes with spectrin (*see* 3.1. item 3), excise the spectrin lane from the protein marker lane after matching each spectrin lane with its specific protein marker lane with pencil marks.
20. Rinsing the membrane strips with deionized water 2–3 times will help remove a bulk of the non-specific antibodies and help reduce the amount of TBST used subsequently and also reduce the number of washes. This wash helps to reduce non-specific binding of NBT/BCIP to the strip. The water, owing to its low ionic strength compared to TBST, will be able to remove contaminants much better than TBST. Water is much cheaper compared to TBST, in terms of money and

labor. Other investigators have found no reduction in detection of specific signals due to washing with water (16).

21. We use paper boards placed in between stacks of ELISA plates in packages of ELISA plates (Costar, Cambridge, MA, USA) for this purpose.

Acknowledgement

This work was supported by NIH grant ARO1844 and Oklahoma Center for the Advancement of Science and Technology to RHS.

References

1. Towbin H, Staehelin T, and Gordon J (1979) Electrophoretic transfer of proteins from polyacrylamide gels to NC sheets: procedure and applications. *Proc Natl Acad Sci USA* 76, 4350–4354. [PubMed: 388439]
2. Kurien BT, and Scofield RH (2006) Western blotting. *Methods* 38, 283–293. [PubMed: 16483794]
3. Renart J, Reiser J, and Stark GR (1979) Transfer of proteins from gels to diazobenzyloxymethyl paper and detection with anti-sera: a method for studying antibody specificity and antigen structure. *Proc Natl Acad Sci USA* 76, 3116–3120. [PubMed: 91164]
4. Gershoni JM, and Palade GE (1983) Protein blotting: principles and applications. *Anal Biochem.* 131, 1–15. [PubMed: 6193725]
5. Peferoen M, Huybrechts R, and De Loof A (1982) Vacuum-blotting: a new simple and efficient transfer of proteins from sodium dodecyl sulfate-polyacrylamide gels to NC. *FEBS Lett.* 145, 369–372.
6. Kurien BT, and Scofield RH (1997) Multiple immunoblots after non-electrophoretic bidirectional transfer of a single SDS-PAGE gel with multiple antigens. *J Immunol Methods* 205, 91–94. [PubMed: 9236919]
7. Protein blotting and detection. Ed: Kurien BT and Scofield RH Humana Press, 2009.
8. MacPhee DJ (2010) Methodological considerations for improving Western blot analysis. *J Pharmacol Toxicol Methods* 61, 171–177. [PubMed: 20006725]
9. Smith GE, and Summers MD (1980) The bidirectional transfer of DNA and RNA to nitrocellulose or diazobenzyloxymethyl-paper. *Anal Biochem.* 109, 123–129. [PubMed: 6162401]
10. Reinhardt MP, and Malamud D (1982) Protein transfer from isoelectric focusing Gels: the native blot. *Anal Biochem.* 123, 229–235. [PubMed: 6181706]
11. Knisley KA, and Rodkey LS. (1986) Affinity immunoblotting. High resolution isoelectric focusing analysis of antibody clonotype distribution. *J Immunol Methods* 95, 79–87. [PubMed: 3537139]
12. Scofield RH, Zhang F, Kurien BT, Anderson CJ, Reichlin M, Harley JB, et al. (1996) Development of the anti-Ro autoantibody response in a patient with systemic lupus erythematosus. *Arthritis Rheum.* 39, 1664–1668. [PubMed: 8843856]
13. Deutscher SL, Harley JB, and Keene JD (1988) Molecular analysis of the 60 kD human Ro ribonucleoprotein. *Proc Natl Acad Sci USA* 85, 9479–9483. [PubMed: 3200833]
14. Ben-Chetrit E, Gandy BJ, Tan EM, and Sullivan KF (1989) Isolation and characterization of a cDNA clone encoding the 60 kDa component of the human SS-A/Ro ribonucleoprotein autoantigen. *J Clin Invest.* 83, 1284–1292. [PubMed: 2649513]
15. Dodge JT, Mitchell C, and Hanahan DJ (1963) The preparation and chemical characteristics of hemoglobin-free ghosts of human erythrocytes. *Arch Biochem Biophys.* 100, 119–130. [PubMed: 14028302]
16. Electrophoresis. Appendix I In: Harlow E, and Lane D, Ed. *Antibodies. A laboratory manual.* Cold Spring Harbor Laboratory, New York, 1988, 638.
17. Electrophoresis. Appendix I In: Harlow E, and Lane D, Ed. *Antibodies. A laboratory manual.* Cold Spring Harbor Laboratory, New York, 1988, 636.

18. Wu M, Stockley PG, and Martin WJ 2nd. (2002) An improved Western blotting technique effectively reduces background. *Electrophoresis* 23, 2373–2376. [PubMed: 12210190]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1:

Gel membrane assembly for the non-electrophoretic transfer of proteins from SDS-PAGE gels to nitrocellulose membranes to obtain up to 12 blots. The polyacrylamide gel is sandwiched between 2 membranes, filter paper and glass plates and incubated at 37°C for varying periods of time to obtain up to 12 blots. (Reproduced from Ref 2 with permission from Elsevier).

Figure 2:

Ro 60 immunoblots obtained following non-electrophoretic transfer for 30 min or 2 h. **Fig. 2A:** One of two Ro 60 blots obtained by the first incubation of nitrocellulose membranes on either side of the gel at 37°C for 30 min. **Fig. 2B:** A blot from the second set (blots 3 and 4) obtained from the same gel following incubation at 37°C for 2 h. (Reproduced from Ref 6 with permission from Elsevier).

Figure 3:

Immunoblots obtained using spectrin as antigen. Lane 1 in A-F shows prestained SDS-PAGE molecular weight standards. Lane 2 shows spectrin probed with either preimmune (B,C,F) or anti-spectrin rabbit sera (A,D,E). **Fig. 3A** shows one of the first two blots obtained and has been probed with anti-spectrin. **Fig. 3B** shows the second of the two blots of the first set obtained from the reverse side of the gel and was probed with pre-immune sera. **Fig. 3C, 3D** show the fifth set (blots 9 and 10) of immunoblots probed with preimmune and anti-spectrin sera, respectively. **Fig. 3E and 3F** show the sixth set (blots 11 and 12) of blots probed with anti-spectrin and preimmune sera, respectively. (Reproduced from Ref 6 with permission from Elsevier).