Peri-operative management of frailty in the orthopedic patient

Pragati H. Mamtora^{a,b}, Michelle A. Fortier^{a,c,d,e}, Sheila R. Barnett^f, Lauren N. Schmid^a, Zeev N. Kain^{a,c,g,h,*}

^a UCI Center on Stress and Health, School of Medicine, University of California-Irvine, Irvine, CA, USA

^b Chicago Medical School, Rosalind Franklin University of Medicine and Science, North Chicago, IL, USA

^c Department of Anesthesiology and Perioperative Care, University of California-Irvine, Irvine, CA, USA

^d Sue & Bill Gross School of Nursing, University of California-Irvine, Irvine, CA, USA

^e Department of Pediatric Psychology, Children's Hospital of Orange County, Orange, CA, USA

^f Department of Anesthesiology, Pain Medicine, and Critical Care, Beth Israel Deaconess Medical Center, Boston, MA, USA

^g Department of Pediatrics, Children's Hospital of Orange County, Orange, CA, USA

^h Yale Child Study Center, Yale University School of Medicine, New Haven, CT, USA

ARTICLE INFO

Keywords:

Anesthesiology
Orthopedic surgery
Frailty
Geriatrics
Perioperative management

ABSTRACT

The purpose of this investigation was to summarize current research on diagnosis, outcomes, and management of frail patients undergoing orthopedic surgery. Because frail patients are at increased risk of negative post-operative outcomes including increased 30-day mortality and post-operative complications including infections and delirium, such a review is timely. Strategies including supervised exercise training programs before surgery, early identification of frailty, prophylactic antibiotics, regular drug chart review, regular monitoring of electrolytes, and other strategies to prevent post-operative delirium are helpful in the management of frail orthopedic patients. It is important for surgeons and anesthesiologists to take action in attempt to alleviate adverse post-operative outcomes in frail patients. Ultimately, more research is needed to identify new strategies and to evaluate whether pre-operative optimization can effectively mitigate post-operative outcomes in large-scale randomized controlled trials.

1. Introduction

With medical and technological advances, global life expectancy has increased significantly in the past few decades and by year 2030, at least 30% of the population in the United States is expected to be older than 65 years.¹ To keep pace with the increasing lifespan, surgical and anesthetic services will need to be adapted to provide safe and specialized care for the older population. Accordingly, there has been heightened attention focused on the concept of frailty in the older adult surgical population in the past decade.

It was only in 2001 that Fried and colleagues defined frailty as a biological condition with its own symptoms and functional impairments that is not synonymous to advanced age.² An international panel in 2011 concluded that frailty is a multidimensional concept comprised of 6 domains including physical performance, gait speed, mobility, nutritional status, mental health, and cognition.³ We and others conclude that ultimately frailty is a measure of decreased physiologic reserve across multiple organ systems.⁴ Frailty has been reported to be associated with an increased perioperative complication rate in older adults

undergoing elective and emergency surgeries^{4–7} and mortality rates are higher among frail patients undergoing surgery.^{8–11} Total Hip Arthroplasty (THA) and Total Knee Arthroplasty (TKA) surgeries are specifically associated with increased morbidity and mortality in the frail population^{12,13}.

As a direct consequence of the growing aging population, the demand for orthopedic surgical procedures is increasing. A recent study predicted the demand for THA to increase by 174%–572,000 procedures per year and the demand for TKA to increase by 673% to 3.48 million procedures per year by the year 2030. The growing number of orthopedic procedures in the older adult population, high prevalence of frailty in the surgical population, and increased risk for morbidity and mortality in frail older adult surgical patients all highlight the importance of increased focus on frailty in the orthopedic space. Accordingly, in this review we seek to summarize the current research and recommendations on frailty in patients undergoing orthopedic surgery. We incorporate the classical biomedical approach to disease that will include a discussion of risk factors, outcomes, diagnosis and management of frailty in this patient population.

* Corresponding author. UCI Center on Stress and Health, Department of Anesthesiology and Perioperative Care, University of California-Irvine, Orange, CA, USA.
E-mail address: zkain@uci.edu (Z.N. Kain).

<https://doi.org/10.1016/j.jor.2020.05.024>

Received 15 April 2020; Received in revised form 25 April 2020; Accepted 5 May 2020

Available online 11 June 2020

0972-978X/ © 2020 Professor P K Surendran Memorial Education Foundation. Published by Elsevier B.V. All rights reserved.

Table 1

Percent Increase in 30-Day Mortality, Re-Operation, Readmission, Clavien-Dindo Class IV Complications and Any Complications with Increase in mFI Score by Type of Orthopedic Surgery.

Orthopedic Surgery Type	30-Day Mortality (%) increase)	Re-operation (%) increase)	Readmission (%) increase)	Clavien-Dindo Class IV Complications (%) increase)	Any Complications (%) increase)
Spine	9.7%	10%	N/A	N/A	25%
Orthopedic Trauma	10.5%	1.3%	13.3%	9.6%	8.5%
THA	4.08%	3.19%	11.28%	N/A	14.8%
TKA	1.49%	N/A	9.45%	N/A	11.27%
HA	11%	N/A	N/A	4.9%	N/A

THA = total hip arthroplasty; TKA = total knee arthroplasty, HA = hemiarthroplasty, N/A = not assessed; mFI = modified frailty index.

2. How to diagnose and screen frailty?

Unfortunately, there is not one single and simple way to diagnose frailty. Rather, there have been at least 27 frailty scales developed to date¹⁴ and the screening tools that exist are either scoring systems based on physical, mental, and functional abilities or single measures of frailty based on functional capabilities.^{2, 15–19} Each screening tool has its own advantages and disadvantages; therefore, caution must be used in selecting the optimal screening tool based on both the patient and the context.

Fried developed an early standardized phenotype for frailty using data from over 5300 adults over age 65 in the Cardiovascular Heart Study.² This screening method is called the Frailty Phenotype (FP) and predicts falls, disability, worsening mobility, future hospitalization, and death, but it fails to account for changes in cognition and mood,²⁰ which are crucial components to frailty (Appendix A). However, this measure has methodological flaws as diagnosis of frailty via detection of two or more characteristics has limited theoretical grounds and there is no reliability or validity data.²¹

Since the development of the Frailty Phenotype, researchers developed the Frailty Index (FI),²² which consists of 44 items. The FI has been found to be an effective measure for frail patients undergoing major elective orthopedic surgeries including total hip and knee replacement and lumbar, sacral, or cervical laminectomy.²³ After the FI was first developed, it was revised to become the Modified Frailty Index (mFI).²⁴ This modified scale provides an even simpler measure of frailty, as the mFI consists of only 11 items along with information about functional status from the medical record to stratify patients into risk categories and predict post-operative outcomes (Appendix B).

In terms of orthopedic surgeries, the mFI has also been proven to be an effective risk assessment tool for both THA and TKA in addition to being easier to implement.^{17, 18, 25, 26} It has been shown to be a stronger predictor for readmission, post-operative complications, re-operation, and post-operative mortality.¹⁶ Throughout the past few years, the mFI has been shrunk to a 5-factor mFI (mFI-5).²⁷ A recent study comparing the credibility of the mFI-5 compared to the original 11 factor mFI has found it an equally credible predictor of frailty in all surgical subspecialties with a correlation coefficient of above 0.9 for all surgical subspecialties except cardiac and vascular surgery.²⁷ Several recent studies have found the 5-mFI a strong predictor of frailty and post-operative morbidity and mortality in a wide variety of surgeries including primary hip and knee arthroplasty, kyphoplasty vertebral augmentation, posterior lumbar fusion, distal radius fractures repair.^{28–31}

The Frailty Phenotype has five domains and all domains have to be collected specifically for the purpose of the phenotype. Some domains also require the use of specialized equipment and training for the collection of data such as weakness and slow gait domains, which require the measurement of grip strength and gait speed, for example. Furthermore, the FI has its own challenges as it has 42 variables and thus requires the use of an algorithm to extract information from individual electronic medical records and administrative databases.³²

Ultimately, the FP, FI, and the 5 factor mFI are all useful for

assessing and screening for frailty pre-operatively for patients undergoing major elective orthopedic surgery³² but differ in the level of complexity. The mFI-5 has the benefit of simplicity with only 5 variables to measure and without the need for specialized equipment and therefore may be the most clinically useful. In the research setting, the original FI may have the most utility as it is more detailed in its inclusion of 70 clinical deficits in a wide variety of domains to more precisely stratify individuals into risk categories. However, both the mFI and mFI-5 have been proven useful in identifying frailty before surgery; therefore, selection of the optimal measure for pre-operative frailty will depend upon feasibility considerations such as available data, logistical constraints, and provider preference³².

3. What are the outcomes for frailty?

Previous research has reported that frailty is highly associated with post-operative mortality.^{12, 25, 26, 33, 34} Frailty is associated with *post-operative mortality* in patients undergoing surgery for pelvis and lower extremity fractures, adult spinal deformity, femoral neck fracture, primary hip arthroplasty, and primary knee arthroplasty.^{12, 25, 26, 33, 34} Higher mFI scores are associated with increased post-operative mortality and are shown to be a stronger predictor of post-operative mortality compared to age, obesity class, and ASA class.^{12, 25, 26, 33, 34} Various studies have shown different results as to increased post-operative mortality for spine surgery, orthopedic trauma, THA, TKA, and HA (Table 1).

There is also an increased risk of post-operative complications, length of stay, and readmission in frail patients undergoing orthopedic surgery.¹² Specifically, as can be seen in Table 1, there are a significant number of *post-operative complications* that are associated with frailty. There is an increased risk of Clavien-Dindo Class IV complications and hospital acquired conditions (surgical-site infections, pneumonia, venous thromboembolism, and urinary tract infections) with increased mFI score.^{12, 13, 25, 26, 33–37} for patients undergoing spine surgery, orthopedic trauma, THA, TKA, and HA (Table 1).

Several recent studies have found an association with frailty with the occurrence of post-operative delirium.^{38–40} Frailty is also associated with higher delirium severity as measured by the Confusion Assessment Method (CAM).⁴¹ Furthermore, for post-operative delirium in frail patients, it has been shown that there is an association with delirium and longer length of hospitalization, lower baseline functional status, and pre-operative cognitive impairment.^{42–44}

4. How to manage the orthopedic frail patient?

Excellent pain control, optimizing sleep environment, minimizing tethers, cognitive reorientation with clocks, clear communication, early mobilization, and good nutrition have been reported to improve the outcome in this group of patients.^{45, 46} Identification and management of triggering factors such as sepsis, dehydration, electrolyte imbalance, and substance withdrawal is paramount.⁴⁷ The European Society of Anesthesiology recommends implementing fast-track surgery to prevent post-operative delirium in high-risk patients such as those who are

frail.^{45, 46} Specific suggestions for anesthesia management include avoiding potentially inappropriate medications such as benzodiazepines for pre-medication and monitoring of anesthesia depth to avoid excessive depth.^{45, 46}

Optimization strategies for functional status for frail patients include referring patients to obtain formal physical therapy evaluations, obtaining assistive devices, and planning for in-hospital and post discharge rehabilitative therapy. Pre-operative exercise programs often referred to as pre-habilitation to improve strength and mobility have shown minimal to no improvement in post-operative outcomes including reduced risk of discharge to a rehabilitation facility post-discharge, improved strength, and functional ability.⁴⁸ A recent systematic review by Wang et al. that included 33 studies found that only two studies reported a significant reduction in several different pain scores at 6 and 12 weeks post-operatively as a function of pre-habilitation.⁴⁸ Only 25% of the studies reported significant improvement of post-operative function which was measured by various ADL scales and various ranges of motion on the operated joint.^{48, 49} Furthermore, length of stay, cost, and quality of life were not significantly changed following these interventions.⁴⁸

To prevent common post-operative infections including respiratory, wound, and urinary infections, antibiotics and standard infection protocols should be given and followed.⁵⁰ Frail patients are also at increased risk for opportunistic infections such as MRSA (methicillin-resistant staph aureus) or clostridium difficile and iatrogenic problems that arise from unnecessary medication as there is a high rate of unnecessary prescribing in hospitals.⁵⁰ Even when drugs are prescribed appropriately, polypharmacy and altered pharmacodynamics and kinetics can lead to systemic side effects including nausea, lethargy, confusion, anorexia, dizziness, constipation, and electrolyte imbalance. Therefore, regular drug chart review and rationalization of each medication is important.⁵⁰

Appendix A. Frailty Phenotype Criteria

Frailty phenotype criteria	Measurement
Weakness	Grip strength (lowest 20% by sex, body mass index)
Slowness	Walking time per 15 feet (slowest 20% by sex, height)
Low level of physical activity	Kcal/week for lowest 20% Males 383 kcal/week Females: 270 kcal/week
Exhaustion/poor endurance	Self-reported exhaustion
Weight Loss	> 10 lbs. lost unintentionally in prior year

Appendix B. Modified Frailty Index Factors

1) Overall functional status (partial or total dependency)	+1
2) Impaired sensorium	+1
3) Diabetes mellitus (with or without insulin treatment)	+1
4) Chronic or acute lung disease (history of COPD, current pneumonia, etc.)	+1
5) Myocardial Infarction (history of myocardial infarction within 6 months)	+1
6) History of congestive heart failure	+1
7) History of angina, percutaneous coronary intervention, or prior cardiac surgery	+1
8) Hypertension requiring medications	+1
9) History of transient ischemic attacks	+1
10) History of cerebrovascular accidents/strokes with neurologic deficits	+1
11) Revascularization/amputation for peripheral vascular disease, rest pain, or gangrene	+1

COPD = chronic obstructive pulmonary disease; the Modified Frailty Index is graded on a Likert scale with a summative score out of 11.

References

- U. C. Bureau, "The Older Population in the United States: 2010 to 2050."
- Fried LP, Tangen CM, Walston J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol Ser A Biol Sci Med Sci*. 2001;56(3):M146–M157.
- Rodríguez-Mañas L, Féart C, Mann G, et al. Searching for an operational definition of frailty: a delphi method based consensus statement. The frailty operative definition-consensus conference project. *J Gerontol Ser A*. Jan. 2013;68(1):62–67.
- Shah R, Attwood K, Arya S, et al. Association of frailty with failure to rescue after low-risk and high-risk inpatient surgery. *JAMA Surg*. May 2018;153(5):e180214.
- Seib CD, Rochefort H, Chomsky-Higgins K, et al. Association of patient frailty with

- increased morbidity after common ambulatory general surgery operations. *JAMA Surg.* Feb. 2018;153(2):160.
6. Augustin T, Burnstein MD, Schneider EB, et al. Frailty predicts risk of life-threatening complications and mortality after pancreatic resections. *Surgery.* Oct. 2016;160(4):987–996.
 7. Joseph B, Zangbar B, Pandit V, et al. Emergency general surgery in the elderly: too old or too frail? *J Am Coll Surg.* May 2016;222(5):805–813.
 8. Affilo J, Motillo S, Eisenberg MJ, et al. Addition of frailty and disability to cardiac surgery risk scores identifies elderly patients at high risk of mortality or major morbidity. *Circ Cardiovasc Qual Outcomes.* Mar. 2012;5(2):222–228.
 9. Sündermann S, Dademasch A, Praetorius J, et al. Comprehensive assessment of frailty for elderly high-risk patients undergoing cardiac surgery. *Eur J Cardio Thorac Surg.* Jan. 2011;39(1):33–37.
 10. Hodari A, Hammoud ZT, Borgi JF, Tsiouris A, Rubinfeld IS. Assessment of morbidity and mortality after esophagectomy using a modified frailty index. *Ann Thorac Surg.* Oct. 2013;96(4):1240–1245.
 11. Karam J, Tsiouris A, Shepard A, Velanovich V, Rubinfeld I. Simplified frailty index to predict adverse outcomes and mortality in vascular surgery patients. *Ann Vasc Surg.* Oct. 2013;27(7):904–908.
 12. McIsaac DI, Beaulé PE, Bryson GL, Van Walraven C. The impact of frailty on outcomes and healthcare resource usage after total joint arthroplasty. *Bone Joint Lett J.* Jun. 2016;98-B(6):799–805.
 13. Dayama A, Olorunfemi O, Greenbaum S, Stone ME, McNelis J. Impact of frailty on outcomes in geriatric femoral neck fracture management: an analysis of national surgical quality improvement program dataset. *Int J Surg.* Apr. 2016;28:185–190.
 14. Bouillon K, Kivimäki M, Hamer M, et al. Measures of frailty in population-based studies: an overview. *BMC Geriatr.* Dec. 2013;13(1):64.
 15. Rolfsen DB, Majumdar SR, Tsuyuki RT, Tahir A, Rockwood K. Validity and reliability of the edmonton frail scale. *Age Ageing.* Sep. 2006;35(5):526–529.
 16. Partridge JSL, Harari D, Dhisi JK. Frailty in the older surgical patient: a review. *Age Ageing.* 2012;41(2):142–147.
 17. Rockwood K, Mitnitski A. Frailty in relation to the accumulation of deficits. *Journals Gerontol. Ser. A Biol. Sci. Med. Sci.* Jul. 2007;62(7):722–727.
 18. de Vries NM, Staal JB, van Ravensberg CD, Hobbelen JSM, Olde Rikkert MGM, Nijhuis-van der Sanden MWG. Outcome instruments to measure frailty: a systematic review. *Ageing Res Rev.* Jan. 2011;10(1):104–114.
 19. Ravaglia G, Forti P, Lucicesare A, Pisacane N, Rietti E, Patterson C. Development of an easy prognostic score for frailty outcomes in the aged. *Age Ageing.* 2008;37(2):161–166.
 20. Chan SP, Ip KY, Irwin MG. Peri-operative optimisation of elderly and frail patients: a narrative review. *Anaesthesia.* Jan. 2019;74:80–89.
 21. Sutton JL, Gould RL, Daley S, et al. Psychometric properties of multicomponent tools designed to assess frailty in older adults: a systematic review. *BMC Geriatr.* Dec. 2016;16(1):55.
 22. Rockwood K, Song X, MacKnight C, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ (Can Med Assoc J).* Aug. 2005;173(5):489–495.
 23. Ali R, Schwalb J, Nerenz D, Antoine H, Rubinfeld I. Use of the modified frailty index to predict 30-day morbidity and mortality from spine surgery. *J Neurosurg.* 2016;25(537):541.
 24. Tsiouris A, Hammoud ZT, Velanovich V, Hodari A, Borgi J, Rubinfeld I. A modified frailty index to assess morbidity and mortality after lobectomy. *J Surg Res.* 2013;183(1):40–46. <https://doi.org/10.1016/j.jss.2012.11.059>.
 25. Runner RP, Bellamy JL, Vu CCL, Erens GA, Schenker ML, Guild GN. Modified frailty index is an effective risk assessment tool in primary total knee arthroplasty. *J Arthroplasty.* Sep. 2017;32(9):S177–S182.
 26. Bellamy JL, Runner RP, Vu CCL, Schenker ML, Bradbury TL, Roberson JR. Modified frailty index is an effective risk assessment tool in primary total hip arthroplasty. *J Arthroplasty.* Oct. 2017;32(10):2963–2968.
 27. Subramaniam S, Aalberg JJ, Soriano RP, Divino CM. New 5-factor modified frailty index using American college of surgeons NSQIP data. *J Am Coll Surg.* 2018;226(2):173–181. <https://doi.org/10.1016/j.jamcollsurg.2017.11.005>.
 28. Segal DN, Wilson JM, Staley C, Michael KW. The 5-item modified frailty index is predictive of 30-day postoperative complications in patients undergoing kyphoplasty vertebral augmentation. *World Neurosurg.* 2018;116:e225–e231. <https://doi.org/10.1016/j.wneu.2018.04.172>.
 29. Traven SA, Reeves RA, Sekar MG, Slone HS, Walton ZJ. New 5-factor modified frailty index predicts morbidity and mortality in primary hip and knee arthroplasty. *J Arthroplasty.* 2019;34(1):140–144. <https://doi.org/10.1016/j.arth.2018.09.040>.
 30. Weaver DJ, Malik AT, Jain N, Yu E, Kim J, Khan SN. The modified 5-item frailty index: a concise and useful tool for assessing the impact of frailty on postoperative morbidity following elective posterior lumbar fusions. *World Neurosurg.* 2019;S1878-8750(19):30035–30038. <https://doi.org/10.1016/j.wneu.2018.12.168>.
 31. Wilson JM, Holzgrefe RE, Staley CA, Schenker ML, Meals CG. Use of a 5-item modified frailty index for risk stratification in patients undergoing surgical management of distal radius fractures. *J. Hand Surg. Am.* 2018;43(8):701–709. <https://doi.org/10.1016/j.jhsa.2018.05.029>.
 32. Cooper Z, Rogers Jr SO, Ngo L, et al. Comparison of frailty measures as predictors of outcomes after orthopedic surgery. *J Am Geriatr Soc.* Dec. 2016;64(12):2464–2471.
 33. Leven DM, Lee NJ, Kothari P, et al. Frailty index is a significant predictor of complications and mortality after surgery for adult spinal deformity. *Spine Phila. Pa.* 1976;41 23, pp. E1394–E1401, Dec. 2016.
 34. Vu CPCL, Runner RP, Reisman WM, Schenker ML. The frail fail: increased mortality and post-operative complications in orthopaedic trauma patients. *Injury.* 2017;48(11):2443–2450. <https://doi.org/10.1016/j.injury.2017.08.026>.
 35. Schmucker AM, Hupert N, Mandl LA. The impact of frailty on short-term outcomes after elective hip and knee arthroplasty in older adults: a systematic review. *Geriatr. Orthop. Surg. Rehabil.* Jan. 2019;10. <https://doi.org/10.1177/2151459319835109> 2151459319835109.
 36. Shin JI, Keswani A, Lovy AJ, Moucha CS. Simplified frailty index as a predictor of adverse outcomes in total hip and knee arthroplasty. *J Arthroplasty.* 2016;31(11):2389–2394.
 37. Folbert EC, Hegeman JH, Gierfeld J, et al. Complications during hospitalization and risk factors in elderly patients with hip fracture following integrated orthogeriatric treatment. *Arch Orthop Trauma Surg.* 2017;137(4):507–515.
 38. Bansal R, Singh S, Chatterjee P, Malhotra R, Dey AB. Impact of frailty on post-operative outcome of major orthopaedic surgery in older Indian. *J Indian Acad Geosci.* 2018;14 180–180.
 39. Kua J, Ramasom R, Rajamoney G, Chong MS. Which frailty measure is a good predictor of early post-operative complications in elderly hip fracture patients? *Arch Orthop Trauma Surg.* May 2016;136(5):639–647.
 40. Sheehan KJ, Guerrero EM, Tainter D, et al. Prognostic factors of in-hospital complications after hip fracture surgery: a scoping review. *Osteoporos Int.* Apr. 2019;1–13.
 41. Racine AM, Jones ScD RN, Gou MA Y, et al. Preoperative frailty is associated with higher postoperative delirium incidence and severity. *Innov Aging. Nov.* 2018;2(suppl_1) pp. 205–205.
 42. Kistler EA, Nicholas JA, Kates SL, Friedman SM. Frailty and short-term outcomes in patients with hip fracture. *Geriatr Orthop Surg Rehabil.* Sep. 2015;6(3):209–214.
 43. Culley DJ, Flaherty D, Fahey MC, et al. Poor performance on a preoperative cognitive screening test predicts postoperative complications in older orthopedic surgical patients. *Anesthesiology.* Nov. 2017;127(5):765–774.
 44. Tahir M, Malik SS, Ahmed U, Kozdryk J, Naqvi SH, Malik A. Risk factors for onset of delirium after neck of femur fracture surgery: a prospective observational study. *SICOT-J.* 2018;4:27.
 45. Aldecoa C, Bettelli G, Bilotta F, et al. European Society of Anaesthesiology evidence-based and consensus-based guideline on postoperative delirium. *Eur J Anaesthesiol.* Apr. 2017;34(4):192–214.
 46. Chan SP, Ip KY, Irwin MG. Peri-operative optimisation of elderly and frail patients: a narrative review. *Anaesthesia.* Jan. 2019;74:80–89.
 47. Inouye SK, Robinson T, Blaum C, et al. Postoperative delirium in older adults: best practice statement from the American geriatrics society. *J Am Coll Surg.* Feb. 2015;220(2) 136–148.e1.
 48. Wang L, Lee M, Zhang Z, Moodie J, Cheng D, Martin J. Does preoperative rehabilitation for patients planning to undergo joint replacement surgery improve outcomes? A systematic review and meta-analysis of randomised controlled trials. *BMJ Open.* Feb. 2016;6(2) e009857.
 49. Shem Tov L, Matot I. Frailty and anesthesia. *Curr Opin Anaesthesiol.* Jun. 2017;30(3):409–417.
 50. Oliver D. Medical input, rehabilitation and discharge planning for patients with hip fracture: why traditional models are not fit for purpose and how things are changing. *Curr Anaesth Crit Care.* Feb. 2005;16(1):11–22.