Published in final edited form as:

J Clin Pediatr Dent. 2020 ; 44(1): 28–34. doi:10.17796/1053-4625-44.1.5.

Identification of *non-Streptococcus mutans* bacteria from predente infant saliva grown on mitis-salivarius-bacitracin agar

Yan Zeng^{1,2}, Moustafa Youssef¹, Lin Wang^{1,3}, Naemah Alkhars¹, Marie Thomas⁴, Rita Cacciato¹, Shuang Qing⁵, Oriana Ly-Mapes¹, Jin Xiao¹

¹Eastman Institute for Oral Health, University of Rochester Medical Center, Rochester, NY, USA

²Department of Forensic Medicine, North Sichuan Medical College, Nanchong, Sichuan, China

³Perking University School of Stomatology, Beijing, China

⁴Department of Family Medicine, University of Rochester Medical Center, Rochester, NY, USA

⁵University of Rochester, Rochester, NY, USA

Abstract

Objective: Although mitis-salivarius-bacitracin (MSB) agar is a commonly used selective medium for detecting *Streptococcus mutans* in clinical studies, non-*S. mutans* microorganisms are cultivatable on MSB agar. Since few studies have identified non-*S. mutans* bacteria grown on MSB, this study aimed to identify and differentiate MSB-grown non-*S. mutans* bacteria from predente infants' oral cavity.

Methods: The saliva from 51 predente infants were plated on MSB agars. Bacteria colonies were characterized based on their morphology under direct visualization and light microscopic observation. Colony PCR targeting *S. mutans htrA* locus and 16S rRNA DNA sequencing were used for further bacteria identification.

Results: Overall, 80% of the predente infants had oral bacteria grown on the MSB agar. Nine bacteria were identified, including *S. mutans*, *Staphylococcus epidermidis*, *Klebsiella quasi-pneumoniae*, *Klebsiella pneumoniae*, *Enterobacter kobei*, *Enterococcus faecalis*, *Staphylococcus hominis*, *Streptococcus anginosus* and *Phytobacter*. The most frequently detected bacteria were *S. epidermidis* (41.5%), followed by *E. kobei* (24.4%), *K. pneumoniae* (17.1%) and *S. mutans* (9.8%).

Conclusions: Multiple non-*S. mutans* bacteria from infants' oral cavity could grow on MSB agar. Caution should be exercised in counting the colony forming units of *S. mutans* from oral samples on MSB agar to avoid overestimation by assuming that all colonies on the MSB agar are

Corresponding author: Jin Xiao, DDS, PhD, Address: 625 Elmwood Ave, Rochester, NY, 14618, Phone: +1 (585) 273 1957, Fax: +1 (585) 273 1237, jin_xiao@urmc.rochester.edu.

Author contributions

Y Zeng and J Xiao contributed to the conception, design, data acquisition, analysis, and interpretation, drafted and critically revised the manuscript; S Qing contributed to data acquisition; M Youssef, L Wang, N Alkhars, M Thomas, R Cacciato and O Ly-Mapes contributed to data acquisition and interpretation and critically revised the manuscript. All authors gave final approval and agree to be accountable for all aspects of the work.

Declaration of interests

The authors declare no conflict of interests.

S. mutans. Using the colony morphological guide we summarized, these non-*S. mutans* bacteria could be distinguished from *S. mutans*. Our study provides a key reference to pediatric cariology clinical-epidemiological studies that commonly use MSB to identify/quantify *S. mutans* in infants and young children.

Keywords

Oral bacteria; Mitis-salivarius-bacitracin agar; Predente infants; Saliva; *Streptococcus mutans*; Colony morphology

Introduction

Dental caries is a chronic infectious disease, initiated from the virulent dental biofilms/plaque formed on tooth surfaces¹. Within the dental biofilms/plaque, oral cariogenic bacteria metabolize dietary carbohydrate, produce acid and initiate demineralization of tooth enamel, and further leads to dental caries². *Streptococcus mutans* has been considered as a well-known culprit for dental caries³ including early childhood caries (ECC)^{4–10}, due to its acidogenicity, aciduricity and capability of synthesizing dental plaque extracellular matrix^{4,11–14}.

Various culture-independent methods have been used to identify and quantify oral *S. mutans* carriage, including biochemical tests¹⁵, DNA probes¹⁶, polymerase chain reaction-restriction (PCR)¹⁷ and 16S rDNA sequencing comparison methods¹⁸. Although culture-independent methods have been widely used to identify the presence and measure relative abundance of *S. mutans* in oral samples, the critical quantitative caries risk indicator remains to be that individual with more than 10⁵ colony forming unit (CFU) /ml of salivary *S. mutans* are considered as high risk for caries^{19–26}. Therefore, isolation, identification and quantification of the *S. mutans* from oral samples (saliva, plaque and swab) using culture-dependent methods still present as essential components for clinical and epidemiological studies, in addition to other culture independent assays^{27–30}.

Mitis-salivarius-bacitracin (MSB) agar is a commonly used selective medium for isolating and quantifying *S. mutans* in many cariology clinical and epidemiological studies^{27,31–34}. However, previous reports have discussed the non-specificity of isolating *S. mutans* using MSB agar^{35–37}. For instance, *S. sobrinus* could also be isolated from the oral cavity of young children using MSB³⁶. Nevertheless, few studies have described and identified other non-*S. mutans* species isolated from MSB^{38,39}. Moreover, our prospective birth-cohort study that examined the colonization of *S. mutans* in infants have observed multiple bacteria that grew on the MSB, in addition to *S. mutans*. Therefore, the aim of this study was to characterize, identify and differentiate these non-*S. mutans* bacteria isolated from predente infants saliva samples using MSB agar, which could serve as a key reference to cariology clinical-epidemiological studies that commonly use MSB to identify and quantify *S. mutans* in infants and young children.

Materials and Methods

Study Subject Enrolment and Clinical Sample Collection

This study was approved by the University of Rochester Institutional Review Board (#67191). Infants met the following conditions were excluded: born pre-term (<37 weeks of gestation at birth), with low birth weight (<2500), with Orofacial deformity (cleft lip, cleft palate, oral-pharyngeal mass), with Down syndrome, immediately admitted into the Neonatal Intensive Care Unit (NICU) after birth and stayed in NICU less than two weeks, or received antifungal oral and/or systematic treatment before the study visit.

The saliva samples from 51 healthy infants (0-6 months) were collected using SalivaBio Infant's Swab (SIS, Salimetrics, Inc. Carlsbad CA, USA). A trained dentist securely held one end of the SIS swab and placed the other end under the infant's tongue (when possible) or at the corners of the mouth. Infants were allowed to chew on the swab. Saliva were collected in intervals by resting the swab inside the mouth for 20 sec, and then re-introducing into the infants' mouth for another 20 sec. Saliva collection continued until the lower third of the swab is saturated (some participants required up to 90 seconds of total collection time). After the saliva collection, the SIS was stored in a sterilized tube and kept on ice and transferred to the laboratory located at the University of Rochester Center for Oral Biology within two hours.

MSB Agar Preparation and Infant Saliva Sample Processing

To prepare the MSB agar, 54 g mitis salivarius agar and 90 g sucrose were added in 600 ml distilled H₂O. The mixture was autoclaved. After the temperature cooled down to 50 °C, 700 µl of 1% tellurite solution and 700 µl of 3 mg/ml bacitracin were added. Poured agar plates were left at room temperature overnight. Infant saliva samples were centrifuged for 15 minutes at 1500 g to obtain infant saliva. In general, 400-1000 µl of saliva was harvested from the centrifuging process. 50 µl of saliva suspensions with no dilution, and with 10⁻¹, 10⁻², and 10⁻³ dilutions were plated onto MSB plates in duplicates and incubated in 5% CO₂ at 37 °C for 72 hours.

Bacteria Identification

Three methods were used to identify bacteria isolated from MSB agar, including colony morphology differentiation⁴⁰, colony PCR with *S. mutans* specific primers and 16s rRNA gene sequencing. *S. mutans* UA159 (ATCC® 700610™) and *S. sobrinus* (ATCC® 33478™) were used as controls.

1) Colony Morphology Differentiation—After 72-hour incubation, bacteria colonies isolated from the MSB plates were observed with naked eyes and under the light microscope (40×, Olympus, Japan). Direct photos and light microscope images were obtained.

2) Colony PCR Targeting *S. mutans*—Three to five colonies were picked from MSB agar plates and added into the PCR strip tube with a mixture of GoTaq® colorless master mix (Promega, USA) and primer pair Sm479F/R (Sm479F: 5'-TCGCGAAAAAGATAAACAACA-3' and Sm479R: 5'-

GCCCCTTCACAGTTGGTTAG-3')⁴¹. This primer set has been used previously to identify *S. mutans* by targeting the segment comprises a portion of the *htrA* locus and a part of an intergenic locus of the *S. mutans* genome⁴¹. PCR were performed on a Thermocycler (Thermo Fisher Scientific, USA). The PCR reaction conditions were: 1) 5 min at 95°C, 2) 35 cycles of: 15 sec at 95°C, 30 sec at an annealing temperature 55°C and 1 min at 72°C, 3) followed by 10 min at 72°C. The PCR amplicon product (479 bp) was imaged using an Invitrogen™ E-Gel™ Power Snap Electrophoresis Device and 2% EX gel (Thermo Fisher Scientific, USA).

3) 16S rRNA Gene Sequencing—Bacteria genomic DNA was extracted from the overnight bacteria liquid culture medium using MasterPure DNA Extraction Kit (Epicentre, USA). Extracted DNA was evaluated for quality and quantity using the NanoDrop Spectrophotometer (Thermo Fisher Scientific, USA). Total DNA was used as a template for PCR amplification of 1.4 kbp fragment of the 16S ribosomal RNA gene. Standard 16S ribosomal RNA PCR primers (27F and 1492R) were used. Water was used as a negative control. All PCR fragments were analyzed for quantity and quality by agarose gel electrophoresis. All PCR fragments were sequenced using the 27F and 1492R primers. All sequencing was performed with the BigDye Terminator Cycle Sequencing Kit, V 3.1, and analyzed on an ABI 3730XL Genetic Analyzer. All sequence reactions were then evaluated and aligned. Consensus sequences were generated for each fragment and compared with the existing species database (NCBI) by the BLAST search algorithm.

Results

Non-*S. mutans* bacteria strains grown on MSB agar

Overall, 80% (41/51) of the predente infants had bacteria grown on the MSB agar. Nine bacteria strains including *S. mutans* and another eight non-*S. mutans* were isolated from predente infants' saliva samples using MSB agar. These bacteria were further identified using 16S rRNA gene sequencing, including *S. mutans*, *Staphylococcus epidermidis*, *Klebsiella quasi-pneumoniae*, *Klebsiella pneumoniae*, *Enterobacter kobei*, *Enterococcus faecalis*, *Staphylococcus hominis*, *Streptococcus anginosus* and *Phytobacter*. The colony morphology from direct visualization and light microscopic observation are summarized in Figure 1 and Table 1. Bacterial characteristics, results from colony PCR that targeted *S. mutans htrA* locus and 16S rRNA gene sequencing identification and clinical relevance were also presented in Table 1. Regarding the colony morphology, *S. mutans* (A-9 and B-9) and *S. sobrinus* (A-10 and B-10) shared similar colony characteristics; both of them had unique morphology of frosted glass colonies which rooted into the agar. The colonies of *S. sobrinus* were surrounded by more distinct polysaccharide.

Despite the similarity between *S. mutans* and *S. sobrinus* colonies, all other non-*S. mutans* bacteria that grew on the MSB agar were distinguishable based on their colony morphology using our summarized colony morphological guide in Figure 1 and Table 1. The colony size (diameter) of the MSB isolated oral bacteria ranged from 0.5 mm to 4 mm. *S. epidermidis*, *E. kobei* and *S. hominis* tended to form larger colonies, with an average size 2-4 mm. The colonies of *S. mutans*, *S. sobrinus*, *S. anginosus* and *Phytobacter* were on the smaller scale,

with an average size of 0.5-1 mm. Majority of the colonies were dark black (except for *K. pneumonia* and *E. kobei*). Several bacteria colonies were accompanied by distinct smell, e.g. strong sour fermentation smell (*E. kobei*) and sweet smell (*K. pneumonia*). Four bacteria did not have distinct smell (*S. hominis*, *S. anginosus*, *S. mutans* and *S. sobrinus*).

The clinical relevance of these MSB-isolated bacteria was summarized in Table 1. *S. mutans* and *S. sobrinus* were considered as cariogenic strains. Several other bacteria were considered as pathogens for nosocomial infection. For example, *K. pneumonia* was closely associated with pneumonias, urinary tract infections, bacteremias, and liver abscesses ⁴², and *E. faecalis* could cause endocarditis, bacteremia, urinary tract infections, meningitis, and other fatal forms of systemic and local infection ⁴³.

Bacteria Detection Frequency and Quantity

The bacteria detection frequency and quantity were shown in Table 2. Eighty percent (41/51) of the infants' saliva had bacteria that grew on MSB agar. Among the isolated bacteria grown on MSB, *S. epidermidis* was the most frequently detected bacteria (41.5%), followed by *E. kobei* (24.4%), *K. pneumoniae* (17.1%) and *S. mutans* (9.8%). *E. kobei* was the most abundant bacteria grown on MSB (CFU/ml $1.7 \pm 4.4 \times 10^7$), whereas *S. hominis* was the least abundant bacteria grown on MSB.

Discussions

Culture-dependent methods that use MSB medium to isolate, cultivate, identify and quantify *S. mutans* are commonly used in cariology clinical and epidemiology studies. In contrast to the popularity of using MSB agar in these studies, limited selectivity of using MSB to isolated/identify *S. mutans* from predente infants' saliva had not been reported. Moreover, limited study have characterized and identified non-*S. mutans* microorganisms isolated from MSB. To fill the gap, our study detailed the characteristics and differentiation of MSB-isolated non-*S. mutans* bacteria, which could be used as important references for cariology clinical and epidemiological studies. Our study identified eight non-*S. mutans* bacteria isolated from infants' saliva using MSB. Several bacteria shared similar appearance with *S. mutans* on the MSB. For instance, *S. sobrinus*, *S. anginosus* and *Phytobacter*. Without proper observation and differentiation, these colonies could be falsely recognized as *S. mutans*, which might lead to inflated positive *S. mutans* detection in clinical studies.

The mutans streptococci constitute a group of oral cariogenic species, with *S. mutans* and *S. sobrinus* being the most frequently detected strains from the oral cavity of individuals at high risk for dental caries ⁴⁴⁻⁴⁶. *S. mutans* has been detected from ~80% to 90% of adults' and ~50% to 100% of preschool children's oral cavity ²⁷, but much less frequent (~10%) in the oral cavity of infants ^{47,48}. The oral colonization by *S. mutans* is considered to take place around 26 months ⁴⁹. Li *et al* showed the *S. mutans* detection rate at 70-80% from old children aged three and five ⁵⁰. Our study showed that *S. mutans* presented in predente infants' oral cavity, with a detection rate of 9.8%. Interestingly, *S. sobrinus* was not detected from any of the 51 predente infants.

The oral cavity is considered as sterile at birth, and quickly colonized by multiple microorganisms through infants' breathing, feeding and contacting with family, medical personals and care-giver⁵¹. Understanding the development and diversity of human oral microorganisms in early infancy is important to understanding the factors that relate to oral health and disease from a microbiological standpoint. In this study, we showed that *S. epidermidis*, *E. kobei* and *K. pneumoniae* were colonized in predente infants' mouth. As reported previously, *E. faecalis*, *E. coli*, *Enterobacter cloacae*, *K. pneumoniae*, *S. epidermidis* and *Staphylococcus haemolyticus* were found as predominant species in infants' intestinal microflora⁵². These three bacteria are commonly recognized as nosocomial pathogens,^{42,53,54} *S. epidermidis* is found ubiquitously on healthy human skin and mucosal surfaces⁵⁵, readily colonizing newborns⁵⁶ and remaining part of the normal microflora throughout life⁵⁷, and it rarely causes infections in healthy tissue⁵⁸. *E. kobei* is a new species of the family *Enterobacteriaceae* resembling *Enterobacter cloacae*⁵⁹, which has been reported as an important opportunistic and multi-resistant bacterial pathogen for humans in hospital wards⁶⁰. *K. pneumoniae*'s virulence factors are closely associated with pneumonia infections⁴². It is speculated that most of the eight non-*S. mutans* bacteria isolated from MSB might be transient species due to the particularities of neonatal innate immunity.

In summary, eight non-*S. mutans* bacteria strains from 51 predente infants' saliva grew on the MSB agar and were identified using 16S rRNA sequencing. Caution should be exercised in counting the colony forming units of *S. mutans* from oral samples on MSB agar to avoid overestimation by assuming that all colonies on the MSB agar are *S. mutans*. Overall, considering the non-*S. mutans* bacteria colonies were distinguishable from *S. mutans* based on their colony morphological characteristics, the lower cost and better affordability when compared to 16S rRNA sequencing, our study results suggest that MSB agar could still be used for initial identification/quantification of *S. mutans* in clinical and epidemiology study setting, followed by further validation using colony PCR that targets the *S. mutans* specific locus.

Conclusions

Our study indicated that a variety of non-*S. mutans* bacteria from predente infant's saliva could grow on MSB agar, including *S. epidermidis*, *K. quasi-pneumoniae*, *K. pneumoniae*, *E. kobei*, *E. faecalis*, *S. hominis*, *S. anginosus*, *Phytobacter*. Using the colony morphological guide we summarized, these bacteria could be distinguished from *S. mutans*, which provides an important key reference to cariology clinical-epidemiological studies that commonly use MSB selective medium to identify and quantify *S. mutans* in infants and young children.

Acknowledgement

Dr. Xiao's research was supported by the National Institute of Dental and Craniofacial Research grant K23DE027412; Dr. Zeng's research was supported by the North Sichuan Medical College fund CBY14-QD-03. The funding agencies had no role in the study design, data collection, analyses, decision to publish, or preparation of the manuscript.

References:Uncategorized References

1. Tanzer JM. Dental caries is a transmissible infectious disease: the Keyes and Fitzgerald revolution. *J Dent Res.* 1995;74(9):1536–1542. [PubMed: 7560413]
2. Bowen WH. Dental caries - not just holes in teeth! A perspective. *Mol Oral Microbiol.* 2016;31(3):228–233. [PubMed: 26343264]
3. Loesche ea WJ. Association of Streptococcus mutants with human dental decay. *Infect Immun.* 1975;1252–1260.
4. Kanasi E, Johansson I, Lu SC, et al. Microbial risk markers for childhood caries in pediatricians' offices. *J Dent Res.* 2010;89(4):378–383. [PubMed: 20164496]
5. Caufield PW, Cutter GR, Dasanayake AP. Initial acquisition of mutans streptococci by infants: evidence for a discrete window of infectivity. *Journal of dental research.* 1993;72(1):37–45. [PubMed: 8418105]
6. Li Y, Caufield PW, Dasanayake AP, Wiener HW, Vermund SH. Mode of delivery and other maternal factors influence the acquisition of Streptococcus mutans in infants. *Journal of dental research.* 2005;84(9):806–811. [PubMed: 16109988]
7. Zhan L, Tan S, Den Besten P, Featherstone JD, Hoover CI. Factors related to maternal transmission of mutans streptococci in high-risk children-pilot study. *Pediatric dentistry.* 2012;34(4):e86–91. [PubMed: 23014079]
8. Slayton RL. Reducing mutans streptococci levels in caregivers may reduce transmission to their children and lead to reduced caries prevalence. *J Evid Based Dent Pract.* 2011;11(1):27–28. [PubMed: 21420005]
9. Klein MI, Florio FM, Pereira AC, Hofling JF, Goncalves RB. Longitudinal study of transmission, diversity, and stability of Streptococcus mutans and Streptococcus sobrinus genotypes in Brazilian nursery children. *Journal of clinical microbiology.* 2004;42(10):4620–4626. [PubMed: 15472319]
10. Klinker T, Urban M, Luck C, Hannig C, Kuhn M, Kramer N. Changes in Candida spp., mutans streptococci and lactobacilli following treatment of early childhood caries: a 1-year follow-up. *Caries Res.* 2014;48(1):24–31. [PubMed: 24216710]
11. van Houte J Role of micro-organisms in caries etiology. *J Dent Res.* 1994;73(3):672–681. [PubMed: 8163737]
12. Banas JA. Virulence properties of Streptococcus mutans. *Front Biosci.* 2004;9:1267–1277. [PubMed: 14977543]
13. Bowen WH, Koo H. Biology of Streptococcus mutans-derived glucosyltransferases: role in extracellular matrix formation of cariogenic biofilms. *Caries Res.* 2011;45(1):69–86.
14. Kowash MB, Curzon ME, Hart P. Association of salivary Streptococcus mutans with caries in young children: effect of dental health education on salivary levels. *European journal of paediatric dentistry : official journal of European Academy of Paediatric Dentistry.* 2002;3(4):199–204.
15. Beighton D, Russell RRB, Whaley RA. A simple biochemical scheme for the differentiation of streptococcus-mutans and streptococcus-sobrinus. *Caries Res.* 1991;25(3):174–178. [PubMed: 1652359]
16. Cangelosi GA, Iversen JM, Zuo Y, Oswald TK, Lamont RJ. Oligonucleotide probes for mutans streptococci. *Mol Cell Probes.* 1994;8(1):73–80. [PubMed: 8028611]
17. Igarashi T, Yamamoto A, Goto N. Direct detection of Streptococcus mutans in human dental plaque by polymerase chain reaction. *Oral Microbiol Immunol.* 1996;11(5):294–298. [PubMed: 9028253]
18. Kawamura Y, Hou XG, Sultana F, Miura H, Ezaki T. Determination of 16S ribosomal-RNS sequences of streptococcus-mitis and streptococcus-gordonii and Phylogenetic-relationships among members of the genus streptococcus. *Int J Syst Bacteriol.* 1995;45(2):406–408. [PubMed: 7537076]
19. Fan CC, Wang WH, Xu T, Zheng SG. Risk factors of early childhood caries (ECC) among children in Beijing - a prospective cohort study. *BMC Oral Health.* 2019;19:1–9. [PubMed: 30611272]
20. Seki M, Karakama F, Terajima T, et al. Evaluation of mutans streptococci in plaque and saliva: correlation with caries development in preschool children. *J Dent.* 2003;31(4):283–290. [PubMed: 12735923]

21. Parisotto TM, Steiner-Oliveira C, Silva C, Rodrigues LKA, Nobre-dos-Santos M. Early Childhood Caries and Mutans Streptococci: A Systematic Review. *Oral Health Prev Dent*. 2010;8(1):59–70. [PubMed: 20480056]
22. Nase L, Hatakka K, Savilahti E, et al. Effect of long-term consumption of a probiotic bacterium, *Lactobacillus rhamnosus* GG, in milk on dental caries and caries risk in children. *Caries Res*. 2001;35(6):412–420. [PubMed: 11799281]
23. Zhou Y, Lin HC, Lo ECM, Wong MCM. Risk indicators for early childhood caries in 2-year-old children in southern China. *Aust Dent J*. 2011;56(1):33–39. [PubMed: 21332738]
24. Campus G, Cocco F, Carta G, et al. Effect of a daily dose of *Lactobacillus brevis* CD2 lozenges in high caries risk schoolchildren. *Clin Oral Investig*. 2014;18(2):555–561.
25. Fontana M, Catt D, Eckert GJ, et al. Xylitol: Effects on the Acquisition of Cariogenic Species in Infants. *Pediatr Dent*. 2009;31(3):257–266. [PubMed: 19552232]
26. ElSalhy M, Soderling E, Honkala E, et al. Salivary microbiota and caries occurrence in Mutans Streptococci-positive school children. *Eur J Paediatr Dent*. 2016;17(3):188–192. [PubMed: 27759406]
27. Xiao J, Moon Y, Li LH, et al. *Candida albicans* Carriage in Children with Severe Early Childhood Caries (S-ECC) and Maternal Relatedness. *PLoS One*. 2016;11(10):1–16.
28. Ribeiro TR, Alves KSD, Mota ACD, et al. Caries experience, mutans streptococci and total protein concentrations in children with protein-energy undernutrition. *Aust Dent J*. 2014;59(1):106–113. [PubMed: 24494693]
29. Fonteles CSR, Guerra MH, Ribeiro TR, et al. Association of free amino acids with caries experience and mutans streptococci levels in whole saliva of children with early childhood caries. *Arch Oral Biol*. 2009;54(1):80–85. [PubMed: 18774124]
30. Tamami Saito O U, Sachiyo Teramoto, Hisao Oguchi, Shigeru Yanagisawa and Hirro Miyazawa. Bacteriological evaluation of mutans streptococci using modified mitis-salivarius-bacitracin (MSB) agar medium in primary dentition period. *Pediatric Dental Journal*. 2007;17(1):53–57.
31. Gold OG, Jordan HV, Vanhoute J. SELECTIVE MEDIUM FOR STREPTOCOCCUS-MUTANS. *Arch Oral Biol*. 1973;18(11):1357–1364. [PubMed: 4518755]
32. Pinheiro SL, Segatti B, Pucca DS, Dutra PT. Dental acid etchant as a sensitizing agent in photodynamic therapy to reduce S-mutans in dentinal carious lesions. *Lasers Med Sci*. 2019;34(2):305–309. [PubMed: 30027423]
33. Kim J, Lee HJ, Hong SH. Inhibition of streptococcal biofilm by hydrogen water. *J Dent*. 2017;58:34–39. [PubMed: 28087297]
34. Relvas M, Coelho C, Henriques CV, Ramos E. Cariogenic bacteria and dental health status in adolescents: the role of oral health behaviours. *Eur J Paediatr Dent*. 2014;15(3):281–287. [PubMed: 25306145]
35. Hildebrandt GH, Bretz WA. Comparison of culture media and chairside assays for enumerating mutans streptococci. *J Appl Microbiol*. 2006;100(6): 1339–1347. [PubMed: 16696682]
36. Saravia ME, Nelson-Filho P, Silva RA, et al. Recovery of mutans streptococci on MSB, SB-20 and SB-20M agar media. *Archives of oral biology*. 2013;58(3):311–316. [PubMed: 23149153]
37. Wan AK, Seow WK, Walsh LJ, Bird PS. Comparison of five selective media for the growth and enumeration of *Streptococcus* mutans. *Australian dental journal*. 2002;47(1):21–26. [PubMed: 12035953]
38. Yoo SY, Kim PS, Hwang HK, et al. Identification of non-mutans streptococci organisms in dental plaques recovering on mitis-salivarius bacitracin agar medium. *J Microbiol*. 2005;43(2):204–208. [PubMed: 15880098]
39. Kim YH, Lee SY. Identification of non-streptococcal organisms from human dental plaque grown on the *Streptococcus*-selective medium mitis-salivarius agar. *Arch Oral Biol*. 2015;60(2):267–271. [PubMed: 25463904]
40. Little WA, Korts, D. C, Thomson, L. A, & Bowen, W. H Comparative recovery of *Streptococcus* mutans on ten isolation media. *J Clin Microbiol*. 1977;5(6):578–583. [PubMed: 885999]
41. Chen Z, Saxena D, Caufield PW, Ge Y, Wang MQ, Li YH. Development of species-specific primers for detection of *Streptococcus* mutans in mixed bacterial samples. *FEMS Microbiol Lett*. 2007;272(2):154–162. [PubMed: 17521362]

42. Paczosa MK, Mecsas J. *Klebsiella pneumoniae*: Going on the Offense with a Strong Defense. *Microbiol Mol Biol Rev.* 2016;80(3):629–661. [PubMed: 27307579]
43. Huycke. NIAHaMM. Enterococcal Disease, Epidemiology, and Implications for Treatment. In: *Enterococci: From Commensals to Leading Causes of Drug Resistant Infection.* 2014.
44. Law V, Seow WK, Townsend G. Factors influencing oral colonization of mutans streptococci in young children. *Aust Dent J.* 2007;52(2):93–100. [PubMed: 17687953]
45. Kishi M, Abe A, Kishi K, Ohara-Nemoto Y, Kimura S, Yonemitsu M. Relationship of quantitative salivary levels of *Streptococcus mutans* and *S-sobrinus* in mothers to caries status and colonization of mutans streptococci in plaque in their 2.5-year-old children. *Community Dentist Oral Epidemiol.* 2009;37(3):241–249.
46. Barfod MN, Magnusson K, Lexner MO, Blomqvist S, Dahlen G, Twetman S. Oral microflora in infants delivered vaginally and by caesarean section. *Int J Paediatr Dent.* 2011;21(6):401–406. [PubMed: 21702851]
47. Durhan MA, Topcuoglu N, Kulekci G, Ozgentas E, Tanboga I. Microbial Profile and Dental Caries in Cleft Lip and Palate Babies Between 0 and 3 Years Old. *Cleft Palate-Craniofac J.* 2019;56(3):349–356. [PubMed: 29800530]
48. Reed SG, Cunningham JE, Latham TN, Shirer SC, Wagner CL. Maternal Oral Mutans Streptococci (MS) Status, Not Breastfeeding, Predicts Preterm Infant Oral MS Status. *Breastfeed Med.* 2014;9(9):446–449. [PubMed: 25290343]
49. Lamont RJ, Hajishengallis GN, Jenkinson HF. *Oral Microbiology and Immunology.* 2013.
50. Li YH, Saraithong P, Chen Z, Leung E, Pattanaporn K, Dasanayake A. Comparison of Real-Time Quantitative PCR with a Chairside Test for *Streptococcus Mutans* Assessment. *Chin J Dent Res.* 2017;20(4):199–210. [PubMed: 29181457]
51. Simon-Soro A, Mira A. Solving the etiology of dental caries. *Trends Microbiol.* 2015;23(2):76–82. [PubMed: 25435135]
52. Fanaro S, Chierici R, Guerrini P, Vigi V. Intestinal microflora in early infancy: composition and development. *Acta Paediatr.* 2003;92:48–55.
53. Otto M *Staphylococcus epidermidis* - the ‘accidental’ pathogen. *Nat Rev Microbiol.* 2009;7(8):555–567. [PubMed: 19609257]
54. Sanders WE, Sanders CC. *Enterobacter* spp.: Pathogens poised to flourish at the turn of the century. *Clin Microbiol Rev.* 1997;10(2):220–241. [PubMed: 9105752]
55. Aly R SH, Strauss WG. Bacterial adherence to nasal mucosal cells. *Infect Immun.* 1977;17:546–549. [PubMed: 409676]
56. Keyworth N, Millar MR, Holland KT. Development of cutaneous microflora in premature neonates. *Arch Dis Child.* 1992;67(7):797–801. [PubMed: 1519978]
57. Kloos WE M M. Distribution and persistence of *Staphylococcus* and *Micrococcus* species and other aerobic bacteria on human skin. *Appl Microbiol.* 1975;30:381–385. [PubMed: 810086]
58. Refsahl K, Andersen BM. Clinically significant coagulase-negative staphylococci-identification and resistance patterns. *J Hosp Infect.* 1992;22(1):19–31. [PubMed: 1358944]
59. Kosako Y, Tamura K, Sakazaki R, Miki K. *Enterobacter kobei* sp nov, a new species of the family Enterobacteriaceae resembling *Enterobacter cloacae*. *Curr Microbiol.* 1996;33(4):261–265. [PubMed: 8824173]
60. Davin-Regli A, Pages JM. *Enterobacter aerogenes* and *Enterobacter cloacae*; versatile bacterial pathogens confronting antibiotic treatment. *Front Microbiol.* 2015;6:1–10. [PubMed: 25653648]
61. Sorlozano A, Gutierrez J, Martinez T, et al. Detection of new mutations conferring resistance to linezolid in glycopeptide-intermediate susceptibility *Staphylococcus hominis* subspecies *hominis* circulating in an intensive care unit. *Eur J Clin Microbiol Infect Dis.* 2010;29(1):73–80. [PubMed: 19876662]
62. Spanu T, Sanguinetti M, Ciccaglione D, et al. Use of the VITEK 2 system for rapid identification of clinical isolates of staphylococci from bloodstream infections. *J Clin Microbiol.* 2003;41(9):4259–4263. [PubMed: 12958254]
63. Molander A, Reit C, Dahlen G, Kvist T. Microbiological status of root-filled teeth with apical periodontitis. *Int Endod J.* 1998;31(1):1–7. [PubMed: 9823122]

64. Brisse S, Passet V, Grimont PAD. Description of *Klebsiella quasipneumoniae* sp nov., isolated from human infections, with two subspecies, *Klebsiella quasipneumoniae* subsp *quasipneumoniae* subsp nov and *Klebsiella quasipneumoniae* subsp *similipneumoniae* subsp nov., and demonstration that *Klebsiella singaporensis* is a junior heterotypic synonym of *Klebsiella variicola*. *Int J Syst Evol Microbiol*. 2014;64:3146–3152. [PubMed: 24958762]
65. Parkins MD, Sibley CD, Surette MG, Rabin HR. The *Streptococcus milleri* group - An unrecognized cause of disease in cystic fibrosis: A case series and literature review. *Pediatr Pulmonol*. 2008;43(5):490–497. [PubMed: 18383109]
66. Takahashi N, Nyvad B. The Role of Bacteria in the Caries Process: Ecological Perspectives. *J Dent Res*. 2011;90(3):294–303. [PubMed: 20924061]
67. Balakrishnan M, Simmonds RS, Tagg JR. Dental caries is a preventable infectious disease. *Aust Dent J*. 2000;45(4):235–245. [PubMed: 11225524]

Figure 1. Colony morphology of bacteria grown on MSB agar

*Nine bacteria strains (A-1 to A-9 and B-1 to B-9) from predente infants' saliva sample grew on MSB agar. A-10 and B-10 were *Streptococcus sobrinus* (ATCC® 33478™) wild strain.

Colony morphology and description of bacteria from predentine infant oral cavity grown on MSB agar

Bacteria	<i>Staphylococcus epidermidis</i>	<i>Enterobacter kobei</i>	<i>Staphylococcus hominis</i>	<i>Enterococcus faecalis</i>	<i>Klebsiella quasi-pneumoniae</i>	<i>Klebsiella pneumoniae</i>	<i>Streptococcus anginosus</i>	<i>Phytobacter sp.</i>	<i>Streptococcus mutans</i>	<i>Streptococcus sobrinus</i>
Colony size (diameter)	2~4 mm	2~4 mm	2~3 mm	1~1.5 mm	~1 mm	~1 mm	~5 mm	~5 mm	0.5~1 mm	0.5~1 mm
Colony morphology	Dark black, perfect round, regular margin, flat, shiny	Bright blue, iris, bull-eye, mucoid, easy to form connected colonies	Black blue, round, flat, matt-finish, wrinkles on the colony edge	Dark black or gray, round, small	Dark black, shiny, round, tall, easy to form chains	Light blue or gray, small dome	Dark black, small dome, irregular round, textured surface	Dark black, shiny, irregular margin	Frosted glass, irregular margin, black blue, rooted into the agar, starshaped	Frosted glass, irregular margin, black blue, rooted into the agar, surrounded by polysaccharide
Colony smell	Slightly fermented	Fermented, sour	No distinct smell	Foul smell	Slightly fermented	Fermented, slightly sweet	No distinct smell	Slightly sour	No distinct smell	No distinct smell
<i>S. mutans</i> Colony PCR	-	-	-	-	-	-	-	-	+	-
Bacteria characteristics	<ul style="list-style-type: none">• Gram-positive• Facultative anaerobic• One of the most commonly etiological agent of nosocomial infections ⁵³	<ul style="list-style-type: none">• Gram-negative• Oxidase-negative• Fermentative rods• Facultative anaerobic• <i>Enterobacter cloacae</i> complex (<i>E. Koberi</i> included) is one of the important nosocomial pathogens, responsible for 65% - 75% of all infections ⁵⁴	<ul style="list-style-type: none">• Gram-positive• Limited facultative anaerobic• Potentially opportunistic pathogens, could cause bloodstream infections, endocarditis, peritonitis, osteomyelitis, bone and joint infections ⁶¹• Commonly recoverable from the blood of hospitalized patients ⁶²	<ul style="list-style-type: none">• Gram-positive• Facultative anaerobic• One of the most commonly etiological agent of nosocomial infection• Could cause endocarditis, bacteremia, urinary tract infections, meningitis, and other fatal forms of systemic and local infection ⁴³• Frequently found in re-infected, root canal-treated teeth ⁶³	<ul style="list-style-type: none">• Gram-negative• Oxidase-negative• Rod-shaped bacteria with a prominent polysaccharidebased capsule Similar to <i>K. Pneumonia</i> ⁶⁴	<ul style="list-style-type: none">• Gram-negative• Facultative anaerobic• Rod-shaped• Commonly found in the normal flora of the mouth, skin and intestines• Could cause a wide range of infections, including pneumonias, urinary tract infections, bacteremias, and liver abscesses ⁴²	<ul style="list-style-type: none">• Gram-positive• Facultative anaerobic• Commonly found in the oral cavity• Carigiogenic bacteria ^{21,44,51,66}	<ul style="list-style-type: none">• Gram-negative• Facultative anaerobic• Commonly found in the gastrointestinal and urogenital tract• Potential pathogenic role in cystic fibrosis, ⁶⁵	<ul style="list-style-type: none">• Gram-positive• Facultative anaerobic• Commonly found in the oral cavity• Carigiogenic bacteria ^{21,44,51,66}	<ul style="list-style-type: none">• Gram-positive• Anaerobic• Closely related species of <i>S. mutans</i> ^{44,67}• Carigiogenic bacteria ^{44,67}

Table 2

Detection frequency and quantity of bacteria strains isolated by MSB agar from predentate infants (n=51)

Bacteria	<i>Staphylococcus epidermidis</i>	<i>Enterobacter kobei</i>	<i>Staphylococcus hominis</i>	<i>Enterococcus faecalis</i>	<i>Klebsiella quasi-pneumoniae</i>	<i>Klebsiella pneumoniae</i>	<i>Streptococcus anginosus</i>	<i>Phytobacter sp.</i>	<i>Streptococcus mutans</i>	<i>Streptococcus sobrinus</i>
Detection frequency	41.5%	24.4%	2.4%	12.2%	2.4%	17.1%	2.4%	2.4%	9.8%	0
Detection quantity (CFU/ml, Mean ± SD)	6.7 ± 12.1 x10 ²	1.7 ± 4.4 x10 ⁷	1.5 x10 ¹	6.7 ± 8.8 x10 ³	9.7 x10 ¹	3.9 ± 5.5 x10 ²	4.5 x10 ¹	1.1 x10 ⁵	1.7 ± 3.4 x10 ⁵	0