Published in final edited form as:

J Chem Theory Comput. 2019 April 09; 15(4): 2692–2705. doi:10.1021/acs.jctc.8b01312.

Thermodynamics of cation binding to the sarco/endoplasmic reticulum calcium ATPase pump and impacts on enzyme function

Bin Sun, Bradley D. Stewart, Amir N. Kucharski, Peter M. Kekeneshuskey*

Department of Chemistry, University of Kentucky, 505 Rose St., Chemistry-Physics Building, Lexington, KY, USA 40506

Abstract

Sarcoendoplasmic reticulum Ca^{2+} -ATPase (SERCA) is a transmembrane pump that plays an important role in transporting calcium into the sarcoplasmic reticulum (SR). While calcium (Ca^{2+}) binds SERCA with micromolar affinity, magnesium (Mg^{2+}) and potassium (K^+) also compete with calcium (Ca^{2+}) binding. However, the molecular bases for these competing ions' influence on SERCA function and the selectivity of the pump for Ca^{2+} are not well-established. We therefore used *in silico* methods to resolve molecular determinants of cation binding in the canonical site I and II Ca^{2+} binding sites: 1) triplicate molecular dynamics (MD) simulations of Mg^{2+} , Ca^{2+} and K^+ -bound SERCA. 2) mean spherical approximation (MSA) theory to determine the affinity and selectivity of cation binding to the MD-resolved structures and 3) state models of SERCA turnover informed from MSA-derived affinity data. Our key findings are that a) coordination at sites I and II are optimized for Ca^{2+} and to a lesser extent for Mg^{2+} and K^+ , as determined by MD-derived cation-amino acid oxygen and bound water configurations, b) the impaired coordination and high desolvation cost for Mg^{2+} precludes favorable Mg^{2+} binding relative to Ca^{2+} , while K^+ has limited capacity to bind site I, c) Mg^{2+} most likely acts as inhibitor and K^+ as intermediate in SERCA's reaction cycle, based on a best-fit state model of SERCA turnover. These findings provide a quantitative basis for SERCA function that leverages molecular-scale thermodynamic data and rationalize enzyme activity across broad ranges of K^+ , Ca^{2+} and Mg^{2+} concentrations.

Graphical Abstract

*Corresponding authors: pkekeneshuskey@uky.edu (P. Kekeneshuskey).

⁷Supporting Information

Additional information regarding details of method and supportive analysis are provided. This information is available free of charge via the Internet at <http://pubs.acs.org>

1 INTRODUCTION

Sarcoendoplasmic reticulum Ca²⁺-ATPase (SERCA) is a 110-kDa transmembrane cation pump which actively transports Ca²⁺ ions into the SR by utilizing energy released from adenosine triphosphate (ATP) hydrolysis.¹ SERCA has been widely studied for its role in returning intracellular Ca²⁺ to basal levels following stimuli that elevates Ca²⁺ content.² The pump's catalytic cycle is roughly characterized by four states comprising a sequential cycle $E1 \rightarrow E1P.2Ca \rightarrow E2P.2Ca \rightarrow E2$. In E1, the Ca²⁺ binding sites are exposed to the cytosolic space, whereas the E2 conformations expose the low-affinity Ca²⁺ sites toward the SR lumen. The transition between E1 and E2 is driven by ATP hydrolysis at residue Asp 351¹ following Ca²⁺ binding, for which E1P and E2P are the respective phosphorylated states of the enzyme. Accompanying transitions between catalytic states are prominent changes in its ten transmembrane (TM) helices as well as the cytosolic actuator (A) domain, nucleotide-binding domain (N) and phosphorylation domain (P). Many of these changes have been resolved through x-ray crystallography.^{3–6}

Based on available structural models of the protein and a wealth of biochemical studies,^{7–14} reaction schemes linking the E1 and E2 states are beginning to emerge. Utilizing SERCA vesicles coupled with spin label molecules, Inesi et al observed changed electron spin resonance spectrum upon Ca²⁺ binding that revealed conformational changes in the enzyme.¹¹ Dupont and coworkers similarly measured changes in intrinsic fluorescence upon Ca²⁺ binding and further proposed a two-step Ca²⁺ binding process to high affinity sites evidenced by the pumps slow rate of fluorescence changes.^{9,10} Additionally, conformational changes linking the E1 and E2 states were explored by kinetic studies of intrinsic fluorescence changes upon Ca²⁺ binding and release.^{7,8} To probe molecular determinants of Ca²⁺ binding in the pump, an E309Q mutant bound with two Ca²⁺ in its phosphorylated state was determined via x-ray crystallography,¹⁵ which revealed that the altered TM arrangements caused by the mutation leads to impaired pump functionality. With respect to cation binding affinities, Inesi et al measured Ca²⁺ binding and stoichiometry to SERCA vesicles in via chromatography,¹⁶ while others have probed the binding of the non-cognate Mg²⁺ and K⁺ ions via intrinsic fluorescence changes in SERCA.^{17,18}

In complement to experiment, extensive molecular dynamics simulations have uncovered many aspects of cation binding and SERCA function that are difficult to probe via

experimental approaches. Huang et al first explored the Ca^{2+} binding pathways to SERCA TM sites via MD simulation and observed the cooperative binding of two Ca^{2+} ions.¹⁹ Kekenos-Huskey et al performed MD simulations on SERCA with and in the absence of Ca^{2+} to examine gating of Ca^{2+} binding by E309,²⁰ in addition to estimating ion binding free energies and kinetics. Espinoza-Fonseca et al evaluated the interaction energy between $\text{Ca}^{2+}/\text{Mg}^{2+}/\text{K}^{+}$ and SERCA binding sites based on MD trajectories and reported that Ca^{2+} has the most negative interaction energy while K^{+} has the least negative value.²¹ In addition to these initial studies on cation binding to the SERCA pump, more recent studies have probed mechanisms of SERCA function and its modulation by regulatory proteins and drugs.^{22–29}

Although studies have been reported on the roles of Mg^{2+} , K^{+} on SERCA binding, less has been done to provide a thermodynamic basis for their effects on SERCA function. This is of particular importance, as intracellular K^{+} and Mg^{2+} concentrations are orders of magnitude larger than the roughly sub micromolar Ca^{2+} concentrations found in typical cells. Here, simulation studies of cation binding to molecular sites that resemble high affinity, amino acid-based motifs have been informative. Dudev et al for instance constructed cation binding sites using model compounds and calculated cation binding energies via quantum mechanics.³⁰ Ye et al designed a theoretical framework in combination with MD simulations to calculate cation binding free energies and isolate the energetic contribution from geometric and confinement effect.³¹ Implicit models that describe Ca^{2+} binding thermodynamics as via ion density-based formalisms have also been reported, including reference interaction site modeling³² and density functional theory.³³ Besides these methods relied on explicit binding site configurations, Nonner et al developed the MSA model in which the binding site is treated as confined filter filled with coordination oxygens. A variant of density functional theory called the mean spherical approximation (MSA) approach has proven effective in rank-ordering the binding of cationic species to oxygen-rich binding domains such as EF-hands in β parvalbumin (β -PV)³⁴ and Ca^{2+} channel selectivity filters,^{35,36} through describing the electrostatics and hard-sphere contributions to the chemical potential of partitioning ions into oxygen-rich ‘filters’.

Our study has therefore focused on utilizing molecular dynamics (MD) derived data with thermodynamic and state-based models to assess contributions of Mg^{2+} and K^{+} binding on the SERCA turnover rate. Here we performed MD simulations of Ca^{2+} , Mg^{2+} and K^{+} -bound wild-type (WT) SERCA as well as the E309Q and N796A variants. These MD data provided structural information to assess cation binding thermodynamics via MSA to elucidate the molecular basis of SERCA’s preference of Ca^{2+} over Mg^{2+} and K^{+} . Further, we relate these studies of E1 state ion binding to a state-based kinetic model of SERCA pump rate to determine the extents to which Mg^{2+} and K^{+} facilitate or inhibit catalysis. With this approach, we provide a multiscale and molecular basis for cation binding to SERCA and impacts on pump function.

2 RESULTS

We performed triplicate simulations of wild-type SERCA and its variants E309Q and N768A to probe the binding site coordination of the cations Ca^{2+} , Mg^{2+} and K^{+} , each

replica was at least 100 ns in length. Ca^{2+} - and Mg^{2+} -bound configurations of the protein have been determined through x-ray crystallography, but to our knowledge, the binding of K^+ to the Ca^{2+} binding domain has only been resolved via simulation.²¹ In this study, we critically examine the protein atoms and waters that directly coordinate bound cations. However, to ensure that our simulations of the intact protein are consistent with prior studies of the enzyme, we briefly summarize standard analyses of the transmembrane bundles and cytosolic domains in the Supplement (see Sect. S1.4).

2.1 Cation coordination in the Ca^{2+} binding region

In this study, we highlight structural and dynamic contributions of the Ca^{2+} and Mg^{2+} binding domains to the thermodynamics of ion binding. This is analogous to our approach for probing $\text{Ca}^{2+}/\text{Mg}^{2+}$ -binding to the β parvalbumin (β -PV) protein,³⁴ for which we used MD simulation-derived structural data from the cation-bound configurations to parameterize a statistical mechanical model of fluid thermodynamics called mean spherical approximation (MSA). Specifically, we used the radius of the ions' inner coordination sphere and amino acid oxygens comprising the sphere to estimate the binding site volume and coordinating oxygen density for MSA. We use a similar strategy for SERCA in that we assess the coordination of a given ion based on the number of oxygens within six Å of the bound ion. In contrast to our previous study, we additionally include coordinated waters in the MSA calculation that are directly involved in stabilizing the ion.

In Fig. 2 we report radial distribution function (RDF)s of cation-coordinating oxygens from the simulations. These RDFs demonstrate that Ca^{2+} , Mg^{2+} , and K^+ have varying degrees of coordination with amino acid oxygens within the binding site. We summarize the identity of the coordinating amino acids in Sect. S1.4, although from the perspective of MSA theory, only the number of contributed amino acid oxygens is important. The Ca^{2+} within site I was optimally coordinated with 8.5 oxygens on average (including three water oxygens). Ca^{2+} within site II maintained six amino acid oxygens pairings, similar to the x-ray crystal structure, but did not directly coordinate waters. By virtue of having more coordinating oxygens, we speculate that the Ca^{2+} ion in site I is bound more tightly than that found within site II. In contrast, for the N796A mutant, the site I Ca^{2+} had a reduced coordination number of six relative to over eight in the WT structure. Interestingly, for the site II Ca^{2+} in the N796A variant, the loss of coordination to site 796 was compensated by interactions formed with E58 and a bound water molecule (see Fig. S11(b)) to yield a greater coordination number than observed in the WT. For the E309Q mutant, two possible side chain rotamers of Q309 were investigated, as both rotamers were viable starting positions (see Fig. S11(c–d) for illustration of rotamer directions). We found that these rotamers yielded identical Ca^{2+} coordination patterns for the two binding sites: site I had six coordinating oxygens versus seven for site II, while neither included bound waters (see Fig. S11(c–d)). Similar to the N796A site II Ca^{2+} case, E58 in TM1 also participated into the coordination with Ca^{2+} at site II in both rotamers of E309Q. Overall, these simulations reveal that ion/oxygen pairing is remarkably labile between the sites I and II, and can incorporate bound waters to maximize Ca^{2+} coordination.

In contrast to Ca^{2+} , Mg^{2+} binds in a ‘hybrid’ site between sites I and II with an average six coordinating oxygens (including one water molecule, see Fig. S13(a)). K^{+} , on the other hand, binds site I with five coordinating oxygens (including one water molecule, see Fig. S13(c)), while K^{+} at site II is highly dynamic and interchanges with water immobilized in the SERCA interior. Given that these ions are both positively charged and not remarkably different in size relative to Ca^{2+} , we had anticipated that the non-cognate ions might adopt higher coordination numbers in the native Ca^{2+} sites. Clearly the molecular simulations did not reflect this expectation; in Sect. 2.2 we provide a thermodynamic rationale via MSA theory for why these non-cognate ions present impaired coordination numbers.

The differences in cation-oxygen coordination patterns for the cases considered are accompanied by variations in the coordinating residues’ mobilities relative to WT. These mobilities are measured as RMSF values in Fig. 3, for which the upper row represents site I and the lower row, site II. Generally, Ca^{2+} -bound systems have RMSF values for most residues around 0.5 Å, which are the smallest among the ions considered. To a certain extent, the reduced mobility could be interpreted as an indication of tighter and more favorable binding, although this would come at an entropic cost that is not explicitly estimated here. We were, however, surprised to see little change in RMSF for the Ca^{2+} -free (apo) state versus the Ca^{2+} -bound cases. One possible reason is that in apo state, waters fill the binding sites and stabilize residues via a hydrogen bonding network (see Fig. S7) - in this capacity, bound waters might ‘prop’ open the Ca^{2+} -binding domains to promote rapid incorporation of solvated Ca^{2+} ions from the bulk medium. Additionally, we found that WT SERCA and its variants presented negligible differences in RMSF upon Ca^{2+} binding, whereas the non-cognate Mg^{2+} and K^{+} manifest significant RMSF increases across all residues (Mg^{2+} generally above 0.8 Å and K^{+} above 1 Å). It is possible that the greater mobility of coordinating residues for the non-cognate ions are indicative of impaired coordination. We had anticipated that waters could be incorporated into the ions’ binding domain to suppress fluctuations in amino acids contributing to coordination, much as was observed for the apo state. However, it is apparent the the strong electrostatic affinity for these ions with the coordination residues limited the volume within which waters could be incorporated. At a minimum, these data suggest that ion coordination is dynamic, with fluctuations on a nanosecond timescale (see Fig. S6), which ultimately may play a role in selecting Ca^{2+} over non-cognate ions.

2.2 Thermodynamics of ion binding at sites I and II

Molecular dynamics simulations provide qualitative insight to the binding of various ions in the binding sites of SERCA, but alone do not directly predict affinities. Therefore we explored MSA to semi-quantitatively estimate free energies and selectivity of ion binding. MSA predicts chemical potentials of partitioning solvated ions into the SERCA binding domains, assuming a similar approach that was performed for the $\beta\text{-PV}^{34}$ protein. Namely, MSA theory estimates the chemical potential of ion binding, based on the assumption of finite sized ions and chelating oxygens confined to a spherical binding site volume. To utilize this method, we first compute oxygen RDFs about bound ions. These data provide oxygen filter densities and volumes, from which chemical potentials of ion partitioning into the binding site ‘filter’ from the surrounding bulk solution can be estimated. As shown in

Fig. 4, Ca^{2+} at site I of WT case presents the most negative and therefore thermodynamically-favorable MSA- predicted chemical potentials across all cases, corresponding to the largest number of coordination oxygens. For the SERCA variants, site I and site II Ca^{2+} ions have modestly less favourable chemical potentials compared with site I Ca^{2+} from WT cases, as the relative values were approximately 0.3 kcal/mol higher. These are consistent with the comparable Ca^{2+} coordination numbers present among variants. Compared with Ca^{2+} -binding, Mg^{2+} binding at WT and mutant SERCA yielded more significantly disfavored chemical potentials at approximately 2 kcal/mol relative to WT site I Ca^{2+} . Among all cases, the K^+ relative potential was the most positive at 4.2 kcal/mol, indicating that K^+ is the least thermodynamically favored at TM sites. For Mg^{2+} and K^+ , the MSA predicted potentials also correlated with the cation-coordination patterns as these two ions had reduced coordination number when compared with Ca^{2+} (5.2/4.7 for $\text{Mg}^{2+}/\text{K}^+$ versus 8.5 for site I Ca^{2+}). These data show that MSA could capture the key factors governing cation affinities such as coordination number and binding site volume. In addition, with inclusion of coordinating waters in MSA, the predicted potential for WT site I Ca^{2+} is most favourable among all cases, which agrees with experimentally measured affinities. Our MSA results indicate that site I confers greater Ca^{2+} affinity due to extensive inclusion of water coordination. In Sect. S1.4, we utilized Grid Inhomogeneous Solvation Method (GIST) to assess the relative thermodynamics of water binding to the Ca^{2+} binding domain. In general, we found that when bound waters are present in the ions' coordination shells, the predicted free energies are on the order of -12 kcal/mol and thus very thermodynamically favorable.

The membrane into which SERCA is embedded presents a negative surface charge owing to solvent-exposed phosphate head groups.³⁷ The corresponding negative electrical potential can attract positively charged ions and thereby increase their concentration near the transmembrane-bound SERCA, as we previously observed in.²⁰ Since the partitioning of cations into the SERCA binding sites depends on the composition of the surrounding electrolyte, we speculated that the local elevation of cations near the membrane surface would subsequently increase the concentration of bound cations. Additionally, the higher ionic strength could also reduce the desolvation energy and thereby further favor cation binding to SERCA, though this effect would likely be offset by screening electrostatic interactions between cations and the oxygens in the SERCA Ca^{2+} binding domains. To investigate this hypothesis, we determined the effective ion concentration near the membrane using a 1D solution of the linearized Poisson-Boltzmann equation,³⁸ $[i]_{eff} = e^{-\beta Z_i \zeta} [i]_{bath}$ where Z_i is the charge of ion and $\beta = \frac{1}{k_B T}$, $[i]_{bath}$ is ion concentration in the bath and ζ is the membrane potential.

Assuming $\zeta = 25$ mV,³⁹ we predicted that monovalent cation concentrations would be increased by 2.7 fold, anions would decrease by 0.37 fold, and the divalent $\text{Ca}^{2+}/\text{Mg}^{2+}$ ions would increase by 7.4 fold. As shown in Fig. 5, at low bath $[\text{Ca}]$, both Mg^{2+} and K^+ were present in the SERCA Ca^{2+} binding domain with concentrations of 7.7 M and 1.7 M, respectively. As bath $[\text{Ca}]$ was increased, Ca^{2+} partitioned into the binding domain in favor of Mg^{2+} ; at roughly 2×10^{-5} M, the ratio of Mg^{2+} to Ca^{2+} was 1:1. The Ca^{2+} concentration

at which $\text{Mg}^{2+}/\text{Ca}^{2+}$ was 1:1 varied by roughly 0.6 fold under the assumption of charged versus neutral membrane. In other words, according to our model, the local electrostatic environment about the membrane did not significantly impact the Ca^{2+} binding affinity. In contrast, in²⁰ we demonstrated that the negative surface charge densities of SERCA and the lipid enhanced the association rate of Ca^{2+} to the protein, thus the local electrostatic environment may have a greater contribution to ion binding kinetics than steady-state binding.

2.3 Steady-state catalytic activity

Lastly, we relate our predictions of Ca^{2+} and non-cognate ion binding to the SERCA pumping rate. For the complete pumping cycle, two Ca^{2+} ions in the cytoplasm are transported into the SR by first binding SERCA to its E1 state. This binding process was proposed by Inesi et al⁴⁰ to consist of two Ca^{2+} successive binding events via a cooperative mechanism. Subsequent steps include binding of MgATP, a slow conformational transition to the E2 state, release of Ca^{2+} ions into the SR lumen, and a return to the E1 apo state. In practice, by accounting for the transition rates between SERCA conformational states, the time-evolution of each state can be described, which in turn can be related to the pump's cycling rate. However, generally the transitions between 'micro state' conformations within the E1 or E2 'macro' states, such as the Ca^{2+} binding steps in E1, are rapid relative to the slow E1 to E2 transitions. Hence, the micro states comprising the E1 and E2 stages are approximately in steady-state. This allowed us to describe the SERCA pump cycle rate as a two-state model for the E1 and E2 macro states (Eq. 8), which we used to relate experimentally and computation-determined binding constants to SERCA function.

In this state-based model, with exception to the undetermined transition rates between E1 and E2 (k_5^+ , k_5^- , k_6^+ and k_6^-), the resting rate constants and substrate concentrations were taken from.^{40,41} The above unknown rates were left as free parameters that were fitted to reproduce experimentally-determined turnover rates from⁴⁰ (see black data points in Fig. 6a)). This fitting accuracy was assessed as the difference between model predicted rates and experimental values reached minimum, as defined by

$$\text{Diff} \equiv \sqrt{\left(\frac{\sum_{i=1}^N (M_{r,i} - \text{Exp}t_{r,i})^2}{N} \right)} \quad (1)$$

where $M_{r,i}$ is state model predicted pump rate at experimental $[\text{Ca}]$, $\text{Exp}t_{r,i}$ is the experimentally measured rate and N is the total number of data points. As shown in Fig. 6(a), the fitted Ca_{only} model (blue) reproduces the experimentally-determined SERCA turnover rates as a function of cytosolic Ca^{2+} concentration, which validates our state-based model.

2.4 Effects on non-cognate ions on steady-state behavior of SERCA

Since a primary focus of this study was to elucidate to which extent the non-cognate ions Mg^{2+} and K^+ influenced the SERCA transport cycle, we introduced additional microstates representing the Mg^{2+} - and K^+ -bound configurations. The resulting representations are

summarized in Fig. S4 and differ in terms of whether the ions serve as inhibitors or intermediates. Using the fitted model from Fig. 6a, we introduced dissociation constants for the Mg^{2+} and K^+ species. We considered two strategies for defining those constants: 1) using experimentally-determined values from Table S4 and 2) constants determined from rescaling of the MSA-predicted values. For 2), the MSA predicted $\text{Mg}^{2+}/\text{K}^+$ chemical potentials were first converted to dissociation constants via $K_d = e^{\frac{\mu_{\text{MSA}}}{RT}}$. Second, the K_d s were multiplied by a scaling factor, λ , that minimizes the difference between the MSA-predicted and experimentally-measured Ca^{2+} dissociation constants,

$$(Cai_{ref} - \lambda \cdot Cai_{MSA})^2 - (Caii_{ref} - \lambda \cdot Caii_{MSA})^2 = 0 \quad (2)$$

where $Cai_{ref}/Caii_{ref}$ and $Cai_{MSA}/Caii_{MSA}$ are site I/II Ca^{2+} dissociation constants from⁴⁰ (4×10^{-8} M and 4×10^{-6} M) and from MSA calculations (9.55×10^{-6} M and 1.38×10^{-5} M), respectively. Eq. 2 was minimized by $\lambda = 0.17$, thus yielding 4.93×10^{-5} M and 3.3×10^{-3} M for the MSA-predicted values of $K_{d,Mg}$ and $K_{d,K}$ for Mg^{2+} and K^+ (see also Table S3). Predictions of the SERCA cycling rate using experimentally-determined dissociation constants are shown in Fig. 6(b) and the rescaled MSA constants in panel (c). Both approaches indicate that the Mg_+K_- (green) and Mg_-K_+ (yellow) provide the best agreement with the experimentally-measured turnover rates, as these two models have relative smaller normalized difference values than other models, with the MSA-determined dissociation constants yielding the strongest agreement overall. Hence, the cycling rate data reported by Cantilina et al⁴⁰ was sufficient to eliminate two of the four proposed models.

To discriminate between the remaining Mg_+K_- and Mg_-K_+ models, we next assessed the abilities of the respective models to reproduce steady-state Ca^{2+} binding data measured at various Mg^{2+} concentrations by Guillain et al.¹⁷ In these experiments, both the E1.Mg and E1.2Ca states contributed a fluorescence signal indicative of Ca^{2+} saturation, therefore we report in Fig. 7 the combined probabilities of those states,

$$P_{\text{Cabin}} = P_{\text{E1.Mg}} + P_{\text{E1.2Ca}} \quad (3)$$

where $P_{\text{E1.Mg}}$ and $P_{\text{E1.2Ca}}$ are the probabilities of the E1.Mg and E1.2Ca states (see Sect. S1.3). The experimental data (black) shown in Fig. 7 demonstrate that Ca^{2+} saturation naturally increases with increasing cytosolic Ca^{2+} , but importantly, saturation increases as Mg^{2+} is raised from 0 mM (circles) to 20 mM (solid triangles). These data indicate that Mg^{2+} locks SERCA into an E1 state in absence of Ca^{2+} . We plot in Fig. 7(a–b) the predicted P_{Cabin} data for the Mg_-K_+ and Mg_+K_- , respectively, as well as the fit in Fig. 7(c). We find that the Mg_-K_+ model provides the optimal fit with difference of 0.75 normalized to Mg_+K_- model. The Mg_-K_+ model correctly captures the plateau in Ca^{2+} saturation at Ca_2 concentrations below 1×10^{-7} , in contrast to the competing model. We note that as Mg^{2+} is increased to unphysiologically high concentrations (> 5 mM), the slope of the experimentally-determined saturation curves decreases, which is indicative of a loss in Ca^{2+} -binding cooperativity. Our model does not directly consider ion-dependent modulation of Ca^{2+} binding cooperativity, therefore this behavior is not reproduced in our predicted data and accounts for some of the error relative to experiment. Additionally, we predict a greater

population of the Ca^{2+} -bound state at increasing Mg^{2+} levels than is experimentally observed, which accounts for the remainder of the error. Nevertheless, we find that Mg-K_+ model provides the best agreement with experimental data, especially within physiological Mg^{2+} concentrations. Therefore, Mg^{2+} most likely acts as inhibitor and K^+ as an intermediate in the SERCA pumping cycle.

Lastly, as a demonstration of how MSA-predicted Ca^{2+} affinities could be applied to infer changes in SERCA pumping function, we compared the turnover rates between WT and two variants using relative affinity estimates from Sect. 2.2. To be consistent with the procedure of obtaining $\text{Mg}^{2+}/\text{K}^+$ affinity from the MSA potentials, the same scaling factor $\lambda = 0.17$ was applied to the MSA-predicted affinities for the E309Q and N796A variants (Ca^{2+} affinities at site I/II are $2.98 \times 10^{-6} \text{ M}/2.81 \times 10^{-6} \text{ M}$ for N796A and $4.06 \times 10^{-6} \text{ M}/2.25 \times 10^{-6} \text{ M}$ for E309Q). Relative to WT SERCA for which the half-maximal pump rate (approximately 2.5 per second) occurs at $6 \times 10^{-7} \text{ M}$ $[\text{Ca}]$, the weaker affinities exhibited by the E309Q and N796A variants right-shift the half-maximal rate to $\sim 1 \times 10^{-5} \text{ M}$ $[\text{Ca}]$. In other words, the SERCA variants are essentially non-functional within physiological Ca^{2+} concentrations (1×10^{-7} to $1 \times 10^{-6} \text{ M}^{42}$), which is consistent with experimentally-observed decreases in SERCA activity for the E309Q and N796A variants.⁴³ Although we recognize that the experimentally-measured activities arise from a culmination of factors beyond just the Ca^{2+} binding affinity in the binding domains, these data qualitatively indicate that MSA predicted affinities can then be used to rationalize steady-state pump turnover rates estimated from molecular-level simulations.

3 DISCUSSION

3.1 Ion coordination and contributions to cation binding affinity

A key contribution from this study was our use of MSA theory to evaluate trends in Ca^{2+} and non-cognate binding in the SERCA E1 states. By using molecular dynamics simulations of the wild type and two site-directed mutations, we could probe shifts in the binding site configurations - relative to the available crystal structures of the pump - that contribute to chelating cations. Overall, the MSA theory, when informed using molecular simulation data including water distributions, appears to be effective in rank-ordering ions by affinity (approximately -7 to -10 kcal/mol , see Fig. 8). Further, our state-based model of SERCA pumping function correctly captures cycling rates across physiological Ca^{2+} concentrations and predicts functional effects of site-directed mutations (N796A and E309Q).

We first applied this combined MSA/MD technique to an EF-hand containing protein called β parvalbumin to investigate factors contributing to its Ca^{2+} affinity and selectivity.³⁴ In both systems, the MSA/MD approach indicated that high affinity Ca^{2+} binding is afforded through the tight clustering of chelating oxygens with partial negative charges. Importantly, by predicting chemical potentials of ion binding, the MSA theory provided thermodynamic information about the relative competition between ions for the Ca^{2+} binding domains, which is generally unfeasible by MD alone. For instance, we determined that Mg^{2+} , and not K^+ , is thermodynamically more probable to occupy the pump at low Ca^{2+} levels, but is disfavored as Ca^{2+} rises toward micromolar concentrations typical in eukaryotic cells. In principle, this approach could additionally account for variations in electrolyte composition

expected near charged lipid bilayers, but we found scant differences in binding assuming a surface charge of 25 mV (see Fig. 5).

Consistent with our earlier findings for Ca^{2+} -binding to β parvalbumin,³⁴ MSA predicts increasing stability of the Ca^{2+} ion as the number of coordinating oxygens is increased. This trend reversed as the oxygen density increased beyond roughly six per $2.2 \times 10^{-1} \text{ nm}^3$, which is presumably when the volume is insufficient to accommodate all coordinating oxygens. For the E309Q and N796A variants, we potentially reduced the number of oxygens that could directly coordinate Ca^{2+} . Based on the MD simulations for E309Q, however, it appeared that the loss of one carboxylic acid oxygen was compensated by a new interaction with E58 that maintained the same coordination number as the wild-type structure. This suggests that there is some degree of flexibility in coordination partners for the ion, which may facilitate the E309 residue's role in kinetically gating the binding site.²⁰ In contrast, the N796A mutation was not offset by a nearby available oxygen, thus the predicted chemical potential was less favorable. Therefore, our data suggest that the MSA could reasonably rank-order Ca^{2+} binding stabilities based on structures of the immediate coordination shell based on Fig. 4, but is less sensitive to broad ranges of binding affinities reported in the literature (see Fig. 8).

We note that our predictions of favorable binding stabilities for Ca^{2+} at site II for the N796A and E309Q variants are at odds with Inesi et al,^{44,45} who suggested based on SERCA ATPase activity that the site was incapable of binding Ca^{2+} . Surprisingly, a structure of the E309Q variant (PDB ID: 4nab¹⁵) deposited in the protein databank contains Ca^{2+} at both binding sites. Ostensibly, the E309Q variant has a much lower, but nonzero affinity for Ca^{2+} . Thus, it is likely that the MSA model does not sufficiently penalize nonoptimal Ca^{2+} coordination, or reflect changes in internal strain that might disfavor coordination.³⁴

We also found that including water distributions made a modest improvement in chemical potentials, as per Fig. 8. Based on Fig. S5, the narrow binding site volumes presented in sites I and II favor Ca^{2+} and Mg^{2+} over K^+ , while both divalent ions are increasingly stabilized by greater densities of coordinating oxygens and waters. It is interesting that the water-free MSA calculations indicate Ca^{2+} could be more stable in a binding site volume of $2.2 \times 10^{-1} \text{ nm}^3$, which is smaller than the volume presented in the MD simulations, albeit with a fewer number of oxygens (approximately five versus six). However, when water is considered, the stabilities at the smaller binding volumes are disfavored. It is possible that there is a kinetic advantage to having mobile waters involved in coordination, which could favor more rapid exchange of bound ions with the surrounding solvent. It is also possible that the trend can be explained on a thermodynamic basis, namely that by preserving waters that coordinate in the ion in bulk solvent, the desolvation cost upon binding SERCA are reduced, which should increase the apparent binding affinity. An intriguing possibility is that the Ca^{2+} affinity could be indirectly tuned by controlling the density of binding site waters.

The primary function of the SERCA pump is to transport cytosolic Ca^{2+} into the sarcoplasmic or endoplasmic reticulum, therefore we verified that MSA would indeed predict an unfavorable change in chemical potential based on conformational changes induced in sites I and II upon forming the E2 state. Since the focus of this study was on the

E1 Ca^{2+} binding thermodynamics, we did not perform MD simulations of the E2 state. Nevertheless, based on the crystal structure of SERCA/beryllium fluoride complex (PDB ID: 3b9b⁶), which represents the pump's E2P state, we illustrate in Fig. 9 that drastic changes in the Ca^{2+} binding site configuration culminate in a significant reduction in oxygens that could potentially coordinate Ca^{2+} . We further evaluate the binding stability for a hypothetical Ca^{2+} bound between the labeled coordination groups and find that the MSA values are not only more positive than those of the E1 state, but are additionally greater than 0 kcal/mol. The positive values of approximately 1 kcal/mol indicate that Ca^{2+} binding at these position is less thermodynamically favorable than partitioning into the surrounding bulk electrolyte. In other words, when SERCA transitions into the E2 state, it is thermodynamically preferred for Ca^{2+} to vacate the binding site in favor of the reticulum lumen. Along these lines, mutations that alter the free energy difference for the Ca^{2+} sites in the E1 and E2 configurations could affect the efficiency of the ATPase.

It has been suggested that the number of cations (Ca^{2+} , Mg^{2+} , and K^+) bound to SERCA is approximately constant across its many conformational states.⁴⁶ In other words, Ca^{2+} binding is offset by Mg^{2+} and K^+ dissociation. ATP is bound to SERCA in complex with Mg^{2+} ,⁴⁷ but there also exists structural⁴⁸ and activity⁴⁹ data that confirm Mg^{2+} binds in the pump's transmembrane region. Toyoshima et al.,⁴⁸ for example, obtained the x-ray crystal structure of the pump with a single Mg^{2+} bound at a 'hybrid' transmembrane site, for which the ion is coordinated by ~5 oxygens with distances of approximately 2.0 Å. Since Mg^{2+} bears the same charge as Ca^{2+} , but with a modestly smaller radius, it is somewhat surprising that Mg^{2+} preferentially binds at an intermediate site between the canonical Ca^{2+} binding sites I and II. We attribute the thermodynamic preference for Ca^{2+} at those sites based on two factors: the higher desolvation energy for Mg^{2+} relative to Ca^{2+} (779.94 kT versus 648.65 kT³³) and the smaller binding site volume for the intermediate site versus sites I and II. With respect to the latter factor, in principle, Mg^{2+} could reduce the site I and II volumes by pulling the chelating oxygens toward the smaller ion, as we previously observed in.³⁴ For the EF-hand (helix-loop-helix) containing protein, we found that constricting the binding loop region introduced greater strain for Mg^{2+} relative to Ca^{2+} , which we suggested would reduce the overall binding affinity for Mg^{2+} . Although we did not explicitly evaluate strain that could be introduced upon Mg^{2+} -binding for SERCA, we anticipate an even greater cost for reducing the binding site volume, since the chelating amino acid are firmly tethered to relatively immobile transmembrane helices. It is important to emphasize, though, that the non-optimal volumes for Mg^{2+} binding at sites I and II does not preclude the ion from binding, as we demonstrate high binding probabilities at very low Ca^{2+} levels (see Fig. 7).

Consistent with our earlier findings in β -PV, K^+ is disfavored from binding the native Ca^{2+} binding volume based on its significantly larger radius and smaller charge.⁵⁰ However, K^+ and other monovalent cations have been demonstrated to stimulate SERCA function, with K^+ being the most efficient agonist.^{18,46,49} Moreover, based on molecular simulations performed in the presence of transmembrane-bound K^+ , Espinosa-Fonseca et al suggest that K^+ facilitates the pump's E2 to E1 transition through its stabilization of the E1 state.²¹ In our simulations, we found that of the two K^+ ions placed in the Ca^{2+} binding domains, the site I K^+ remained bound with approximately four coordinating oxygens with ion/oxygen distances exceeding 4 Å. Based on the MSA predictions, although K^+ has a favorable ($\mu < 0$)

binding potential that is consistent with its sub-millimolar binding affinity at site I,¹⁸ Mg^{2+} and Ca^{2+} are considerably more likely to be bound (see Fig. 5). Meanwhile, water frequently exchanged with K^+ in site II, which is suggestive of a low affinity for the cation.

Based on our comparison with proteins that selectively bind K^+ , such as the KcsA K^+ channel, selective binding of K^+ over competing Ca^{2+} might best be afforded through placement of carbonyl oxygens at sufficiently large distances to accommodate the K^+ ion's larger volume. Namely, in K^+ channels, oxygens from the backbone or side chains of amino acid forming the selectivity filter are exquisitely arranged to achieve precise pore size control and K^+ -oxygen interaction strength.⁵¹ In contrast, it is likely that K^+ exerts its agonistic effects on SERCA through binding the cytoplasmic domain, as was evidenced in a crystal structure determined by Sorenson et al, based on a bound K^+ ion in the P-domain.⁵ According to the K^+ -oxygen coordination pattern shown in Fig. S10, the MSA predicted potential for the P domain K^+ is -2.38 kcal/mol, which is comparable to the values predicted for K^+ bound to site I (-2.35 kcal/mol). Consistent with this structural evidence, the E2P.2Ca dephosphorylation data indicate that K^+ stimulates the Ca^{2+} -release step in this state, whereby the luminal Ca^{2+} affinity is reduced, rates of exchanging luminal Ca^{2+} with luminal solvated Ca^{2+} are accelerated,⁵² and Ca^{2+} release is enhanced.⁵³ Similarly, dephosphorylation of the E2P state is blunted in the absence of K^+ .⁴⁶

Although the thermodynamics of K^+ binding are unfavorable relative to Mg^{2+} and Ca^{2+} , there remains the possible role of K^+ in shaping the kinetics of SERCA function. It has been speculated, for instance, that K^+ accelerates Ca^{2+} binding by first transiently occupying site I, after which it exchanges quickly with a Ca^{2+} at site II.⁵³ K^+ -facilitated exchange could therefore permit faster incorporation of Ca^{2+} into site I, as opposed to the direct migration of Ca^{2+} into a site that is potentially only partially-formed.⁵³ This interpretation is consistent with our observations from molecular dynamics simulations, and was initially reported in simulations of Ca^{2+} -free SERCA by Musgaard et al.⁵⁴

3.2 Relating cation affinity to ATPase activity

In our opinion, the bridging of molecular-level simulation and MSA thermodynamic data with state-based models represents a significant contribution toward multi-scale modeling of steady-state SERCA activity. Numerical and computational modeling of SERCA activity has spanned phenomenological frameworks, such as Hill-type models^{55–57} to those representing distinct stages of the catalytic process as states.^{40,41,58–60} Our implementation is based on the stepwise binding of Ca^{2+} originally proposed by Inesi et al,⁴⁰ but additionally considers Mg^{2+} - and K^+ -bound states, followed by a reduction scheme to lump the E1 states separately from those comprising the E2 configurations. Significantly, the states defined in our model coincide with SERCA structures determined by x-ray crystallography and are ordered in a manner consistent with assays of SERCA function⁶¹ In contrast, the state model proposed by Tran et al⁴¹ assumed that ATP binding precedes the binding of two Ca^{2+} ions, which has not been experimentally verified.

Existing models of SERCA function have faithfully recapitulated the pump's activity. However, an advantage of our state-based model is that its alignment with experimentally-determined structures permits us to directly investigate how structural modifications might

impact activity. For example, we performed MSA/MD simulations of two mutated SERCA variants that yielded modified Ca^{2+} binding constants that we used to predict SERCA activity (see Fig. 6). While we acknowledge that there are likely myriad changes to the pump's energetics following mutation that could be accounted for in the state-based model, our implementation here is a significant step toward structure-based modeling of SERCA activity. At a minimum, our fitted state model is consistent with steady-state pump rates data collected as a function of Ca^{2+} by Cantilina et al.,⁴⁰ Ca^{2+} -saturated versus Mg^{2+} ,¹⁷ as well as molecular simulations that predict Mg^{2+} inhibition of the pump.²¹ Furthermore, by considering different linkages of K^{+} - and Mg^{2+} -bound states to the reaction scheme, we were able to determine that assumptions of Mg^{2+} inhibition and K^{+} agonist were most consistent with experimental data collected by Guillain et al.¹⁷ In principle, multiscale models of SERCA activity that include structure-derived thermodynamic information could permit in silico investigations of how disease-associated SERCA mutations,^{62,63} post-translational modifications,⁶⁴ and binding of regulatory proteins such as phospholamban⁵⁸ affect pump function.

3.3 Limitations

There are several limitations of our approach that could be addressed in future studies. Our study largely focused on conformational changes and energetics of the cations' immediate coordination sphere. Mg^{2+} and Ca^{2+} -bound structures (PDB ID: 1su4 and 3w5b, respectively) deposited in the Protein Data Bank indicate substantial differences in the conformations of the transmembrane bundle helices and cytosolic domains that will necessarily contribute to the free energies of binding. For this reason, the free energy differences implied in experimental measurements of cation binding affinities reflect contributions from both ion coordination and the pump's different conformations. It is also important to emphasize that the MSA assumptions of a spherical binding volume within which amino acid oxygens are immobilized crudely approximates the actual binding site. Thus, the predicted thermodynamic quantities are most appropriate for rank-ordering different ion/binding site configurations. Additionally, there is evidence^{16,61} that Ca^{2+} binding is cooperative, although here we treat the binding events independently. It may be possible to partially recover some of the cooperative behavior by modeling SERCA with only one bound Ca^{2+} and using end-point methods such as Molecular Mechanics/Poisson Boltzmann Surface Area (MM/PBSA)⁶⁵ to estimate energy changes upon binding a second Ca^{2+} . To our knowledge, however, a half-saturated Ca^{2+} -bound SERCA structure has not been determined, which would challenge efforts to validate predictions. Additionally, in the sequential binding model used by Inesi and others^{6,40} ($E1 \rightarrow E1.Ca \rightarrow E1'.Ca \rightarrow E1'.2Ca$), the apparent kinetics governing the transition between the $E1.Ca$ to $E1'.Ca$ states can vary depending on a variety of factors, including the presence the regulatory protein phospholamban.^{58,66} Thus, extending this model to broader conditions and regulatory proteins would require careful consideration of conformational changes that might accompany the $E1.Ca$ to $E1'.Ca$ transition. Lastly, the Ca^{2+} -binding steps in our model are assumed to be in steady state relative to the substantially slower transitions between the $E1$ and $E2$ states. In the event that the pump is subject to rapid Ca^{2+} oscillations, such as spontaneous Ca^{2+} spiking in glia,⁶⁷ the steady-state approximation may be unreliable.

4 CONCLUSIONS

We utilized molecular dynamics simulations, mean sphere approximation theory and state-based modeling to probe effects of Ca^{2+} , Mg^{2+} and K^{+} binding on the SERCA pump cycle. The MD and MSA approaches indicate that favorable binding of Ca^{2+} in the wild-type SERCA configuration is facilitated through a high degree of coordination by amino acids comprising the binding sites, as well as significant contributions from water coordination. This coordination pattern appears to be impaired in the E309Q and N796A variants; using MSA theory, we found that the chemical potential of Ca^{2+} binding is less favourable relative to wild-type as a result. Hence, mutations near the Ca^{2+} binding domains that alter the coordination number, hydration and binding site volume can be expected to modulate Ca^{2+} affinity in a manner qualitatively described by the MSA theory. Similarly, the coordination patterns exhibited in the Mg^{2+} and K^{+} -bound structures led to less favorable binding estimates from MSA-theory. These findings were qualitatively consistent with measured affinity data reported in the literature.^{15–18,44,53} Additionally, we developed a state-based model of SERCA activity that we used to: 1) relate Ca^{2+} binding affinities to the SERCA pump rate; and 2) determine whether Mg^{2+} and K^{+} are pump agonists or inhibitors. We found that the state model treating Mg^{2+} as an inhibitor and K^{+} as an agonist ($\text{Mg}_{-}\text{K}_{+}$) was most consistent with experimental data. Despite the limitations of the assumptions used for the various models, we believe that the approach provides an attractive framework for evaluating allosteric functional effects of ion binding on SERCA, which may be extendable to other Ca^{2+} transporters, such as P2X4.⁶⁸

5 METHODS

5.1 Construction of the SERCA systems

Molecular dynamics (MD) simulations on wild-type (WT) SERCA and two variants of SERCA: E309Q and N796A were performed; the latter variants were chosen given diminished site II Ca^{2+} -binding.⁴³ For the WT SERCA system, we considered apo (free of bound ions) Ca^{2+} -bound, Mg^{2+} -bound and K^{+} -bound states. Ca^{2+} -bound (PDB ID: 1su4³) and Mg^{2+} -bound (E1.Mg, PDB ID: 3w5b⁴⁸) SERCA X-ray crystal structures were used as the starting structures. The apo SERCA system was constructed based on 1su4 with two Ca^{2+} ions removed as done in.⁶⁹ The structure of the site I or II bound K^{+} SERCA has not been determined, thus we created the structure based on the Mg^{2+} -bound variant. For the N796A mutant, Ca^{2+} -bound and Mg^{2+} -bound cases were considered, based on mutating N796 to alanine. Similarly, for the E309Q mutant, we obtained two rotamers compatible with the binding site as evaluated through UCSF Chimera.⁷⁰ These E309Q rotamers were designated as "E309Q_r1" and "E309Q_r2". All SERCA cases considered in present study are summarized in Table S1.

The cation coordinating acidic residues E309, E771 and D800 were assumed to be deprotonated, while E908 was protonated, consistent with;²¹ further a disulfide bond was introduced between residues C876 and C888. The system was inserted into a homogeneous 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (POPC) lipid bilayer of POPC via the Membrane Builder⁷¹ within the CHARMM-GUI software.⁷² This system was solvated via TIP3P waters⁷³ using a 20 Å margin perpendicular to the membrane. Counterions K^{+} and

chloride (Cl^-) were added into the system via Monte-Carlo method to neutralize the system and maintain an ionic strength of 0.15 M. Both the solvation and ion-adding were performed via the Solvator module within CHARMM-GUI online-server. The final system contained ~255,000 atoms including lipid bilayer with 461 lipids, and ~59,000 TIP3P water molecules and was parameterized by the CHARMM36 force field.^{74,75}

5.2 Molecular dynamics simulations

MD simulations were performed using NAMD.⁷⁶ The system was first subjected to an energy minimization process consisting of 2000 steps' steepest descent (SD) and 2000 steps' adopted basis Newton rRphson (ABNR) algorithm. For each case, the minimized system was heated from 0 K to 303.14 K over 25 ps with 1 fs timestep via the Langevin thermostat, using randomized velocities for three triplicate preparations. For each replica, harmonic constraints were introduced during minimization and heating on protein side chains, protein backbone atoms, lipid heavy atoms and ions. The force constants of constraints were set to 5, 10, 10 and $1.0 \times 10^1 \frac{\text{k}}{\text{mol } \text{\AA}^2}$, respectively, and were gradually reduced to zero over five equilibration steps of 20 ps in duration. 100 ns production simulations was then performed on the equilibrated system with the Shake algorithm,⁷⁷ using 2 fs timesteps.

5.3 Simulation trajectory analysis

Standard trajectory analyses including RMSF and coordination oxygen/ Ca^{2+} distances were computed using Lightweight Object-Oriented Structure Library (LOOS).⁷⁸ Coordination pattern analysis on each cation with oxygen atoms from nearby amino acids were performed in each binding site. This consisted of extracting residues within 20 Å from D800 (the shared residue of the conventional two Ca^{2+} -binding sites in SERCA) based on the Ca^{2+} -bound SERCA crystal structure (PDB ID: 1su4). The water density around cation was computed via the radial command implemented in CPPTRAJ⁷⁹ and analyzed via Grid Inhomogeneous Solvation Method (GIST) (as described in Sect. S1.4).

The MSA is a mean-field model which estimates cation chemical potentials in electrolyte solution with finite-sized ions. In this study, the SERCA cation binding sites were treated as confined filters filled with oxygens from coordinating residues and water molecules. The MSA model calculates cation distributions between bulk electrolyte solution that minimize the chemical potential for partitioning ions into a finite size volume occupied by coordinating oxygens; these volume and number of oxygens were determined by MD as in Kucharski et al.³⁴ The free energy expression in this model was assuming negative-charged spherical domains was based on a chemical potential accounting for electrostatic and hard-sphere interactions between ions inclusive of oxygens, as described in.^{35,80,81} In this representation, which is analogous to the Ca^{2+} binding domain in β -PV and calcium channels,^{33–35} ionic species that have a negative chemical potential in the binding filter are thermodynamically favored to bind. We include in this approach a solvation contribution as estimated via generalized Born theory of ion hydration energies:

$$\Delta G_{GB} = \frac{z^2}{r} \left(\frac{e^2 N_a}{2 \cdot 4\pi\epsilon_0} \right) \left(1 - \frac{1}{\epsilon_r} \right) \quad (4)$$

where z is charge number, r is radii, e is electron charge, N_a is the Avogadro constant, ϵ_0 and ϵ_r are vacuum dielectric and the relative dielectric constant of the solvent. Additional details are elaborated in the supplementary material of.³⁴

In the SERCA system, the cation/SERCA configurations can present differing numbers of coordination oxygen atoms (see Fig. 2 and Table S2) and volumes, as shown previously for β -PV in:³⁴

$$V_f = \frac{4}{3}\pi R^3 \quad (5)$$

$$R = r_{rdf} + r_{oxy} \quad (6)$$

where r_{rdf} is the radii of optimal coordination sphere which can be obtained from the cation-oxygen coordination pattern analysis based on MD simulations (see Fig. 2) while r_{oxy} is the radii of oxygen atom. Charges and radii values of all ions are taken from the Li-Merz work.

⁸² The specific parameters used in the MSA calculations are listed in Table S2.

5.4 State-based model of the SERCA pump cycle

SERCA pumping is characterized by two prominent states, E1 and E2, comprised of microstates corresponding to various conformations or bound states of the pump (Fig. S4). In the E1 state, two Ca^{2+} ions bind to SERCA through cooperative mechanism followed by the binding of MgATP.⁴⁰ We represent each state as by an ordinary differential equation of the form

$$ds_i/dt = \sum k_{ji}s_j - \sum k_{ij}s_i \quad (7)$$

where s_i is state i , and k_{ij} describe rates for transitioning between states i and j . The models tested in this paper are summarized in Fig. S4. While the cooperative Ca^{2+} binding mechanism in the E1 state is well-accepted, the Ca^{2+} release mechanism of the E2 state is not clearly resolved, therefore we use the state representations proposed in Tran et al.⁴¹

Based on the technique proposed by Smith et al.,⁸³ we can applied rapid equilibrium assumption to fast processes (those within E1 or E2) relative to the slow rates of transitions between the E1 and E2 states to simplify the model. As shown in Fig. S4, states in dashed boxes were assumed to be in quasi-steady-state and thus were lumped together. Hence, the original multi-state models were simplified into a two-state model (see bottom Fig. S4) in which new apparent rate constants were derived based on original rate constants. The resulting equations are summarized in the supplement Sect. S1.3.

For the lumped 2-state model, the turn over rate is derived as in Tran's paper:⁴¹

$$V = \frac{\alpha_1^+ \alpha_2^+ - \alpha_1^- \alpha_2^-}{\alpha_1^+ + \alpha_1^- + \alpha_2^+ + \alpha_2^-} \quad (8)$$

where $\alpha_i^\pm$ are apparent rate constants between lumped two states (the expressions are given in Sect. S1.3). The final parameters used are listed in Table S3.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENT

Research reported in this publication was supported by the Maximizing Investigators' Research Award (MIRA) (R35) from the National Institute of General Medical Sciences (NIGMS) of the National Institutes of Health (NIH) under grant number R35GM124977, as well as the award 1F32HL114365-01A1. This work used the Extreme Science and Engineering Discovery Environment (XSEDE), which is supported by National Science Foundation under grant number ACI-1548562. We thank Michael Autry for discussion of this work.

References

- (1). Ma H; Inesi G; Toyoshima C Substrate-induced conformational fit and headpiece closure in the Ca²⁺ ATPase (SERCA). *J. Biol. Chem* 2003, 278, 28938–28943. [PubMed: 12750373]
- (2). Inesi G; Prasad AM; Pilankatta R The Ca²⁺ATPase of cardiac sarcoplasmic reticulum: Physiological role and relevance to diseases. *Biochem. Biophys. Res. Commun* 2008, 369, 182–187. [PubMed: 18068669]
- (3). Toyoshima C; Nakasako M; Nomura H; Ogawa H Crystal structure of the calcium pump of sarcoplasmic reticulum at 2.6 Å resolution. *Nature* 2000, 405, 647–655. [PubMed: 10864315]
- (4). Toyoshima C; Nomura H Structural changes in the calcium pump accompanying the dissociation of calcium. *Nature* 2002, 418, 605–611. [PubMed: 12167852]
- (5). Sørensen TLM; Clausen JD; Jensen AML; Vilsen B; Møller JV; Andersen JP; Nissen P Localization of a K⁺-binding site involved in dephosphorylation of the sarcoplasmic reticulum Ca²⁺-ATPase. *J. Biol. Chem* 2004, 279, 46355–46358. [PubMed: 15383548]
- (6). Olesen C; Picard M; Winther AML; Gyrupe C; Morth JP; Oxvig C; Møller JV; Nissen P The structural basis of calcium transport by the calcium pump. *Nature* 2007, 450, 1036–1042. [PubMed: 18075584]
- (7). Inesi G; Scarpa A Fast kinetics of adenosine triphosphate dependent Ca²⁺ uptake by fragmented sarcoplasmic reticulum. *Biochemistry*. 1972, 11, 356–359. [PubMed: 5059117]
- (8). Froehlich JP; Taylor EW Transient state kinetic effects of calcium ion on sarcoplasmic reticulum adenosine triphosphatase. *J. Biol. Chem* 1976, 251, 2307–2315. [PubMed: 131125]
- (9). Dupont Y Fluorescence studies of the sarcoplasmic reticulum calcium pump. *Biochem. Biophys. Res. Commun* 1976, 71, 544–550. [PubMed: 134709]
- (10). Dupont Y; Leigh JB Transient kinetics of sarcoplasmic reticulum Ca²⁺ATPase studied by fluorescence. *Nature* 1978, 273, 396–398. [PubMed: 149252]
- (11). Coan CR; Inesi G Ca²⁺-dependent effect of acetylphosphate on spin-labeled sarcoplasmic reticulum. *Biochem. Biophys. Res. Commun* 1976, 71, 1283–1288. [PubMed: 184786]
- (12). Meis L; Vianna AL Energy Interconversion by the Ca²⁺-Dependent ATPase of the Sarcoplasmic Reticulum. *Annu. Rev. Biochem* 1979, 48, 275–292. [PubMed: 157714]
- (13). Inesi G Mechanism of Calcium Transport. *Annu. Rev. Physiol* 1985, 47, 573–601. [PubMed: 2581506]
- (14). Mintz E; Guillain F Ca²⁺ transport by the sarcoplasmic reticulum ATPase. *Biochim. Biophys. Acta, Bioenerg* 1997, 1318, 52–70.

- (15). Clausen JD; Bublitz M; Arnou B; Montigny C; Jaxel C; Møller JV; Nissen P; Andersen JP; Le Maire M SERCA mutant E309Q binds two Ca²⁺ ions but adopts a catalytically incompetent conformation. *EMBO J* 2013, 32, 3231–3243. [PubMed: 24270570]
- (16). Inesi G; Kurzmack M; Coan C; Lewis DE Cooperative calcium binding and ATPase activation in sarcoplasmic reticulum vesicles. *J. Biol. Chem* 1980, 255, 3025–3031. [PubMed: 6244305]
- (17). Guillain F; Gingold MP; Champeil P Direct fluorescence measurements of Mg²⁺ binding to sarcoplasmic reticulum ATPase. *J. Biol. Chem* 1982, 257, 7366–7371. [PubMed: 6211442]
- (18). Lee a. G.; Baker K; Khan YM; East JM Effects of K⁺ on the binding of Ca²⁺ to the Ca(2+)-ATPase of sarcoplasmic reticulum. *Biochem. J* 1995, 305 (Pt 1), 225–231. [PubMed: 7826333]
- (19). Huang Y; Li H; Bu Y Molecular dynamics simulation exploration of cooperative migration mechanism of calcium ions in sarcoplasmic reticulum Ca²⁺-ATPase. *J. Comput. Chem* 2009, 30, 2136–2145. [PubMed: 19242958]
- (20). Keken-Huskey PM; Metzger VT; Grant BJ; Andrew McCammon J Calcium binding and allosteric signaling mechanisms for the sarcoplasmic reticulum Ca²⁺ ATPase. *Protein Sci* 2012, 21, 1429–1443. [PubMed: 22821874]
- (21). Espinoza-Fonseca LM; Autry JM; Thomas DD Microsecond molecular dynamics simulations of Mg²⁺- And K⁺-Bound E1 intermediate states of the calcium pump. *PLoS ONE* 2014, 9, DOI: 10.1371/journal.pone.0095979.
- (22). Das A; Rui H; Nakamoto R; Roux B Conformational Transitions and Alternating-Access Mechanism in the Sarcoplasmic Reticulum Calcium Pump. *J. Mol. Biol* 2017, 429, 647–666. [PubMed: 28093226]
- (23). Rui H; Das A; Nakamoto R; Roux B Proton Countertransport and Coupled Gating in the Sarcoplasmic Reticulum Calcium Pump. *J. Mol. Biol* 2018, 430, 5050–5065. [PubMed: 30539761]
- (24). Fernández-De Gortari E; Espinoza-Fonseca LM Structural basis for relief of phospholamban-mediated inhibition of the sarcoplasmic reticulum Ca²⁺-ATPase at saturating Ca²⁺ conditions. *J. Biol. Chem* 2018, 293, 12405–12414. [PubMed: 29934304]
- (25). Ha KN; Masterson LR; Hou Z; Verardi R; Walsh N; Veglia G; Robia SL Lethal Arg9Cys phospholamban mutation hinders Ca²⁺-ATPase regulation and phosphorylation by protein kinase A. *Proc. Natl. Acad. Sci. U.S.A* 2011, 108, 2735–2740. [PubMed: 21282613]
- (26). Bidwell P; Blackwell DJ; Hou Z; Zima AV; Robia SL Phospholamban binds with differential affinity to calcium pump conformers. *J. Biol. Chem* 2011, 286, 35044–35050. [PubMed: 21832088]
- (27). Smolin N; Robia SL A structural mechanism for calcium transporter headpiece closure. *J. Phys. Chem. B* 2015, 119, 1407–1415. [PubMed: 25531267]
- (28). Zhao Y; Ogawa H; Yonekura S-I; Mitsuhashi H; Mitsuhashi S; Nishino I; Toyoshima C; Ishiura S Functional analysis of SERCA1b, a highly expressed SERCA1 variant in myotonic dystrophy type 1 muscle. *Biochim. Biophys. Acta, Mol. Basis Dis* 2015, 1852, 2042–2047.
- (29). Tadini-Buoninsegni F; Smeazzetto S; Gualdani R; Moncelli MR Drug Interactions With the Ca²⁺-ATPase From Sarco(Endo)Plasmic Reticulum (SERCA). *Front. Mol. Biosci* 2018, 5, 36. [PubMed: 29696147]
- (30). Dudev T; Lim C Competition among metal ions for protein binding sites: Determinants of metal ion selectivity in proteins. *Chem. Rev* 2014, 114, 538–556. [PubMed: 24040963]
- (31). Yu H; Noskov SY; Roux B Two mechanisms of ion selectivity in protein binding sites. *Proc. Natl. Acad. Sci. U.S.A* 2010, 107, 20329–20334. [PubMed: 21057111]
- (32). Kiyota Y; Takeda-Shitaka M Molecular recognition study on the binding of calcium to calbindin D9kbased on 3D reference interaction site model theory. *J. Phys. Chem. B* 2014, 118, 11496–11503. [PubMed: 25251030]
- (33). Nonner W; Gillespie D; Henderson D; Eisenberg B Ion accumulation in a biological calcium channel: Effects of solvent and confining pressure. *J. Phys. Chem. B* 2001, 105, 6427–6436.
- (34). Kucharski AN; Scott CE; Davis JP; Keken-Huskey PM Understanding Ion Binding Affinity and Selectivity in -Parvalbumin Using Molecular Dynamics and Mean Spherical Approximation Theory. *J. Phys. Chem. B* 2016, 120, 8617–8630. [PubMed: 27267153]

- (35). Nonner W; Catacuzzeno L; Eisenberg B Binding and selectivity in L-type calcium channels: a mean spherical approximation. *Biophys. J* 2000, 79, 1976–1992. [PubMed: 11023902]
- (36). Gillespie D; Nonner W; Henderson D; Eisenberg RS A physical mechanism for large-ion selectivity of ion channels. *Phys. Chem. Chem. Phys* 2002, 4, 4763–4769.
- (37). Hughes E; Clayton JC; Middleton DA Cytoplasmic residues of phospholamban interact with membrane surfaces in the presence of SERCA: A new role for phospholipids in the regulation of cardiac calcium cycling? *Biochim. Biophys. Acta, Biomembr* 2009, 1788, 559–566.
- (38). Israelachvili JN, *Intermolecular and Surface forces*; Academic Press: 2011.
- (39). Worley JF; McIntyre MS; Spencer B; Mertz RJ; Roe MW; Dukes ID Endoplasmic reticulum calcium store regulates membrane potential in mouse islet β -cells. *J. Biol. Chem* 1994, 269, 14359–14362. [PubMed: 8182038]
- (40). Cantilina T; Sagara Y; Inesi G; Jones LR Comparative studies of cardiac and skeletal sarcoplasmic reticulum ATPases: Effect of a phospholamban antibody on enzyme activation by Ca^{2+} . *J. Biol. Chem* 1993, 268, 17018–17025. [PubMed: 8349590]
- (41). Tran K; Smith NP; Loiselle DS; Crampin EJ A thermodynamic model of the cardiac sarcoplasmic/endoplasmic Ca^{2+} (SERCA) pump. *Biophys. J* 2009, 96, 2029–2042. [PubMed: 19254563]
- (42). Berridge MJ; Bootman MD; Roderick HL Calcium signalling: dynamics, homeostasis and remodelling. *Nat. Rev. Mol. Cell Biol* 2003, 4, 517–529. [PubMed: 12838335]
- (43). Clarke DM; Loo TW; Inesi G; MacLennan DH Location of high affinity Ca^{2+} -binding sites within the predicted transmembrane domain of the sarco-plasmic reticulum Ca^{2+} -ATPase. *Nature* 1989, 339, 476–478. [PubMed: 2524669]
- (44). Inesi G; Zhang Z; Lewis D Cooperative setting for long-range linkage of Ca^{2+} -binding and ATP synthesis in the Ca^{2+} -ATPase. *Biophys. J* 2002, 83, 2327–2332. [PubMed: 12414670]
- (45). Inesi G; Ma H; Lewis D; Xu C Ca^{2+} occlusion and gating function of Glu309 in the ADP-fluoroaluminate analog of the Ca^{2+} -ATPase phosphoenzyme intermediate. *J. Biol. Chem* 2004, 279, 31629–31637. [PubMed: 15150270]
- (46). Shigekawa M; Pearl LJ Activation of calcium transport in skeletal muscle sarcoplasmic reticulum by monovalent cations. *J. Biol. Chem* 1976, 251, 6947–6952. [PubMed: 136443]
- (47). Vianna AL Interaction of calcium and magnesium in activating and inhibiting the nucleoside triphosphatase of sarcoplasmic reticulum vesicles. *Biochim. Biophys. Acta, Enzymol* 1975, 410, 389–406.
- (48). Toyoshima C; Iwasawa S; Ogawa H; Hirata A; Tsueda J; Inesi G Crystal structures of the calcium pump and sarcolipin in the Mg^{2+} -bound E1 state. *Nature* 2013, 495, 260–264. [PubMed: 23455422]
- (49). Jones LR Mg^{2+} and ATP effects on K⁺ activation of the Ca^{2+} -transport ATPase of cardiac sarcoplasmic reticulum. *Biochim. Biophys. Acta, Biomembr* 1979, 557, 230–242.
- (50). Shannon RD; Prewitt CT Effective ionic radii in oxides and fluorides. *Acta Crystallographica Section B* 1969, 25, 925–946.
- (51). Cordero-Morales JF; Cuello LG; Zhao Y; Jogini V; Cortes DM; Roux B; Perozo E Molecular determinants of gating at the potassium-channel selectivity filter. *Nat. Struct. Mol. Biol* 2006, 13, 311–318. [PubMed: 16532009]
- (52). Yamasaki K; Wang G; Daiho T; Danko S; Suzuki H Roles of Tyr122-hydrophobic cluster and K⁺ binding in Ca^{2+} -releasing process of ADP-insensitive phosphoenzyme of sarcoplasmic reticulum Ca^{2+} -ATPase. *J. Biol. Chem* 2008, 283, 29144–29155. [PubMed: 18728008]
- (53). Moutin MJ; Dupont Y Interaction of potassium and magnesium with the high affinity calcium-binding sites of the sarcoplasmic reticulum calcium-ATPase. *J. Biol. Chem* 1991, 266, 5580–5586. [PubMed: 1826001]
- (54). Musgaard M; Thøgersen L; Schiøtt B Protonation states of important acidic residues in the central Ca^{2+} ion binding sites of the Ca^{2+} -ATPase: A molecular modeling study. *Biochemistry*. 2011, 50, 11109–11120. [PubMed: 22082179]
- (55). Sipido KR; Wier WG Flux of Ca^{2+} across the sarcoplasmic reticulum of guinea pig cardiac cells during excitation contraction coupling. *J. Physiol. (Lond.)* 1991, 435, 605–630. [PubMed: 1770453]

- (56). Luo CH; Rudy Y A dynamic model of the cardiac ventricular action potential: I. Simulations of ionic currents and concentration changes. *Circ. Res* 1994, 74, 1071–1096. [PubMed: 7514509]
- (57). Shannon TR; Wang F; Puglisi J; Weber C; Bers DM A mathematical treatment of integrated Ca dynamics within the ventricular myocyte. *Biophys. J* 2004, 87, 3351–3371. [PubMed: 15347581]
- (58). Ceholski DK; Trieber CA; Holmes CFB; Young HS Lethal, hereditary mutants of phospholamban elude phosphorylation by protein kinase A. *J. Biol. Chem* 2012, DOI: 10.1074/jbc.M112.382713.
- (59). Higgins ER; Cannell MB; Sneyd J A buffering SERCA pump in models of calcium dynamics. *Biophys. J* 2006, 91, 151–163. [PubMed: 16617079]
- (60). Koivumäki JT; Takalo J; Korhonen T; Tavi P; Weckström M Modelling sarcoplasmic reticulum calcium ATPase and its regulation in cardiac myocytes. *Philos. Trans. A. Math. Phys. Eng. Sci* 2009, 367, 2181–2202. [PubMed: 19414452]
- (61). Zhang Z; Lewis D; Strock C; Inesi G; Nakasako M; Nomura H; Toyoshima C Detailed characterization of the cooperative mechanism of Ca²⁺ binding and catalytic activation in the Ca²⁺ transport (SERCA) ATPase. *Biochemistry*. 2000, 39, 8758–8767. [PubMed: 10913287]
- (62). Dhitavat J; Dode L; Leslie N; Sakuntabhai A; Lorette G; Hovnanian A Mutations in the sarcoplasmic/endoplasmic reticulum Ca²⁺ ATPase isoform cause Darier's disease. *J. Invest. Dermatol* 2003, 121, 486–489. [PubMed: 12925205]
- (63). Hovnanian A SERCA pumps and human diseases. *Subcell. Biochem* 2007, 45, 337–363. [PubMed: 18193643]
- (64). Xu A; Hawkins C; Narayanan N Phosphorylation and activation of the Ca(2+)-pumping ATPase of cardiac sarcoplasmic reticulum by Ca²⁺/calmodulin-dependent protein kinase. *J. Biol. Chem* 1993, 268, 8394–7. [PubMed: 8386159]
- (65). Kollman PA; Massova I; Reyes C; Kuhn B; Huo S; Chong L; Lee M; Lee T; Duan Y; Wang W; Donini O; Cieplak P; Srinivasan J; Case DA; Cheatham TE Calculating Structures and Free Energies of Complex Molecules: Combining Molecular Mechanics and Continuum Models. *Acc. Chem. Res* 2000, 33, 889–897. [PubMed: 11123888]
- (66). Glaves JP; Trieber CA; Ceholski DK; Stokes DL; Young HS Phosphorylation and mutation of phospholamban alter physical interactions with the sarcoplasmic reticulum calcium pump. *J. Mol. Biol* 2011, 405, 707–723. [PubMed: 21108950]
- (67). Schipke CG; Heidemann A; Skupin A; Peters O; Falcke M; Kettenmann H Temperature and nitric oxide control spontaneous calcium transients in astrocytes. *Cell Calcium* 2008, 43, 285–295. [PubMed: 17698190]
- (68). Chun BJ; Stewart BD; Vaughan DD; Bachstetter AD; Keken-Huskey PM Simulation of P2X-mediated calcium signalling in microglia. *J. Physiol. (Lond.)* 2019, 597, 799–818. [PubMed: 30462840]
- (69). Espinoza-Fonseca LM; Thomas DD Atomic-level characterization of the activation mechanism of serca by calcium. *PLoS ONE* 2011, 6, ed. by Stultz CM, e26936.
- (70). Pettersen EF; Goddard TD; Huang CC; Couch GS; Greenblatt DM; Meng EC; Ferrin TE UCSF Chimera A visualization system for exploratory research and analysis. *J. Comput. Chem* 25, 1605–1612.
- (71). Wu EL; Cheng X; Jo S; Rui H; Song KC; D vila-Contreras EM; Qi Y; Lee J; Monje-Galvan V; Venable RM; Klauda JB; Im W CHARMM-GUI Membrane Builder toward realistic biological membrane simulations. *J. Comput. Chem* 2014, 35, 1997–2004. [PubMed: 25130509]
- (72). Jo S; Kim T; Iyer VG; Im W CHARMM-GUI: A web-based graphical user interface for CHARMM. *J. Comput. Chem* 2008, 29, 1859–1865. [PubMed: 18351591]
- (73). Jorgensen WL; Chandrasekhar J; Madura JD; Impey RW; Klein ML Comparison of simple potential functions for simulating liquid water. *J. Chem. Phys* 1983, 79, 926–935.
- (74). MacKerell AD et al. All-Atom Empirical Potential for Molecular Modeling and Dynamics Studies of Proteins. *J. Phys. Chem. B* 1998, 102, 3586–3616. [PubMed: 24889800]
- (75). MacKerell AD; Feig M; Brooks CL Improved Treatment of the Protein Backbone in Empirical Force Fields. *J. Am. Chem. Soc* 2004, 126, 698–699. [PubMed: 14733527]

- (76). Phillips JC; Braun R; Wang W; Gumbart J; Tajkhorshid E; Villa E; Chipot C; Skeel RD; Kal L; Schulten K Scalable molecular dynamics with NAMD. *J. Comput. Chem* 2005, 26, 1781–1802. [PubMed: 16222654]
- (77). Ryckaert J-P; Ciccotti G; Berendsen HJ Numerical integration of the cartesian equations of motion of a system with constraints: molecular dynamics of n-alkanes. *J. Comput. Phys* 1977, 23, 327–341.
- (78). Romo TD; Leioatts N; Grossfield A Lightweight Object Oriented Structure analysis: Tools for building Tools to Analyze Molecular Dynamics Simulations. *J. Comput. Chem* 2014, 35, 2305–2318. [PubMed: 25327784]
- (79). Roe DR; Cheatham TE PTRAJ and CPPTRAJ: Software for Processing and Analysis of Molecular Dynamics Trajectory Data. *J. Chem. Theory Comput* 2013, 9, 3084–3095. [PubMed: 26583988]
- (80). Høye JS; Blum L The mean spherical model for asymmetric electrolytes: Thermodynamics and the pair correlation function. *Mol. Phys* 1978, 35, 299–300.
- (81). Herrera JN; Blum L Sticky electrolyte mixtures in the Percus Yevick-/mean spherical approximation. *J. Chem. Phys* 1991, 94, 5077–5082.
- (82). Li P; Roberts BP; Chakravorty DK; Merz KM Rational Design of Particle Mesh Ewald Compatible Lennard-Jones Parameters for +2 Metal Cations in Explicit Solvent. *J. Chem. Theory Comput* 2013, 9, 2733–2748. [PubMed: 23914143]
- (83). Smith N; Crampin E Development of models of active ion transport for whole-cell modelling: cardiac sodium potassium pump as a case study. *Prog. Biophys. Mol. Biol* 2004, 85, 387–405. [PubMed: 15142754]

Figure 1:
Comparison of Ca²⁺-bound (PDB ID: 1su4) and Mg²⁺-bound (PDB ID: 3w5b) SERCA crystal structures. (a) Superimpose of two crystal structures with 1su4/3w5b cytosolic domains (A, P and N) colored in red/salmon, blue/slate and green/palegreen. The TM helices are colored in cyan and grey for 1su4 and 3w5b, respectively. (b-c) Mg²⁺ and Ca²⁺ binding site comparison. In this orientation, sites I and II are right and left, respectively. Mg²⁺ and Ca²⁺ are represented as magenta and yellow balls. The distance between coordination oxygen atoms and cation are also shown. The oxygen atoms of crystal water molecules are shown as red balls.

Figure 2:

Radial distribution function (RDF) of amino acid and water oxygen atoms about bound Ca^{2+} , Mg^{2+} , or K^{+} . The distribution around Ca^{2+} for each individual case is shown in Fig. S9 for clarity. The inset bar graphs show the maximum number of coordinating amino acid oxygen and water density around the cation in each case (the bars with black dots represent site II Ca^{2+}). The coordinating waters with site I Ca^{2+} of WT SERCA is also shown. Ca^{2+} ions bound to WT SERCA tend to reflect the highest degree of coordination among the modeled systems.

Figure 3:

Heavy atom RMSF of key coordinating residues at the cation binding sites. The error bars were calculated from the triplicate trajectories for each case. The upper row represents site I residues while lower row were site II residues (E58 was not involved into coordination with cations in the crystal structure, although interactions with bound ions were predicted in our simulations, see Fig. S11). Ca^{2+} -bound configurations generally exhibit lesser mobility compared to the configurations with non-cognate ions Mg^{2+} and K^{+} .

Figure 4:

Mean spherical approximation (MSA)-predicted chemical potentials for each cation based on oxygen-coordination pattern (waters included) and optimal filter volumes derived from MD simulations. Potentials are given relative to Ca^{2+} bound to wild-type SERCA at site I. Ca^{2+} bound cases exhibit more favorable binding interactions compared to the binding of non-cognate ions, with the WT cases reflecting the most favorable potentials relative to the N796A and E309Q variants.

Figure 5:

Cation concentration in the SERCA bind domains (assuming a filter volume = 0.33 nm^3 from our MD simulations), $N_{oxy} = 7$ (coordination number of Ca^{2+} ,³⁰ this value lies between the coordination number of site I, $N_{oxy} = 8.5$, and site II, $N_{oxy} = 6.7$, for Ca^{2+} determined in our simulation of WT SERCA) and with the cation solvation energy from.³³ $[\text{KCl}] = 150 \text{ mM}$ and $[\text{MgCl}_2] = 2 \text{ mM}$. As cytosolic Ca^{2+} is increased, Ca^{2+} displaces Mg^{2+} bound to SERCA and reaches saturation at millimolar Ca^{2+} concentrations. Data are also presented assuming a membrane potential of $\zeta = -25 \text{ mV}$, which locally increases bath cation concentrations by several fold according to Poisson-Boltzmann theory estimates.

Figure 6:

(a) Comparison of pump turnover rate between WT and SERCA variants using the Ca_{only} model with MSA predicted Ca^{2+} affinity for N796A and E309Q variants. The experimental data is from Inesi et al.⁴⁰ (b-c) Comparison of SERCA pump turnover rate of different state models against experimental data at $[\text{Mg}] = 2 \text{ mM}$ and $[\text{K}] = 150 \text{ mM}$. In panel (b) the experimentally measured $K_{d,K}$ and $K_{d,Mg}$ in Table S4 were used. In panel (c) the MSA predicted $K_{d,K}$ and $K_{d,Mg}$ were used. (d) Normalized difference between each state model and experimental data for panels (a-c), as evaluated by Eq. 1. Our state-based models reproduce steady-state WT SERCA pumping rates and predict impaired rates for the N796A and E309Q variants.

Figure 7:

(a-b) Comparison of [Ca] dependence of Ca^{2+} -SERCA binding of Mg₋K₊ and Mg₊K₋ models at varying Mg^{2+} concentrations ([K]= 150 mM) against experimental data from.17 Experimentally measured $K_{d,K}$ and $K_{d,Mg}$ were used. (c) Normalized difference between state models and experimental data as evaluated by Eq. 1. The Mg₋K₊ model, which assumes Mg^{2+} and K^{+} act as inhibitors and agonists, respectively, provide the best fit to experimental data.

Figure 8:

Correlation between MSA predicted chemical potential and experimentally measured binding free energy. The circles and triangles represent MSA results with waters and without waters taken into account, respectively. $RT = 0.593$ kcal/mol at room temperature and Kd_{exp} is experimentally measured dissociation constants in Table S4. The MSA chemical potentials correctly rank-order Ca^{2+} versus non-cognate ion binding.

Figure 9:

Comparison of Ca²⁺ binding sites in E1 state (a-b) and E2 state (c-d) from crystal structures of PDB 1su4 and 3b9b,⁶ respectively. Ca²⁺ and water molecules are shown as yellow and red spheres, respectively. The blue dashed lines in (a-b) outline the overall shape of Ca²⁺ coordinating spheres in E1 state while in (c-d) they depict assumed Ca²⁺ binding sites in E2 state. Panel (e) depicts MSA predicted Ca²⁺ chemical potential for these two sites. In E1 state the number of coordinating oxygens are seven and six for site I and II while these values are assumed as four for both sites in E2 state.