Published in final edited form as:

Org Lett. 2019 July 05; 21(13): 5143–5146. doi:10.1021/acs.orglett.9b01749.

Phosphine-Catalyzed α -Umpolung–Aldol Reaction for the Synthesis of Benzo[*b*]azapin-3-ones

Kui Zhang, Lingchao Cai, Sooji Hong, Ohyun Kwon*

Department of Chemistry and Biochemistry, University of California, Los Angeles, California 90095-1569, United States

Abstract

A novel phosphine-catalyzed intermolecular cyclization between 2-sulfonamidobenzaldehydes and ynones is reported. This methodology serves as a conduit for the construction of benzo[*b*]azepin-3-ones in good to excellent yields under mild conditions. The resulting 2-benzylidene moieties are formed exclusively in the *E*-configuration. Mechanistically, this unusual annulation occurs through a phosphine-catalyzed α -umpolung addition, followed by an aldol reaction. One of the benzo[*b*]azepin-3-one products was converted to the core structure of 3-amino-[*a*]benzazepin-2-one-1-alkanoic acids, many of which function as angiotensin-converting enzyme inhibitors.

Graphical Abstract

Benzoazepines are structural motifs within a wide variety of biologically and pharmacologically significant compounds.¹ In particular, a benzo[*b*]azepine moiety is a core unit of several marketed tricyclic antidepressant drugs, including tienopramine, anafranil, amezepine, and imipramine (Figure 1).² Lotensin, which features a benzoazepine nucleus as the key subunit, is used in the treatment of hypertension, congestive heart failure, and heart attacks, and also to prevent the renal and retinal complications of diabetes.³ Mozavaptan, which also possesses a benzoazepine skeleton, is an effective vasopressin V-2 receptor antagonist.⁴ Despite their promising biological properties, benzo[*b*]azepines have drawn

*Corresponding Author: ohyun@chem.ucla.edu.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.9b01749.

Experimental procedures and characterization data for all new compounds (PDF)

The authors declare no competing financial interest.

relatively little synthetic attention. Nevertheless, several methods have been described for the construction of benzo[*b*]azepines, including approaches through intramolecular Heck coupling,⁵ iridium-catalyzed asymmetric allylic amination,⁶ cyclizations of suitably substituted Morita–Baylis–Hillman adducts,⁷ FeCl₃- and Au-catalyzed cyclizations of alkynes and alkenes,⁸ Pd-catalyzed oxidative annulations of *ortho*-alkenylanilines and allenes,⁹ and Pd- and Au-catalyzed ring-expansions of alkylidenecyclopropanes.¹⁰ Despite the great potential of these methods, they require harsh reaction conditions and/or starting materials that are synthesized in several steps. In this regard, we became interested in developing new pathways for the synthesis of benzoazepines from readily available starting materials under benign reaction conditions.

Nucleophilic phosphine-catalyzed cyclizations have expanded chemical space by allowing access to a variety of carbo- and heterocycles.¹¹ While the use of electron-deficient alkenes, allenes, and alkynoates in those reactions has been explored widely,¹² the application of α,β -ynones in phosphine-catalyzed annulations has drawn scant attention until quite recently. The products of ynone annulations can be synthetically useful and biologically important compounds, including oxazolidines,^{13a} thiazolidines,^{13a} pyrrolidines,^{13a} dihydropyrrolopyridines,^{13b} benzimidazoline,^{13b} benzomorpholines,^{13b} tetrahydroquinolines, 3-oxanones, 3-oxepanones, spirooxazolines, spirooxindoles,^{13e} spirocyclopentanones,^{13f} spiro-cyclopentanone pyrazolones,^{13g} cyclopent[*b*]annulated heteroarenes,^{13h} hydropyridazines,¹³ⁱ cyclopentanone-fused benzosultams,^{13j} and spirobarbiturate-cyclopentanones.^{13k} In almost all of these examples, the α,β -ynone provides two- or three-carbon atoms for the formation of common five- or six-membered rings; to the best of our knowledge, there has been only one report of seven-membered ring formation.¹⁴

Considering these precedents and based on our previously described tandem γ -umpolung addition–Wittig olefination,¹⁵ we envisioned the possibility of synthesizing the dihydroquinoline **3'** through α -umpolung addition–Wittig alkenylation (Scheme 1). We found, however, that a solution of *o*-(*p*-toluenesulfonamido)benzaldehyde (**1a**) and the α,β -ynone **2a** in toluene produced, in the presence of 1 equiv of PPh₃, the benzo[*b*]azepin-3-one **3a** in 38% yield. We identified the *E*-configuration of this benzyldene moiety through NOESY analysis of the product **3a**.

As displayed in Scheme 2, we suspect that the mechanism begins with nucleophilic addition of PPh₃ to the ynone **2a** to generate the zwitterion **I**. The intermediate **I** subsequently deprotonates the sulfonamide **1a**, with α -umpolung addition of the resulting amide anion onto the vinylphosphonium species producing the ylide **II**. Proton transfer delivers the enolate **III**, which is ready for the aldol reaction with the aldehydic unit to construct the benzoazapin-3-one ring **IV**. Alkoxide-to-enolate proton transfer and subsequent β -elimination of the catalyst (PPh₃) furnishes the product **3a**. The exclusive formation of the *E*-benzyldene moiety is due, we believe, to steric repulsion between the *p*-toluenesulfonyl group and the phenyl ring when forming the alternative *Z*-isomer. Although there have been reported α -umpolung additions¹⁶ and tandem reactions involving α -umpolung addition,¹⁷ to the best of our knowledge there are no previous reports of phosphine-catalyzed α -umpolung-aldol processes. Herein, we describe our study of this cascade process for the

synthesis of benzo[*b*]azapin-3-ones through a phosphine-catalyzed α -umpolung-aldol reaction.

Recognizing that the α -umpolung-aldol reaction is catalytic in PPh₃, we tested various catalysis conditions for the α -umpolung-aldol reaction, engaging ethyl *o*-(*p*-toluenesulfonamido)benzaldehyde (**1a**) and the ynone **2a** as model substrates and employing various tertiary phosphines as catalysts at 20 mol%. We first screened various solvents in the presence of 20 mol % of PPh₃ at room temperature (Table 1). The product benzo[*b*]azepin-3-one **3a** was obtained in 10–42% yields in toluene, CH₂Cl₂, CHCl₃, and benzene (entries 1–4), but no product formed in tetrahydrofuran (THF), MeCN, or dioxane (entries 5–7). Noting that CHCl₃, with its acidic proton, was a particularly efficacious medium, we tested other protic solvents—alcohols, which could be deprotonated by the zwitterionic phosphonium enolate intermediates to promote the aldol reaction—for this reaction.¹⁸ Interestingly, the Michael addition product **3''** was generated in 83% yield when using MeOH as the solvent (entry 8).¹⁹ In contrast, EtOH and *n*-propanol enhanced the yield of the α -umpolung-aldol reaction to 58 and 69%, respectively (entries 9 and 10). *n*-Butanol could be used as the solvent without increasing the yield, but no significant product formed when running the reaction in isopropanol or *tert*-butanol (entries 11 and 13). Furthermore, we examined the effects of other phosphine catalysts (entries 14–17). Performing the reaction in the presence of tributylphosphine, a more nucleophilic catalyst, did not provide any product, but the yield increased to 74% when employing ethyldiphenylphosphine as the catalyst. Attempts to modulate the electronic properties of PPh₃ by using tris(*p*-fluorophenyl)-phosphine or tris(*p*-tolyl)phosphine had no significant benefits, supplying the benzo[*b*]azapin-3-one in yields of 69 and 54%, respectively.

Hereafter, employing ethyldiphenylphosphine as the catalyst in *n*-propanol as the solvent, our focus shifted to applying various other *o*-sulfonamidobenzaldehydes **1** and ynones **2** for the synthesis of a range of benzoazapinones **3** (Table 2). Initially, we studied the effects of varying the substituents on the benzene ring of the *o*-sulfonamidobenzaldehydes in reactions with the ynone **2a**. Electron-donating 5-methoxy and 5-methyl substituents on the benzene ring generated the desired products **3b** and **3c** in yields of 82 and 77%, respectively, while *N*-(3-formylnaphth-2-yl)-4-methylbenzenesulfonamide provided the benzo[*b*]azapin-3-one **3d** in 70% yield (entries 2–4). Electron-withdrawing groups, namely, 5-iodo, 5-bromo, and 5-chloro units, on the benzene ring induced the formation of the desired products **3e–g** in yields of 70–75% (entries 5–7). When we employed the 3-methyl-substituted 2-sulfonamidobenzaldehyde as the pronucleophile, no reaction occurred, presumably because of steric effects (entry 8). Performing the reaction with 4-methyl-2-sulfonamidobenzaldehyde delivered the benzoazapin-3-one **3h** in 72% yield (entry 9). Next, we probed the effects of various protecting groups of the aniline functionality. *N*-(2-Formylphenyl)benzenesulfonamide, *N*-(2-formylphenyl)-4-methoxybenzenesulfonamide, and *N*-(2-formylphenyl)-4-bromobenzenesulfonamide produced their desired products **3i–k**, respectively, in moderate yields (59–63%, entries 10–12).

We examine a range of ynones for the ethyldiphenylphosphine-catalyzed α -umpolung-aldol reaction. An electron-donating group on the benzene ring of the ynone **2** seemed crucial for a highly efficient reaction. When 4-(4-methoxyphenyl)but-3-yn-2-one was involved in the

annulation with **1a**, the desired product was obtained in 94% yield (entry 13). When run on 1 mmol scale, the reaction produced the product **3l** in 83% yield. The combination of electron-rich *o*-sulfonamidobenzaldehydes and 4-(4-methoxyphenyl)but-3-yn-2-one produced the desired benzo[*b*]azapin-3-ones **3m** and **3n** in the highest yields of this study: 95 and 99%, respectively (entries 14 and 15). Following this trend, 4-(4-methylphenyl)but-3-yn-2-one, when reacted with *N*-(2-formylphenyl)-4-methylbenzenesulfonamide and *N*-(2-formyl-4-methoxyphenyl)-4-methylbenzenesulfonamide, also provided high yields of the benzoazapin-3-ones **3o** and **3p** (91 and 93%, respectively; entries 16 and 17). Conducting the reactions of 4-(3-methylphenyl)but-3-yn-2-one with *N*-(2-formylphenyl)-4-methylbenzenesulfonamide and *N*-(2-formyl-4-methoxyphenyl)-4-methylbenzenesulfonamide gave the desired products **3q** and **3r** in slightly lower yields (85 and 81%, respectively; entries 18 and 19). When we employed 4-(2-methylphenyl)but-3-yn-2-one, however, no reaction occurred, presumably because the *o*-methyl group on the benzene ring of the alkynone blocked the nucleophilic addition of ethyldiphenylphosphine to the β -carbon atom of the α,β -ynone (entry 20). When an electron-withdrawing chlorine atom was positioned on the phenyl ring of the ynone, the corresponding product **3s** was furnished in 75% yield (entry 21). Interestingly, when 1,4-diphenylbut-3-yn-2-one was subjected to the standard reaction conditions, we obtained the 1,4-dihydroquinoline **4** instead of the benzo[*b*]azapin-3-one (eq 1).²⁰

(1)

Applying this α -umpolung–aldol technology, we realized the rapid synthesis of the 3-amino-benzo[*b*]azepin-2-one skeleton—the core unit of 3-amino-[*a*]benzazepin-2-one-1-alkanoic acids that are known angiotensin-converting enzyme (ACE) inhibitors.²¹ Starting from the benzo[*b*]azapin-3-one **3a**, diastereoselective reductive amination generated the *syn*-aminoalcohol **5**, whose benzyldene motif was readily converted to a C=O group through ozonolysis to obtain the 3-amino-[*a*]benzazepin-2-one **6** (Scheme 3).

In conclusion, we have developed a phosphine-catalyzed intermolecular cyclization of *o*-sulfonamidobenzaldehydes and ynones to yield highly functionalized (*E*)-benzo[*b*]azapin-3-ones under mild reaction conditions at room temperature. This α -umpolung–aldol process provides a range of benzo[*b*]azapin-3-ones **3** in synthetically useful yields and with exclusive *E*-selectivity. In the further functionalization of **3a**, we found that compound **6** could be constructed rapidly through reductive amination and ozonolysis, providing an alternative means toward the skeleton of 3-amino-[*a*]benzazepin-2-one-1-alkanoic acids that are useful ACE inhibitors.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

We thank the NIH (GM071779) for financial support.

REFERENCES

- (1). (a) Albright JD; Reich MF; Delos Santos EG; Dusza JP; Sum FW; Venkatesan AM; Coupet J; Chan PS; Ru X; Mazandarani H; Bailey TJ Med. Chem 2000, 43, 4388(b) Le Diguarher T; Ortuno JC; Shanks D; Guilbaud N; Pierre A; Raimbaud E; Fauchere JL; Hickman JA; Tucker GC; Casara P Bioorg. Med. Chem. Lett 2004, 14, 767 [PubMed: 14741286] (c) Failli AA; Shumsky JS; Steffan RJ; Caggiano TJ; Willoams DK; Trybulski EJ; Ning X; Lock Y; Tanikella T; Hartmann D; Chan PS; Park CH Bioorg. Med. Chem. Lett 2006, 16, 954. [PubMed: 16297621]
- (2). Singh H; Gupta N; Kumar P; Dubey SK; Sharma PK Org. Process Res. Dev 2009, 13, 870.
- (3). (a) Hou F; Zhang X; Zhang G; Xie D; Chen P; Zhang W; Jiang J; Liang M; Wang G; Liu Z; Geng RN N. Engl. J. Med 2006, 354, 131 [PubMed: 16407508] (b) O'Grady MR; O'Sullivan ML; Minors SL; Horne RJ Vet. Intern. Med 2009, 23, 977.
- (4). (a) Decaux G; Soupart A; Vassart G Lancet 2008, 371, 1624 [PubMed: 18468546] (b) Robertson GL Nat. Rev. Endocrinol 2011, 7, 151 [PubMed: 21283124] (c) Ghose AK; Herbertz T; Hudkins RL; Dorsey BD; Mallamo JP ACS Chem. Neurosci 2012, 3, 50. [PubMed: 22267984]
- (5). Qadir M; Cobb J; Sheldrake WP; Whittall N; White AJP; Hill KK; Horton PN; Hursthouse MB J. Org. Chem 2005, 70, 1545. [PubMed: 15730272]
- (6). (a) He H; Liu W; Dai L; You S Angew. Chem. Int. Ed 2010, 49, 1496(b) Ye K; Dai L; You S Org. Biomol Chem 2012, 10, 5932. [PubMed: 22441972]
- (7). (a) Singh V; Batra S Eur. J. Org. Chem. 2007, 2007, 2970(b) Zhan G; Shi M; He Q; Du W; Chen Y Org. Lett 2015, 17, 4750. [PubMed: 26359687]
- (8). (a) Cui L; Zhang G; Peng Y; Zhang L Org. Lett 2009, 11, 1225 [PubMed: 19220014] (b) Jalal S; Bera K; Sarkar S; Paul K; Jana U Org. Biomol. Chem 2014, 12, 1759 [PubMed: 24500306] (c) Undeela S; Ravikumar G; Nanubolu JB; Singarapu KK; Menon RS Chem. Commun 2016, 52, 4824.
- (9). (a) Wu L; Meng Y; Ferguson J; Wang L; Zeng FJ Org. Chem 2017, 82, 4121(b) Dhandabani GK; Mutra MR; Wang J Adv. Synth. Catal 2018, 360, 4754.
- (10). (a) Yu L; Zhu Z; Hu X; Tang X; Shi M Chem. Commun 2016, 52, 6581(b) Yu L; Xu Q; Tang X; Shi M ACS Catal. 2016, 6, 526.
- (11). For selective examples, see:(a) Andrews IP; Kwon O Chem. Sci 2012, 3, 2510 [PubMed: 22798981] (b) Cai L; Zhang K; Kwon O J. Am. Chem. Soc 2016, 138, 3298 [PubMed: 26910382] (c) Li J; Zhang W; Zhang F; Chen Y; Li A J. Am. Chem. Soc 2017, 139, 14893 [PubMed: 28956924] (d) Chen X; Zhang H; Yang X; Lv H; Shao X; Tao C; Wang H; Cheng B; Li Y; Guo J; Zhang J; Zhai H Angew. Chem. Int. Ed 2018, 57, 947.
- (12). For selected reviews, see:(a) Lu X; Zhang C; Xu Z Acc. Chem. Res 2001, 34, 535 [PubMed: 11456471] (b) Valentine DH Jr.; Hillhouse JH Synthesis 2003, 2003, 317(c) Methot JL; Roush WR Adv. Synth. Catal 2004, 346, 1035(d) Ye L-W; Zhou J; Tang Y Chem. Soc. Rev 2008, 37, 1140 [PubMed: 18497927] (e) Cowen BJ; Miller SJ Chem. Soc. Rev 2009, 38, 3102 [PubMed: 19847345] (f) Lopez F; Mascarenas JL Chem. - Eur. J 2011, 17, 418 [PubMed: 21207554] (g) Zhao Q-Y; Lian Z; Wei Y; Shi M Chem. Commun 2012, 48, 1724(h) Fan YC; Kwon O Chem. Commun 2013, 49, 11588(i) Xu S; He Z RSC Adv. 2013, 3, 16885(j) Wang Z; Xu X; Kwon O Chem. Soc. Rev 2014, 43, 2927 [PubMed: 24663290] (k) Li W; Zhang J Chem. Soc. Rev 2016, 45, 1657 [PubMed: 26776280] (l) Wei Y; Shi M Org. Chem. Front 2017, 4, 1876(m) Li H; Lu Y Asian J. Org. Chem 2017, 6, 1130(n) Ni H; Chan W; Lu Y Chem. Rev 2018, 118, 9344 [PubMed: 30204423] (o) Guo H; Fan YC; Sun Z; Wu Y; Kwon O Chem. Rev 2018, 118, 10049. [PubMed: 30260217]
- (13). (a) Sriramurthy V; Kwon O J. Am. Chem. Soc 2007, 129, 12928 [PubMed: 17924625] (b) Sriramurthy V; Kwon O Org. Lett 2010, 12, 1084 [PubMed: 20143856] (c) Saha J; Lorenc C; Surana B; Peczu MW J. Org. Chem 2012, 77, 3846 [PubMed: 22428530] (d) Yang L; Xie P; Li E; Li X; Huang Y; Chen R Org. Biomol. Chem 2012, 10, 7628 [PubMed: 22903685] (e)

Ramachary DB; Venkaiah C; Krishna PM *Org. Lett* 2013, 15, 4714 [PubMed: 24015947] (f) Liang L; Li E; Xie P; Huang Y *Chem. -Asian J* 2014, 9, 1270 [PubMed: 24678009] (g) Li J; Du D *Adv. Synth. Catal* 2015, 357, 3986(h) Raghu M; Grover J; Ramasastry SSV *Chem. -Eur. J* 2016, 22, 18316 [PubMed: 27731920] (i) Liang L; Huang Y *Org. Lett* 2016, 18, 2604 [PubMed: 27218675] (j) Ramachary DB; Krishna PM; Reddy TP *Org. Biomol. Chem* 2016, 14, 6413 [PubMed: 27305594] (k) Gao X; Li Z; Yang W; Liu Y; Chen W; Zhang C; Zheng L; Guo H *Org. Biomol. Chem* 2017, 15, 5298. [PubMed: 28598480]

- (14). Sun L; Wei Y; Shi M *Adv. Synth. Catal* 2017, 359, 3176.
- (15). Zhang K; Cai L; Yang Z; Houk KN; Kwon O *Chem. Sci* 2018, 9, 1867. [PubMed: 29732112]
- (16). (a) Trost BM; Dake GR *J. Am. Chem. Soc* 1997, 119, 7595(b) Zheng F; Leung T-F; Chan K-W; Sung HHY; Williams ID; Xie Z; Jia G *Chem. Commun* 2016, 52, 10767.
- (17). (a) Liu B; Davis R; Joshi B; Reynolds DW *J. Org. Chem* 2002, 67, 4595 [PubMed: 12076163] (b) Lu C; Lu X *Org. Lett* 2002, 4, 4677 [PubMed: 12489959] (c) Kuroda H; Tomita I; Endo T *Org. Lett* 2003, 5, 129 [PubMed: 12529122] (d) Yavari I; Moradi L *Tetrahedron Lett.* 2006, 47, 1627(e) Yavari I; Soury S; Sirouspour M; Djahaniani H *Synthesis* 2006, 2006, 3243(f) Gabillet S; Lecercle D; Loreau O; Carboni M; Dezard S; Gomis J-M; Taran F *Synthesis* 2007, 2007, 515(g) Gabillet S; Lecercle D; Loreau O; Carboni M; Dezard S; Gomis J-M; Taran F *Org. Lett* 2007, 9, 3925 [PubMed: 17803312] (h) Alizadeh A; Sheikhi E *Tetrahedron Lett.* 2007, 48, 4887(i) Carboni M; Gomis J-M; Loreau O; Taran F *Synthesis* 2008, 2008, 417(j) Wilson JE; Sun J; Fu GC *Angew. Chem., Int. Ed* 2010, 49, 161(k) Nasiri F; Bayzidi M; Zolali A *Mol. Diversity* 2012, 16, 619(l) Lian Z; Shi M *Eur. J. Org. Chem* 2012, 2012, 581.(m) Anary-Abbasinejad M; Farashah HD; Hassanabadi A; Anaraki-Ardakani H; Shams N *Synth. Commun* 2012, 42, 1877.(n) Zhou Q-F; Chu X-P; Ge F-F; Li C; Lu T *Mol. Diversity* 2013, 16, 563(o) Gabillet S; Loreau O; Specklin S; Rasalofonjatovo E; Taran F *J. Org. Chem* 2014, 79, 9894 [PubMed: 25238600] (p) Li J; Du D *Adv. Synth. Catal* 2015, 357, 3986(q) Zhang J; Zhang M; Li Y; Liu S; Miao Z *RSC Adv.* 2016, 6, 107984(r) Ametovski J; Dutta U; Burchill L; Maiti D; Lupton DW; Hooper JF *Chem. Commun* 2017, 53, 13071.
- (18). Thalji RK; Roush WR *J. Am. Chem. Soc* 2005, 127, 16778. [PubMed: 16316211]
- (19). (a) Inanaga J; Baba Y; Hanamoto T *Chem. Lett* 1993, 22, 241(b) Stoddard RL; Luo J; van der Wal N; O'Rourke NF; Wulff JD; McIndoe JS *New J. Chem* 2014, 38, 5382(c) Wende M; Meier R; Gladysz JA *J. Am. Chem. Soc* 2001, 123, 11490. [PubMed: 11707132]
- (20). (a) Khong SN; Kwon O *J. Org. Chem* 2012, 77, 8257 [PubMed: 22928667] (b) Khong SN; Kwon O *Asian J. Org. Chem* 2014, 3, 453. [PubMed: 26207200]
- (21). Watthey JWH 3-Amino-[1]-benzazepin-3-one-1-alkanoic acids. US 4575503 A, 1986.

Figure 1.
Representative marketed drugs containing benzo[*b*]azepine core structures.

Scheme 1.
Discovery of α -Umpolung-Aldol Reaction

Scheme 2.
Proposed Mechanism for the Formation of the Benzo[*b*]azepin-3-one 3a

Scheme 3.
Synthetic Application of the α -Umpolung–Aldol Reaction

Table 1.

optimization of the Reaction Conditions^a

entry	phosphine	solvent	temp. (°C)	yield (%) ^b
1	Ph ₃ P	toluene	rt	36
2	Ph ₃ P	CH ₂ Cl ₂	rt	10
3	Ph ₃ P	CHCl ₃	rt	42
4	Ph ₃ P	benzene	rt	14
5	Ph ₃ P	THF	rt	NR
6	Ph ₃ P	MeCN	rt	NR
7	Ph ₃ P	dioxane	rt	NR
8	Ph ₃ P	MeOH	rt	3'' (83%) ^c
9	Ph ₃ P	EtOH	rt	58
10	Ph ₃ P	<i>n</i> -propanol	rt	69
11	Ph ₃ P	isopropanol	rt	NR
12	Ph ₃ P	<i>n</i> -butanol	rt	40
13	Ph ₃ P	<i>tert</i> -butanol	rt	NR
14	<i>n</i> -Bu ₃ P	<i>n</i> -propanol	rt	NR
15	EtPh ₂ P	<i>n</i> -propanol	rt	74
16	(<i>p</i> -FC ₆ H ₄) ₃ P	<i>n</i> -propanol	rt	69
17	(<i>p</i> -MeC ₆ H ₄) ₃ P	<i>n</i> -propanol	rt	54

^aReaction conditions: **1a** (0.2 mmol), **2a** (0.3 mmol), phosphine (0.04 mmol), and solvent (2 mL) at room temperature.^bIsolated yield of **3a**.^c**3''** = 4-methoxy-4-phenyl-3-buten-2-one.

Table 2.

Synthesis of Benzo[*b*]azapin-3-ones^a

entry	R ¹	R ²	R ³	yield (%) ^b
1	H	Ts	H	3a , 74
2	5-MeO	Ts	H	3b , 82
3	5-Me	Ts	H	3c , 77
4	1d ^c	Ts	H	3d , 70
5	5-I	Ts	H	3e , 75
6	5-Br	Ts	H	3f , 73
7	5-Cl	Ts	H	3g , 70
8	3-Me	Ts	H	NR
9	4-Me	Ts	H	3h , 72
10	H	PhSO ₂	H	3i , 63
11	H	(4-MeOC ₆ H ₄)SO ₂	H	3j , 65
12	H	(4-BrC ₆ H ₄)SO ₂	H	3k , 59
13	H	Ts	4-MeO	3l , 94 (83) ^d
14	5-MeO	Ts	4-MeO	3m , 99
15	5-Me	Ts	4-MeO	3n , 95
16	H	Ts	4-Me	3o , 91
17	5-MeO	Ts	4-Me	3p , 93
18	H	Ts	3-Me	3q , 85
19	5-MeO	Ts	3-Me	3r , 81
20	H	Ts	2-Me	NR
21	H	Ts	4-Cl	3s , 75

^aReaction conditions: **1** (0.2 mmol), **2** (0.3 mmol), phosphine (0.04 mmol), and *n*-propanol (2 mL) at room temperature.^bIsolated yields.^c**1d** = *N*-(3-formylnaphth-2-yl)-4-methylbenzenesulfonamide.^dReaction run on 1 mmol scale.