Published in final edited form as:

J Parkinsons Dis. 2020 ; 10(2): 693–705. doi:10.3233/JPD-191793.

Well Water and Parkinson Disease in Medicare Beneficiaries: A Nationwide Case-Control Study

Maya R. Silver^a, Brad A. Racette^{a,b}, Umber Dube^a, Irene M. Faust^a, Susan Searles Nielsen^a

^aDepartment of Neurology, Washington University School of Medicine, St. Louis, Missouri, USA

^bSchool of Public Health, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

Abstract

Background—Well water frequently is considered a risk factor for Parkinson disease (PD), but few studies were designed appropriately to test whether geographic factors affect PD risk.

Objective—To determine the risk of PD in relation to residential use of private well water.

Methods—In a nationwide, population-based case-control study, we identified all incident PD cases (N=89,790) and all comparable controls (N=21,549,400) age 66–90 who solely relied on Medicare coverage in the U.S. in 2009. We estimated the probability of use of private well water using zip code of residence at diagnosis/reference and U.S. Census data on household water source. We modeled this exposure linearly in logistic regression to calculate the odds ratio (OR) and 95% confidence interval (CI) of PD risk in relation to well water use. We adjusted for age, sex and race/ethnicity, and verified that smoking and use of medical care did not confound results. We repeated analyses with a 2-year exposure lag and separately within each U.S. state.

Results—Use of well water was inversely associated with PD risk (OR=0.87, 95% CI 0.85–0.89). We confirmed this association in a Cox survival analysis in which we followed controls for 5 years, death or PD diagnosis. There was little evidence that well water use increased risk of PD in any individual state.

Conclusions—Although it remains possible that exposures in well water in more narrow geographic regions increase PD risk, in general these results suggest that exposures more common in urban/suburban areas might also be relevant.

Keywords

Case-Control Studies; Drinking Water; Epidemiology; Geographic Factors; Ground Water; Medicare; Parkinson Disease

Address correspondence to Susan Searles Nielsen, PhD, Washington University School of Medicine, Department of Neurology, Campus Box 8111, 660 South Euclid Avenue St. Louis, MO 63110 USA. Telephone: (314) 362-5291. Fax: (314) 747-3274. sn Nielsen@wustl.edu.

Conflict of Interest: The authors have no conflicts of interest to report.

Supplementary Material: Supplementary Data include tables of characteristics associated with use of well water, PD-well water ORs by decile of exposure and state-specific PD-well water ORs and use of well water).

Introduction

Although numerous genes contribute to PD risk [1–3], studies in twins indicate that environmental risk factors play a prominent role [4–6]. As such, identification of environmental risk factors for PD is an essential step in reducing disease burden through primary prevention.

One potential risk factor that has received considerable attention is residential use of private well water as the primary source of water for drinking, cooking and bathing. Several epidemiologic studies have concluded that the use of well water might increase the risk of PD [7–24], presumably due to water contaminants. However, other studies observed an inverse association [25] or no association [26–41]. While geographic differences in the quality of well water in different regions makes comparison across studies difficult, the inconsistency in results is also likely due to differences in study design. Most of the studies either were very small, or were not designed to test the hypothesis that PD risk is associated with a geographically varying exposure and therefore may have been biased. In particular, most well water studies were clinic-based and included prevalent PD cases. Only one study, conducted in the Central Valley of California, was adequately sized, population-based, restricted to incident PD cases and avoided geographic matching of controls to cases [19]. In that study, PD risk was increased in relation to well water potentially contaminated with the insecticide diazinon. Nonetheless, there was no clear association between PD and ever using well water in general during one's lifetime (odds ratio [OR]=1.2, 95% confidence interval [CI] 0.82–1.80).

In order to address the limitations of most other prior studies of PD and well water, we utilized United States (U.S.) Medicare data to conduct a large, population-based case-control study with incident PD cases and controls (comparable to the cases yet selected without geographic matching). Based on the results from the study in California [19], we hypothesized that any positive association between well water and PD would be modest and would vary depending on potential well water contaminants and hence geographic location.

Materials and Methods

Protection of Human Subjects/Data

This study was approved by the Centers for Medicare and Medicaid Services (CMS) and Washington University's Human Research Protection Office. CMS de-identified all data prior to release to Washington University. Informed consent was not required because this was a records-based study only using de-identified data.

Overall Study Design

We conducted a population-based case-control study using U.S. Medicare data. We used comprehensive administrative claims data and beneficiary annual summary (base) files from 2004–2009 to identify all cases and controls [42]. In summary, in the year 2009 all cases and controls lived in the U.S., were 66–90 years of age and had Medicare Parts A and/or B, but had no health maintenance organization coverage (the main exclusion in age-eligible beneficiaries).

Case Ascertainment

We included as cases (N=89,790) all Medicare beneficiaries in 2009 who met all above criteria and who had 1 International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9) diagnosis code 332 or 332.0 in any file (inpatient/skilled nursing facility, outpatient, carrier, home health care, or durable medical equipment) in 2009 but not in 2004–2008. Cases did not have a code potentially indicative of an atypical parkinsonism or Lewy body dementia.

Control Selection

We included as controls (N=21,549,400) all Medicare beneficiaries in 2009 who met all of the same criteria as cases, but who had no ICD-9 code for PD and who had survived to their randomly assigned reference date in 2009. We included all eligible controls to maximize statistical power. We avoided matching controls to cases to ensure that we did not overmatch with regard to geography. Instead, we addressed potential confounding analytically, and we verified that we had excellent statistical power. Matching is most important when gains in statistical efficiency are necessary [43], but even without matching we had >99.9% power to detect an OR as modest as 1.09.

Determination of Type of Water Source

Medicare base files contain each beneficiary's residential zip code for the respective benefit year. We assigned each beneficiary's residential zip code as of the diagnosis/reference year (2009) to the 2010 ZIP Code Tabulation Area (ZCTA) using a lookup table [44]. We then assigned each beneficiary to the latitude and longitude coordinate of the population centroid of the ZCTA weighted for households with residents age 65 using ArcMap [45]. Next, we spatially joined each coordinate to the corresponding 1990 U.S. Census block group [46], which is the most recent year for which the U.S. Census collected data on household water source [47]. From this Census data, we estimated the percent of households in each block group that used private well water by dividing the number of households served by individual drilled or dug wells by the total number of households. In most areas the remaining households use water from a "public system or private company" that originates from deeper wells, or surface water sources, monitored for compliance with federal drinking water standards [48]. We thus assigned the percentage of households that use water from individual drilled or dug wells (hereafter "wells") to each beneficiary, such that exposure was expressed probabilistically, ranging from 0% to 100%. We were able to estimate exposure for all but 85 (0.09%) cases and 22,375 (0.10%) controls.

We validated our exposure assessment approach by applying a similar approach to estimate the probability of well water for 107 residences sampled from Washington State through random digit dialing [49]. Residents reported their source of drinking water and their mailing and street addresses; 18 (17%) used an individual or shared private well and the remainder used water from a publicly monitored source. Even when we applied our exposure assessment approach using only the 5-digit mailing zip code, we observed good agreement between probability of well water use and our gold standard measure, self-reported water source (biserial correlation coefficient=0.45, $p=0.0001$). As a secondary validation we compared our probability of well water use variable, which was based on 1990

Census data, to a variable that was assessed at the time of PD diagnosis/reference, a county level variable from 2010 from the U.S. Geological Survey [50]. Specifically, we calculated the percent of the population in the county whose “domestic” water was “self-supplied” in that year. This county level variable also agreed well with our primary (zip code level) variable despite the different geographical scale and 20-year time gap (Spearman’s $\rho=0.49$).

Determination of Covariates

We obtained beneficiary date of birth (age), sex and race/ethnicity from the Medicare base file for 2009. Medicare base files also include individual level summaries of use of medical care, including the number of physician visits (visits to physician offices as well as outpatient institutional visits). We obtained the number of physician visits from the 2008 base file so that we did not count post-diagnosis visits. We also had complete Medicare claims data from 2004–2009 for all cases and all 118,095 controls in a 0.5% random sample of the full control group. In these cases and controls, we calculated the total count of unique diagnosis codes prior to PD diagnosis/reference (as another indicator of use of medical care) and three other covariates previously associated with PD. These included ever/never tobacco smoking, [51, 52] physical activity [53–55] and constipation [56]. We determined constipation from ICD-9 codes 564.0x. For physical activity, we were interested in assessing lifetime physical activity [53, 54]. In addition, individuals in rural areas might have more opportunity for some types of outdoor physical activity, including from an outdoor occupation, which itself has been associated with a reduced risk of PD [57, 58], presumably due to greater physical activity and/or exposure to ultraviolet light (UV). Therefore, we used UV-related cancers as a proxy for lifetime outdoor physical activity including from an outdoor occupation. We classified the following cancers as UV-related: lip cancer (ICD-9 140.xx); squamous, basal cell or melanoma skin cancer (ICD-9 172.xx, 173.xx); ocular melanoma/conjunctival squamous cell carcinoma (ICD-9 190.3, 190.5, 190.6, 190.8, 190.9) [59].

We estimated the probability (0% to 100%) of “ever smoking” tobacco using a multifaceted approach [42, 60]. We assigned a probability of 100% to any beneficiary with a diagnosis or procedure code specific to tobacco use, primarily ICD-9 305.1 (tobacco use disorder) or ICD-9 V15.82 (history of tobacco use). Otherwise, we estimated the probability from a logistic regression model that we created based on sex, race/ethnicity, birth cohort and >600 other diagnosis or procedure codes. We have detailed the development and validation of this predictive model previously [42].

We used publicly available data to create variables for selected environmental exposures likely to be associated with use of well water. Overall, well water use is more common in areas with intensive use of agricultural pesticides than in urban areas without any well water use. Given the suspected association between residential exposure to agricultural pesticides and PD [61–63], we obtained county level data on kilograms of agricultural pesticides applied in each county of the U.S. in 2009 [64]. Specifically, we calculated the midpoint between the lower and upper estimates for each pesticide and then summed across all pesticides. Then we divided this total by the land area (acres) in the county to standardize agricultural pesticide use in terms of intensity of pesticide use. We also obtained the

prevalence of daily smoking in the county in 2009 [65]. We used this as a proxy for exposure to environmental tobacco smoke (ETS), since some studies suggest that ETS, not just active smoking, is associated with a reduced risk of PD [66, 67].

Statistical Analysis

We performed all statistical analysis using Stata [68]. We used logistic regression to estimate the OR and 95% CIs for well water use and PD. Probability of well water use ranged from 0% to 100%. We retained this as a continuous variable that ranged from 0 to 1, so that a one unit change (i.e. the OR) can be interpreted as the risk of PD among beneficiaries with vs. without well water (100% vs. 0% probability of well water). This also allowed the substantial number of beneficiaries (51.0% of cases, 52.4% of controls) with intermediate probabilities of well water use to contribute to risk estimation.

In all models we adjusted *a priori* for age (as 2 linear splines with a knot at age 85), sex and race/ethnicity (in seven categories unless noted). We then examined the effect of adjustment for the following variables: geographic region (state) of residence, agricultural pesticide use (kg/acre) and ETS. We also examined the effect of adjustment for several individual level variables, including use of medical care. We adjusted for use of medical care because we previously found that greater use of care is associated with a greater PD case ascertainment in the present study and can bias associations [60]. In the present analysis, our concern was that use of medical care could be a nuisance mediator. In this context, a nuisance mediator should be handled as a confounder to factor out any effect on the PD-well water association if beneficiaries in more rural areas tend to have fewer encounters with physicians or receive fewer ICD-9 codes on average than their counterparts from more urban areas (Figure 1, Supplementary Table 1). Our primary indicator of medical care was the total number of physician visits in 2008 for each individual. This was our primary indicator because it was available in the full case-control sample. Among beneficiaries for whom we had complete Medicare claims data, we alternatively considered the total number of unique diagnosis codes up to the PD diagnosis/reference date. Among this same subset of beneficiaries we examined the effect of adjusting for smoking and UV-related cancer, as well, with the latter as a proxy for outdoor occupations and outdoor physical activity in general. Finally, we checked whether constipation modified the association between use of well water and PD, given that constipation long precedes PD diagnosis and could affect drinking water habits. Except where noted, we modeled continuous variables linearly, and coded all percentage variables on the 0 to 1 scale, so that the OR could be interpreted easily as described above.

We conducted four sensitivity analyses. First, since it is possible that residence could change during the prodromal period of PD, due to PD symptoms, we lagged exposure by 2 years. Specifically, we used the probability of well water according to the beneficiary's zip code in 2007, the first year that residential zip code was usually recorded at the 9-digit (vs. 5-digit) zip code level. Second, we restricted to those cases and controls who had the same 5-digit zip code in 2004 as in 2009. Third, we restricted to beneficiaries with either 0% or 100% probability of well water use. This allows for a traditional analysis with a simple 0/1 dichotomous exposure variable, and also restricts to the cases and controls whose exposure

in 2009 was known with the greatest certainty. Fourth, we restricted our analysis to PD cases who received a PD diagnosis code from a neurologist or who had 3 PD diagnosis codes.

We considered a 2-sided p-value of <0.05 significant in the nationwide analyses, as evidenced by 95% CIs that excluded the value of one. We repeated all statistical analyses for each of the 50 U.S. states individually. For this state-by-state analysis, we applied a Bonferroni correction to account for multiple comparisons (2-sided $\alpha=0.001$).

Data Sharing

Per our agreement with CMS, we are not allowed to share individual beneficiaries' data.

Results

Characteristics of Cases and Controls

The demographic characteristics of cases and controls were consistent with known risk factors for PD (Table 1). Half of cases were male, compared to 43% of controls. Proportionally more white and Hispanic beneficiaries had PD than other beneficiaries. On average, cases were older than controls; the mean age was 78.8 years (standard deviation [SD] 6.1) for cases and 76.0 years (SD 6.2) for controls. Among beneficiaries with claims data, PD was inversely associated with smoking (OR=0.70, 95% CI 0.67–0.72) after accounting for the above demographic differences and greater use of medical care among cases than controls. After accounting for the same variables, as well as (active) smoking and state, PD also appeared to be inversely associated with ETS (OR=0.87, 95% CI 0.63–1.19) and UV-related cancer (OR=0.86, 95% CI 0.84–0.87), although the former was non-significant.

Well Water and PD Risk

A total of 10,579 (11.8%) of cases and 2,824,319 (13.1%) of controls had a $>50\%$ probability of using well water. In the nationwide sample we observed an overall inverse association between use of well water and PD (OR=0.87, 95% CI 0.85–0.89) (Table 2, Supplementary Table 2). This OR attenuated by $<5\%$ with adjustment for any of the individual potential confounders that we considered, with the greatest effect through adjustment for total number of unique diagnosis codes. In the subset of cases and controls for whom we could calculate this indicator of use of medical care, further adjustment for all potential confounders resulted in a modestly attenuated OR that remained significant (OR=0.91, 95% CI 0.88–0.95). The inverse association between well water and PD also attenuated slightly but clearly remained in each of the sensitivity analyses. Risk of PD in relation to use of well water was 0.90 (95% CI 0.88–0.92) when applying a 2-year lag; 0.88 (95% CI 0.85–0.90) when restricting to beneficiaries who had the same 5-digit zip code in 2004 as in 2009; 0.89 (95% CI 0.84–0.95) when only retaining beneficiaries with 0% or 100% probability of using well water; and 0.89 (95% CI 0.83–0.94) when restricting to the most certain PD cases. The PD-well water association was inverse among beneficiaries with or without constipation (respectively, OR=0.84, 95% CI 0.79–0.90; OR=0.88, 95% CI 0.85–0.92).

Because the inverse association contrasted with a majority of previous studies, we conducted a *post hoc* Cox proportional hazards survival analysis to attempt to replicate our results using a different analytical approach. Using complete Medicare claims data and base files for 2010–2014, we followed all controls for whom we had claims data and who survived through 2009 (N=115,492) for up to 5 years, from January 1, 2010 until death, PD diagnosis (N=2,326) or December 31, 2014. Again we observed an inverse association between use of well water and PD (hazard ratio=0.83, 95% CI 0.72–0.97). Similarly, in another *post hoc* analyses given our unexpected findings, we examined the association between PD and the probability of well water use as assessed at the county level in 2010, i.e. at the time of diagnosis/reference. Here again we observed a significant inverse association despite the broader geographic scale when estimating exposure (OR=0.94, 95% CI 0.89–0.99, adjusted for age, sex, race/ethnicity, smoking and number of diagnosis codes). Finally, we explored whether the inverse association between well water use and PD could occur due to greater recognition or coding of PD in the most urban areas. We recategorized controls as PD cases if they had each of the following: tremor (ICD-9 333.1 or 781.0), gait/coordination abnormality (ICD-9 781.2 or 781.3) and at least one additional condition associated with PD (constipation [ICD-9 564.0×], REM sleep behavior disorder [ICD-9 327.42] or smell/taste disturbance [ICD-9 781.1]) The association between our primary well water variable and PD remained inverse (OR=0.92, 95% CI 0.90–0.95) when we treated the 794 (0.7%) of controls from the control group who met these criteria as PD cases who each represented 200 cases nationwide. We applied this weight (in logistic regression) because our controls were a 0.5% random sample of all eligible controls.

Well Water and PD Risk by State

Use of well water was not associated with greater risk of PD in any individual state after Bonferroni correction (Figure 2). In addition, all OR point estimates >1 were relatively modest. ORs were >1.10 in only four states: Alaska, Connecticut, Massachusetts and Tennessee. The association between well water use and PD in Tennessee represented the largest positive association (1.33, 95% CI 1.07–1.65, $p=0.01$). This association weakened (OR=1.24, 95% CI 0.99–1.54) when we excluded Memphis, which is both populous and served by ground water rather than surface water. However, the risk of PD in relation to well water use in Alaska was very similar in models with or without Fairbanks, which shares these relatively unique features with Memphis.

Five states had a significant inverse association between well water use and PD after Bonferroni correction: California, Florida, Maine, Michigan and New York (Figure 2). These were among the most populous states (Figure 2) or among those with above average use of well water (Supplementary Table 3).

Discussion

In this large, population-based study of well water and PD risk across the entire U.S, we observed a modest inverse association between use of private well water and PD, in contrast with several previous studies [7–24]. The large number and geographic distribution of residences allowed for a statistically powerful nationwide analysis and an ability to explore

results in disparate regions. We did not observe an increased risk of PD nationwide or, after correction for multiple comparisons, in any individual state. There are several reasons our results might differ from previous studies of the association between well water and PD risk. Of particular note, most prior studies have been clinic-based interview studies, and it is very plausible that PD cases were more willing than controls to travel farther to reach the clinic in order to receive care or participate in the study. In general, this would bias the PD-well water OR upward, and thus potentially even create a spurious positive association when the true association is inverse. Our study was records-based and therefore not subject to this bias. In addition, we identified incident cases and highly comparable controls within a well-defined population. Accordingly, our PD study demonstrated all known demographic associations, as well as the known associations for smoking, constipation and REM sleep behavior disorder [42]. These expected results provide evidence of the strength of our design and the generalizability of the results. Of note, a meta-analysis of PD and well water use did observe an OR of 1.0 when restricting to the highest quality studies [69]. The present work further calls into question the generally accepted notion that the use of well water increases PD risk.

In a state-by-state analysis, the states with a significant inverse association between use of well water and PD either had either a particularly large number of Medicare beneficiaries or above average use of well water, and therefore relatively good statistical power. While no states demonstrated a strong positive association between well water and PD, our analysis does not rule out such an association in smaller geographical areas. Specifically, our study was not designed to test for highly localized geographic effects of specific contaminants. For example, one well-designed case-control study determined that exposure to PD risk was increased in relation to well water potentially contaminated with diazinon in an agricultural region of California [19]. We did not consider geographic regions smaller than states, which is a limitation since many states are diverse in terms of land use. We observed an inverse PD-well water association for California overall, despite some agricultural regions within California potentially having an increased risk of PD among well water users. Similarly, we did not systematically explore whether any particular environmental exposure might affect PD risk. Selected pesticides and solvents are among the most common contaminants in private wells in the U.S. [70], and a recent geographic study in France confirmed that both groups of chemicals might include agents that increase PD risk [71]. However, in private wells in the U.S. pesticides and solvents are typically present at levels below federal drinking water standards [70] and “well water” is likely a poor indicator of exposure to these classes of chemicals from drinking water. In addition, a complete examination would require consideration of multiple sources of exposure, not just water.

Overall, the inverse association we observed between the use of well water and PD nationwide was robust. We also found little evidence for confounding by a variety of potential confounders, or for reverse causation, whereby individuals with prodromal PD might move to urban areas as they begin to manifest symptoms. However, we cannot fully discount either of these possibilities, or other effects of prodromal PD, given the limited ability to apply exposure lagging or to account for potential confounders using individual level data. At the same time, we cannot rule out the possibility that PD risk is increased by environmental exposures that are more prevalent in regions served by public water. For example, disinfection byproducts are more common in public tap water than in water from

individual private wells, although there is currently no evidence that exposure to these chemicals is associated with a greater risk of PD. In addition, air quality is poorer in urban areas than in rural areas, particularly with respect to fine particulate matter (PM_{2.5}) [72]. Air pollution, particularly from traffic, including PM_{2.5}, was associated with increased risk of PD in a recent meta-analysis [73]. Such an association is biologically plausible because urban air pollution has been associated with neuroinflammation and cerebrospinal fluid oligomeric α -synuclein in humans [74, 75]. Similarly, a variety of other exposures might be more common in non-rural areas. For example, exposure to agricultural pesticides is not the only source of pesticide exposures. It is possible that as compared to rural areas, which often are non-agricultural areas, residential use of pesticides in and around homes might be more common in urban and suburban areas. We were unable to consider this potential confounder in the present records-based study. Given a 42% increased risk of PD in relation to pesticide exposure [76], we estimate that the difference in exposure to pesticides would need to be very marked – for example a 15% prevalence of pesticide exposure among controls who use well water but a 55% prevalence of pesticide exposure among controls who do not use well water – in order to have biased an OR of 1.00 down to the OR we observed.

We also acknowledge some additional limitations. There are other potential confounders that we could not address that also could have biased ORs downward, such as some types of physical activity and education/socioeconomic status. Another important potential limitation is that PD ascertainment was based upon claims data that may contain some inaccuracies. Nonetheless, we observed a clear, significant inverse association even when restricting to our most certain PD cases, or when we explored the effect of broadening our case criteria to include controls with evidence of PD, based upon motor signs identified by ICD-9 codes, but not yet identified as having PD. Similarly, there will be misclassification of exposure because we only considered water source for the most recent residence; however, some beneficiaries lived in other regions earlier in life, and early life exposures might be relevant [6]. In addition, we relied on a probabilistic measure of exposure. We note, though, that use of well water in our nationwide Medicare-aged sample from 2009 is consistent with the overall estimate that 15% of homes in the U.S. used well water in 2007 [77]. More importantly, results were unchanged when we restricted to beneficiaries with probabilities of either 0% or 100%, i.e. essentially with water source known. Finally, some beneficiaries likely drank bottled or filtered water, so their assigned water source might not accurately reflect exposures from ingesting water. However, insofar as exposure measurement error did not differ by PD case status, this error likely biased our results toward null. Therefore, exposure measurement error is an unlikely cause of the inverse association between well water and PD.

In summary, reliance on well water does not appear to increase risk of PD in general. While this does not preclude the possibility that specific contaminants in well water increase PD risk in more focused geographic regions, it does provide substantial reassurance that well water is not a risk factor for PD. This contradicts the generally accepted notion that use of well water increases PD risk, and the inverse association we found suggests potentially that some future environmental epidemiologic studies might focus on urban risk factors for PD.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements including sources of support

We thank Dr. Beth A. Mueller of the Fred Hutchinson Cancer Research Center for assisting us with the primary validation study component of the present work. We appreciate the assistance of the American Academy of Family Physicians, which provided the Uniform Data System (UDS) Mapper (www.udsmapper.org).

The results reported herein correspond to specific aims of grant K01ES028295 to Susan Searles Nielsen from NIH – NIEHS. This work was also supported by grants K24ES017765 from NIH – NIEHS; grants from the Michael J. Fox Foundation; grants from the American Parkinson Disease Association (APDA); an APDA Summer Student Fellowship (PDF-APDA-SFW-1664); the Washington University School of Medicine Dean's Fellowship and R03CA94735 from NIH – NCI. The funding organizations played no role in the study design, analysis or decision to publish.

References

- [1]. Nalls MA, Pankratz N, Lill CM, Do CB, Hernandez DG, Saad M, DeStefano AL, Kara E, Bras J, Sharma M, Schulte C, Keller MF, Arepalli S, Letson C, Edsall C, Stefansson H, Liu X, Pliner H, Lee JH, Cheng R, International Parkinson's Disease Genomics C, Parkinson's Study Group Parkinson's Research: The Organized GI, andMe, GenePd, NeuroGenetics Research C, Hussman Institute of Human G, Ashkenazi Jewish Dataset I, Cohorts for H, Aging Research in Genetic E, North American Brain Expression C, United Kingdom Brain Expression C, Greek Parkinson's Disease C, Alzheimer Genetic Analysis G, Ikram MA, Ioannidis JP, Hadjigeorgiou GM, Bis JC, Martinez M, Perlmutter JS, Goate A, Marder K, Fiske B, Sutherland M, Xiomerisiou G, Myers RHLN, Stefansson K, Hardy JA, Heutink P, Chen H, Wood NW, Houlden H, Payami H, Brice A, Scott WK, Gasser T, Bertram L, Eriksson N, Foroud T, Singleton AB (2014) Large-scale meta-analysis of genome-wide association data identifies six new risk loci for Parkinson's disease. *Nat Genet* 46, 989–993. [PubMed: 25064009]
- [2]. Lill CM, Roehr JT, McQueen MB, Kavvoura FK, Bagade S, Schjeide BM, Schjeide LM, Meissner E, Zauft U, Allen NC, Liu T, Schilling M, Anderson KJ, Beecham G, Berg D, Biernacka JM, Brice A, DeStefano AL, Do CB, Eriksson N, Factor SA, Farrer MJ, Foroud T, Gasser T, Hamza T, Hardy JA, Heutink P, Hill-Burns EM, Klein C, Latourelle JC, Maraganore DM, Martin ER, Martinez M, Myers RH, Nalls MA, Pankratz N, Payami H, Satake W, Scott WK, Sharma M, Singleton AB, Stefansson K, Toda T, Tung JY, Vance J, Wood NW, Zabetian CP, 23andMe, The Genetic Epidemiology of Parkinson's Disease Consortium, The International Parkinson's Disease Genomics Consortium, The Parkinson's Disease GWAS Consortium, The Wellcome Trust Case Control Consortium 2, Young P, Tanzi RE, Khoury MJ, Zipp F, Lehrach H, Ioannidis JP, Bertram L (2012) Comprehensive research synopsis and systematic meta-analyses in Parkinson's disease genetics: The PDGene database. *PLoS Genet* 8, e1002548. [PubMed: 22438815]
- [3]. Nalls MA, Blauwendraat C, Vallerga CL, Heilbron K, Bandres-Ciga S, Chang D, Tan M, Kia DA, Noyce AJ, Xue A, Bras J, Young E, von Coelln R, Simon-Sanchez J, Schulte C, Sharma M, Krohn L, Pihlstrom L, Siitonen A, Iwaki H, Leonard H, Faghri F, Gibbs JR, Hernandez DG, Scholz SW, Botia JA, Martinez M, Corvol JC, Lesage S, Jankovic J, Shulman LM, Sutherland M, Tienari P, Majamaa K, Toft M, Andreassen OA, Bangale T, Brice A, Yang J, Gan-Or Z, Gasser T, Heutink P, Shulman JM, Wood NW, Hinds DA, Hardy JA, Morris HR, Gratten J, Visscher PM, Graham RR, Singleton AB, 23andMe Research Team, System Genomics of Parkinson's Disease Consortium, International Parkinson's Disease Genomics Consortium (2019) Identification of novel risk loci, causal insights, and heritable risk for Parkinson's disease: a meta-analysis of genome-wide association studies. *Lancet Neurol* 18, 1091–1102. [PubMed: 31701892]
- [4]. Wirdefeldt K, Gatz M, Schalling M, Pedersen NL (2004) No evidence for heritability of Parkinson disease in Swedish twins. *Neurology* 63, 305–311. [PubMed: 15277625]
- [5]. Wirdefeldt K, Gatz M, Reynolds CA, Prescott CA, Pedersen NL (2011) Heritability of Parkinson disease in Swedish twins: a longitudinal study. *Neurobiol Aging* 32, 1923.e0001–e0008.

- [6]. Goldman SM, Marek K, Ottman R, Meng C, Comyns K, Chan P, Ma J, Marras C, Langston JW, Ross GW, Tanner CM (2019) Concordance for Parkinson's disease in twins: A 20-year update. *Ann Neurol* 85, 600–605. [PubMed: 30786044]
- [7]. Zorzon M, Capus L, Pellegrino A, Cazzato G, Zivadinov R (2002) Familial and environmental risk factors in Parkinson's disease: a case-control study in north-east Italy. *Acta Neurol Scand* 105, 77–82. [PubMed: 11903115]
- [8]. Wright JM, Keller-Byrne J (2005) Environmental determinants of Parkinson's disease. *Arch Environ Occup Health* 60, 32–38. [PubMed: 16961006]
- [9]. Vljajinac HD, Sipetic SB, Maksimovic JM, Marinkovic JM, Dzoljic ED, Ratkov IS, Kostic VS (2010) Environmental factors and Parkinson's disease: a case-control study in Belgrade, Serbia. *Int J Neurosci* 120, 361–367. [PubMed: 20402575]
- [10]. Tsai CH, Lo SK, See LC, Chen HZ, Chen RS, Weng YH, Chang FC, Lu CS (2002) Environmental risk factors of young onset Parkinson's disease: a case-control study. *Clin Neurol Neurosurg* 104, 328–333. [PubMed: 12140099]
- [11]. Smargiassi A, Mutti A, De Rosa A, De Palma G, Negrotti A, Calzetti S (1998) A case-control study of occupational and environmental risk factors for Parkinson's disease in the Emilia-Romagna region of Italy. *Neurotoxicology* 19, 709–712. [PubMed: 9745932]
- [12]. Sanyal J, Chakraborty DP, Sarkar B, Banerjee TK, Mukherjee SC, Ray BC, Rao VR (2010) Environmental and familial risk factors of Parkinson's disease: case-control study. *Can J Neurol Sci* 37, 637–642. [PubMed: 21059511]
- [13]. De Palma G, Mozzoni P, Mutti A, Calzetti S, Negrotti A (1998) Case-control study of interactions between genetic and environmental factors in Parkinson's disease. *Lancet* 352, 1986–1987. [PubMed: 9872254]
- [14]. Morano A, Jimenez-Jimenez FJ, Molina JA, Antolin MA (1994) Risk-factors for Parkinson's disease: case-control study in the province of Caceres, Spain. *Acta Neurol Scand* 89, 164–170. [PubMed: 8030397]
- [15]. Koller W, Vetere-Overfield B, Gray C, Alexander C, Chin T, Dolezal J, Hassanein R, Tanner C (1990) Environmental risk factors in Parkinson's disease. *Neurology* 40, 1218–1221. [PubMed: 2381528]
- [16]. Jimenez-Jimenez FJ, Mateo D, Gimenez-Roldan S (1992) Exposure to well water and pesticides in Parkinson's disease: a case-control study in the Madrid area. *Mov Disord* 7, 149–152. [PubMed: 1584237]
- [17]. James KA, Hall DA (2015) Groundwater pesticide levels and the association with Parkinson disease. *Int J Toxicol* 34, 266–273. [PubMed: 25939349]
- [18]. Hertzman C, Wiens M, Snow B, Kelly S, Calne D (1994) A case-control study of Parkinson's disease in a horticultural region of British Columbia. *Mov Disord* 9, 69–75. [PubMed: 8139607]
- [19]. Gatto NM, Cockburn M, Bronstein J, Manthripragada AD, Ritz B (2009) Well-water consumption and Parkinson's disease in rural California. *Environ Health Perspect* 117, 1912–1918. [PubMed: 20049211]
- [20]. Firestone JA, Smith-Weller T, Franklin G, Swanson P, Longstreth WT Jr., Checkoway H (2005) Pesticides and risk of Parkinson disease: a population-based case-control study. *Arch Neurol* 62, 91–95. [PubMed: 15642854]
- [21]. De Michele G, Filla A, Volpe G, De Marco V, Gogliettino A, Ambrosio G, Marconi R, Castellano AE, Campanella G (1996) Environmental and genetic risk factors in Parkinson's disease: a case-control study in southern Italy. *Mov Disord* 11, 17–23. [PubMed: 8771062]
- [22]. Das K, Ghosh M, Nag C, Nandy SP, Banerjee M, Datta M, Devi G, Chatterjee G (2011) Role of familial, environmental and occupational factors in the development of Parkinson's disease. *Neurodegener Dis* 8, 345–351. [PubMed: 21346317]
- [23]. Chen ML, Lin CH, Lee MJ, Wu RM (2014) BST1 rs11724635 interacts with environmental factors to increase the risk of Parkinson's disease in a Taiwanese population. *Parkinsonism Relat Disord* 20, 280–283. [PubMed: 24342025]
- [24]. Behari M, Srivastava AK, Das RR, Pandey RM (2001) Risk factors of Parkinson's disease in Indian patients. *J Neurol Sci* 190, 49–55. [PubMed: 11574106]

- [25]. Taylor CA, Saint-Hilaire MH, Cupples LA, Thomas CA, Burchard AE, Feldman RG, Myers RH (1999) Environmental, medical, and family history risk factors for Parkinson's disease: a New England-based case control study. *Am J Med Genet* 88, 742–749. [PubMed: 10581500]
- [26]. Chan DK, Woo J, Ho SC, Pang CP, Law LK, Ng PW, Hung WT, Kwok T, Hui E, Orr K, Leung MF, Kay R (1998) Genetic and environmental risk factors for Parkinson's disease in a Chinese population. *J Neurol Neurosurg Psychiatry* 65, 781–784. [PubMed: 9810958]
- [27]. Dick FD, De Palma G, Ahmadi A, Scott NW, Prescott GJ, Bennett J, Semple S, Dick S, Counsell C, Mozzoni P, Haite N, Wettinger SB, Mutti A, Otelea M, Seaton A, Soderkvist P, Felice A, Geoparkinson Study Group (2007) Environmental risk factors for Parkinson's disease and parkinsonism: the Geoparkinson study. *Occup Environ Med* 64, 666–672. [PubMed: 17332139]
- [28]. Dong JQ, Zhang ZX, Zhang KL (2003) Parkinson's disease and smoking: an integral part of PD's etiological study. *Biomed Environ Sci* 16, 173–179. [PubMed: 12964791]
- [29]. Elbaz A, Clavel J, Rathouz PJ, Moisan F, Galanaud JP, Delebotte B, Alperovitch A, Tzourio C (2009) Professional exposure to pesticides and Parkinson disease. *Ann Neurol* 66, 494–504. [PubMed: 19847896]
- [30]. Gorell JM, Johnson CC, Rybicki BA, Peterson EL, Richardson RJ (1998) The risk of Parkinson's disease with exposure to pesticides, farming, well water, and rural living. *Neurology* 50, 1346–1350. [PubMed: 9595985]
- [31]. Kuopio AM, Marttila RJ, Helenius H, Rinne UK (1999) Environmental risk factors in Parkinson's disease. *Mov Disord* 14, 928–939. [PubMed: 10584666]
- [32]. Liou HH, Tsai MC, Chen CJ, Jeng JS, Chang YC, Chen SY, Chen RC (1997) Environmental risk factors and Parkinson's disease: a case-control study in Taiwan. *Neurology* 48, 1583–1588. [PubMed: 9191770]
- [33]. Marder K, Logroschino G, Alfaro B, Mejia H, Halim A, Louis E, Cote L, Mayeux R (1998) Environmental risk factors for Parkinson's disease in an urban multiethnic community. *Neurology* 50, 279–281. [PubMed: 9443493]
- [34]. McCann SJ, LeCouteur DG, Green AC, Brayne C, Johnson AG, Chan D, McManus ME, Pond SM (1998) The epidemiology of Parkinson's disease in an Australian population. *Neuroepidemiology* 17, 310–317. [PubMed: 9778597]
- [35]. Park J, Yoo CI, Sim CS, Kim HK, Kim JW, Jeon BS, Kim KR, Bang OY, Lee WY, Yi Y, Jung KY, Chung SE, Kim Y (2005) Occupations and Parkinson's disease: a multi-center case-control study in South Korea. *Neurotoxicology* 26, 99–105. [PubMed: 15527877]
- [36]. Park J, Yoo CI, Sim CS, Kim JW, Yi Y, Jung KY, Chung SE, Kim Y (2004) Occupations and Parkinson's disease: a case-control study in South Korea. *Ind Health* 42, 352–358. [PubMed: 15295907]
- [37]. Preux PM, Condet A, Anglade C, Druet-Cabanac M, Debrock C, Macharia W, Couratier P, Boutros-Toni F, Dumas M (2000) Parkinson's disease and environmental factors. Matched case-control study in the Limousin region, France. *Neuroepidemiology* 19, 333–337. [PubMed: 11060508]
- [38]. Seidler A, Hellenbrand W, Robra BP, Vieregge P, Nischan P, Joerg J, Oertel WH, Ulm G, Schneider E (1996) Possible environmental, occupational, and other etiologic factors for Parkinson's disease: a case-control study in Germany. *Neurology* 46, 1275–1284. [PubMed: 8628466]
- [39]. Stern M, Dulaney E, Gruber SB, Golbe L, Bergen M, Hurtig H, Gollomp S, Stolley P (1991) The epidemiology of Parkinson's disease. A case-control study of young-onset and old-onset patients. *Arch Neurol* 48, 903–907. [PubMed: 1953412]
- [40]. Tanner CM, Chen B, Wang W, Peng M, Liu Z, Liang X, Kao LC, Gilley DW, Goetz CG, Schoenberg BS (1989) Environmental factors and Parkinson's disease: a case-control study in China. *Neurology* 39, 660–664. [PubMed: 2710356]
- [41]. Werneck AL, Alvarenga H (1999) Genetics, drugs and environmental factors in Parkinson's disease. A case-control study. *Arq Neuropsiquiatr* 57, 347–355. [PubMed: 10450337]
- [42]. Searles Nielsen S, Warden MN, Camacho-Soto A, Willis AW, Wright BA, Racette BA (2017) A predictive model to identify Parkinson disease from administrative claims data. *Neurology* 89, 1448–1456. [PubMed: 28864676]

- [43]. Kupper LL, Karon JM, Kleinbaum DG, Morgenstern H, Lewis DK (1981) Matching in epidemiologic studies: validity and efficiency considerations. *Biometrics* 37, 271–291. [PubMed: 7272415]
- [44]. Health Resources and Services Administration (HRSA) (2017) Uniform Data System (UDS) Mapper, <https://www.udsmapper.org>, John Snow Inc., American Academy of Family Physicians, Accessed May 15, 2018.
- [45]. ArcMap, version 10.6, Esri, Redlands, CA.
- [46]. Manson S, Schroeder J, Van Riper D, Ruggles S (2018) IPUMS National Historical Geographic Information System: Version 13.0, 10.18128/D050.V13.0, University of Minnesota, Minneapolis, MN, Accessed May 15, 2018.
- [47]. U.S. Census Bureau (1990) U.S. Census Summary Tape File 3, <https://www.census.gov/data/datasets/1990/dec/summary-file-3.html>, U.S. Department of Commerce, Economic and Statistics Administration, Accessed May 15, 2018.
- [48]. EPA (2016) Safe Drinking Water Act (SDWA) Compliance Monitoring, <https://www.epa.gov/compliance/safe-drinking-water-act-sdwa-compliance-monitoring>, Accessed May 7, 2018.
- [49]. Searles Nielsen S, Kuehn CM, Mueller BA (2010) Water quality monitoring records for estimating tap water arsenic and nitrate: a validation study. *Environ Health* 9, 4. [PubMed: 20109206]
- [50]. U.S. Geological Survey (USGS) (2014) Estimated Use of Water in the United States County-Level Data for 2010, Accessed October 23, 2019.
- [51]. Hernan MA, Takkouche B, Caamano-Isorna F, Gestal-Otero JJ (2002) A meta-analysis of coffee drinking, cigarette smoking, and the risk of Parkinson's disease. *Ann Neurol* 52, 276–284. [PubMed: 12205639]
- [52]. Ritz B, Ascherio A, Checkoway H, Marder KS, Nelson LM, Rocca WA, Ross GW, Strickland D, Van Den Eeden SK, Gorell J (2007) Pooled analysis of tobacco use and risk of Parkinson disease. *Arch Neurol* 64, 990–997. [PubMed: 17620489]
- [53]. Shih IF, Liew Z, Krause N, Ritz B (2016) Lifetime occupational and leisure time physical activity and risk of Parkinson's disease. *Parkinsonism Relat Disord* 28, 112–117. [PubMed: 27177695]
- [54]. Shih IF, Starhof C, Lassen CF, Hansen J, Liew Z, Ritz B (2017) Occupational and recreational physical activity and Parkinson's disease in Denmark. *Scand J Work Environ Health* 43, 210–216. [PubMed: 28319247]
- [55]. Chen H, Zhang SM, Schwarzschild MA, Hernan MA, Ascherio A (2005) Physical activity and the risk of Parkinson disease. *Neurology* 64, 664–669. [PubMed: 15728289]
- [56]. Abbott RD, Petrovitch H, White LR, Masaki KH, Tanner CM, Curb JD, Grandinetti A, Blanchette PL, Popper JS, Ross GW (2001) Frequency of bowel movements and the future risk of Parkinson's disease. *Neurology* 57, 456–462. [PubMed: 11502913]
- [57]. Kenborg L, Lassen CF, Ritz B, Schernhammer ES, Hansen J, Gatto NM, Olsen JH (2011) Outdoor work and risk for Parkinson's disease: a population-based case-control study. *Occup Environ Med* 68, 273–278. [PubMed: 20884793]
- [58]. Kwon E, Gallagher LG, Searles Nielsen S, Franklin GM, Littell CT, Longstreth WT Jr., Swanson PD, Checkoway H (2013) Parkinson's disease and history of outdoor occupation. *Parkinsonism Relat Disord* 19, 1164–1166. [PubMed: 24044947]
- [59]. International Agency for Research on Cancer (IARC) (2012) in IARC Monographs on the Evaluation of Carcinogenic Risks to Humans: Radiation IARC, Lyon, France.
- [60]. Gross A, Racette BA, Camacho-Soto A, Dube U, Searles Nielsen S (2018) Use of medical care biases associations between Parkinson disease and other medical conditions. *Neurology* 90, e2155–e2165. [PubMed: 29743207]
- [61]. Ritz B, Yu F (2000) Parkinson's disease mortality and pesticide exposure in California 1984–1994. *Int J Epidemiol* 29, 323–329. [PubMed: 10817132]
- [62]. Costello S, Cockburn M, Bronstein J, Zhang X, Ritz B (2009) Parkinson's disease and residential exposure to maneb and paraquat from agricultural applications in the Central Valley of California. *Am Journal of Epidemiol* 169, 919–926. [PubMed: 19270050]
- [63]. Wang A, Costello S, Cockburn M, Zhang X, Bronstein J, Ritz B (2011) Parkinson's disease risk from ambient exposure to pesticides. *Eur J Epidemiol* 26, 547–555. [PubMed: 21505849]

- [64]. U.S. Geological Survey (USGS) (2018) Estimated Annual Agricultural Pesticide Use: County-Level Data, Accessed 10/18/2019.
- [65]. Dwyer-Lindgren L, Mokdad AH, Srebotnjak T, Flaxman AD, Hansen GM, Murray CJ (2014) Cigarette smoking prevalence in US counties: 1996–2012. *Popul Health Metr* 12, 5. [PubMed: 24661401]
- [66]. Searles Nielsen S, Gallagher LG, Lundin JI, Longstreth WT Jr., Smith-Weller T, Franklin GM, Swanson PD, Checkoway H (2012) Environmental tobacco smoke and Parkinson's disease. *Mov Disord.* 27, 293–296. [PubMed: 22095755]
- [67]. O'Reilly EJ, Chen H, Gardener H, Gao X, Schwarzschild MA, Ascherio A (2009) Smoking and Parkinson's disease: using parental smoking as a proxy to explore causality. *Am J Epidemiol* 169, 678–682. [PubMed: 19131566]
- [68]. Stata MC, version 14.2, StataCorp LP, College Station, TX.
- [69]. Pezzoli G, Cereda E (2013) Exposure to pesticides or solvents and risk of Parkinson disease. *Neurology* 80, 2035–2041. [PubMed: 23713084]
- [70]. DeSimone LA, Hamilton PA, Gilliom RJ (2009) Quality of Water from Domestic Wells in Principal Aquifers of the United States, 1991–2004 Overview of Major Findings In National Water-Quality Assessment Program, Circular 1332, U.S. Geological Survey, <https://pubs.usgs.gov/circ/circ1332/includes/circ1332.pdf>, Pembroke Publishing Service Center, Reston, Virginia, Accessed May 1, 2019.
- [71]. Vlaar T, Kab S, Schwaab Y, Frery N, Elbaz A, Moisan F (2018) Association of Parkinson's disease with industry sectors: a French nationwide incidence study. *Eur J Epidemiol* 33, 1101–1111. [PubMed: 29730746]
- [72]. Strosnider H, Kennedy C, Monti M, Yip F (2017) Rural and Urban Differences in Air Quality, 2008–2012, and Community Drinking Water Quality, 2010–2015 - United States. *MMWR Surveill Summ* 66, 1–10.
- [73]. Kasdagli MI, Katsouyanni K, Dimakopoulou K, Samoli E (2019) Air pollution and Parkinson's disease: A systematic review and meta-analysis up to 2018. *Int J Hyg Environ Health* 222, 402–409. [PubMed: 30606679]
- [74]. Block ML, Calderon-Garciduenas L (2009) Air pollution: mechanisms of neuroinflammation and CNS disease. *Trends Neurosci* 32, 506–516. [PubMed: 19716187]
- [75]. Calderon-Garciduenas L, Avila-Ramirez J, Calderon-Garciduenas A, Gonzalez-Heredia T, Acuna-Ayala H, Chao CK, Thompson C, Ruiz-Ramos R, Cortes-Gonzalez V, Martinez-Martinez L, Garcia-Perez MA, Reis J, Mukherjee PS, Torres-Jardon R, Lachmann I (2016) Cerebrospinal fluid biomarkers in highly exposed PM2.5 urbanites: the risk of Alzheimer's and Parkinson's diseases in young Mexico City residents. *J Alzheimers Dis* 54, 597–613. [PubMed: 27567860]
- [76]. Allen MT, Levy LS (2013) Parkinson's disease and pesticide exposure--a new assessment. *Crit Rev Toxicol* 43, 515–534. [PubMed: 23844699]
- [77]. U.S. Census Bureau (2008) American Housing Survey for the United States: 2007 In Current Housing Reports, Series H150/07, U.S. Department of Housing and Urban Development, Office of Policy Development and Research, <https://www.census.gov/prod/2008pubs/h150-07.pdf>, U.S. Government Printing Office, Washington, DC, Accessed May 1, 2019.

Figure 1. Factors potentially affecting the association between use of well water and Parkinson disease (PD)

Figure legend: Several variables might act as potential confounders on the association between use of well water and PD because of the strong positive association between residence in a rural area and use of well water. These include age, sex and race/ethnicity, which might bias the PD-well water odds ratios upward, and tobacco smoking and environmental tobacco smoke (ETS) exposure, which might bias the PD-well water odds ratios downward (see Supplementary Table 1). Several additional variables likewise might mediate the association between PD and well water, but should be treated analytically as confounders to remove their potential effect on the PD-well water odds ratios. These potential “nuisance” mediators include use of medical care, agricultural pesticides, outdoor occupation, lifetime physical activity and other environmental and lifestyle factors. Education and socioeconomic status (SES) could act as a confounder and/or mediator.

Figure 2. Risk of Parkinson disease in relation to use of well water, by state, U.S. Medicare 2009

Figure Legend: Use of well water was not significantly positively associated with Parkinson disease in any individual state after Bonferroni correction ($p < 0.001$). Use of well water means the probability of well water from an individual dug or drilled well, based on residential zip code in the PD diagnosis/control reference year. Probability of well water use is a continuous variable ranging from 0 to 1 (0% to 100%), such that a one unit change, and hence the OR, compares beneficiaries with 100% probability of well water use vs. beneficiaries with 0% probability of well water use, i.e. similar to a dichotomous exposure variable. All ORs and 95% CIs are adjusted for age (as 2 linear splines with a knot at age 85) and sex. ORs and 95% CIs are adjusted for race/ethnicity in as much detail as possible for each state (Supplementary Table 3). Abbreviations: OR, odds ratio; CI, confidence interval.

Table 1.

Characteristics of Parkinson disease cases and controls, U.S. Medicare 2009

	Cases N=89,790	Controls N=21,549,400
	n (%)	n (%)
Sex ^a		
Male	44,684 (49.8)	9,287,842 (43.1)
Female	45,106 (50.2)	12,261,500 (56.9)
Race/ethnicity		
White	79,697 (88.8)	18,616,341 (86.4)
Black	5,386 (6.0)	1,635,170 (7.6)
Hispanic	1,904 (2.1)	399,355 (1.9)
Asian	1,565 (1.7)	460,214 (2.1)
Native American	285 (0.3)	92,978 (0.4)
Pacific Islander/other	872 (1.0)	324,397 (1.5)
Unknown	81 (0.1)	20,945 (0.1)
Smoking index ^{b,c} median	38,492 (42.9)	65,451 (55.4)
UV-related cancer ^d	19,560 (21.8)	19,997 (16.9)
Age, years		
Mean (SD)	78.8 (6.1)	76.0 (6.2)
Minimum	66	66
Median	79	75
Maximum	90	90
Number of physician visits		
Mean (SD)	14.7 (12.4)	10.5 (10.6)
Minimum	0	0
Median	12	8
Maximum	289	655
Number of diagnoses ^{c,e}		
Mean (SD)	100.9 (54)	65.6 (47.1)
Minimum	0	0
Median	94	58
Maximum	441	441
Agricultural pesticides, kg/acre ^f		
Mean (SD)	0.26 (0.54)	0.26 (0.54)
Minimum	0	0
Median	0.08	0.08
Maximum	6.12	6.12
Environmental tobacco smoke, % ^g		
Mean (SD)	16.2 (4.2)	16.3 (4.2)
Minimum	5.7	5.7

	Cases N=89,790	Controls N=21,549,400
	n (%)	n (%)
Median	16.3	16.3
Maximum	31.3	31.3

Abbreviations: SD, standard deviation; International Classification of Diseases Ninth Revision Clinical Modification, ICD-9; kg, kilograms.

^aNumbers and percentages of controls exclude 58 (0.0003%) controls with unknown sex.

^bPredicted probability of ever smoking tobacco divided by the person's total number of unique diagnosis codes (or 1 for 292 cases and 6,227 controls without any diagnosis codes).

^cAvailable for all cases and for all controls in a randomly selected subset of 118,095 controls for whom claims data were available.

^dLip, skin and selected eye cancers[59] as a proxy for lifetime outdoor physical activity, including from outdoor occupations.

^eTotal number of unique ICD-9 diagnosis codes up to the PD diagnosis/control reference date.

^fTotal kilograms of agricultural pesticides applied in 2009 in the county of residence in 2009, divided by the total number of agricultural and non-agricultural acres in the county. Excludes 1268 (1.4%) of cases and 327,323 (1.5%) of controls.

^gPercent of individuals in the county of residence who smoked daily in 2009.

Table 2.

Risk of Parkinson disease in relation to use of well water, U.S. Medicare 2009

Adjustment in addition to age, sex and race/ethnicity ^d	All Cases and Controls 89,705 ^a cases/21,526,967 ^b controls	All with Claims Data ^b 89,705 ^a cases/117,971 ^b controls
	OR (95% CI) ^{d,e}	OR (95% CI) ^{d,e}
None	0.87 (0.85–0.89)	0.86 (0.83–0.89)
Geographic region (state)	0.89 (0.87–0.92)	0.88 (0.85–0.91)
Agricultural pesticides (county), kg/acre	0.87 (0.85–0.89)	0.86 (0.83–0.89)
Number of physician visits ^f	0.88 (0.85–0.90)	0.86 (0.83–0.89)
Number of diagnoses ^g	N/A	0.90 (0.87–0.94)
Smoking ^h	N/A	0.86 (0.83–0.89)
Number of diagnoses ^g and smoking ^h	N/A	0.91 (0.88–0.94)
Number of diagnoses, ^g smoking, ^h ETS, ⁱ UV-related cancer, ^j agricultural pesticides and state	N/A	0.91 (0.88–0.95)

Abbreviations: OR, odds ratio; CI, confidence interval; International Classification of Diseases Ninth Revision Clinical Modification, ICD-9; kg, kilogram; UV, ultraviolet light; ETS, environmental tobacco smoke.

^aExcludes 85 (0.09%) of cases without data on well water use.

^bExcludes 22,375 (0.10%) of controls without data on well water use (including 124 [0.10%] of controls with claims data), and 58 (0.0002%) additional controls with unknown sex. Sex was known for all subjects with claims data.

^cAll cases and a random sample of 0.5% of controls, who represent all cases and controls for whom complete Medicare claims data for 2004–2009, and therefore data on number of diagnoses and smoking, were available.

^dAdjusted for age (as 2 linear splines with a knot at age 85), sex and race/ethnicity (in 7 categories); and, if noted, additional covariates with all continuous variables modeled linearly.

^eProbability of well water use is a continuous variable ranging from 0 to 1 (0% to 100%), such that a one unit change, and hence the OR, compares beneficiaries with 100% probability of well water use vs. beneficiaries with 0% probability of well water use, i.e. similar to a dichotomous exposure variable. Use of well water means the probability of well water from an individual dug or drilled well, based on residential zip code in the PD diagnosis/control reference year.

^fNumber of physician visits (outpatient institutional visits and visits to physician offices combined) in 2008 (prior to PD diagnosis/control reference).

^gTotal number of unique ICD-9 diagnosis codes up to PD diagnosis/control reference date.

^hProbability of ever/never tobacco smoking, determined based on tobacco-specific ICD-9 diagnosis codes and a probabilistic model based on sex, race/ethnicity, birth cohort and >600 other diagnosis and procedure codes.

ⁱPercent of county residents who smoke tobacco daily.

^jAny UV-related cancer[59] according to ICD-9 diagnosis code, as a proxy for lifetime outdoor physical activity.