Brazilian Journal of Physical Therapy

<https://www.journals.elsevier.com/brazilian-journal-of-physical-therapy>

ORIGINAL RESEARCH

Baseline characteristics did not identify people with low back pain who respond best to a Movement System Impairment-Based classification treatment

Daniel Camara Azevedo^{a,b,*}, Paulo Henrique Ferreira^c,
Henrique de Oliveira Santos^b, Daniel Ribeiro Oliveira^b, Joao Victor Leite de Souza^b,
Leonardo Oliveira Pena Costa^{a,d}

^a Programa de Mestrado e Doutorado em Fisioterapia, Universidade Cidade de São Paulo, São Paulo, SP, Brazil

^b Departamento de Fisioterapia, Pontifícia Universidade Católica de Minas Gerais, Belo Horizonte, MG, Brazil

^c Faculty of Health Sciences, University of Sydney, Sydney, Australia

^d Musculoskeletal Division, The George Institute for Global Health, Sydney, NSW, Australia

Received 25 November 2018; received in revised form 14 May 2019; accepted 31 May 2019

Available online 13 June 2019

KEYWORDS

Musculoskeletal;
Pain;
Disability;
Rehabilitation

Abstract

Study design: Secondary analysis of data from a randomized controlled trial.

Background: Treatment based on the Movement System Impairment-Based classification for chronic low back pain results in the same benefit when compared to other forms of exercise. It is possible that participant's characteristics measured at baseline can identify people with chronic low back pain who would respond best to a treatment based on the Movement System Impairment model.

Objectives: To assess if specific characteristics of people with chronic low back pain measured at baseline can modify the effects of a treatment based on the Movement System Impairment model on pain and disability.

Methods: Four variables assessed at baseline that could potentially modify the treatment effects of the treatment based on the Movement System Impairment model were selected (age, educational status, physical activity status and STarT back tool classification). Separate univariate models were used to investigate a possible modifier treatment effect of baseline participant's characteristics on pain and disability after the treatment. Findings of interaction values above 1 point for the outcome mean pain intensity or above 3 points for disability (Roland Morris questionnaire) were considered clinically relevant.

Results: Linear regression analyses for the outcomes of pain and disability did not show interaction values considered clinically relevant for age, educational status, physical activity status and STarT back tool classification.

* Corresponding author at: Pontifícia Universidade Católica de Minas Gerais – Departamento de Fisioterapia, Dom Jose Gaspar, 500, CEP: 30535-901, Belo Horizonte, MG, Brazil.

E-mail: danielazevedo@pucminas.br (D.C. Azevedo).

Conclusion: Age, educational status, physical activity status and STarT back tool classification did not modify the effects of an 8-week treatment based on the Movement System Impairment model in patients with chronic low back pain.

Registered at www.clinicaltrials.gov: NCT02221609 (<https://clinicaltrials.gov/ct2/show/NCT02221609>).

© 2019 Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia. Published by Elsevier Editora Ltda. All rights reserved.

Introduction

Low back pain (LBP) is the most prevalent musculoskeletal disorder¹ and the leading cause of disability worldwide.² LBP is recurrent in nature^{3–6} although the exact estimates of risk of LBP recurrence are unknown.⁷ Exercise is usually a consistent recommendation for treating chronic LBP across different guidelines⁸ resulting however in only modest effects on pain and disability.⁹ Several classification systems have been described to guide the treatment of LBP trying to increase the treatment effects.^{10–15}

The Movement System Impairment based classification (MSI) model is a classification system available to guide the treatment of different musculoskeletal disorders including chronic LBP.^{14,16,17} MSI classification procedure involve interpreting different tests of alignments and movements. Data from timing and magnitude of movement and presence of pain during each test allows the therapist to classify the patient into one of the MSI syndromes. After the classification procedure, treatment is based on correcting the alignment and movement impairments associated with each specific MSI syndrome through patient education, analysis and correction of daily activities and prescription of specific exercises.^{14,16,17}

The efficacy of a treatment based on the MSI model for chronic LBP has been tested in some randomized controlled trials resulting however in the same benefit when compared to other forms of exercise.^{18–20} In a recent clinical trial,^{20,21} treatment based on the MSI model against a treatment based on stretching and strengthening exercise in people with chronic LBP were compared in a high-quality randomized controlled trial. The results of this study showed that patients with chronic LBP had similar improvements in pain, disability and global impression of recovery after participating in a treatment based on the MSI model or a treatment based on stretching and strengthening exercises with a 6-month follow-up. However, this study²⁰ did not investigate if participant's characteristics measured at baseline could identify people with chronic LBP who would respond best to a treatment based on the MSI model. This information might help physical therapists, health care providers and people with chronic LBP in choosing between a treatment based on the MSI model or a treatment based on stretching and strengthening exercises.

The objective of the current study was to assess if specific characteristics of people with chronic LBP measured at baseline can modify the effects of a treatment based on

the MSI model on pain and disability when compared to a treatment based on stretching and strengthening exercises.

Methods

Study design

This study was a secondary analysis of data from a randomized controlled trial comparing the efficacy of a treatment based on the MSI model with a treatment based on stretching and strengthening exercises people with chronic LBP.^{20,21}

Participants

A total of 148 participants of both sexes, between 18 and 65 years of age with non-specific LBP for more than three months and a current minimum pain intensity of three on an 11-point numeric pain rating scale participated in this study. Inclusion criteria also included to be able to stand and walk independently and be literate in Portuguese. Participants were excluded if they presented any contraindications to exercise according to the guidelines of the American College of Sports Medicine,²² serious cardio-respiratory diseases, previous spinal surgery, serious spinal pathologies (fractures, tumors or inflammatory pathologies), nerve root compromise (diagnosed by clinical examination of sensibility, power and reflexes), major depression,^{23,24} pregnancy or if they could not be classified into one of the five categories of the MSI model.^{14,25}

All participants received information about the study and signed the informed consent form. The study was approved by the Ethics Committee of the Pontificia Universidade Catolica de Minas Gerais, Belo Horizonte, MG, Brazil (CAAE 17660913.0.0000.5137) and it was prospectively registered at www.clinicaltrials.gov (NCT02221609).

Randomization and interventions

Participants were randomly allocated into one of the two treatment groups (treatment based on the MSI model or treatment based on stretching and strengthening exercises) by using a randomization schedule performed in Excel for Windows. Both groups participated in an 8-week treatment program consisting of 12 physical therapy sessions supervised by two trained physical therapists (one physical therapist for each group).

Treatment based on the MSI model^{14,16,17} included patient education, analysis and modification of performance of daily activities and specific exercises based on the classification of the participants into one of five possible categories (flexion, extension, rotation, flexion and rotation or extension and rotation) according to the MSI model. The treatment also included a home exercise program that should be performed at least once a day when no treatment sessions were scheduled.

Treatment based on stretching and strengthening exercises^{26,27} included stretching exercises addressing the lumbar, abdominal and lower limbs muscles and strengthening exercises involving the abdominal and paraspinal muscles.^{28,29} The treatment also included a home exercise program that should be performed one or two days a week, in addition to the treatment session days.

Outcome measures

Mean pain intensity and disability after the treatment (eight weeks after randomization) were selected as the main outcome measures of this study. Mean pain intensity was assessed using a numeric pain rating scale (NRS).^{30,31} Disability was assessed using the Roland Morris Disability Questionnaire.³⁰⁻³³ The outcome measures have been translated and cross-culturally adapted into Brazilian Portuguese³⁰⁻³² and are considered as the most important outcome measures for back pain patients.³⁴

Variables of interest

Four variables assessed in baseline that could potentially modify the treatment effects of the treatment based on the MSI model were selected based on theoretical rationale: age, educational level, physical activity status and STarT back screening tool classification.³⁵⁻³⁷

Age

Treatment based on MSI model is based on modification of impaired movements and postures that are supposed to be associated with LBP symptoms. The motor skills required to perform the exercises and activities in the MSI based treatment may be considered more challenging when compared to the exercises prescribed in the treatment based on stretching and strengthening exercises. Because motor skills are expected to gradually deteriorate with age,^{38,39} the hypothesis of this study was that younger people would respond better to the treatment based on MSI model compared to treatment based on stretching and strengthening exercises.

Educational status

Because cognitive-motor learning is influenced positively by educational status,⁴⁰ the hypothesis of this study was that people with higher educational status would respond better to the treatment based on the MSI model. Cognitive-motor learning seems more important to perform exercises and activities associated to the treatment based on MSI model

compared to the performance of the exercises associated with treatment based on stretching and strengthening exercises.

Physical activity status

Different forms of exercise can induce hypoalgesia and improve function through conditioned pain modulation,^{41,42} reduction of systemic inflammation found in chronic pain conditions,⁴³⁻⁴⁵ reduced fear avoidance beliefs, and interrupting the cycle of pain, disuse and deconditioning.⁴⁶ The hypothesis of this study was that people who were physically active before treatment would respond better to treatment based on the MSI model because a different treatment component (modification of impaired movements and postures) was added. On the contrary, people who were physically active before treatment may respond worst to the treatment based on stretching and strengthening exercises because this treatment program may be just adding more exercise to a person who is already involved in some form of exercise.

STarT back screening classification

The STarT back screening tool allows the identification of prognostic indicators in people with LBP in primary care. It categorizes people with LBP into one of three categories, suggesting the best management options for each subgroup: low risk (analgesia, advice and education), medium risk (evidence-based physical therapy) and high risk (combination of physical therapy and cognitive-behavioral therapy).³⁵⁻³⁷ Because treatment based on the MSI model is more tailored to a kinesiotherapeutic approach^{14,16,17} than to a psychosocial approach, the hypothesis of this study was that the participants classified as a low or medium risk according to the STarT back screening tool would respond better to the treatment based on the MSI model compared to treatment based on stretching and strengthening exercises.

Statistical analysis

Separate univariate models were used to assess a possible modifier treatment effect of four baseline participant's characteristics (age, educational level, physical activity status and STarT back screening classification) on pain and disability after a treatment based on the MSI model and treatment based on stretching and strengthening exercises. This subgroup analysis was performed according to the approach recommended by previous research.^{47,48}

Age was dichotomized by using the median split as performed in a previous study.⁴⁹ The other variables were dichotomized as follows: educational level (university and graduate degree versus elementary degree and high school), physical active status (physically active versus physically inactive), STarT back screening classification (low and medium risk versus high risk).

Because subgroup analysis usually requires larger sample sizes compared to randomized controlled trials investigating main effects,⁵⁰ the main interest of this study was to look at the estimated effect sizes rather than statistical significance as suggested by other studies.^{49,51} Findings of

Table 1 Participants' characteristics and potential effect modifiers at baseline.

	Stretching and strengthening exercises (<i>n</i> = 74)	Movement System Impairment based treatment (<i>n</i> = 74)
Age (y)	40.4 ± 13.4	43.4 ± 12.5
Body Mass Index (kg/m ²)	25.5 ± 5.3	25.5 ± 4.4
Sex		
Male	31 (41.9%)	26 (35.1%)
Female	43 (58.1%)	48 (64.9%)
History of low back pain (months)	75.2 ± 74.0	94.9 ± 96.7
^a Mean pain intensity (0–10)	6.5 ± 1.9	6.6 ± 1.8
^b Disability (0–24)	10.2 ± 5.0	10.7 ± 5.1
Effect modifiers		
Age ≤ 43	41 ± 55.4	33 ± 44.6
Higher educational status	21 (28.4%)	21 (28.4%)
Physically active	30 (40.5%)	25 (33.8%)
^c STarT back tool classification (Low and medium risk)	57 (77.0%)	57 (77.0%)

Continuous variables are expressed as mean ± standard deviation.

Categorical variables are expressed as *n* (%).

^a Numeric Pain Rating Scale (NRS).^{30,31}

^b Roland Morris Disability Questionnaire.^{30–32}

^c STarT back screening tool.^{35–37}

interaction values above 1 point for the outcome mean pain intensity (NRS) or above 3 points for disability (Roland Morris questionnaire) were considered clinically relevant.⁴⁹ In this case, the analysis was followed by calculating the marginal means for the subgroups.⁵² IBM SPSS 19 statistical package for Windows (IBM Corp, Armonk, New York) was used for the analyses.

Results

A total of 231 participants volunteered to participate in the study. After the initial assessment, 83 participants were excluded. A total of 148 participants were assessed at baseline and participated in this study. After the treatment, 145 participants completed the 2-month follow-up assessment (treatment based on the MSI model, *n* = 71 and treatment based on stretching and strengthening exercises, *n* = 74). Participants' characteristics and potential effect modifiers at baseline are presented in Table 1.

Linear regression analyses for the outcomes of pain and disability did not show interaction values considered clinically relevant (>1 point for pain and >3 for disability) for any of the potential effect modifiers: age, educational status, physical activity status and STarT back tool classification (Tables 2 and 3).

Discussion

The objective of this study was to investigate if specific baseline characteristics in patients with chronic LBP (age, educational status, physical activity status and STarT back tool classification) could modify the effects of a treatment based on the MSI model compared to a treatment based on

stretching and strengthening exercise on pain and disability. We did not observe any clinically relevant influence⁵³ of those baseline characteristics on the effects of a treatment based on the MSI model after an 8-week treatment program.

Because the current study was underpowered to perform a subgroup analysis, the interpretation of the results was based on the treatment effect modifier sizes represented by the results of interaction tests rather than statistical significance. Although interaction effect values were not exactly equal to treatment effect, the size of interaction can quantify the additional benefit the participants in a specific subgroup would get from treatment when they were compared with those not in this subgroup.⁵² The size of interaction found in all analyses were very small with all interaction values being close to zero (–0.25 to 0.38) (Tables 2 and 3).

The hypothesis of this study was that younger people, with higher educational status, who are physically active and with a STarT back tool classification of low or medium risk would benefit more from a treatment based on the MSI model compared to a treatment based on stretching and strengthening exercises. Although several other baseline characteristics were assessed in baseline, we only included a few number of variables based on theoretical rationale or indirect evidence supporting the hypothesized interaction as suggested by other studies.^{47,48}

There are limited number of randomized controlled trials investigating the effect of a treatment based on the MSI model in people with chronic LBP.^{18–20} To the best of our knowledge, this is the first study to investigate if baseline characteristics of people with chronic LBP could influence the effects of a treatment based on the MSI model on pain and disability. Van Dillen et al.¹⁸ showed that higher adherence to daily activity performance is associated with

Table 2 Linear regression analysis for pain.

Variables	Beta coefficient	<i>p</i>	95% Confidence Interval
<i>Age</i>			
Treatment	−0.06	0.64	−1.53, 0.94
Age <43	−0.25	0.35	−4.06, 1.45
Interaction: treatment × age	0.38	0.16	−0.51, 3.00
Constant	2.99	0.00	0.99, 5.00
<i>Educational level</i>			
Treatment	0.08	0.45	−0.64, 1.44
Educational status (university and graduate degree)	0.10	0.71	−2.49, 3.66
Interaction: treatment × educational status	−0.14	0.61	−2.47, 1.44
Constant	2.24	0.01	0.61, 3.88
<i>Physical activity status</i>			
Treatment	0.05	0.63	−0.84, 1.39
Physically active	0.00	0.99	−2.87, 2.82
Interaction: treatment × physically active	−0.02	0.93	−1.91, 1.75
Constant	2.43	0.01	0.65, 4.21
<i>STarT Back tool classification</i>			
Treatment	−0.02	0.90	−1.91, 1.68
STarT (low and medium risk)	−0.27	0.31	−4.91, 1.57
Interaction: treatment × STarT	0.14	0.65	−1.58, 2.53
Constant	3.71	0.01	0.87, 6.55

Table 3 Linear regression analysis for disability.

Variables	Beta coefficient	<i>p</i>	95% Confidence Interval
<i>Age</i>			
Treatment	0.11	0.36	−1.44, 3.94
Age <43	0.27	0.32	−2.97, 9.05
Interaction: treatment × age	−0.25	0.35	−5.63, 2.03
Constant	2.90	0.19	−1.47, 7.28
<i>Educational status</i>			
Treatment	0.08	0.45	−1.38, 3.10
Educational status (university and graduate degree)	0.15	0.56	−4.67, 8.56
Interaction: treatment × educational level	−0.24	0.37	−6.13, 2.30
Constant	3.97	0.03	0.45, 7.50
<i>Physical activity status</i>			
Treatment	0.01	0.91	−2.27, 2.54
Physically active	−0.10	0.71	−7.31, 4.98
Interaction: treatment × physically active	0.05	0.85	−3.59, 4.32
Constant	5.02	0.01	1.19, 8.86
<i>STarT Back tool classification</i>			
Treatment	0.10	0.57	−2.74, 4.98
STarT (low and medium risk)	−0.07	0.79	−7.93, 6.00
Interaction: treatment × STarT	−0.14	0.63	−5.49, 3.33
Constant	5.29	0.09	−0.81, 11.40

greater effect of a treatment based on the MSI model on function but the authors did not assess the influence of baseline characteristics in their analyses. Other studies have investigated the influence of age, educational status and STarT back screening classification as a potential treatment modifier on different outcomes and conditions. Garcia et al.⁴⁹ showed that older people with chronic LBP respond

better to Mechanical Diagnosis and Treatment compared to Back School. Being younger, however, resulted in greater improvement after a cognitive-behavioral treatment for LBP.⁵⁴ According to Oliveira et al.,⁵⁵ age influenced pain intensity and disability in people with chronic LBP after four weeks of treatment. However, this association was not clinically important. Jain et al.⁵⁶ reported better outcomes

of nonoperative treatment of rotator cuff tears when people have at least a college education. Patients with acute LBP classified as high risk by the STarT Back screening tool respond better to early physical therapy intervention than usual care (only patient education).⁵⁷

Based on the results of the current study, we cannot recommend choosing between treatment based on the MSI model and treatment based on stretching and strengthening exercises for people with chronic LBP by using baseline characteristics of age, educational status, physical activity status and STarT back tool classification. Because treatment based on the MSI model for chronic LBP has been shown to result in the same improvement when compared to other forms of exercise,^{18–20} choosing between both treatments should be based on patient and therapist preferences.

Limitations of the present study included not having equally sized groups positive and negative for all the potential effect modifiers and the low statistical power associated with secondary analysis of data from randomized controlled trials.^{51,52} Results of the primary analysis of the current trial showed that the mean differences in pain and disability between groups were very close to zero.²⁰ This is a potential limitation of the secondary analysis performed in the current study as the chances of detecting treatment effect modification when the main effect achieves the null hypothesis are very unlikely.⁵² However, other studies have described baseline characteristics in people with chronic LBP that can modify the effect of different treatments even when the main effect of treatment was close to zero.^{49,58} Future studies with larger sample sizes should investigate whether other characteristics of people with chronic LBP can modify the effects of a treatment based on the MSI model.

Conclusion

Age, educational status, physical activity status and STarT back tool classification did not modify the effects of an 8-week treatment based on the MSI model in patients with chronic LBP.

Acknowledgements

At the time of the study, the first author was a doctoral candidate funded by *Coordenação de Aperfeiçoamento de Pessoal de Nível Superior* (CAPES). This study was funded by *Conselho Nacional de Desenvolvimento Científico e Tecnológico*/Brazil (CNPQ grant number 470273/2013-5).

Conflicts of interest

The authors declare no conflicts of interest.

References

1. March L, Smith EU, Hoy DG, et al. Burden of disability due to musculoskeletal (MSK) disorders. *Best Pract Res Clin Rheumatol*. 2014;28(3):353–366.
2. Vos T, Barber RM, Bell B, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2015;386(9995):743–800.
3. Andersson GB. Epidemiological features of chronic low-back pain. *Lancet*. 1999;354(9178):581–585.
4. Bergquist-Ullman M, Larsson U. Acute low back pain in industry: a controlled prospective study with special reference to therapy and confounding factors. *Acta Orthop Scand*. 1977;48(suppl 170):1–117.
5. Carey TS, Garrett JM, Jackman A, Hadler N. Recurrence and care seeking after acute back pain: results of a long-term follow-up study. *Med Care*. 1999;37(2):157–164.
6. Hoy D, Brooks P, Blyth F, Buchbinder R. The Epidemiology of low back pain. *Best Pract Res Clin Rheumatol*. 2010;24(6):769–781.
7. da Silva T, Mills K, Brown BT, Herbert RD, Maher CG, Hancock MJ. Risk of recurrence of low back pain: a systematic review. *J Orthop Sports Phys Ther*. 2017;47(5):305–313.
8. O'Connell NE, Cook CE, Wand BM, Ward SP. Clinical guidelines for low back pain: a critical review of consensus and inconsistencies across three major guidelines. *Best Pract Res Clin Rheumatol*. 2016;30(6):968–980.
9. Saragiotto BT, Maher CG, Traeger AC, Li Q, McAuley JH. Dispelling the myth that chronic pain is unresponsive to treatment. *Br J Sports Med*. 2017;51(13):986–988.
10. Delitto A, Erhard RE, Bowling RW. A treatment-based classification approach to low back syndrome: identifying and staging patients for conservative treatment. *Phys Ther*. 1995;75(6):470–485, discussion 485–479.
11. McKenzie RA. *The Lumbar Spine: Mechanical Diagnosis and Therapy*. Waikanae: Spinal; 2003.
12. O'Sullivan P. Diagnosis and classification of chronic low back pain disorders: maladaptive movement and motor control impairments as underlying mechanism. *Man Ther*. 2005;10(4):242–255.
13. Petersen T, Olsen S, Laslett M, et al. Inter-tester reliability of a new diagnostic classification system for patients with non-specific low back pain. *Aust J Physiother*. 2004;50(2):85–94.
14. Sahrman SA. *Diagnosis and Treatment of Movement Impairment Syndromes*. 1st ed. St Louis: Mosby; 2001.
15. Hall H, McIntosh G, Boyle C. Effectiveness of a low back pain classification system. *Spine J*. 2009;9(8):648–657.
16. Sahrman S. *Movement System Impairment Syndromes of the Extremities, Cervical and Thoracic Spines*. Elsevier Health Sciences; 2010.
17. Sahrman S, Azevedo DC, Dillen LV. Diagnosis and treatment of movement system impairment syndromes. *Braz J Phys Ther*. 2017;21(6):391–399.
18. Van Dillen LR, Norton BJ, Sahrman SA, et al. Efficacy of classification-specific treatment and adherence on outcomes in people with chronic low back pain. A one-year follow-up, prospective, randomized, controlled clinical trial. *Man Ther*. 2016;24:52–64.
19. Henry SM, Van Dillen LR, Ouellette-Morton RH, et al. Outcomes are not different for patient-matched versus non-matched treatment in subjects with chronic recurrent low back pain: a randomized clinical trial. *Spine J*. 2014;14(12):2799–2810.
20. Azevedo DC, Ferreira PH, Santos HO, Oliveira DR, de Souza JVL, Costa LOP. Movement system impairment-based classification treatment versus general exercises for chronic low back pain: randomized controlled trial. *Phys Ther*. 2018;98(1):28–39.
21. Azevedo DC, Van Dillen LR, Santos Hde O, Oliveira DR, Ferreira PH, Costa LO. Movement system impairment-based classification versus general exercise for chronic low back

- pain: protocol of a randomized controlled trial. *Phys Ther.* 2015;95(9):1287–1294.
22. ACSM. *ACSM's Guidelines for Exercise Testing and Prescription*. Baltimore: Williams and Wilkins; 1995.
 23. Lovibond SH, Lovibond PF. *Manual for the Depression Anxiety Stress Scales*. 2nd ed. Sydney: Psychology Foundation; 1995.
 24. Vignola RC, Tucci AM. Adaptation and validation of the depression, anxiety and stress scale (DASS) to Brazilian Portuguese. *J Affect Disord.* 2014;155:104–109.
 25. Van Dillen LR, Sahrmann SA, Norton BJ, Caldwell CA, McDonnell MK, Bloom NJ. Movement system impairment-based categories for low back pain: stage 1 validation. *J Orthop Sports Phys Ther.* 2003;33(3):126–142.
 26. Airaksinen O, Brox JI, Cedraschi C, et al. Chapter 4. European guidelines for the management of chronic nonspecific low back pain. *Eur Spine J.* 2006;15(suppl 2):S192–S300.
 27. Pillastriani P, Gardenghi I, Bonetti F, et al. An updated overview of clinical guidelines for chronic low back pain management in primary care. *Jt Bone Spine.* 2012;79(2):176–185.
 28. Hayden JA, van Tulder MW, Malmivaara A, Koes BW. Exercise therapy for treatment of non-specific low back pain. *Cochrane Database Syst Rev.* 2005;(3). CD000335.
 29. Rainville J, Hartigan C, Martinez E, Limke J, Jouve C, Finno M. Exercise as a treatment for chronic low back pain. *Spine J.* 2004;4(1):106–115.
 30. Costa LOP, Maher CG, Latimer J, et al. Clinimetric testing of three self-report outcome measures for low back pain patients in Brazil. Which one is the best? *Spine.* 2008;33(22):2459–2463.
 31. Costa LOP, Maher CG, Latimer J, Ferreira PH, Pozzi GC, Ribeiro RN. Psychometric characteristics of the Brazilian-Portuguese versions of the Functional Rating Index and the Roland Morris Disability Questionnaire. *Spine.* 2007;32(17):1902–1907.
 32. Nusbaum L, Natour J, Ferraz MB, Goldenberg J. Translation, adaptation and validation of the Roland Morris questionnaire – Brazil Roland Morris. *Braz J Med Biol Res.* 2001;34(2):203–210.
 33. Roland M, Morris R. A study of the natural history of low-back pain. Part II: Development of guidelines for trials of treatment in primary care. *Spine (Phila Pa 1976).* 1983;8(2):145–150.
 34. Chiarotto A, Boers M, Deyo RA, et al. Core outcome measurement instruments for clinical trials in nonspecific low back pain. *Pain.* 2018;159(3):481–495.
 35. Hill JC, Dunn KM, Lewis M, et al. A primary care back pain screening tool: identifying patient subgroups for initial treatment. *Arthritis Rheum.* 2008;59(5):632–641.
 36. Pilz B, Vasconcelos RA, Marcondes FB, Lodovichi SS, Mello W, Grossi DB. The Brazilian version of STarT Back Screening Tool – translation, cross-cultural adaptation and reliability. *Braz J Phys Ther.* 2014;18(5):453–461.
 37. Pilz B, Vasconcelos RA, Teixeira PP, et al. Construct and discriminant validity of STarT Back Screening Tool – Brazilian version. *Braz J Phys Ther.* 2017;21(1):69–73.
 38. Ketcham CJ, Seidler RD, Van Gemmert AW, Stelmach GE. Age-related kinematic differences as influenced by task difficulty, target size, and movement amplitude. *J Gerontol B: Psychol Sci Soc Sci.* 2002;57(1):P54–P64.
 39. Shimoda N, Lee J, Kodama M, Kakei S, Masakado Y. Quantitative evaluation of age-related decline in control of preprogrammed movement. *PLOS ONE.* 2017;12(11):e0188657.
 40. Voos MC, Piemonte ME, Mansur LL, Caromano FA, Brucki SMD, Valle L. Educational status influences cognitive-motor learning in older adults: going to university provides greater protection against aging than going to high school. *Arq Neuropsiquiatr.* 2017;75(12):843–849.
 41. Ellingson L, Cook D. Exercise induces hypoalgesia through conditioned pain modulation. *J Pain.* 2011;12(4):P37.
 42. Wonders KY, Drury DG. Exercise intensity as a determinant of exercise induced hypoalgesia. *J Exerc Physiol.* 2011;14(4).
 43. Irwin MR, Olmstead R. Mitigating cellular inflammation in older adults: a randomized controlled trial of Tai Chi Chih. *Am J Geriatr Psychiatry.* 2012;20(9):764–772.
 44. Loprinzi PD. Frequency of moderate-to-vigorous physical activity (MVPA) is a greater predictor of systemic inflammation than total weekly volume of MVPA: Implications for physical activity promotion. *Physiol Behav.* 2015;141:46–50.
 45. Stewart LK, Flynn MG, Campbell WW, et al. The influence of exercise training on inflammatory cytokines and C-reactive protein. *Med Sci Sports Exerc.* 2007;39(10):1714–1719.
 46. Nelson N, Churilla JR. Physical activity, fear avoidance, and chronic non-specific pain: a narrative review. *J Bodyw Mov Ther.* 2015;19(3):494–499.
 47. Sun X, Briel M, Busse JW, et al. Credibility of claims of subgroup effects in randomised controlled trials: systematic review. *BMJ.* 2012;344:e1553.
 48. Sun X, Briel M, Walter SD, Guyatt GH. Is a subgroup effect believable? Updating criteria to evaluate the credibility of subgroup analyses. *BMJ.* 2010;340:c117.
 49. Garcia AN, Costa Lda C, Hancock M, Costa LO. Identifying patients with chronic low back pain who respond best to mechanical diagnosis and therapy: secondary analysis of a randomized controlled trial. *Phys Ther.* 2016;96(5):623–630.
 50. Brookes ST, Whitely E, Egger M, Smith GD, Mulheran PA, Peters TJ. Subgroup analyses in randomized trials: risks of subgroup-specific analyses; power and sample size for the interaction test. *J Clin Epidemiol.* 2004;57(3):229–236.
 51. Kamper SJ, Maher CG, Hancock MJ, Koes BW, Croft PR, Hay E. Treatment-based subgroups of low back pain: a guide to appraisal of research studies and a summary of current evidence. *Best Pract Res Clin Rheumatol.* 2010;24(2):181–191.
 52. Hancock M, Herbert RD, Maher CG. A guide to interpretation of studies investigating subgroups of responders to physical therapy interventions. *Phys Ther.* 2009;89(7):698–704.
 53. Armijo-Olivo S. The importance of determining the clinical significance of research results in physical therapy clinical research. *Braz J Phys Ther.* 2018;22(3):175–176.
 54. Underwood M, Mistry D, Lall R, Lamb S. Predicting response to a cognitive-behavioral approach to treating low back pain: secondary analysis of the BeST data set. *Arthritis Care Res (Hoboken).* 2011;63(9):1271–1279.
 55. Oliveira IS, Costa LOP, Garcia AN, Miyamoto GC, Cabral CMN, Costa L. Can demographic and anthropometric characteristics predict clinical improvement in patients with chronic non-specific low back pain? *Braz J Phys Ther.* 2018;22(4):328–335.
 56. Jain NB, Ayers GD, Fan R, et al. Predictors of pain and functional outcomes after the nonoperative treatment of rotator cuff tears. *Orthop J Sports Med.* 2018;6(8), 2325967118788531.
 57. Magel J, Fritz JM, Greene T, Kjaer P, Marcus RL, Brennan GP. Outcomes of patients with acute low back pain stratified by the STarT back screening tool: secondary analysis of a randomized trial. *Phys Ther.* 2017;97(3):330–337.
 58. Macedo LG, Maher CG, Hancock MJ, et al. Predicting response to motor control exercises and graded activity for patients with low back pain: preplanned secondary analysis of a randomized controlled trial. *Phys Ther.* 2014;94(11):1543–1554, <http://dx.doi.org/10.2522/ptj.20140014>.