Published in final edited form as:

Methods Mol Biol. 2019 ; 2034: 83–95. doi:10.1007/978-1-4939-9658-2_7.

Ultrastructural Analyses of Microglial Interactions with Synapses

Marie-Ève Tremblay^{1,2}, Ania K. Majewska^{3,4}

¹Axe Neurosciences, Centre de Recherche du CHU de Québec-Université Laval, Québec, QC, Canada.

²Département de Médecine Moléculaire, Faculté de Médecine, Université Laval, Québec, QC, Canada.

³Center for Visual Science, University of Rochester, Rochester, NY, USA.

⁴Department of Neuroscience, University of Rochester, Rochester, NY, USA.

Abstract

Immunohistochemical electron microscopy (EM) allows the identification of microglial cell bodies and processes, which are otherwise difficult to recognize based on their ultrastructural features. The technique has been essential in defining, at high spatial resolution, microglial interactions with neurons and synapses, thus providing, among other discoveries, important insights into their roles in synaptic pruning and stripping. In this protocol, we describe the preparation of mouse brain tissue for EM, the immunocytochemical staining against ionized calcium binding adaptor molecule 1, the imaging of microglial cell bodies and processes, and the analysis of microglial relationships with the synaptic neuropil.

Keywords

Immunohistochemistry; Electron Microscopy; Mouse; Brain; Imaging; Ultrastructural Analysis; Microglia; Synapses

1 Introduction

In the mid-twentieth century, the development of electron microscopy as a tool to study the nervous system allowed neuroscientists to view neurons and other cell types at unprecedented resolution in situ. Being able to resolve membranes and intracellular organelles precipitated the understanding of synaptic transmission and illuminated the role that glial cells play in the brain [1]. A number of early and more recent studies have described microglial cell bodies in electron micrographs [2–6], as these can be distinguished from those of other cells because of their small size and a characteristically electron-dense cytoplasm [7–9]. However, while most neuronal and glial processes have unique identifying ultrastructural characteristics, microglial processes have been difficult to characterize based

purely on their morphological properties [10]. Here, we describe an immunohistochemical method to label microglia using the marker ionized calcium binding adaptor molecule 1 (IBA1) for electron microscopic studies. This method allows the delineation of microglial cell bodies, along with large and small processes, and is compatible with ultrastructural analyses of other cell types in order to determine how microglia interact with other brain elements [11, 12]. This technique has allowed the examination of microglial behaviors in the context of normal brain development [11], experience-dependent plasticity [13–15], and disease [6, 16–18], and will prove important in the technical arsenal that will be needed to explore the growing list of functions that microglia have been proposed to play in physiological and disease processes [10].

2 Materials

Use double-distilled water prepared by purifying deionized water, to attain a sensitivity of 18 MΩ-cm at 25 °C. Prepare and store all reagents at room temperature (RT) unless indicated otherwise. Strictly follow the waste disposal regulations and safety guidelines provided by the manufacturers and your institution.

2.1 Rodent Perfusion

The proposed volumes allow the perfusion of six adult mice. Phosphate buffer (PB), phosphate-buffered saline (PBS), and paraformaldehyde (PFA) are prepared the day before perfusion. Acrolein is prepared on the day of perfusion.

1. PB: 100 mM, pH 7.4. Weigh 23.48 g of Na_2HPO_4 and 4.8 g of $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ to prepare a 2 l solution in double-distilled water.
2. PBS: 0.9% NaCl, 50 mM PB, pH 7.4. Weigh 5.87 g of Na_2HPO_4 , 1.2 g of $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$, and 9 g of NaCl to prepare a 1 l solution in double-distilled water.
3. PFA: 4%. Heat PB to 60 °C and add 40 g of electron microscopy grade granular PFA under a fume hood to prepare a 1 l solution. Stir until the solution clears, with the PFA completely dissolved, then cool to RT.
4. Acrolein: 3.5%. Measure 19.45 ml of acrolein 90% (from Sigma; catalog number 110221) in PB under a fume hood to prepare a 500 ml solution (see Note 1).
5. Filter the 3.5% acrolein and 4% PFA solutions using coarse filter paper of 25 µm particle retention. Store at 4 °C.
6. Peristaltic pump with tubing of adequate length to hold approximately 15 ml of liquid.
7. 27½-gauge needle.
8. 25-gauge winged blood collection needle.
9. Surgical tools.

2.2 Brain Sectioning and Immunostaining for IBA1

1. PBS: 0.9% NaCl in 50 mM PB, pH 7.4. Weigh 5.87 g of Na_2HPO_4 , 1.2 g of $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$, and 9 g of NaCl in double-distilled water to prepare a 1 l solution.
2. Vibratome.
3. Cryoprotectant: 40% PBS, 30% ethylene glycol, and 30% glycerol.
4. Tris-buffered saline (TBS): 50 mM, pH 7.4. Weigh 6.05 g of Tris and 8.75 g of NaCl in double-distilled water to prepare a 1 l solution. Adjust the pH to 7.4.
5. Fine paintbrush.
6. Sodium borohydride.
7. Blocking buffer: 10% fetal bovine serum, 3% bovine serum albumin, and 0.01% Triton X-100 in TBS.
8. Rabbit anti-IBA1 primary antibody from Wako (#019–19741).
9. Goat antirabbit secondary antibody conjugated to biotin.
10. 0.05% diaminobenzidine (DAB) and 0.015% hydrogen peroxide in TBS.
11. Orbital rotator.
12. Light microscope.

2.3 Post-fixation and Embedding

1. Osmium tetroxide 1%: Under a fume hood, dissolve 4% osmium tetroxide in PB within a glass vial.
2. Durcupan resin: 20 g of component A, 20 g of component B, 0.6 g of component C, and 0.4 g of component D. Combine the components together in a disposable beaker and mix them well using a 10 ml serological pipette until a uniform color is obtained.
3. Ethanol.
4. Propylene oxide.
5. Aluminum weighing dishes.
6. PCTFE film sheets.
7. Superglue and razor blades.
8. Resin blocks. Prepare resin blocks by pouring Durcupan resin into molds. Polymerize by placing the molds in the oven at 55–60 °C for 3 days.
9. Stereomicroscope.

2.4 Ultrathin Sectioning

1. Ultramicrotome.

2. Glass knife.
3. Diamond knife.
4. Section pick-up tool (such as Perfect Loop from Electron Microscopy Sciences).
5. Toluidine blue: Weigh 2 g sodium borate and 0.2 g toluidine blue to prepare a 200 ml solution in double-distilled water.
6. Heating plate.
7. Light microscope.
8. Filter paper.
9. Mesh grids and grid storage boxes.
10. Transmission electron microscope.

3 Methods

3.1 Mouse Brain Perfusion

1. Use a peristaltic pump to deliver approximately 15 ml of PBS, 75 ml of acrolein, and 150 ml of PFA into the blood circulation, at a flow rate of 25 ml/min (*see* Notes 2 and 3).
2. To setup the pump, insert one end of the tubing in the PBS solution, fill the tubing with PBS and fix a 25-gauge winged blood collection needle to the other end.
3. Insert the tubing in the acrolein solution.
4. Anesthetize one mouse at a time with a 90 mg/kg of bodyweight dose of sodium pentobarbital injected intraperitoneally with a 27½-gauge needle. Assess responses to tail/toe pinches. Proceed only if the mouse is unresponsive to such aversive, painful stimuli.
5. Secure the mouse in the supine position and carefully expose the heart without damaging other organs. Work quickly after puncturing the diaphragm.
6. Hold the heart steady with blunt forceps, incise the right atrium, and start the infusion of PBS. Immediately insert the needle into the left ventricle.
7. Perfuse the mouse with PBS (inside of the tubing) followed by acrolein solution for 3 min (corresponding to 75 ml) then stop the pump and delicately switch the tubing from the acrolein to the PFA solutions for 6 min (corresponding to 150 ml) (*see* Note 4).
8. Extract the fixed brain and post-fix in 4% PFA within a glass vial for 2 h at 4 °C (*see* Note 2).

3.2 Brain Sectioning

1. Wash the fixed brain 3 times with chilled PBS. With a razor blade, remove the cerebellum and the olfactory bulb (unless these regions are to be investigated) and cut the remaining tissue transversally into two pieces of equal height, all of which can be sectioned simultaneously to accelerate the procedure.
2. Glue the brain pieces vertically onto the specimen plate secured into the tray. Add enough PBS solution to submerge completely the entire brain surface.
3. Place the tray inside the vibratome, adjust the sectioning speed to 0.5 mm/s, frequency to 90 Hz, and feed thickness to 50 μ m to yield 50 μ m thick sections. Transfer with a fine paintbrush the sections into 20 ml glass vials containing cryoprotectant solution for storage at -20°C without ultrastructural degradation (*see Note 5*).

3.3 Pre-embedding Immunostaining for IBA1

The following steps can be conducted in a multi-well plate or individual glass vials. All the incubations, except with sodium borohydride, are performed on an orbital rotator.

1. Select the sections to be processed (*see Note 6*).
2. Thoroughly wash the sections with approximately 1 ml of PBS, 3 times for 10 min each.
3. Block endogenous peroxidases with 0.3% hydrogen peroxide in PBS for 10 min.
4. Wash the tissue in PBS, 3 times for 10 min each.
5. Incubate the sections in 0.1% sodium borohydride in PBS for 30 min (*see Note 7*).
6. Wash the tissue in PBS, 3 times for 10 min each, making sure to remove all the bubbles.
7. Incubate the sections in blocking buffer for 1 h (*see Note 8*).
8. Remove the blocking buffer from the tissue and incubate in primary antibody solution (rabbit anti-IBA1, diluted [1:1000] prepared in blocking buffer) at 4°C overnight.
9. Wash in TBS, 3 times for 10 min each.
10. Incubate in secondary antibody (goat antirabbit conjugated to biotin) [1:200] in 0.05 M TBS for 90 min.
11. Wash in TBS, 5 times for 5 min each.
12. Incubate in Avidin–Biotin complex solution (Avidin [1:100], Biotin [1:100]) in TBS solution for 1 h. This solution should be prepared 30 min before incubation.
13. Wash in TBS, 5 times for 5 min each.
14. Reveal the staining with 0.05% DAB and 0.015% hydrogen peroxide in TBS for approximately 5 min (*see Notes 9 and 10*).

15. Stop the reaction by washing the sections in chilled PB, 5 times for 5 min each.

3.4 Processing for Electron Microscopy

The following steps can be performed in multi-well plates or glass vials, except the dehydration step with propylene oxide, which requires glass vials.

1. Prepare 1% osmium tetroxide solution (*see* Note 11).
2. Remove the PB from the brain sections and spread them flat in a multi-well plate using a fine paintbrush.
3. Immerse the sections in osmium tetroxide for 30 min at RT, adding one drop of osmium tetroxide at a time with a transfer pipette to prevent the sections from folding (*see* Note 12).
4. While sections are in osmium tetroxide, prepare the Durcupan resin.
5. Transfer the prepared mixture to aluminum weighing dishes that will receive the sections once they have been dehydrated.
6. Dehydrate the sections in increasing concentrations of ethanol for 5 min each in the following order: 35%, 35%, 50%, 70%, 80%, 90%, 100%, 100%, and 100%. Stop the reaction by washing the sections in PB, 3 times for 10 min each.
7. To remove residual ethanol, immerse the sections in propylene oxide 3 times for 5 min each (*see* Note 13).
8. Use a fine paintbrush to transfer the sections from propylene oxide into the resin and leave overnight for infiltration at RT (*see* Note 14).
9. The following day, place the aluminum weighing dishes containing the specimens into a 55–60 °C oven for 10–15 min.
10. Using a fine paintbrush, paint a thin layer of resin onto one PCTFE film sheet.
11. Move one section of tissue at a time from the aluminum weighing dish to the PCTFE sheet, gently removing excess resin.
12. After moving the sections to the PCTFE sheet, place a second PCTFE sheet over the first, creating a sandwich of tissue and resin (*see* Note 15).
13. Polymerize the resin by placing the sandwiches in the oven for 3 days at 55–60 °C.

3.5 Ultrathin Sectioning

1. After polymerization, remove the light weighing dishes and PCTFE sheet covering the sections. Under a binocular microscope, select square areas of interest (about 2 × 2 mm) and carefully excise them from the PCTFE sheet with a razor blade (*see* Note 16).
2. Glue the areas of interest at the tip of resin blocks using Superglue and cure in the oven at 55–60 °C for at least 1 h.

3. In preparation for sectioning, trim the resin block to the shape of an isosceles trapezoid with a razor blade.
4. Using an ultramicrotome and a glass knife, remove the glue and resin at the surface of the tissue. Fill the boat of the glass knife with double-distilled water and cut a few semi-thin sections to remove the resin (0.5–1 μm thick).
5. Transfer the sections to a SuperFrost slide using a section pick-up tool. Dry the sections by placing the slide on a heating plate at 80 °C for 1 min. Cover with a few drops of 0.1% toluidine blue stain for 1 min. Rinse the excess stain with double-distilled water.
6. Examine the sections under a light microscope to visualize the stained tissue and determine whether the sample is ready for ultrathin sectioning, or whether semithin sectioning should be resumed to go deeper into the tissue.
7. For ultrathin sectioning, replace the glass knife with a diamond knife and fill the boat with double-distilled water. Cut silver ultrathin (60–70 nm) sections and carefully collect them onto mesh grids using a section pick-up tool. Dry the grids on a filter paper and store them in grid storage boxes. The samples are then ready for transmission electron microscopy imaging.

3.6 Imaging and Analysis

In addition to their immunostaining for IBA1, microglia are distinguished from oligodendrocytes by their nuclear heterochromatin pattern, narrower space between nuclear membranes, frequent association with pockets of extracellular space, accumulation of lipid bodies and cellular debris, among other types of phagocytic inclusions, as well as frequent contacts with synaptic elements (Fig. 1).

1. To analyze microglial cell bodies, randomly acquire pictures of cells immunopositive for IBA1 at 6800 $\times$, making a mosaic in Illustrator for each cell. A lower magnification picture at 2900 $\times$ showing the entire cell body is also useful for presentation purpose and creating the mosaic (*see* Notes 17–19, as well as Fig. 1).
2. To analyze microglial processes, randomly acquire pictures of processes immunopositive for IBA1 at 9300 $\times$ (*see* Notes 17–19, as well as Fig. 2).
3. Images are analyzed using ImageJ by an observer blinded to the experimental conditions (*see* Note 20).
4. To analyze area, perimeter, and morphology (circularity, solidity), the “Area”, “Perimeter”, and “Shape descriptors” measurements are selected in Image J, using the “Analyze” and “Set Measurements” tabs (Fig. 3).
5. Select the “ROI Manager” under the “Analyze” and “Tools” tabs.
6. Trace profiles of microglial cell bodies or processes, using the “Freehand selections” tool in ImageJ (Fig. 4).

7. To analyze contacts with synapses, identify all the elements surrounding individual microglial cell bodies or processes. Dendritic spines are recognized by their postsynaptic density, axon terminals by their 40-nm synaptic vesicles, while perisynaptic astrocytic processes have a clear cytoplasm and acute angles, as they go in between the other elements of neuropil. Considering that most of the time, not all elements can be identified in ultrathin section, a separate “unknown” category can be created for those. The different types of elements can be annotated directly in ImageJ using the “Text” tool (Figs. 1 and 2).
8. Direct microglial contacts with pockets of extracellular space, recognized by their absence of delineating membrane, and often accumulation of cellular debris, are determined in the same manner (Figs. 1 and 2).

4 Notes

1. Take special care when working with acrolein, it is corrosive and toxic to researchers and the environment. Also use suitable glassware to prepare acrolein solutions.
2. Perform perfusion and brain extraction strictly inside a fume hood to avoid being exposed to PFA and acrolein.
3. Keep the perfusion solutions on ice throughout the procedure.
4. During perfusion, ensure that the tubing is free from air bubbles. If air bubbles form, gently stop the peristaltic pump, disconnect the needle from the tubing extremity, making sure that the needle position inside of the heart is preserved, start again the pump to let the bubbles pass, stop, then reconnect the tubing, and resume the perfusion.
5. When storing sections in cryoprotectant, make sure to wait until the sections have sunk to the bottom of the vials, that is, impregnated the cryoprotectant, before placing them at -20°C . Section freezing would be highly detrimental to the ultrastructure due to the formation of ice crystals.
6. When selecting brain sections, make sure to use the same Bregma levels across all animals and experimental conditions to allow for comparison.
7. Sodium borohydride is important to reduce any remaining aldehydes from the fixation step, and is especially important to prevent non-specific staining when using acrolein for fixation.
8. In the immunostaining, the low concentration of Triton X-100 allows slight permeabilization of membranes for better staining, and is low enough to preserve most ultrastructural features.
9. Be careful when using DAB, as it is a known carcinogen.
10. The timing of DAB reaction should be determined by carefully monitoring the sections under a light microscope. For IBA1, the cell bodies and processes of IBA1-stained microglia should be visible at $20\times$.

11. Perform osmium tetroxide post-fixation one well at a time to avoid the sections from drying out. Be sure to flatten the sections immediately before adding osmium tetroxide, as any folds in the sections will become permanent and attempting to flatten tissue after osmium fixation will only break the sections. Because osmium tetroxide is photosensitive, protect from light throughout the experiment.
12. As osmium tetroxide fixes lipids, brain sections will appear dark after osmium after fixation. If the sections fail to darken, the procedure should be repeated with a fresh osmium solution. However, it should be noted that neuronal and glial cells bodies, as well as myelin tracts, will still be visible on the sections treated with osmium, under the binocular microscope, allowing to select the regions of interest after embedding.
13. Always use glass vials when working with propylene oxide, as it dissolves plastic in addition to being hazardous.
14. Be careful not to dilute the resin with propylene oxide.
15. When preparing the sandwiches for embedding, evenly distribute light weights (e.g., the plastic caps of glass vials) on top of the sandwich to facilitate the resin spreading. Placing the sandwich between large glass slides (as used for monkey brain tissue) also helps to flatten the brain sections during polymerization.
16. On the embedded sections, make sure to select the same region of interest in all the animals and experimental conditions to allow for comparison.
17. During imaging, make sure to capture images at the tissue-resin border, where the penetration of antibodies and staining intensity is maximal.
18. The provided magnifications were established using a FEI Tecnai Spirit G2 microscope equipped with an ORCA-HR Hamamatsu (10MP) camera.
19. Also make sure to image the same number of microglial cell bodies and processes in all the animals and experimental conditions to allow for comparison.
20. When doing analysis, trace all profiles of cell bodies or processes in a picture before moving to the next, to avoid introducing selection bias.

Acknowledgements

We are grateful to Kanchan Bisht, Julie C. Savage and Hassan El Hajj for their help developing the protocol and providing pictures. We also thank Julie-Christine Lévesque at the Bio-Imaging platform of the Infectious Disease Research Centre of CRCHU de Québec-Université Laval, funded by an equipment and infrastructure grant from the Canadian Foundation Innovation. M.E.T. holds a Canada Research Chair of *Neuroimmune plasticity in health and therapy*. This work was also supported by NIH grants EY019277, AA027111, NS099973 and NSF grant 1557971 to A. K.M.

References

1. Peters A, Palay SL, Webster HF (1976) The fine structure of the nervous system: the neurons and supporting cells. Saunders, New York

2. Schultz RL, Maynard EA, Pease DC (1957) Electron microscopy of neurons and neuroglia of cerebral cortex and corpus callosum. *Am J Anat* 100(3):369–407. 10.1002/aja.1001000305 [PubMed: 13458116]
3. Blinzinger K, Kreutzberg G (1968) Displacement of synaptic terminals from regenerating motoneurons by microglial cells. *Z Zellforsch Mikrosk Anat* 85:145–157 [PubMed: 5706753]
4. Murabe Y, Sano Y (1982) Morphological studies on neuroglia. V. Microglial cells in the cerebral cortex of the rat, with special reference to their possible involvement in synaptic function. *Cell Tissue Res* 223(3):493–506 [PubMed: 6124316]
5. Peinado MA, Quesada A, Pedrosa JA, Torres MI, Martinez M, Esteban FJ, Del Moral ML, Hernandez R, Rodrigo J, Peinado JM (1998) Quantitative and ultrastructural changes in glia and pericytes in the parietal cortex of the aging rat. *Microsc Res Tech* 43(1):34–42. 10.1002/(SICI)1097-0029(19981001)43:1<34::AID-JEMT6>3.0.CO;2-G [PubMed: 9829457]
6. Ohmi K, Greenberg DS, Rajavel KS, Ryazantsev S, Li HH, Neufeld EF (2003) Activated microglia in cortex of mouse models of mucopolysaccharidoses I and IIIB. *Proc Natl Acad Sci U S A* 100(4):1902–1907. 10.1073/pnas.252784899 [PubMed: 12576554]
7. Garcia-Cabezas MA, John YJ, Barbas H, Zikopoulos B (2016) Distinction of neurons, glia and endothelial cells in the cerebral cortex: an algorithm based on cytological features. *Front Neuroanat* 10:107 10.3389/fnana.2016.00107 [PubMed: 27847469]
8. Mori S, Leblond CP (1969) Identification of microglia in light and electron microscopy. *J Comp Neurol* 135(1):57–80. 10.1002/cne.901350104 [PubMed: 4181000]
9. Shapiro LA, Perez ZD, Foresti ML, Arisi GM, Ribak CE (2009) Morphological and ultrastructural features of Iba1-immunolabeled microglial cells in the hippocampal dentate gyrus. *Brain Res* 1266:29–36. 10.1016/j.brainres.2009.02.031 [PubMed: 19249294]
10. Savage JC, Picard K, Gonzalez-Ibanez F, Tremblay ME (2018) A brief history of microglial ultrastructure: distinctive features, phenotypes, and functions discovered over the past 60 years by electron microscopy. *Front Immunol* 9:803 10.3389/fimmu.2018.00803 [PubMed: 29922276]
11. Tremblay ME, Lowery RL, Majewska AK (2010) Microglial interactions with synapses are modulated by visual experience. *PLoS Biol* 8(11):e1000527 10.1371/journal.pbio.1000527 [PubMed: 21072242]
12. Tremblay ME, Riad M, Majewska A (2010) Preparation of mouse brain tissue for immunoelectron microscopy. *J Vis Exp* (41). 10.3791/20212021
13. Lowery RL, Tremblay ME, Hopkins BE, Majewska AK (2017) The microglial fractalkine receptor is not required for activity-dependent plasticity in the mouse visual system. *Glia* 65 (11):1744–1761. 10.1002/glia.23192 [PubMed: 28836393]
14. Sipe GO, Lowery RL, Tremblay ME, Kelly EA, Lamantia CE, Majewska AK (2016) Microglial P2Y12 is necessary for synaptic plasticity in mouse visual cortex. *Nat Commun* 7: ARTN10905 10.1038/ncomms10905
15. Tremblay ME, Zettel ML, Ison JR, Allen PD, Majewska AK (2012) Effects of aging and sensory loss on glial cells in mouse visual and auditory cortices. *Glia*. 10.1002/glia.22287
16. Trapp BD, Wujek JR, Criste GA, Jalabi W, Yin X, Kidd GJ, Stohlman S, Ransohoff R (2007) Evidence for synaptic stripping by cortical microglia. *Glia* 55(4):360–368 [PubMed: 17136771]
17. Stalder M, Phinney A, Probst A, Sommer B, Staufenbiel M, Jucker M (1999) Association of microglia with amyloid plaques in brains of APP23 transgenic mice. *Am J Pathol* 154 (6):1673–1684. 10.1016/S0002-9440(10)65423-5 [PubMed: 10362792]
18. Tischer J, Krueger M, Mueller W, Staszewski O, Prinz M, Streit WJ, Bechmann I (2016) Inhomogeneous distribution of Iba-1 characterizes microglial pathology in Alzheimer's disease. *Glia* 64(9):1562–1572. 10.1002/glia.23024 [PubMed: 27404378]

Fig. 1.

Example of microglial cell body stained for IBA1 (pseudo-colored in purple). The staining with DAB produces an electron-dense precipitate. The cell body was captured at 6800× in the hippocampus CA1 *stratum radiatum* of an aged APP-PS1 mouse, a model of Alzheimer's disease. *a* astrocytic process, *d* dendrite, *dn* dystrophic neurite, *er* endoplasmic reticulum, *g* Golgi apparatus, *lb* lipid body, *mt* mitochondrion, *my* myelin debris, *n* nucleus, *s* dendritic spine, *t* axon terminal. Extracellular space pockets are shown by asterisks

Fig. 2. Microglial processes stained for IBA1 (pseudo-colored in purple). The processes were captured at 9800× in the hippocampus CA1 *stratum lacunosum-moleculare* of an aged APP-PS1 mouse, a model of Alzheimer's disease. Annotations as in Fig. 1. *Ma* myelinated axon

Fig. 3.
Measurements selected in ImageJ for the ultrastructural analysis of microglial cell bodies and processes

Fig. 4.
Example of microglial process tracing using the Freehand selections tool in Image J, from the picture shown in Fig. 2