Published in final edited form as:

Evolution. 2019 November ; 73(11): 2333–2342. doi:10.1111/evo.13853.

An interaction between host and microbe genotypes determines colonization success of a key bumble bee gut microbiota member

Logan A. Sauers*, Ben M. Sadd

School of Biological Sciences, Illinois State University, Normal, IL, USA

Abstract

There has been a proliferation of studies demonstrating an organism's health is influenced by its microbiota. However, factors influencing beneficial microbe colonization and the evolution of these relationships remain understudied relative to host-pathogen interactions. Vertically transmitted beneficial microbes are predicted to show high levels of specificity in colonization, including genotype matching, which may transpire through coevolution. We investigate how host and bacterial genotypes influence colonization of a core coevolved microbiota member in bumble bees. The hindgut colonizing *Snodgrassella alvi* confers direct benefits, but, as an early colonizer, also facilitates the further development of a healthy microbiota. Due to predominantly vertical transmission promoting tight evolution between colonization factors of bacteria and host lineages, we predict that genotype-by-genotype interactions will determine successful colonization. Germ-free adult bees from seven bumble bee colonies (host genotypic units) were inoculated with one of six genetically distinct strains of *S. alvi*. Subsequent colonization within host-genotype and microbe-genotype combinations ranged from zero to one hundred percent, and an interaction between host and microbe genotypes determined colonization success. This novel finding of a genotype-by-genotype interaction determining colonization in an animal host-beneficial microbe system has implications for the ecological and evolutionary dynamics of host and microbe, including associated host-fitness benefits.

Keywords

Bumble bee; *Snodgrassella alvi*; microbiota; genotype-by-genotype interactions; host colonization

Introduction

Recent advances have shed light on the important contributions of microbial communities for the health and evolutionary fitness of their hosts. These microbiota have implications for host defense (Dong et al. 2009; Oliver et al. 2012; Parker et al. 2017), development (Visick et al. 2000; Palm et al. 2015), and physiology (Kikuchi et al. 2012; Brune 2014; Kešnerová

*Corresponding author; lsauers@ilstu.edu address: Julian 210, Campus Box 4120, Normal, IL. 61790-4120, fax number: 309-438-3722.

Author Contributions: LAS and BMS developed the study design. LAS collected data, conducted the analysis, and wrote the first draft of the manuscript. BMS advised on the development of data analysis and contributed to all subsequent manuscript drafts.

et al. 2017). The role of an organism's microbiota in determining fitness-relevant host traits necessitates an understanding of ecological and evolutionary determinants of the composition of these microbial communities, including a focus on factors that drive the colonization and establishment of particular microbes.

Colonization represents an essential component in host-beneficial microbe systems (Visick et al. 2000; Vivas et al. 2003; Lee et al. 2013), with the beneficial effects dependent upon successful microbial colonization. Many factors, such as environmental influences (Yellowlees et al. 2008; Bourne et al. 2009; Kikuchi and Yumoto 2013), host-species specificity (Kwong and Moran 2015; La Rivière et al. 2015), and the host or symbiont genotypes (Chong and Moran 2016; Goodrich et al. 2016; Parker et al. 2017; Wang et al. 2017), may determine colonization success. The influence of these factors on the success of colonization also depends upon the route of microbe transmission between hosts (Bright and Bulgheresi 2010). Where symbionts are environmentally acquired, abiotic factors can serve as the primary driving force behind microbial colonization (Thursz et al. 1999; Sison-Mangus et al. 2018). Such symbionts are facultative and experience a free-living stage between hosts, making survival of both their host and external environment paramount. The environmental influences in these systems are two-fold, as the environment serves as the source of these microbial organisms, but also abiotic factors influence survival and colonization success. Abiotic factors, such as temperature and salinity, can influence microbial growth within hosts and when free-living (Soto et al. 2009). Although environmental influences are involved, systems where microbes are acquired through vertical transmission are predicted to show a greater degree of coevolution and host-microbe specificity (Poisot et al. 2011).

The vertical transmission of beneficial microbial symbionts from mother to offspring facilitates interactions that can shape host specificity (Moeller et al. 2018; Moran et al. 1995; Poisot et al. 2011; Powell et al. 2014; Schardl et al. 1997). Relationships between hosts and microbes may become highly specialized, with genotype matching of hosts and vertically transmitted microbes (Poisot et al. 2011). Such specificity may have consequences for subsequent ecological and evolutionary interactions related to host and symbiont fitness (Gundel et al. 2012). In animals, studies of the effects of host genotype on colonization include examples in multiple systems, such as aphids (Chong & Moran 2016), mice (Moeller et al. 2018), stink bugs (Hosokawa et al. 2016), and humans (Blekham et al. 2015). Furthermore, reciprocal genotype-level specificity (Poisot et al. 2011) is predicted to lead to host-genotype by microbe-genotype interactions, but if host and bacterial genotypes interact to determine colonization success in animal host and beneficial microbe systems has not been investigated.

Despite a lack of research investigating the role of host and beneficial microbe genotypes in determining colonization, there are studies examining genotype-dependent functional outcomes. Functional outcomes, including pathogen defense, digestion, and host development, may depend upon host genotype, beneficial microbe genotype, or interactions between the two. Host genotype affects the formation of microbe induced nitrogen fixing nodules in plants (Smith and Goodman 1999) and how host immune function determines mammalian gut microbiota composition (Spor et al. 2011). Additionally, beneficial

microbial genotypes may differentially determine host defense against pathogens (Ford et al. 2017), ecological invasion success (Gueguen et al. 2010; Rudgers et al. 2010), and resistance to pesticides (Kikuchi et al. 2012, Hosokawa et al. 2016). Furthermore, interactions between host and beneficial microbial genotypes in aphids can mediate pathogen susceptibility (Parker et al. 2017) and the fitness costs of possessing defensive symbionts (Vorburger and Gouskov 2011). Although the interactions between genotypes have been studied in regard to such functional outcomes, the ability for these interactions to determine the successful colonization of beneficial symbionts remains uninvestigated outside of plant systems (Wang et al. 2017).

The contribution of host and symbiont genotypes to colonization have not been well studied in host-beneficial microbe systems, but work in host-parasite systems provides potential insights into the factors involved, and the implications of genotype-level specificity. Within host-parasite systems, host genotypes are implicated in differential immune responses (Lazzaro et al. 2006) and susceptibility to infection (Samuel 2002). However, the influences of genotypes in colonization is not constrained to hosts, as parasite genotypes differ in their abilities to infect (de Roode et al. 2008). Furthermore, the differential immune expression of host individuals in response to different parasite genotypes is evidence of the importance of parasite genotype in these systems (Barribeau and Schmid-Hempel 2013). In systems with diverse host and parasite genotypes, infection outcomes may be determined by the combination of host and parasite genotypes, and there are numerous examples of such genotype-by-genotype interactions (Carius et al. 2001; Lambrechts et al. 2005; Sadd and Schmid-Hempel 2009; Barribeau et al. 2014). In these host-parasite systems, genotype-level specificity is a pre-requisite for the existence of such genotype-by-genotype interactions. It is thought that these interactions are the consequence of coevolution, but they will also feedback on coevolutionary dynamics (Sadd and Schmid-Hempel 2009).

Social bees are a useful model system for understanding host-microbe evolution and its outcomes, and corbiculate bees possess a relatively simple gut bacterial community with 11-15 core species (Martinson et al. 2011). These core species have been implicated in host defense (Koch and Schmid-Hempel 2011; Schwarz et al. 2016), digestion (Kešnerová et al. 2017; Zheng et al. 2017) and development (Näpflin and Schmid-Hempel 2016; Kwong et al. 2017). Further underlining the vital function of these microbes are the negative consequences following their perturbation by pathogens (Schwarz et al. 2016), pesticides (Kakumanu et al. 2016), and antibiotics (Raymann et al. 2017). Vertical transmission of many of these microbes creates a situation where microbial fitness is intricately linked with host health and sets the stage for potential coevolution between host-lineages and their microbes (Powell et al. 2016). This is expected to be greatest where genetic variability within host lineages is low, such as in many bumble bee species with high relatedness of worker bees from a single colony (Hines 2008), and sociality may amplify vertical transmission and the potential for coevolution (Koch et al. 2013). Specificity and genetic differences within hosts and microbes influencing colonization success could result.

Snodgrassella alvi is a key gut colonizer within social bees, including bumble bees. This microbe is highly prevalent in the hindguts of adult bees and is the first microbial species to begin settling (Li et al. 2015; Lim et al. 2015; Powell et al. 2014). Because of this, direct

interactions between the host and *S. alvi* are highly relevant to understand. The biofilm that *S. alvi* forms, directly on the host epithelium (Katsnelson 2015), mediates many environmental hindgut conditions (Zheng et al. 2017). These modifications may be critical in allowing other gut symbionts to colonize the nutrient poor hindgut.

The importance of *S. alvi* for the health of their pollinator hosts make the benefits of studying *S. alvi* colonization two-fold. First, studies in this system will provide novel insights into the principal factors of beneficial symbiont colonization more generally and highlight potential implications for host fitness and subsequent evolutionary dynamics. Additionally, understanding the patterns and factors influencing successful colonization can shed light upon potential implications for pollinator health.

Previous research has demonstrated that *S. alvi* can display host-genus level specificity, where *S. alvi* isolated from specific host genera, bumble bee (*Bombus*) and honey bee (*Apis*), do not colonize hosts of the foreign genus as well as their native hosts (Kwong et al. 2014). The limited number of bumble bee strains used in the study were from a different species than the inoculated host and phylogenetic analyses suggest some sharing of strains across species, but many do appear more narrowly specialized (Powell et al. 2016). Transmission of *S. alvi* in social bumble bees may facilitate evolution with host lineages and potentially genotype-level matching for certain strains (Poisot et al. 2011), which would result in colonization being restricted to related host types with the matching factors relevant for colonization. However, the extent of specificity in this system has not been experimentally examined on the level of host and bacterial genotypes.

We ask if the colonization success of the beneficial gut symbiont *S. alvi* into its bumble bee host *Bombus impatiens* is influenced by the bacterial strain genotype and the genotype of the host, delimited by the host genotypic units of different colonies of origin. We hypothesize that some *S. alvi* strains exhibit narrow specificity to host genotypes, which will stem from interactions between host genotypes and their vertically transmitted bacteria. Thus, we predict that genotype-by-genotype interactions will determine successful colonization.

Materials and Methods

1. Overall Experimental Design

Three *B. impatiens* colonies (A, B and C) were raised from wild collected queens from the Mackinaw River Study Area (Lexington, IL., U.S.A.). Queens were collected with the permission of the ParkLands Foundation (<http://www.parklandsfoundation.org/>) in April 2018. Three additional colonies (D, E, F) were raised from lab-reared and mated queens originally derived from distinct parental commercial colonies from Koppert Biological Systems. One colony (G) was obtained directly from Koppert Biological Systems. Colonies were held under red-light at 26±1.5°C and provided with sugar water (1g cane sugar, 1mL boiled tap water, and 0.1% cream of tartar [potassium bitartate] to partially invert the sugars) and pollen (Brushy Mountain Bee Farms, Moravian Falls, NC). Founding queens and a subset of workers were checked by fecal screening and confirmed free of detectable infections of known gut pathogens. Six strains of *S. alvi* (three from colonies of commercial origin [C7, C11, C14] and three from colonies of wild origin [I1, I2, I49]) were used to

create inoculums. Gut-microbe-free adult worker bees from each colony were assigned to one of seven treatments. All combinations of host genotypic background and *S. alvi* strain were created, with five replicates per genotype treatment combination for a total sample size of 210. An additional 35 bees across all colonies were used to check the gut microbe-free status of uninoculated bees. All strains used were foreign to the host colonies. Native strains could not be used due to restrictions of the colony life-cycle and the time intensive nature of isolating and confirming *S. alvi* identity. However, *S. alvi* strains were isolated from three of the colonies before worker production ceased and a limited number of inoculations showed that colonization in native hosts was achievable, as would be predicted.

2. *Snodgrassella alvi* Strain Origin, Culturing, and Identification

Original hosts (*B. impatiens*) of wild origin strains were sourced from Central Illinois as described above. Original hosts of strains designated commercial were from colonies obtained from Koppert Biological Systems. Bees were isolated for 24 hours after collection and fed gamma irradiated pollen and sterile sugar water *ad libitum*. This isolation period ensures any environmentally acquired microbes just passing through the bees' guts are cleared, preventing environmental contaminants from overgrowing *S. alvi* on agar plates and hindering *S. alvi* isolation. After the 24-hour isolation period, bees were chilled on ice, their guts removed aseptically, and hindguts homogenized for 30 seconds in 200µL of ringer saline with 2.4mm metal beads in a Bead Ruptor (Omni International) on high. Serial dilutions of homogenized samples were spread plated on brain heart infusion agar (Sigma-Aldrich [53286]) supplemented with food coloring for visualization. Plates were incubated at 37°C and 10% CO₂ for 48 hours (Kwong and Moran 2013). Isolated colonies from plates were inoculated into brain heart infusion broth and grown at 37°C and 10% CO₂ for 48 hours. 400µL of each culture was mixed with 400µL of sterile 50% glycerol solution in a 1.5mL screw top tube, and these stock cultures were slow frozen (1°C per minute) and stored at -80°C.

16S RNA gene sequences were used to confirm isolates as *S. alvi*. Briefly, cells were cultured as above, pelleted (8,000rpm, 10 minutes, Micro 200R microcentrifuge, Hettich Zentrifugen), and suspended in lysis buffer (20mM Tris-Cl, 2mM EDTA, 1% Triton X-100, 20mg lysozyme/mL) for 30 minutes at room temperature. Following lysis, DNA was extracted using an IBI Bacterial DNA Extraction Kit following the manufacturer's protocol. 16S RNA gene sequences were amplified with the primers 27F and 1492R (Kwong and Moran 2013). amplicon purity checked using gel electrophoresis, and sample concentration and quality quantified with using a µDrop plate in a Thermo Scientific MultiSkan GO. Sanger sequencing was performed at the University of Illinois Roy J Carter Biotechnology Center, Urbana-Champaign, Illinois. Sequences were manually curated using Sequencher software (Gene Codes Corp, Ann Arbor, MI), and Blastn (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) used to confirm isolates as *S. alvi*.

Partial single copy genes *minD* (Powell et al. 2016) and *phage repressor protein C* (525bp, locus SALWKB2_RS03510 of GenBank accession [CP007446](#)) were used to assess *S. alvi* strain relatedness. Primers RND1_MinDF1 and RND1_MinDR2wbl2, minus Illumina platform primers, were used for *minD* (product 514bp, Powell et al. 2016), with initial

denaturing of 5 minutes at 95°C, 35 cycles of 95°C for 45 seconds, 54°C for 30 seconds, and 72°C for 60 seconds, followed by a final 5 minute extension at 72°C. The reported parameters vary from the previous study (Powell et al. 2016) because of the exclusion of the deep-sequencing portion of amplification. *Phage repressor protein C* (product 593bp) was amplified with novel primers PRPFor (ATGACTATGGCCGGTTTCAG) and PRPRev (CGGCCAAACCACAATACTTT), with PCR as for *minD* except annealing at 52°C. Sequencing was as above, with alignments of trimmed sequences (492bp for *minD* and 526bp for *phage repressor protein C*) carried out using MUSCLE (Edgar 2004) through the Sequencher version 5.4 DNA-Seq tools plugin. Sequences have been deposited in the GenBank under the Accession numbers [MN385152](#) to [MN385157](#) for *minD* and [MN385146](#) to [MN385151](#) *phage repressor protein C*.

3. Preparation of *Snodgrassella* strains for inoculation

Inoculation cultures were initiated from stocks using agar and broth culture parameters as above. After 48 hours in liquid culture, tubes were shaken vigorously to resuspend cells, and optical density (OD) at 600nm measured in four technical replicates to estimate cell number from pre-determined OD-cell number relationships. Dilutions were performed to create stock inoculums of 300µL containing an estimated 10⁶ cells of *S. alvi* per 10µL. As cell to optical density measurements can be variable (Francois et al. 2005), a 1:10,000 dilution of 10µL of each stock solution was plated to verify realized stock cell concentrations through counting the colony forming units (CFUs) and multiplying by the dilution factor. The mean realized inoculation size was 1.26×10^6 cells per 10 µL (standard deviation = 0.46×10^6). We used the realized inoculum size as a covariate in analyses to control for any variation caused by varying inoculum sizes.

Stocks were held at 4°C to limit cell division and reductions in the inoculum viability over time. However, as low temperatures can change bacterial gene expression, stress responses, and metabolism (Liu et al. 2002), fresh stocks were recreated every two to three days to reduce any influence of longer refrigeration on *S. alvi* colonization. However, ages of the stock solutions were tracked and also included in analyses.

4. Rearing and Inoculation of Germ-Free Bees

Cohorts of developing bumble bee larvae will usually be spatially grouped within the colony, with clumps of well-defined individual pupae. Clumps of late stage pupae were identified, removed, and housed in sterilized individual containers until a single worker emerged. Containers had been soaked in 10% bleach for 30 minutes, rinsed with ultrapure water, and dried at 60°C. The emergence of a single worker was used to age the isolated clump, as once a single worker has emerged the remaining individuals should be late stage pupae or pharate adults and individuals in these stages are better able to survive the germ-free treatment. Aged clumps were soaked in a 3% bleach solution for 90 seconds (Näpflin and Schmid-Hempel 2016), sterilizing the outer pupal casing. After soaking, pupal clumps were dried with kimwipes, placed into new germ-free containers, and checked daily for emergence. During the pupal stage, bees shed their gut lining, which, combined with casing sterilization, creates adult bees lacking gut microbiota. This has been confirmed in other studies (Näpflin and Schmid-Hempel 2016) and replicated in our lab.

Emerged adult workers were isolated in individual vials processed using the same procedures as the sterile housing. After an hour, bees were presented with 10 μ L of sugar water with the estimated 10⁶ *S. alvi* cells. This same technique is used for parasite exposures in this system (Sadd 2011). Inoculums were created by pelleting stock inoculums (2,500rpm for 5 minutes), removing the supernatant, and resuspending cells in sugar water. Workers were monitored to ensure inoculum consumption and isolated in sterile housing with autoclaved sugar water and gamma irradiated honey bee pollen provided *ad libitum*.

After four days, bees were chilled on ice, their guts removed aseptically, and hindguts homogenized in 200 μ L of ringer saline. Following homogenization, guts were serially diluted to 1:100, 1:10,000, and 1:1,000,000, plated onto brain heart infusion agar, and grown for four days as above. The four-day cut-off allowed sufficient time for visible colonies to form, even from slower growing strains. Plates were imaged (Canon EOS 7D and Canon Remote Capture software) and CFUs counted as a proxy for viable cell number on the dilution plate allowing individual CFUs to be discerned from one another (i.e. were non-overlapping). This dilution was then included in the conversion of counted CFUs on that plate to the viable number of cells per bee hindgut. Forewings were also removed from each bee and the radial cell measured as a surrogate for body size to use as a covariate in analyses (Palmer-Young et al. 2018).

5. Statistical Analysis

Analyses were carried out in R 3.4.2. (R Core Team 2017). Initially, estimated viable cells per hindgut were used, but these count data contained a large proportion of zeros and models designed to address such distributions, zero-inflated and hurdle, did not converge. Therefore, to further analyze the data, cell estimates were converted to prevalence data, with 0 being individuals with no apparent bacterial colonization and 1 being individuals with *S. alvi* colonization. Absence represents samples that were actually zero or below the detection threshold of 100 cells. A generalized linear model with a binomial distribution was used to analyze presence/absence data. Wing size, inoculum size, and stock age were included in the model as covariates. The initial model included microbe strain and host colony as fixed effects, with an interaction. Terms were sequentially removed from the model and hypothesis testing, with likelihood ratios, used to detect significant differences between models due to removed terms. Additionally, AICc values were calculated for each of the models. The final model was determined using the combination of this hypothesis testing and information criteria, so that the final model did not differ significantly from the full model, while also having the lowest associated AICc (Table 1). The final binomial model was considered appropriate as no overdispersion (dispersion value = 0.96) was indicated by the R package *msme* (Hilbe and Robinson 2013).

Results

Uninoculated control bees across all colonies exhibited no evidence of *S. alvi* colonization, indicating the effectiveness of the germ-free treatment (Figure 1A).

Based on pairwise distances (Figure 1B) between partial sequences of the *minD* and *phage repressor protein C* gene the six *S. alvi* isolates represent six distinct *S. alvi* strains. The

colonization levels of *S. alvi* strains at four days post-inoculation ranged from zero (below the detection limit of 100 cells) to 252,000,000 viable cells in a single bee's hindgut (Figure 1). Due to 66% of samples having zero cell counts, colonization was analyzed as presence or absence for each sample. Coded in this way, colonization ranged from 0 to 100% across replicates of host colony and bacterial strain combinations (Figure 2). The best fitting model for the binomial colonization data included strain, colony, realized inoculum size, and the strain by colony interaction (Table 1). Realized inoculum size had an unsurprising positive effect on colonization (Table 2). Indicative of a genotype-by-genotype interaction, the colonization of *S. alvi* was also determined by a significant interaction between strain and host genotypic unit (Table 2).

Discussion

Although predominant vertical transmission of mutualistic symbionts is predicted to result in high levels of specificity and genotype matching (Poisot et al. 2011), the interaction between host and microbial genotypes in determining colonization success within animal-host and beneficial-microbe systems has remained largely unexplored. The few studies investigating potential interactions in these systems do not explicitly test the influence of genotypic interactions in microbial colonization in an experimental manner (Gomez et al. 2016; Moeller et al. 2016). Our results provide evidence for a genotype-by-genotype interaction determining the colonization success of the beneficial bumblebee symbiont *S. alvi* in its *B. impatiens* host. This shows that specificity can be at a much finer scale than the previously reported genus level specificity (Kwong et al. 2014). The demonstration of a genotype-by-genotype interaction in colonization is also intriguing as previously the effects of host and microbial genotype have mostly been studied within the context of host-parasite systems (Carius et al. 2001; Barribeau et al. 2014). Such genotype-by-genotype specificity will have important implications for the ecological and evolutionary outcomes of interactions in host and beneficial microbe systems.

The vertical transmission of microbes across generations is widespread (Funkhouser and Bordenstein 2013), making understanding the evolutionary processes and patterns in these systems of critical importance. Social bees and their gut microbiota offer an amenable system in which to do this (Kwong and Moran 2016), including studies of specificity of interactions and fitness-relevant outcomes of differential colonization. It is hypothesized that vertical transmission and associated coevolution of hosts and microbes will lead to narrow specificity in some strains, with sociality potentially amplifying the potential for coevolution (Koch et al. 2013). Although this study does not directly test coevolution between host and microbe, the finding of a genotype-by-genotype interaction may be indicative of its action (Poisot et al. 2011). However, it is important to acknowledge that other evolutionary processes can determine symbiont genome evolution, including genetic drift (McCutcheon and Moran 2012), and, much like in host-parasite systems (Antonovics et al. 2013), specificity may not only be the result of coevolution. A determinant of colonization that is non-mutually exclusive of the explanations above and could contribute to colonization patterns observed is the presence of generalist and specialist strains of *S. alvi*. Previous work highlights the presence of some *S. alvi* strains able to colonize across host species (Kwong et al. 2014; Kwong et al. 2017) and clusters of both narrow species specialists and more

generalists based on phylogenetic data (Powell et al. 2016). Our results, however, indicate that even if some strains are generalists across host species, a level of within-species specificity could be common in bumble bee *S. alvi* interactions. In fact, only one strain (C7) successfully colonized in at least one replicate bee across all host backgrounds.

Ideally, native strains to the bee colonies would've been used as one of the inoculation treatments, but this was not possible due to logistical constraints. Therefore, only a small number of native inoculations took place to confirm they do indeed colonize. Under the scenario of coevolution between host and microbe, in comparison to the foreign strains used, native strains would be predicted to be the best colonizers in general. However, even without these native combinations, the described genotype-by-genotype interaction shows specificity, with colonization determined by an apparent matching of factors between host and microbe. These same factors could be those associated with the factors determining colonization success of native strains.

In addition to the mechanisms behind these interactions, there are implications for the presence of genotype-by-genotype interactions that extend beyond beneficial symbiont colonization phenotypes into the functional and evolutionary outcomes. Different patterns of *S. alvi* diversity have been found between honey bee and bumble bee individuals and colonies (Powell et al. 2016). The different ecologies of these bees along with the genotype-by-genotype interactions demonstrated here could contribute to these patterns. Bumble bee species typically possess a single strain of *S. alvi*, while honeybees more often possess multiple strains (Powell et al. 2016). This observation was previously attributed to colony founding events. Bumble bees have an annual social lifecycle and colonies are founded by a single queen after hibernation (Goulson et al. 2008), while honey bees found colonies through swarming of a queen and many workers (Villa 2004). While these founding events may contribute to the maintenance of strain diversity within a colony, genotype-by-genotype interactions may also be involved. Genetic variation within honey and bumble bee colonies will typically differ as a result of different levels of polyandry. Bumble bee queens are predominantly singly mated (Estoup et al. 1995), although this varies by species, whereas honeybee queens can mate upwards of ten times (Tarpy and Nielsen 2002; Mattila and Seeley 2007). Genotype matching between *S. alvi* strains and genotypes of their bee hosts could subsequently drive the observed strain variation within colonies. Polyandry of honeybees increases the genetic variation between workers in a colony (Mattila and Seeley 2007) and multiple strains may be maintained due to differential colonization across the multiple host genotypes present. In bumble bees, the low number of starting strains due to colony foundation by a single queen (Powell et al. 2016) and high level of relatedness between bumble bee workers within a colony (Hines 2008) would restrict *S. alvi* strain diversity.

In aphids, symbiont titers determined by host-genotype have been shown to be correlated with host-fitness (Chong & Moran 2016). For the bumble bee and *S. alvi* system, further studies are required to elucidate fitness consequences of differential colonization. If an association with host-fitness exists, as the benefits of colonization with *S. alvi* elude to (Kwong and Moran 2016), a potential implication of the genotype-by-genotype interaction relates to compatibility mismatches, and hence negative consequences for host fitness

(Gundel et al. 2012). As *S. alvi* is acquired through vertical transmission (Martinson et al. 2011; Koch et al. 2013), with the mother queen passing the bacteria to her daughter offspring, matings creating incompatible offspring genotypes could reduce *S. alvi* colonization success in the subsequent colony, depriving offspring of the bacteria's benefits. These offspring, with low or absent levels of *S. alvi* colonization, would experience negative functional outcomes (Raymann et al. 2017), such as increased pathogen susceptibility or reduced levels of nutrition. These effects could result directly from *S. alvi* abundance, but also indirectly due to overall microbiota dysbiosis because of the foundational role of *S. alvi* in the adult bee gut (Zheng et al. 2017). The influence of host outcrossing creating incompatible host-genotypes has been demonstrated in plant-symbiont systems (Gundel et al. 2012; Sneek et al. 2019) and has been suggested as a mechanism underlying imperfect vertical transmission resulting in the loss of symbionts in certain host lineages (Sneek et al. 2019). There could also be consequences for beneficial microbe colonization and bumble bees due to the introgression of genes from commercial bumble bee colonies into wild populations (Kraus et al. 2011). The importance of genotype-by-genotype interactions and their implications in determining colonization patterns in social bees demonstrates the need for future studies aimed at disentangling underlying mechanisms.

Genotype-by-genotype interactions within a host-beneficial microbe colonizer system have potential consequences for host and microbe evolution. These interactions can determine the beneficial outcomes a host receives from vertically transmitted microbial species, and hence evolutionary fitness, as discussed above. Additionally, as outlined for host-parasite interactions, genotype-by-genotype interactions can provide a basis for local adaptation (Lively and Dybdahl 2000; Zhan et al. 2002). They have also been shown to not only be a consequence of coevolution, but also to feedback into evolutionary dynamics (Thompson and Burdon 1992; Lambrechts et al. 2006). Furthermore, these genotype-by-genotype interactions can result in differing host biotypes (Chu et al. 2011), with the potential to lead to future speciation (Brucker and Bordenstein 2012). These elements require further thought concerning the implications of genotype-by-genotype interactions for host-beneficial microbe evolution.

By demonstrating the importance of host and beneficial symbiont genotypes in determining colonization, the results presented here advance our understanding of host-beneficial microbial colonization and support predictions relating to genotype matching in such systems (Poisot et al. 2011). The further implications of such interactions for ecological and evolutionary processes, and the crucial roles of beneficial microbes in host health, highlight the need for subsequent studies to understand the underlying factors determining such patterns.

Acknowledgements:

We would like to thank Earon Grinage for assistance with bacterial culturing, and Kylie Hampton for help with sample collection. Research reported in this publication was supported by the National Institute of General Medical Sciences of the National Institutes of Health under award number R15GM129681 to BMS, an Illinois State University Pre-tenure Faculty Initiative Grant to BMS, and an Illinois State University Phi Sigma Honors Society R.D. Weigel Grant to LAS.

References

- Antonovics J, Boots M, Ebert D, Koskella B, Poss M and Sadd BM 2013 The origin of specificity by means of natural selection: evolved and nonhost resistance in host–pathogen interactions. *Evolution* 67:1–9. [PubMed: 23289557]
- Barribeau SM, and Schmid-Hempel P. 2013 Qualitatively different immune response of the bumblebee host, *Bombus terrestris*, to infection by different genotypes of the trypanosome gut parasite, *Crithidia bombi*. *Infect. Genet. Evol* 20:249–256. [PubMed: 24055962]
- Barribeau SM, Sadd BM, du Plessis L, and Schmid-Hempel P. 2014 Gene expression differences underlying genotype-by-genotype specificity in a host–parasite system. *Proc. Natl. Acad. Sci. USA* 111:3496–3501. [PubMed: 24550506]
- Blekham R, Goodrich JK, Huang K, Sun Q, Bukowski R, Bell JT, Spector TD, Keinan A, Ley RE, Gevers D, and Clark AG 2015 Host genetic variation impacts microbiome composition across human body sites. *Genome Biol.* 16:191. [PubMed: 26374288]
- Bourne DG, Garren M, Work TM, Rosenberg E, Smith GW, and Harvell CD. 2009 Microbial disease and the coral holobiont. *Trends Microbiol.* 17:554–562. [PubMed: 19822428]
- Bright M and Bulgheresi S 2010 A complex journey: transmission of microbial symbionts. *Nat. Rev. Microbiol* 8:218. [PubMed: 20157340]
- Brucker R and Bordenstein S. 2012 Speciation by symbiosis. *Trends in Ecol. Evol* 27:443–451.
- Brune A 2014 Symbiotic digestion of lignocellulose in termite guts. *Nat. Rev. Microbiol* 12:168–180. [PubMed: 24487819]
- Carius HJ, Little TJ, and Ebert D. 2001 Genetic variation in a host-parasite association: Potential for coevolution and frequency-dependent selection. *Evolution* 55:1136–1145. [PubMed: 11475049]
- Chandler JA, Eisen JA, and Kopp A. 2012 Yeast communities of diverse *Drosophila* species: Comparison of two symbiont groups in the same hosts. *Appl. Environ. Microbiol* 78:7327–7336. [PubMed: 22885750]
- Chong RA and Moran NA. 2016 Intraspecific genetic variation in hosts affects regulation of obligate heritable symbionts. *Proc. Natl. Acad. Sci. USA* 113:13114–13119. [PubMed: 27799532]
- Chu D, Gao C, DeBarro P, Zhang Y, Wan F, & Khan I 2011 Further insights into the strange role of bacterial endosymbionts in whitefly, *Bemisia tabaci*: Comparison of secondary symbionts from biotypes B and Q in China. *Bull. Entomol. Res* 101:477–486. [PubMed: 21329550]
- de Roode JC, Yates AJ, and Altizer S. 2008 Virulence-transmission trade-offs and population divergence in virulence in a naturally occurring butterfly parasite. *Proc. Natl. Acad. Sci. USA* 105:7489–7494. [PubMed: 18492806]
- Dong Y, Manfredini F, and Dimopoulos G. 2009 Implication of the mosquito midgut microbiota in the defense against malaria parasites. *PLoS Pathog.* 5:1–10.
- Edgar RC 2004 MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32:1792–1797. [PubMed: 15034147]
- Estoup A, Scholl A, Pouvreau A, and Solignac M. 1995 Monoandry and polyandry in bumble bees (Hymenoptera: Bombinae) as evidenced by highly variable microsatellites. *Mol. Ecol* 4:89–94. [PubMed: 7711957]
- Ford SA, Williams D, Paterson S, and King KC. 2017 Co-evolutionary dynamics between a defensive microbe and a pathogen driven by fluctuating selection. *Mol. Ecol* 26:1778–1789. [PubMed: 27862515]
- Francois K, Devlieghere F, Standaert R, Geeraerd H, Cools I, Van Impe JF, and Debevere J. 2005 Environmental factors influencing the relationship between optical density and cell count for *Listeria monocytogenes*. *J. Appl. Microbiol* 99:1503–1515. [PubMed: 16313423]
- Funkhouser LJ and Bordenstein SR. 2013 Mom knows best: the universality of maternal microbial transmission. *PLoS Biol.* 11:e1001631. [PubMed: 23976878]
- Gomez A, Petrzalkova KJ, Burns MB, Yeoman CJ, Amato KR, Vickova K, Modry D, Todd A, Jost Robinson CA, Remis MJ, Torralba MG, Morton E, Umana JD, Carbonero F, Gaskins HR, Nelson KE, Wilsos BA, Stumpf RM, White BA, Leigh SR, and Blekham R 2016 Gut microbiome of coexisting BaAka pygmies and Bantu reflects gradients of traditional subsistence patterns. *Cell. Rep* 14:2142–2153. [PubMed: 26923597]

- Goodrich JK, Davenport ER, Waters JL, Clark AG and Ley RE 2016 Cross-species comparisons of host genetic associations with the microbiome. *Science* 352:532–535. [PubMed: 27126034]
- Goulson D, Lye GC, and Darvill B. 2008 Decline and conservation of bumblebee. *Annu. Rev. Entomol* 53:191–208. [PubMed: 17803456]
- Gueguen G, Vavre F, Gnankine O, Peterschmitt M, Charif D, Chiel E, Gottlieb Y, Ghanim M, Zchori-Fein E, and Fleury F. 2010 Endosymbiont metacommunities, mtDNA diversity and the evolution of the *Bemisia tabaci* (Hemiptera: Aleyrodidae) species complex. *Mol. Ecol* 19:4365–4376. [PubMed: 20723069]
- Gundel PE, Martínez-Ghersa MA, Omacini M, Cuyeu R, Pagano E, Ríos R and Ghersa CM 2012 Mutualism effectiveness and vertical transmission of symbiotic fungal endophytes in response to host genetic background. *Evol. Appl* 5:838–849 [PubMed: 23346228]
- Hilbe JM and Robinson AP 2013 *Methods of statistical model estimation*. Chapman and Hall/CRC Press London.
- Hines HM 2008 Historical biogeography, divergence times, and diversification patterns of bumble bees (Hymenoptera: Apidae: *Bombus*). *Syst. Biol* 57:58–75. [PubMed: 18275002]
- Hosokawa T, Ishii Y, Nikoh N, Fujie M, Satoh N, and Fukatsu T 2016 Obligate bacterial mutualists evolving from environmental bacteria in natural insect populations. *Nat. Microbiol.* 1:15011. [PubMed: 27571756]
- Huth-Schwarz A, León A, Vandame R, Moritz RF, and Kraus FB. 2011 Workers dominate male production in the neotropical bumblebee *Bombus wilmattae* (Hymenoptera: Apidae). *Front. Zool* 8:13. [PubMed: 21651814]
- Kakumanu ML, Reeves AM, Anderson TD, Rodrigues RR, and Williams MA. 2016 Honey bee gut microbiome is altered by in-hive pesticide exposures. *Front. Microbiol* 7:1255. [PubMed: 27579024]
- Katsnelson A 2015 Microbiome: The puzzle in a bee's gut. *Nature* 521:S56. [PubMed: 25992673]
- Kešnerová L, Mars RAT, Ellegaard KM, Troilo M, Sauer U, and Engel P. 2017 Disentangling metabolic functions of bacteria in the honey bee gut. *PLoS Biol.* 15:1–28.
- Kikuchi Y, Hosokawa T, and Fukatsu T. 2007 Insect-microbe mutualism without vertical transmission: A stinkbug acquires a beneficial gut symbiont from the environment every generation. *Appl. Environ. Microbiol* 73:4308–4316. [PubMed: 17483286]
- Kikuchi Y, Hayatsu M, Hosokawa T, Nagayama A, Tago K, and Fukatsu T 2012 Symbiont-mediated insecticide resistance. *Proc. Natl. Acad. Sci. USA* 109:8618–8622. [PubMed: 22529384]
- Kikuchi Y, and Yumoto I. 2013 Efficient colonization of the bean bug *Riptortus pedestris* by an environmentally transmitted *Burkholderia* symbiont. *Appl. Environ. Microbiol* 79:2088–2091. [PubMed: 23291548]
- Koch H, and Schmid-Hempel P. 2011 Socially transmitted gut microbiota protect bumble bees against an intestinal parasite. *Proc. Natl. Acad. Sci. USA* 108:19288–92. [PubMed: 22084077]
- Koch H, Abrol DP, Li J, and Schmid-Hempel P. 2013 Diversity and evolutionary patterns of bacterial gut associates of corbiculate bees. *Mol. Ecol* 22:2028–2044. [PubMed: 23347062]
- Kraus FB, Szentgyörgyi H, Roj E, Rhode M, Morio D, Woyciechowski M and Moritz RF 2011 Greenhouse bumblebees (*Bombus terrestris*) spread their genes into the wild. *Conser. Genet* 12:187–192
- Kwong WK, and Moran NA. 2013 Cultivation and characterization of the gut symbionts of honey bees and bumble bees: Description of *Snodgrassella alvi* gen. nov., sp. nov., a member of the family Neisseriaceae of the betaproteobacteria, and *Gilliamella apicola* gen. nov., sp. nov., *Int. J. Syst. Evol. Microbiol* 63:2008–2018. [PubMed: 23041637]
- Kwong WK, Engel P, Koch H, and Moran NA. 2014 Genomics and host specialization of honey bee and bumble bee gut symbionts. *Proc. Natl. Acad. Sci. USA* 111:11509–11514. [PubMed: 25053814]
- Kwong WK, and Moran NA. 2015 Evolution of host specialization in gut microbes: The bee gut as a model. *Gut Microbes* 6:214–220 [PubMed: 26011669]
- Kwong WK and Moran NA, 2016 Gut microbial communities of social bees. *Nat. Rev. Microbiol* 14:374. [PubMed: 27140688]

- Kwong WK, Mancenido AL, and Moran NA. 2017 Immune system stimulation by the native gut microbiota of honey bees. *R. Soc. Open Sci* 4:170003. [PubMed: 28386455]
- Kwong WK, Medina LA, Koch H, Sing K, Soh EY, Ascher JS, Jaffé R, and Moran NA, 2017 Dynamic microbiome evolution in social bees. *Sci. Adv* 3:e1600513. [PubMed: 28435856]
- La Rivière M, Garrabou J, and Bally M. 2015 Evidence for host specificity among dominant bacterial symbionts in temperate gorgonian corals. *Coral Reefs* 34:1087–1098.
- Lambrechts L, Halbert J, Durand P, Gouagna LC, and Koella JC. 2005 Host genotype by parasite genotype interactions underlying the resistance of anopheline mosquitoes to *Plasmodium falciparum*. *Malar. J* 4:1–8. [PubMed: 15638948]
- Lambrechts L, Fellous S, and Koella JC. 2006 Coevolutionary interactions between host and parasite genotypes. *Trends Parasitol.* 22:12–16. [PubMed: 16310412]
- Lazzaro BP, Sackton TB, and Clark AG. 2006 Genetic variation in *Drosophila melanogaster* resistance to infection: A comparison across bacteria. *Genetics* 174:1539–1554. [PubMed: 16888344]
- Lee SM, Donaldson GP, Mikulski Z, Boyajian S, Ley K, and Mazmanian SK. 2013 Bacterial colonization factors control specificity and stability of the gut microbiota. *Nature* 501:426–429. [PubMed: 23955152]
- Li J, Powell JE, Guo J, Evans J, Wu J, Williams P, Lin Q, Moran N, and Zhang Z 2015 Two gut community enterotypes recur in diverse bumblebee species. *Curr. Biol* 25:R652–R653. [PubMed: 26241138]
- Lim HC, Chu CC, Seufferheld MJ, and Cameron SA 2015 Deep sequencing and ecological characterization of gut microbial communities of diverse bumble bee species. *PloS ONE* 10:e0118566. [PubMed: 25768110]
- Liu S, Graham JE, Bigelow L, Morse PD, and Wilkinson BJ. 2002 Identification of *Listeria monocytogenes* genes expressed in response to growth at low temperature. *Appl. Environ. Microbiol* 68:1697–1705. [PubMed: 11916687]
- Lively CM, and Dybdahl MF. 2000 Parasite adaptation to locally common host genotypes. *Nature* 405:679–681 [PubMed: 10864323]
- Martinson VG, Danforth BN, Minckley RL, Rueppell O, Tingek S, and Moran NA. 2011 A simple and distinctive microbiota associated with honey bees and bumble bees. *Mol. Ecol* 20:619–628. [PubMed: 21175905]
- Mattila HR, and Seeley TD. 2007 Genetic diversity in honey bee colonies enhances productivity and fitness. *Science* 317:362–364. [PubMed: 17641199]
- McCutcheon JP and Moran NA 2012 Extreme genome reduction in symbiotic bacteria. *Nat. Rev. Microbiol* 10:13.
- Moeller AH, Caro-Quintero A, Mjungu D, Georgiev A, Lonsdorf EV, Muller MN, Pusey AE, Peeters M, Hahn BH, and Ochman H 2016 Cospeciation of gut microbiota with hominids. *Science* 353:380–382. [PubMed: 27463672]
- Moeller AH, Suzuki TA, Phifer-Rixey M, and Nachman M 2018 Transmission modes of the mammalian gut microbiota. *Science* 362:453–457. [PubMed: 30361372]
- Moran NA, vonDohlen CD, and Baumann P. 1995 Faster evolutionary rates in endosymbiotic bacteria than in cospeciating insect hosts. *J. Mol. Evol* 41:727–731.
- Näpflin K, and Schmid-Hempel P. 2016 Immune response and gut microbial community structure in bumblebees after microbiota transplants. *Proc. R. Soc. B Biol. Sci* 283:1–8.
- Nyholm SV, and McFall-Ngai MJ. 2004 The winnowing: Establishing the squid- *Vibrio symbiosis*. *Nat. Rev. Microbiol* 2:632–642. [PubMed: 15263898]
- Oliver KM, Noge K, Huang EM, Campos JM, Becerra JX, and Hunter MS. 2012 Parasitic wasp responses to symbiont-based defense in aphids. *BMC Biol.* 10:1–10. [PubMed: 22214525]
- Palm NW, deZoete MR, and Flavell RA. 2015 Immune-microbiota interactions in health and disease. *Clin. Immunol.* 159:122–127. [PubMed: 26141651]
- Palmer-Young EC, Calhoun AC, Mirzayeva A, and Sadd BM. 2018 Effects of the floral phytochemical eugenol on parasite evolution and bumble bee infection and preference. *Sci. Rep.* 8:1–11. [PubMed: 29311619]

- Parker BJ, Hretek J, McLean AHC, and Godfray HCJ. 2017 Genotype specificity among hosts, pathogens, and beneficial microbes influences the strength of symbiot-mediated protection. *Evolution* 71:1222–1231. [PubMed: 28252804]
- Poisot T, Bever JD, Nemri A, Thrall PH and Hochberg ME 2011 A conceptual framework for the evolution of ecological specialisation. *Ecol. Lett* 14:841–851. [PubMed: 21699641]
- Powell E, Martinson VG, Urban-Mead K, and Moran NA 2014 Routes of acquisition of the gut microbiota of the honey bee *Apis mellifera*. *Appl. Environ. Microbiol* 80:7378–7387. [PubMed: 25239900]
- Powell E, Ratnayeke N, and Moran NA. 2016 Strain diversity and host specificity in a specialized gut symbiont of honey bees and bumble bees. *Mol. Ecol.* 1:4461–4471.
- Core Team R. 2017 R: A language and environment for statistical computing R Foundation for Statistical Computing. Vienna, Austria.
- Raymann K, Shaffer Z, and Moran NA. 2017 Antibiotic exposure perturbs the gut microbiota and elevates mortality in honeybees. *PLoS Biol.* 15:1–22.
- Rudgers JA, Fischer S, and Clay K. 2010 Managing plant symbiosis: Fungal endophyte genotype alters plant community composition. *J. Appl. Ecol* 47:468–477.
- Sadd BM and Schmid-Hempel P 2009 Ecological and evolutionary implications of specific immune responses. 225–240 In Rolff J and Reynolds SE. *Insect infection and immunity: evolution, ecology, and mechanisms*. Oxford Univ. Press Oxford.
- Sadd BM 2011 Food-environment mediates the outcome of specific interactions between a bumblebee and its trypanosome parasite. *Evolution* 65:2995–3001. [PubMed: 21967438]
- Samuel CE 2002 Host genetic variability and West Nile virus susceptibility. *Proc. Natl. Acad. Sci. USA* 99:11555–11557. [PubMed: 12192094]
- Schardl CL, Leuchtmann A, Chung KR, Penny D, and Siegel MR. 1997 Coevolution by common descent of fungal symbionts (*Epichloe spp.*) and grass hosts. *Mol. Biol. Evol* 14:133–143.
- Schwarz RS, Moran NA, and Evans JD. 2016 Early gut colonizers shape parasite susceptibility and microbiota composition in honey bee workers. *Proc. Natl. Acad. Sci. USA* 113:9345–9350. [PubMed: 27482088]
- Sison-Mangus MP, Metzger CMJA, and Ebert D. 2018 Host genotype-specific microbiota do not influence the susceptibility of *D. magna* to a bacterial pathogen. *Sci. Rep* 8:9407. [PubMed: 29925845]
- Smith KP, and Goodman RM. 1999 Host variation for interactions with beneficial plant-associated microbes. *Annu. Rev. Phytopathol* 37:473–491. [PubMed: 11701832]
- Sneck ME, Rudgers JA, Young CA and Miller TE, 2019 Does host outcrossing disrupt compatibility with heritable symbionts? *Oikos*. Online early
- Soto W, Gutierrez J, Remmenga MD, and Nishiguchi MK. 2009 Salinity and temperature effects on physiological responses of *Vibrio fischeri* from diverse ecological niches. *Microb. Ecol* 57:140–150. [PubMed: 18587609]
- Spor A, Koren O, and Ley R. 2011 Unravelling the effects of the environment and host genotype on the gut microbiome. *Nat. Rev. Microbiol* 9:279–290. [PubMed: 21407244]
- Tarpy DR, and Nielsen DI. 2002 Sampling error, effective paternity, and estimating the genetic structure of honey bee colonies (Hymenoptera: Apidae). *Ann. Entomol. Soc. Am* 95:513–528.
- Thompson JN, and Burdon JJ. 1992 Gene-for-gene coevolution between plants and parasites. *Nature* 360:121–125.
- Thursz M, Yallop R, Goldin R, Trepo C, and Thomas HC. 1999 Influence of MHC class II genotype on outcome of infection with hepatitis C virus. *Lancet* 354:2119–2124. [PubMed: 10609818]
- Villa JD 2004 Swarming behavior of honey bees (Hymenoptera: Apidae) in southeastern Louisiana. *Ann. Entomol. Soc. Am* 97:111–116.
- Visick KL, Foster J, Doi J, McFall-Ngai M, and Ruby EG. 2000 *Vibrio fischeri* lux genes play an important role in colonization and development of the host light organ. *J. Bacteriol* 182:4578–4586. [PubMed: 10913092]
- Vivas A, Vörös I, Biró B, Campos E, Barea JM, and Azcón R. 2003 Symbiotic efficiency of autochthonous arbuscular mycorrhizal fungus (*G. mosseae*) and *Brevibacillus sp.* isolated from

cadmium polluted soil under increasing cadmium levels. Environ. Pollut 126:179–189. [PubMed: 12927489]

Vorburger C, and Gouskov A. 2011 Only helpful when required: A longevity cost of harbouring defensive symbionts. J. Evol. Biol 24:1611–1617. [PubMed: 21569156]

Wang Q, Yang S, Liu J, Terecskei K, Ábrahám E, Gombár A, Domonkos Á, Szűcs A, Körmöczi P, Wang T and Fodor L 2017 Host-secreted antimicrobial peptide enforces symbiotic selectivity in *Medicago truncatula*. Proc. Natl. Acad. Sci. USA 114:6854–6859. [PubMed: 28607058]

Yellowlees D, Rees TAV, and Leggat W. 2008 Metabolic interactions between algal symbionts and invertebrate hosts. Plant Cell Environ. 31:679–694. [PubMed: 18315536]

Zhan J, Mundt CC, Hoffer ME, and McDonald BA. 2002 Local adaptation and effect of host genotype on the rate of pathogen evolution: An experimental test in a plant pathosystem. J. Evol. Biol 15:634–647.

Zheng H, Powell E, Steele M, Dietrich C, and Moran N. 2017 Honeybee gut microbiota promotes host weight gain via bacterial metabolism and hormonal signaling. Proc. Natl. Acad. Sci. USA 114:4775–4780. [PubMed: 28420790]

Figure 1.
A) Viable cells of *S. alvi* colonizing bee guts 4 days following inoculation of different *S. alvi* genotypes (x-axes) across multiple host colonies (genotypic units, panels A-G). Different *S. alvi* strains used to inoculate them are on the x-axis. Control individuals (first column in each panel) received the germ-free bee protocol, but were not given *S. alvi*. **B) Pairwise between strain distance matrix for *minD* (below diagonal, unshaded) and *phage repressor protein C* (above diagonal, shaded).** Dissimilarity was calculated with *seqinr* package in R and represents the square root of the pairwise proportional distance.

Figure 2. Proportion of inoculated bees colonized by *S. alvi* based on the host genotypic unit (colony) and *S. alvi* strain genotype.

There were five samples per treatment type combination.

Table 1.
Hypothesis testing model selection.

Table showing the degrees of freedom, residual variance, and p-value for tested models during the model selection process. Models were tested for the null hypothesis of no significant difference until the final model was obtained.

Model	Residual d.f.	Residual Deviance	AICc	P-value
Full	158	162.06	278.43	
Body Size Removed	159	162.06	275.13	0.9793
Inoculum Age Removed	159	164.96	278.03	0.0885
Inoculum Size Removed	159	168.89	281.96	0.0089
Final Model	160	164.98	274.78	0.0875

Table 2.
Factors in the final binomial generalized linear model of *S. alvi* colonization.

Significant terms ($p < 0.05$) are bolded.

Effect	d.f.	LR χ^2	P-value
Strain	5	12.50	0.0285
Colony	6	9.58	0.1435
Inoculum Size	1	6.93	0.0084
Strain x Colony	30	44.51	0.0428