Conservation paradox: Large-scale mining waste in protected areas in two global hotspots, southeastern Brazil

Luciana Hiromi Yoshino Kamino , Eric Oliveira Pereira,
Flávio Fonseca do Carmo

Received: 15 November 2019 / Revised: 14 February 2020 / Accepted: 17 February 2020 / Published online: 16 March 2020

Abstract Irreversible losses of natural areas and damage to ecosystem services are caused by large-scale mining. These effects are prominent in Minas Gerais, which is home to a mining industry with hundreds of tailings dams concentrated in global hotspots: the Cerrado and the Atlantic Forest. We present an overview of the spatial overlap between protected areas (PAs), priority conservation areas (PCAs), tailings dams, and their influence areas. A total of 218 dams accumulated environmental costs and liability that are represented by 2.24 thousand million m³ of tailings and industrial waste. Most dams ($\approx 70\%$) are located in PCAs, 45 dams in conservation units, 58 in their respective buffer zones, and 51 in special protection areas. We have identified an urgent need to implement adequate public policies to establish management and monitoring mechanisms in PAs and PCAs and to address a scenario of high potential for socioenvironmental damage associated with tailings dams.

Keywords Ecological disaster · Environmental footprints · Governance · Megadiversity · Tailings Dam

INTRODUCTION

Brazil has the largest biological heritage in the world; that is, it is home to approximately 20% of all known species (Ministério do Meio Ambiente, MMA 2019a). Minas Gerais stands out among the Brazilian states because it contains two global hotspots (the Cerrado and the Atlantic Forest), which contain a wide diversity of vegetation types, including forest formations (ombrophilous, semideciduous, and deciduous), cerrado (Brazilian savanna), and plant communities associated with rock outcrops (The Brazil Flora Group, BFG 2015). However, biodiversity is not

homogeneously distributed and tends to concentrate in unique natural areas due to historical, evolutionary, and ecological factors (Carvalho 2009), such as ancient metal-rich ecosystems—ironstone ranges and canga outcrops (Jacobi et al. 2011; Carmo et al. 2018). Ironstone ranges contain one of the largest concentrations of caves in Brazil (Centro Nacional de Pesquisa e Conservação de Cavernas, CECAV 2018) and harbor rare invertebrate communities, such as the troglotauna, which present physiological and ecological specializations to the underground environment (Ferreira et al. 2015).

Due to this heterogeneous scenario, methods such as systematic conservation planning (Margules and Pressey 2000) are utilized for the identification of priority conservation areas (PCAs), which represent territorial planning units for the rational use of natural resources and the conservation of species and ecosystems. In Brazil, PCA are referenced for the implementation or formulation of public policies, programs, projects, and activities that are aimed at improving biodiversity protection management (Minas Gerais 2002; Ministério do Meio Ambiente, MMA 2018). Among the most important actions specified from the identification of a PCA is the indication for the creation of protected areas (PAs) in the regions where the objects of conservation are concentrated. Creating PAs is globally recognized as the most effective strategy to contain biodiversity degradation, improve human well-being, maintain the quality of environmental services, and stimulate the rational use of natural resources (Naughton-Treves et al. 2005; Gaston et al. 2008). However, the effectiveness of this strategy depends on both adequate management within PAs and the type and intensity of human land use outside its borders (Hansen and DeFries 2007).

One of the main land uses in Minas Gerais is large-scale mining (Jacobi et al. 2011) because the state contains one

of the largest geological reserves of metal minerals in the world and is the largest producer of iron, gold, zinc, and niobium in Brazil (Agência Nacional de Mineração, ANM 2018). Irreversible losses of natural areas and damage to unique terrestrial and aquatic ecosystems are caused globally by the exploitation of mineral commodities, for example, due to the removal of substrates, changes to aquifers, destruction of springs, and concentration of pollutants in watercourses (Palmer et al. 2010; Murguía et al. 2016). These effects are prominent in Minas Gerais (which translates to “general mines”), where intense mining is conducted in some of the most important metal-rich regions in the world (Salgado and Carmo 2015). Thus, mining activities, even when located at considerable distances, are potential threats to PAs (Durán et al. 2013).

With hundreds of tailings dams in operation and a history of environmental disasters, the National Dam Safety Policy (PNSB, its abbreviation in Portuguese) was established in Brazil in 2010. The PNSB regulates all structures with at least one of the following characteristics: dam height ≥ 15 m; total reservoir capacity $\geq 3\,000\,000\text{ m}^3$; reservoir that contains hazardous waste; and high- or medium-associated damage potential (DPA, its abbreviation in Portuguese). The dam safety policy defined DPA based on the loss of human lives and the economic, social, and environmental impacts from a dam failure and established that the population should be informed of preventive and emergency actions and participation and social control mechanisms (Brasil 2010). All dams with a high DPA must have an emergency action plan that encompasses the geographical delimitation of potentially affected areas. The plan should include a flood map, which is an important planning and management tool, and be based on scenarios generated from a hypothetical breach of the dam, which is technically known as a “dam break” (Agência Nacional de Mineração, ANM 2017).

The need for awareness, participation, and social control in decision-making processes of environmental issues is explained both in international treaties, such as Principle 10, which was adopted in 1992 as a part of the Rio Declaration (United Nations Environment Programme, UNEP 2019), and in Brazilian public policies, as defined in the National Environmental Policy, National System of Conservation Units (SNUC, its abbreviation in Portuguese), and National Strategic Plan for Protected Areas (PNAP, its abbreviation in Portuguese) (Brasil 1981, 2000, 2007). When decision-making processes involve specialized knowledge and technical language, the lay public faces a relevant obstacle for their participation in democratic-participatory processes (Fonseca et al. 2012). This obstacle is evident in environmental licensing procedures and disaster risk management of engineering structures, such as tailings dams. Thus, Principle 10 highlights the importance of communities with

adequate access to information about hazardous materials and activities, including judicial and administrative mechanisms such as compensation and reparation for environmental damage. The objectives of the SNUC and PNAP include increasing public awareness for participation and social control over the creation, implementation and management of conservation units (CUs), and the adoption of precautionary measures in areas with indications of high environmental sensitivity and under threat.

From this perspective, we provide the following: (1) the scenario of environmental costs and liability of tailings dams in Minas Gerais State, which highlights the structures with the greatest damage potential, from the total deposited volume and the type of tailings; (2) the spatial overlap between tailings dams and their influence areas on PAs and PCAs; and (3) an overview of the cases of overlap, which indicate whether the PAs have a specific level of management to monitor or evaluate the damage potential associated with tailings dams.

MATERIALS AND METHODS

Minas Gerais is the fourth largest state in Brazil, has 586.5 thousand km^2 , which is an area slightly smaller than France, and harbors some of the main watersheds in southeastern Brazil. The state contains a megadiversity represented by more than 11 200 species of vascular plants, including the largest number of endemic plants (2245 spp.) and rare plants cataloged in the country (550 spp., sensu Giulietti et al. 2009), 43% of bird species, 36% of mammal species, 32% of reptile species, and 25% of amphibian species that occur in Brazil (Drummond et al. 2009; The Brazil Flora Group, BFG 2015).

We used official databases to obtain the following data to support our analyses: (1) data on PAs and PCAs, such as location, protection category, priority actions, geographic boundaries, buffer zones, and management plans (Instituto Chico Mendes de Conservação da Biodiversidade, ICMBio 2019; Instituto Estadual de Florestas, IEF 2019; Sistema Estadual de Meio Ambiente, SISEMA 2019). These data are available for public consultation via agencies responsible for information management in these areas (Box 1); and (2) data regarding the general characteristics of tailings dams, such as location, size, volume, and type of tailings deposited in the dam. These data are available for public consultation via regulatory and inspection agencies of mining activities throughout the country (Agência Nacional de Mineração, ANM 2019a; Fundação Estadual do Meio Ambiente, FEAM 2019).

For each tailings dam regulated by the PNSB (i.e., those with at least one of the characteristics: dam height ≥ 15 m; total reservoir capacity $\geq 3\,000\,000\text{ m}^3$; reservoir that

Box 1 Definition of the six types of priority conservation areas (PCAs) and protected areas (PAs) in Minas Gerais, southeastern Brazil	
State PCA	Establishes guidelines and criteria for the creation and management of PAs; recommendations for the revision of fiscal and financial instruments of conservation and proposals for the creation of new instruments; and the identification of lines of action that are aimed at improving the management of public policies for the protection of biodiversity (Minas Gerais 2002)
Federal PCA	Federal regulation recognizes the PCA for the formulation and implementation of public policies, programs, projects, and activities under the responsibility of the federal government (Ministério do Meio Ambiente, MMA 2018)
Conservation units: sustainable use	Federal law establishes that the basic objective is to reconcile nature conservation with the sustainable use of a portion of its natural resources (Brasil 2000). The categories identified in this study were Environmental Protection Area (APA, its abbreviation in Portuguese; equivalent to category V of the IUCN system) and Private Natural Heritage Reserve (RPPN, its abbreviation in Portuguese; equivalent to category VI of the IUCN system (Ministério do Meio Ambiente, MMA 2007a, 2019b)
Conservation units: strict protection	Federal law establishes that the basic objective is to preserve nature and that only the indirect use of natural resources is allowed. (Brasil 2000). The categories identified in this study were State/National Park, Natural Monument and Ecological Station, which are equivalent to categories II, III, and Ia, respectively, of the IUCN system (Ministério do Meio Ambiente, MMA 2007a, 2019b)
Buffer zone of conservation units	Federal law establishes that the area surrounding a conservation unit, where human activities are subject to specific norms and restrictions, serves the purpose of minimizing negative impacts on the PA (Brasil 2000). Conservation units, except for the APA and RPPN categories, must have a buffer zone
Special protection area (SPA)	State regulation establishes important areas for the conservation of biodiversity and water sources for water harvesting to ensure the perpetuity of the qualitative and quantitative characteristics of the water supply sources (Companhia de Saneamento de Minas Gerais, COPASA 2019a)

contains hazardous waste; and high or medium DPA), we generated an “influence area,” which consists of a territorial unit for the analysis of overlap with the PA and PCA. The area of influence was based on the concept of watersheds, which delimits landscape elements that are connected via water-mediated transfer (Pringle 2001). In a geographic information system (GIS) environment, the coordinates of the dams were superimposed on the drainage network, and from the point of contact with the watercourse, we selected a stretch with an extension of approximately 10 km downstream of the dam. We used a distance of 10 km due to a certain correspondence between the associated damage potential of a dam and the concept of a self-saving zone expressed in federal regulations on the management of mining dam safety (Agência Nacional de Mineração, ANM 2017). According to these regulations, a self-saving zone corresponds to the region downstream of the dam, which in emergencies, such as a dam failure, the time for intervention by competent authorities would be insufficient. The delimitation of the self-saving zone must adopt the greater of the following distances: an arrival time of the flood wave within 30 min or 10 km downstream of the dam. The 10 km stretch of the watercourse was used to select the areas of *ottobacias*, which was our territorial unit of analysis. Each tailings dam and “influence area” were spatially superimposed on the limits of the PA and PCA to contextualize the potential damage to biodiversity, natural areas, and formally recognized ecosystem services. Thus, the total number of dams (N) and their “influence area” (*ottobacias* in ha) for each of the six types of PCA and PA and their respective categories (Box 1) were taken into

account, considering the legal and administrative complexity of Brazil, where PAs under different regulations (municipal, state, and federal) may occur in a given territory (Delelis et al. 2010).

DISCUSSION

Magnitude of environmental costs and liability

Minas Gerais has 218 PNSB tailings dams, which is almost half of the total PNSB tailings dams registered in Brazil. In these 218 PNSB dams, 2.24 thousand million m³ of tailings, mainly those generated from the treatment and processing of iron ore (52%), gold (26%), phosphate (16%), and bauxite (2%), are deposited (refer to Fig. 1).

Dams are usually built in natural valleys, causing, in many cases, changes to the landscape via the suppression of forest fragments and cutting of soil and rocks, which are associated with drainage networks. In Brazil, the most common ore processing method consists of washing and sieving, which generate waste known as ‘slurry’ with an appearance similar to mud that is deposited in tailings dams (Instituto Brasileiro de Mineração, IBRAM 2016). The industrial waste generated during the processing of minerals causes severe impacts, mainly to aquatic ecosystems, which can persist for decades and spread over dozens of kilometers (Mebane et al. 2015; Cross et al. 2017).

The largest dams have heights greater than 80 m—the highest dam reaches 175 m—and store volumes greater than 10 million m³—the largest dam currently stores 400

Fig. 1 Total volume stored in the 218 tailings dams included in the National Dam Safety Policy (PNSB), Minas Gerais, southeastern Brazil

million m³. As an indicator of the scale of the socioenvironmental damage potential of these mega-structures, the greatest technological disaster of the mineral industry, which was caused by the Fundão tailings dam failure (Carmo et al. 2017), spilled “only” 43 million m³, which is less than 2% of the accumulated volume of mining tailings and industrial waste in the Quadrilátero Ferrífero (QF) region, Mariana Municipality, and Minas Gerais. This disaster caused 19 deaths and polluted 668 km of watercourses from the Doce River Basin to the Atlantic Ocean (Carmo et al. 2017; Queiroz et al. 2018). The socio-cultural effects of the disaster were equally catastrophic due to the loss of cultural heritage (century-old buildings and ways of life typical of the region), interruption of the public water supply, displacement of hundreds of families, loss of jobs, and strong decrease in tax collection in some municipalities, with consequent decline in the local economy (Sánchez et al. 2018; Freitas et al. 2019).

Acute changes in river morphology caused by tailing tsunamis, pollution of freshwater and saltwater, loss of biota and ecosystems, and loss of living and working places are listed among the main direct impacts caused by Fundão tailings dam failure (Sánchez et al. 2018). Some disasters generated by mining dam breaks have caused environmental catastrophes in important PAs, such as Doñana National Park (Spain), which is one of Europe’s most important wetlands and a UNESCO World Heritage Site (Pain et al. 1998), and the disaster related to the Fundão tailing dam affected Brazilian-protected coastal zones, such as Santa Cruz Wildlife Refuge and Comboios Biological Reserve, which are important locations for spawning sea turtles (Carmo et al. 2017).

Tailings dams: Environmental footprints

Best practices in engineering, geotechnics, safety, and socioenvironmental risk management issues addressed by the PNSB have the goal of avoiding new catastrophic events generated by dam failures. However, the high pollutant potential during the “regular” operation of dams should also be considered. Over decades of tailings deposition, silent and chronic contamination (Fig. 2) has progressed via hundreds of kilometers of watercourses. This contamination is caused by numerous reported leakage events, which are described as “accidents” or “anomalies” by regulatory agencies (Ávila 2012). Among the most frequent impact associated with tailings dams is the pollution of groundwater, rivers, and soil by metals and trace elements (such as, Cr, Fe, Mn, and Zn) due to the easy transport of very small tailings particles (< 150 µm), alteration of the turbidity of the water column, and silting of stretches of streams located downstream of the dams (Fig. 2). In addition, thousands of tons of chemical products, such as caustic soda, sodium hypochlorite, ether amines, fatty diamines, polyphenols, cyanide, and hydrogen peroxide, are employed annually for the treatment and processing of minerals. Some of these metals, trace elements, and chemical reagents are capable of severely altering the natural conditions of water bodies, soil, biota, ecosystems, and human health (refer to revision in Dudka and Adriano 1997; Araujo et al. 2005; Pearse 2005; Veado et al. 2006).

Fig. 2 Decades of deposition of tailings in dams caused silent and chronic hydric contamination. **a** Entry of a river (yellow arrow) that drains a watershed containing 12 PNSB tailings dams into a public water supply. **b** and **c** Situations of silting in watercourses. 2019 image. Adapted from *Google Earth Pro*

Conservation paradox

Of the 218 PNSB dams, 189 overlap with PCAs and PAs. In some situations, a given dam and/or its “influence area” overlapped with more than one PA/PCA (Fig. 3). The concentration of dams in the central region of the state, which is known as the QF, is intense. This region is a PCA

that is recognized at the state and federal levels as one of the most important PCAs for the conservation of biodiversity and ecosystem services, mainly because it harbors water sources used for public water supplies, hundreds of caves and dozens of rare plant species, and invertebrates associated with ironstone ranges and canga outcrops

Fig. 3 Concentration of tailings dams in Quadrilátero Ferrífero, central region of Minas Gerais State, and overlap between Protect areas (PAs) and Priority conservation areas (PCAs). Adapted from *Google Earth Pro*

(Ministério do Meio Ambiente, MMA 2007b; Ferreira et al. 2015; Carmo et al. 2018).

Financial and human resources were directed to ensure that the PCAs are public policy instruments with the objective of implementing appropriate measures for in situ conservation, including the indication of areas for the creation of CUs, the recovery of degraded areas and species overexploited or threatened with extinction, and the sustainable use of biodiversity components (Brasil 2007). However, most of the 189 PNSB tailings dams ($\approx 70\%$) and their influence areas are inserted precisely in priority areas for biodiversity conservation. These dams are always associated with other structures, such as open pit mines, wastes dump, industrial plants, and access roads, which represent numerous industrial complexes with high polluting potential.

A total of 45 dams located within CUs were observed, with three dams classified as strict protection (i.e., equivalent to IUCN Category Ia), which demonstrates a situation that is incompatible with both the conservation objectives of the PA and the premises of federal public policy (Brasil 2000). A total of 58 tailings dams exist in the buffer zones

Table 1 Priority conservation areas (PCAs) and protect areas (PAs) types and overlap between tailings dam (*N*) and influence areas (*otobacias* em ha). The limits of the different areas may overlap in some regions. %: relative to the 218 PNSB dams

PCA and PA types (<i>N</i>)	Overlap	
	Tailings dam <i>N</i> (%)	Influence areas (ha)
State PCAs (15)	157 (72)	58 755
Federal PCAs (17)	147 (67)	50 744
Conservation units: sustainable use/ IUCN, IV–VI (26)	42 (19)	13 830
Conservation units: strict protection/ IUCN, Ia–III (09)	3 (1)	1558
Buffer zone of conservation units (27)	58 (27)	14 649
Special protection areas, SPAs (06)	51 (23)	14 270

of these CUs, and 51 of these structures are located within a Special Protection Area (SPA) (Table 1). Even when the tailings dams are located outside the limits of the PA, 26 CUs and 5 SPAs are directly under the influence of the environmental impacts generated by these structures. This

situation occurs when a dam is located upstream of the PA; thus, pollutant materials are transported inside its boundary and follow the flow of the numerous watercourses within its influence area.

The dams accumulate extraordinary environmental costs and liability represented by approximately 2.24 thousand million m³ of tailings and industrial waste. However, the agencies in charge of the PAs have not developed adequate policies to establish management mechanisms and integrated monitoring to address a scenario of high socioenvironmental potential damage associated with tailings dams. This worrying scenario was also observed in a way for the SPAs, which provide essential ecosystem services, such as water recharge and storage, that contribute to the public supply of the third largest metropolitan region of Brazil, with an estimated population of approximately six million people. For the SPAs, no open access information that recognizes the conflicting use, monitoring of water risks associated with the accumulation of tailings, and preventive management of water sources located downstream from the dams was obtained. An exception was the publication of information related to the environmental damage caused by the recent B1 tailings dam failure, primarily because it prevented water harvesting on the day of the disaster, January 25, 2019 (Companhia de Saneamento de Minas Gerais, COPASA 2019b). For 9 of the 24 CUs, management plans and management councils existed. In only two management plans, information, albeit basic, about tailings dams and some associated risks, conflicting use with the PA, and some guidelines regarding the need to intensify inspection and monitoring of the impacts generated by the mining activity was provided to managers. Information related to the emergency action plan of the dams was not available in any of the cases. This situation may be the result of some interdependent factors, such as the historical power relations, particularly in Minas Gerais, among the mining industry, politicians, and decision makers (Meira et al. 2016; Milanez et al. 2019); the difficulty in establishing effective actions in the PA based on ecosystem data and explicit objectives for the management and reduction of impacts (Lima et al. 2005; Vieira et al. 2019); which are related to the giant gap between conservation science and policy decisions based on poor-quality data in Brazil (Azevedo-Santos et al. 2017).

In recent decades, the numerous environmental disasters in Brazil caused by mine tailings dam failures have caused unprecedented damage to natural, social, and economic heritage. Two of the largest dam failures occurred in Minas Gerais and became known worldwide. The first failure occurred in 2015, as reported in this paper. The second disaster occurred in 2019 and caused 259 fatalities; 11 people are still missing. 13 million m³ of iron ore tailings spilled from the B1 tailings dam, Córrego do Feijão Mine

in Brumadinho. The tailings plume traveled at least 110 km along the Paraopeba River, which is a tributary of the São Francisco River (Ministério Público de Minas Gerais, MPMG 2020), and an estimated 424 communities were affected, including indigenous, quilombolas, foresters, and artisanal fishers, whose living and working conditions were affected in the different territories (Freitas et al. 2019). Both disasters affected several PCAs and PAs, which created urgency regarding the incorporation of the experiences and lessons learned into conservation planning. However, this process can be very complicated because it involves changes in both the mindset of individuals and the management of the institutions involved (Grantham et al. 2010).

Lessons (not yet) learned

Even after the environmental disasters caused by the break of tailings dams, the socioenvironmental risk situation remains critical in Brazil, especially in Minas Gerais. Currently, the Federal Regulatory Agency interdicted 54 Brazilian dams that did not submit or certify a Statement of Dam Stability Conditions, which is a mandatory document that should be prepared by the entrepreneur. Of the Brazilian states, Minas Gerais has the highest risk situation because it contains 33 (61%) interdicted dams. Of these dams, 19 have emergency levels between 1 and 3, which represent dams with “potential compromised safety” to dams at “imminent risk” of failure (Agência Nacional de Mineração, ANM 2019b; Ministério Público de Minas Gerais, MPMG 2020). Considering the critical scenario, we have identified an urgent need to implement effective actions and propose the following:

- public disclosure of the Emergency Action Plans for Dams, including versions understandable by the lay public to instruct and train managers and advisory councils, communities, city halls, and civil defenses;
- integrated assessment of the effectiveness of the PA and PCA as a strategy to reduce the loss of biodiversity and ecosystems services, based on the integration of information generated by regulatory agencies, licensors, managers, scientists, and communities. For the evaluation of effectiveness, the high potential of socioenvironmental damage related to the hundreds of tailings dams should be considered;
- development of an open access database to support a risk management program and strategic action plan in the case of a tailings dam break. The database should include the location of the scenarios generated from a hypothetical dam break, which is technically known as a “Dam Break.” The database should also include the environmental attributes and objects of conservation in

the PA, including its buffer zone, and the PCAs located downstream from the dams;

- update the mitigation and compensation actions provided in the Management Plans;
- revision or creation of fiscal and administrative instruments for allocating resources (material, financial, and human) to perform regular technical–operational training to address emergency situations that involve managers and advisory councils, communities, city halls, and civil defenses.

Recalling Principle 10 established in the Rio Declaration on Environment and Development in 1992 and the Brazilian objectives established in the SNUC and PNAP, only through raising public awareness regarding participation in the processes of creation, implementation, and management of PAs will it be possible to advance the development of measures to contain threats to biodiversity and ecosystem services and mitigate negative impacts, with cumulative and synergistic effects, generated by large-scale mining.

Acknowledgements We especially thank the anonymous reviewers and donors.

REFERENCES

- Agência Nacional de Mineração, ANM. 2017. *Ordinance n° 70.389, 17 May, 2017* (in Portuguese). <http://www.anm.gov.br/acao-a-informacao/legislacao/portarias-do-diretor-geral-do-dnmp/portarias-do-diretor-geral/portaria-70-389-de-2017/view>. Accessed 15 Nov 2019.
- Agência Nacional de Mineração, ANM. 2018. *Brazilian Mineral Yearbook: Main Metallic Substances* (in Portuguese). http://www.anm.gov.br/dnmp/publicacoes/serie-estatisticas-e-economia-mineral/anuario-mineral/anuario-mineral-brasileiro/amb_2018.pdf. Accessed 15 Nov 2019.
- Agência Nacional de Mineração, ANM. 2019a. *National Register of Mining Dams* (in Portuguese). <http://www.anm.gov.br/assuntos/barragens/pasta-cadastro-nacional-de-barragens-de-mineracao/cadastro-nacional-de-barragens-de-mineracao>. Accessed 20 Sep 2019.
- Agência Nacional de Mineração, ANM. 2019b. *ANM bans 54 dams for lack of stability condition* (in Portuguese). <http://www.anm.gov.br/noticias/anm-interdita-54-barragens-por-falta-de-condicao-de-estabilidade>. Accessed 15 Nov 2019.
- Araujo, A.C., P.R.M. Viana, and A.E.C. Peres. 2005. Reagents in iron ores flotation. *Minerals Engineering* 18: 219–224. <https://doi.org/10.1016/j.mineng.2004.08.023>.
- Ávila, J.P. 2012. *Tailings dams in Brazil*. Rio de Janeiro: CBDB (in Portuguese).
- Azevedo-Santos, V.M., P.M. Fearnside, C.S. Oliveira, A.A. Padial, F.M. Pelicice, D.P. Lima Jr., D. Simberloff, T.E. Lovejoy, et al. 2017. Removing the abyss between conservation science and policy decisions in Brazil. *Biodiversity Conservation* 26: 1745–1752. <https://doi.org/10.1007/s10531-017-1316-x>.
- Brasil. 1981. *Law n° 6.938, 31 August, 1981* (in Portuguese). <https://www2.camara.leg.br/legin/fed/lei/1980-1987/lei-6938-31-agosto-1981-366135-norma-actualizada-pl.pdf>. Accessed 20 Jan 2020.
- Brasil. 2000. *Law n° 9.985, 18 July, 2000* (in Portuguese). http://www.planalto.gov.br/ccivil_03/LEIS/L9985.htm. Accessed 15 Nov 2019.
- Brasil. 2007. *Ordinance n° 9, 1 December, 2007* (in Portuguese). <http://pesquisa.in.gov.br/imprensa/jsp/visualiza/index.jsp?data=24/01/2007&jornal=1&pagina=55&totalArquivos=64>. Accessed 15 Nov 2019.
- Brasil. 2010. *Law n° 12.334, 20 September, 2010* (in Portuguese). http://www.planalto.gov.br/ccivil_03/_Ato2007-2010/2010/Lei/L12334.htm. Accessed 15 Nov 2019.
- Carmo, F.F., L.H.Y. Kamino, R. Tobias Junior, I.C. Campos, F.F. Carmo, G. Silvino, K.J.S.X. Castro, M.L. Mauro, et al. 2017. Fundão tailings dam failures: The environment tragedy of the largest technological disaster of Brazilian mining in global context. *Perspectives in Ecology and Conservation* 15: 145–151. <https://doi.org/10.1016/j.pecon.2017.06.002>.
- Carmo, F.F., R.C. Mota, L.H.Y. Kamino, and C.M. Jacobi. 2018. Check-list of vascular plant communities on ironstone ranges of south-eastern Brazil: Dataset for conservation. *Biodiversity Data Journal* 6: e27032. <https://doi.org/10.3897/BDJ.6.e27032>.
- Carvalho, C.J.B. 2009. Endemism patterns and biodiversity conservation. *Megadiversidade* 5: 77–86 (in Portuguese).
- Centro Nacional de Pesquisa e Conservação de Cavernas, CECAV. 2018. *Statistical Yearbook of Brazilian Speleological Heritage*. Brasília: Instituto Chico Mendes de Conservação da Biodiversidade (in Portuguese). <http://www.icmbio.gov.br/cecaav/destaques/90-anuario-estatistico-do-patrimonio-espeleologico-brasileiro-2018.html>. Accessed 15 Nov 2019.
- Companhia de Saneamento de Minas Gerais, COPASA. 2019a. *Protection areas* (in Portuguese). <http://www.copasa.com.br/wps/portal/internet/meio-ambiente/areas>. Accessed 14 Feb 2019.
- Companhia de Saneamento de Minas Gerais, COPASA. 2019b. *RMBH water supply situation* (in Portuguese). <http://www.copasa.com.br/wps/portal/internet/imprensa/noticias/informacoes-sobre-abastecimento/2019/setembro19co/coper-rmbh-ie14549/>. Accessed 14 Sep 2019.
- Cross, A.T., J.C. Stevens, and K.W. Dixon. 2017. One giant leap for mankind: Can ecopoiesis avert mine tailings disasters? *Plant and Soil* 421: 1–5.
- Delelis, C.J., T. Rehder, and T.M. Cardoso. 2010. Mosaics of protected areas: Reflections and proposals of Franco-Brazilian cooperation. Brasília: Ministério do Meio Ambiente, MMA; Embaixada da França no Brasil; CDS UnB (in Portuguese). https://www.mma.gov.br/estruturas/240/_arquivos/livro_mosaicos_portugues_montado_240.pdf. Accessed 20 Jan 2020.
- Drummond, G.M., C.S. Martins, M.B. Greco, and F. Vieira. 2009. *Minas Biota: Diagnosis of knowledge about biodiversity in the State of Minas Gerais—Minas subsidy to the Biota Program*. Belo Horizonte: Fundação Biodiversitas (in Portuguese).
- Dudka, S., and D.C. Adriano. 1997. Environmental impacts of metal ore mining and processing: A review. *Journal of Environmental Quality* 26: 590–602. <https://doi.org/10.2134/jeq1997.00472425002600030003x>.
- Durán, A.P., J. Rauch, and K.J. Gaston. 2013. Global spatial coincidence between protected areas and metal mining activities. *Biological Conservation* 160: 272–278. <https://doi.org/10.1016/j.biocon.2013.02.003>.
- Ferreira, R.L., M.P.O. Oliveira, and M.S. Silva. 2015. Underground biodiversity in iron geosystems. In *Iron geosystems of Brazil: Priority areas for the conservation of geological and biological diversity, cultural heritage and environmental services*, ed. F.F. Carmo, and L.H.Y. Kamino, 195–231. Belo Horizonte: 3i Editora (in Portuguese). <https://www.institutopristino.org.br/>

- [wp-content/uploads/2016/03/Geossistemas-ferruginosos-no-Brasil-CD.pdf](#). Accessed 15 Nov 2019.
- Fonseca, I.F.D., M. Bursztyn, and A.M.M.D. Moura. 2012. Technical knowledge, public policy and participation: The case of the National Council on the Environment. *Revista de Sociologia e Política* 20: 183–198. <https://doi.org/10.1590/S0104-44782012000200013> (in Portuguese).
- Freitas, C.M., C. Barcellos, C.I.R.F. Asmus, M.A. Silva, and D.R. Xavier. 2019. From Samarco in Mariana to Vale in Brumadinho: Mining dam disasters and Public Health. *Reports in Public Health* 35: e00052519. <https://doi.org/10.1590/0102-311x00052519>.
- Fundação Estadual do Meio Ambiente, FEAM. 2019. *Dam management* (in Portuguese). <http://www.feam.br/monitoramento/gestao-de-barragens>. Accessed 15 Nov 2019.
- Gaston, K.J., S.F. Jackson, L. Cantu-Salazar, and G. Cruz-Piñón. 2008. The ecological performance of protected areas. *Annual Review of Ecology, Evolution and Systematics* 39: 93–113. <https://doi.org/10.1146/annurev.ecolsys.39.110707.173529>.
- Giulietti, A., A. Rapini, M. Andrade, and L.E.S. Queiroz. 2009. *Rare plants in Brazil*. Belo Horizonte: Conservação Internacional-Universidade Estadual de Feira de Santana (in Portuguese).
- Graham, H.S., M. Bode, E. McDonald-Madden, E.T. Game, A.T. Knight, and H.P. Possingham. 2010. Effective conservation planning requires learning and adaptation. *Frontiers in Ecology and the Environment* 8: 431–437. <https://doi.org/10.1890/080151>.
- Hansen, A.J., and R. DeFries. 2007. Ecological mechanisms linking protected areas to surrounding lands. *Ecological Applications* 17: 974–988. <https://doi.org/10.1890/05-1098>.
- Instituto Brasileiro de Mineração, IBRAM. 2016. *Mining Waste Management and Monitoring*. Brasília: IBRAM (in Portuguese). <http://www.ibram.org.br/sites/1300/1382/00006222.pdf>. Accessed 20 Sep 2019.
- Instituto Chico Mendes de Conservação da Biodiversidade, ICMBio. 2019. *List of protected areas with management plan* (in Portuguese). <http://www.icmbio.gov.br/portal/planosmanejo>. Accessed 10 Sep 2019.
- Instituto Estadual de Florestas, IEF. 2019. *Management Councils* (in Portuguese). <http://www.ief.mg.gov.br/unidades-de-conservacao/gestao/1725-conselhos>. Accessed 12 Sep 2019.
- Jacobi, C.M., F.F. Carmo, and I.C. de Campos. 2011. Soaring extinction threats to endemic plants in Brazilian metal-rich regions. *Ambio* 40: 540–543. <https://doi.org/10.1007/s13280-011-0151-7>.
- Lima, G.S., G.A. Ribeiro, and W. Gonçalves. 2005. Management effectiveness of conservation units with integral protection in Minas Gerais. *Revista Árvore* 29: 647–653. <https://doi.org/10.1590/S0100-67622005000400017> (in Portuguese).
- Margules, C.R., and R.L. Pressey. 2000. Systematic conservation planning. *Nature* 405: 243–253. <https://doi.org/10.1038/35012251>.
- Mebane, C.A., R.J. Eakins, B.G. Fraser, and W.J. Adams. 2015. Recovery of a mining-damaged stream ecosystem. *Elementa: Science of the Anthropocene* 3: 000042. <https://doi.org/10.12952/journal.elementa.000042>.
- Meira, R.M., A.L. Peixoto, M.A. Coelho, A.P. Ponzo, V.G. Esteves, M.C. Silva, P.E.A.S. Câmara, and J.A. Meira-Neto. 2016. Brazil's mining code under attack: Giant mining companies impose unprecedented risk to biodiversity. *Biodiversity and Conservation* 2: 407–409. <https://doi.org/10.1007/s10531-016-1050-9>.
- Milanez, B., L. Magno, and R.G. Pinto. 2019. From weak politics to private policies: The role of the mining sector in the transformations of environmental policy in Minas Gerais State, Brazil. *Cadernos de Saúde Pública* 35: e00051219. <https://doi.org/10.1590/0102-311x00051219>.
- Minas Gerais. 2002. *Normative Deliberation COPAM n° 55, 13 June, 2002* (in Portuguese). <http://www.siam.mg.gov.br/sla/download.pdf?idNorma=137>. Accessed 10 July 2019.
- Ministério do Meio Ambiente, MMA. 2007a. *National Report on Protected Areas in Brazil*. Brasília: Secretaria de Biodiversidade e Florestas, Departamento de Áreas Protegidas (in Portuguese). https://www.mma.gov.br/estruturas/sbf2008_dap/_publicacao/149_publicacao16122010110837.pdf. Accessed 1 Oct 2019.
- Ministério do Meio Ambiente, MMA. 2007b. *Priority areas for the conservation, sustainable use and benefit sharing of Brazilian biological diversity*. https://www.mma.gov.br/estruturas/chm/_arquivos/Priority_Area_Book.pdf. Accessed 10 July 2019.
- Ministério do Meio Ambiente, MMA. 2018. *Ordinance n° 463, 18 December, 2018* (in Portuguese). http://www.in.gov.br/materia/-/asset_publisher/Kujrw0TZC2Mb/content/id/55881195/do1-2018-12-19-portaria-n-463-de-18-de-dezembro-de-2018-55880954. Accessed 20 July 2019.
- Ministério do Meio Ambiente, MMA. 2019a. *Brazilian Biodiversity* (in Portuguese). <https://www.mma.gov.br/biodiversidade/biodiversidade-brasileira.html>. Accessed 10 July 2019.
- Ministério do Meio Ambiente, MMA. 2019b. *National Register of Conservation Units* (in Portuguese). <https://www.mma.gov.br/areas-protegidas/cadastro-nacional-de-ucs>. Accessed 20 July 2019.
- Ministério Público de Minas Gerais, MPMG. 2020. *Dams situation in Minas Gerais* (in Portuguese). <https://www.mpmg.mp.br/comunicacao/noticias/mpmg-acompanha-situacao-da-barragem-de-mineracao-em-brumadinho.htm>. Accessed 25 Jan 2020.
- Murguía, D.I., S. Bringezu, and R. Schaldach. 2016. Global direct pressures on biodiversity by large-scale metal mining: Spatial distribution and implications for conservation. *Journal of Environmental Management* 180: 409–420. <https://doi.org/10.1016/j.jenvman.2016.05.040>.
- Naughton-Treves, L., M.B. Holland, and K. Brandon. 2005. The role of protected areas in conserving biodiversity and sustaining local livelihoods. *Annual Review of Environment and Resources* 30: 219–252. <https://doi.org/10.1146/annurev.energy.30.050504.164507>.
- Pain, D.J., A. Sanchez, and A.A. Meharg. 1998. The Doñana Ecological Disaster: Contamination of a world heritage estuarine marsh ecosystem with acidified pyrite mine waste. *Science of the Total Environment* 222: 45–54. [https://doi.org/10.1016/S0048-9697\(98\)00290-3](https://doi.org/10.1016/S0048-9697(98)00290-3).
- Palmer, M.A., E.S. Bernhardt, W.H. Schlesinger, K.N. Eshleman, E. Foufoula-Georgiou, M.S. Hendryx, A.D. Lemly, G.E. Likens, et al. 2010. Mountaintop mining consequences. *Science* 327: 148–149. <https://doi.org/10.1126/science.1180543>.
- Pearse, M.J. 2005. An overview of the use of chemical reagents in mineral processing. *Minerals Engineering* 18: 139–149. <https://doi.org/10.1016/j.mineng.2004.09.015>.
- Pringle, C.M. 2001. Hydrologic connectivity and the management of biological reserves: A global perspective. *Ecological Applications* 11: 981–998. <https://doi.org/10.2307/3061006>.
- Queiroz, H.M., G.N. Nóbrega, T.O. Ferreira, L.S. Almeida, T.B. Romero, S.T. Santaella, A.F. Bernardino, and X.L. Otero. 2018. The Samarco Mine Tailing Disaster: A possible time-bomb for heavy metals contamination? *Science of the Total Environment* 637–638: 498–506. <https://doi.org/10.1016/j.scitotenv.2018.04.370>.
- Salgado, A.A.R., and F.F. Carmo. 2015. Quadrilátero Ferrífero: A beautiful and neglected landscape between the gold and iron ore reservoirs. In *Landscapes and Landforms of Brazil*, ed. B.C. Vieira, A.A.R. Salgado, and L. Santas, 319–330. London: Springer. https://doi.org/10.1007/978-94-017-8023-0_29.

- Sánchez, L.E., K. Alger, L. Alonso, G. Barbosa, M.C.W. Brito, F.V. Laureano, P. May, H. Roeser, and Y. Kakabadse. 2018. *Impacts of the Fundão Dam failure. A pathway to sustainable and resilient mitigation*. Rio Doce Panel Thematic Report No. 1. Gland: IUCN. <https://doi.org/10.2305/IUCN.CH.2018.18.en>.
- Sistema Estadual de Meio Ambiente, SISEMA. 2019. *Spatial Data Infrastructure of the State Environment and Water Resources System*. Belo Horizonte: IDE-Sisema (in Portuguese). <http://www.idesisema.meioambiente.mg.gov.br>. Accessed 16 Oct 2019.
- The Brazil Flora Group, BFG. 2015. Growing knowledge: An overview of seed plant diversity in Brazil. *Rodriguésia* 66: 1085–1113. <https://doi.org/10.1590/2175-7860201566411>.
- United Nations Environment Programme, UNEP. 2019. *Principle 10*. <https://www.unenvironment.org/civil-society-engagement/partnerships/principle-10>. Accessed 10 July 2019.
- Veado, M.A.R.V., I.A. Arantes, A.H. Oliveira, M.R.M.G. Almeida, R.A. Miguel, M.I. Severo, and H.L. Cabaleiro. 2006. Metal pollution in the environment of Minas Gerais State-Brazil. *Environmental Monitoring and Assessment* 117: 157–172. <https://doi.org/10.1007/s10661-006-8716-9>.
- Vieira, R.R.S., R.L. Pressey, and R. Loyola. 2019. The residual nature of protected areas in Brazil. *Biological Conservation* 233: 152–161. <https://doi.org/10.1016/j.biocon.2019.02.010>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

AUTHOR BIOGRAPHIES

Luciana Hiromi Yoshino Kamino (✉) PhD, is a Coordinator of Research and Development Projects at the Instituto Prístino. Her research interests include conservation biology, botany, and environmental analysis.

Address: Instituto Prístino, Rua Santa Maria Goretti, 86, Barreiro, Belo Horizonte, Minas Gerais CEP 30642-020, Brazil.

e-mail: luciana@institutopristino.org.br

Eric Oliveira Pereira PhD, is a Geographer at the Instituto Prístino. His research interests include environmental systems modeling and environmental analysis.

Address: Instituto Prístino, Rua Santa Maria Goretti, 86, Barreiro, Belo Horizonte, Minas Gerais CEP 30642-020, Brazil.

e-mail: ericpereiraufmg@gmail.com

Flávio Fonseca do Carmo PhD, is a Coordinator of Research and Development Projects at the Instituto Prístino. His research interests include conservation biology, ecology, and environmental governance.

Address: Instituto Prístino, Rua Santa Maria Goretti, 86, Barreiro, Belo Horizonte, Minas Gerais CEP 30642-020, Brazil.

e-mail: flavio@institutopristino.org.br