Published in final edited form as:

Nat Neurosci. 2020 July ; 23(7): 842–853. doi:10.1038/s41593-020-0628-4.

Tumor Necrosis Factor Overcomes Immune Evasion in p53-Mutant Medulloblastoma

Alexandra Garancher¹, Hiromichi Suzuki², Svasti Haricharan³, Lianne Q. Chau¹, Meher Beigi Masihi¹, Jessica M. Ruser¹, Paula S. Norris³, Florent Carrette³, Megan M. Romero⁴, Sorana A. Morrissy², Patryk Skowron², Florence M.G. Cavalli², Hamza Farooq², Vijay Ramaswamy⁵, Steven J.M. Jones⁶, Richard A. Moore⁶, Andrew J. Mungall⁶, Yussanne Ma⁶, Nina Thiessen⁶, Yisu Li⁶, Alaide Morcavallo⁷, Lin Qi^{8,9}, Mari Kogiso⁸, Yuchen Du^{8,9}, Patricia Baxter⁸, Jacob J. Henderson¹⁰, John R. Crawford¹¹, Michael L. Levy¹², James M. Olson¹³, Yoon-Jae Cho¹⁰, Aniruddha J. Deshpande¹, Xiao-Nan Li^{8,9}, Louis Chesler⁷, Marco A. Marra⁶, Harald Wajant¹⁴, Oren J. Becher⁴, Linda M. Bradley³, Carl F. Ware³, Michael D. Taylor^{2,15}, Robert J. Wechsler-Reya^{1,*}

¹Tumor Initiation & Maintenance Program, NCI-Designated Cancer Center, Sanford Burnham Prebys Medical Discovery Institute, La Jolla, California, USA

²Developmental and Stem Cell Biology Program, The Hospital for Sick Children, Toronto, Ontario, Canada

³Tumor Microenvironment and Cancer Immunology Program, NCI-Designated Cancer Center and the Infectious and Inflammatory Disease Center, Sanford Burnham Prebys Medical Discovery Institute, La Jolla, California, USA

⁴Department of Pediatrics, Northwestern University, Chicago, Illinois, USA

⁵Division of Haematology/Oncology and Department of Paediatrics, Hospital for Sick Children, Toronto, Ontario, Canada.

⁶Canada's Michael Smith Genome Sciences Centre, BC Cancer, Vancouver, British Columbia, Canada

⁷Division of Clinical Studies, The Institute of Cancer Research, London, UK

⁸Brain Tumor Program, Texas Children's Cancer Center, Department of Pediatrics, Baylor College of Medicine, Houston, Texas, USA

* Address correspondence to rwreya@sbpdiscovery.org.

AUTHOR CONTRIBUTIONS

A.G. and R.J.W.-R. conceived and designed experiments. A.G. and L.Q.C., M.B.M. J.M.R. performed experiments. A.G. H.S., V.R., S.H., S.A.M., P.S., F.M.G.C., H.F., S.J.M.J., R.A.M., A.J.M., Y.M., N.T., Y.L., M.M. analyzed data. L.Q., M.K., Y.D., P.B., J.J.H., J.R.C., M.M.R., O.J.B., M.L.L., J.M.O., Y.-J.C., A.M., L.C., A.J.D. and X.-N.L. provided patient samples, cell lines or xenografts and associated data. P.S.N., C.F.W. and H.W. provided reagents and protocols. P.S.N., F.C., L.M.B., C.F.W., M.D.T. discussed results and strategy. R.J.W.-R. supervised the study. A.G. and R.J.W.-R. wrote the manuscript, which was revised and approved by all authors.

COMPETING INTERESTS

Sanford Burnham Prebys Medical Discovery Institute has filed a patent application with R.J.W.-R., A.G. and C.F.W. listed as inventors. PCT application serial number PCT/US2018/048916 was filed on August 30, 2018, and is currently pending. This application covers the potential of p53 as a biomarker for responsiveness to T cell based immunotherapy, and the use of TNF receptor superfamily ligands to increase MHC expression and enhance immunotherapy.

⁹Program of Precision Medicine PDOX Modeling of Pediatric Tumors, Ann & Robert H. Lurie Children's Hospital of Chicago, Department of Pediatrics, Northwestern University, Chicago, Illinois, USA

¹⁰Department of Pediatrics, Oregon Health & Science University, Portland, Oregon, USA

¹¹Departments of Pediatrics and Neurosciences, University of California San Diego – Rady Children's Hospital, San Diego, California, USA

¹²Department of Neurosurgery, University of California San Diego – Rady Children's Hospital, San Diego, California, USA

¹³Clinical Research Division, Fred Hutchinson Cancer Research Center, Seattle, Washington, USA

¹⁴Division of Molecular Internal Medicine Department of Internal Medicine II, University Hospital Würzburg, Würzburg, Germany

¹⁵Division of Neurosurgery, Hospital for Sick Children, Toronto, Ontario, Canada.

Abstract

Many immunotherapies act by enhancing the ability of cytotoxic T cells to kill tumor cells. Killing depends on T cell recognition of antigens presented by class I major histocompatibility complex (MHC-I) proteins on tumor cells. Here we show that medulloblastomas lacking the p53 tumor suppressor do not express surface MHC-I, and are therefore resistant to immune rejection. Mechanistically, this is because p53 regulates expression of the peptide transporter Tap1 and the aminopeptidase Erp1, which are required for MHC-I trafficking to the cell surface. In vitro, tumor necrosis factor (TNF) or lymphotoxin- β receptor agonist (LT β Rag) can rescue expression of Erp1, Tap1 and MHC-I on p53-mutant tumor cells. In vivo, low doses of TNF prolong survival and synergize with immune checkpoint inhibitors to promote tumor rejection. These studies identify p53 as a key regulator of immune evasion, and suggest that TNF could be used to enhance sensitivity of tumors to immunotherapy.

Medulloblastoma is the most common malignant brain tumor in children. Surgery, radiation and chemotherapy have improved outcomes, but approximately 30% of patients remain incurable, and survivors suffer severe long-term side effects from these therapies. Improved approaches to treating medulloblastoma are therefore critical.

One marker associated with poor prognosis in medulloblastoma is p53 (encoded by the *TP53* gene in humans and the *Trp53* gene in mice). Newly diagnosed medulloblastoma patients whose tumors exhibit Sonic hedgehog (SHH) pathway activation and *TP53* mutations have nearly 100 per cent mortality ¹. *TP53* mutations are even more common in recurrent medulloblastoma, and patients with MYC amplification and p53 pathway defects at relapse at all die of rapidly progressive disease ^{2,3}. Thus, novel therapies would be particularly beneficial for medulloblastoma patients with *TP53* mutations.

Immunotherapy has emerged as a powerful approach to treating cancer. Antagonists of immune checkpoint regulators, T lymphocytes engineered to recognize tumor antigens, and

vaccines that amplify tumor-specific lymphocytes are being tested against a variety of human malignancies⁴⁻⁷. Although some remarkable responses have been reported, only a subset of patients benefit from these therapies, and the mechanisms that underlie resistance are poorly understood⁸. As immunotherapies begin to undergo clinical testing for medulloblastoma^{9,10}, it would be valuable to identify biomarkers of responsiveness and strategies for overcoming resistance in this tumor. With this in mind, we have begun to characterize anti-tumor immune responses in genetically engineered mouse models of medulloblastoma.

RESULTS

p53-mutant medulloblastomas are resistant to T cell-mediated rejection

We recently created animal models of medulloblastoma by isolating neural stem cells from the neonatal cerebellum, infecting them with viruses encoding oncogenes, and transplanting them into the cerebellum of naïve mice. One model, termed MP, uses stem cells expressing **Myc** and a dominant-negative form of **p53** (DNp53, a fragment of p53 that binds to the wild type protein and interferes with multimerization)¹¹⁻¹³; another, termed MG, uses cells transduced with **Myc** and **Gfi1**, a transcription factor whose expression is activated by enhancer hijacking in a subset of medulloblastoma patients¹⁴. In previous studies these models were established in immunodeficient (**NOD-SCID** interleukin-2 receptor-gamma-deficient, or **NSG**) mice, but in order to use them for studies of immunotherapy, we transplanted them into an immunocompetent strain, albino **C57BL/6** (**aB6**). **aB6** mice are identical to conventional **C57BL/6** mice except for the presence of a tyrosinase mutation, which results in an absence of pigmentation in skin and hair¹⁵; this lack of pigmentation facilitates bioluminescent imaging of tumor growth. **MP** tumors grew with similar kinetics and 100% penetrance in both immunodeficient (**NSG**) and immunocompetent (**aB6**) hosts (Figure 1a-b). In contrast, **MG** tumors were only able to grow efficiently in immunodeficient (**NSG**) hosts : only 2/45 (4.4%) **aB6** mice transplanted with **MG** tumor cells went on to develop tumors, and these developed with much longer latency than those in **NSG** mice (Figure 1c-d).

To determine if the failure of **MG** tumors to grow in **aB6** mice was mediated by the immune system, we isolated tissue from mice during the first week after transplantation and analyzed the number of tumor cells and infiltrating immune cells. As shown in Figure 1e, when mice were transplanted with **MP** tumors, the number of tumor cells increased over the first 7 days, but when mice were transplanted with **MG** tumors the number of tumor cells steadily decreased. There were no marked differences in the proportions of **B220+** B cells, **CD11b+** macrophages or **NK1.1+** natural killer cells infiltrating **MP** and **MG** tumors (Extended Data 1a-c). However, there was a significant difference in the proportion of **CD3+** and **CD8+** T cells (Figure 1f and Extended Data 1d): whereas mice transplanted with **MP** tumors showed no change in T cell numbers, mice transplanted with **MG** tumors exhibited a sharp spike in **CD3+** and **CD8+** T cells 2-5 days after transplantation. Consistent with these findings, depletion of T cells using anti-Thy-1 antibodies allowed **MG** tumors to grow in **aB6** mice (Figure 1g-i). Similar results were seen following depletion of cytotoxic T cells using anti-

CD8 antibodies (Extended Data 1e-g) These studies suggest that the failure of MG tumors to grow in immunocompetent mice is due to rejection by T cells.

Inactivation of p53 results in loss of cell surface MHC-I

The difference in ability of MP and MG tumors to grow in immunocompetent mice could reflect an enhanced ability of MG tumors to activate T cells or a decreased tendency of MP tumors to do so. Since both tumors are generated by transplantation of stem cells engineered to express Myc and one other protein (DNp53 or Gfi1), we tested the consequences of transducing MP tumors with Gfi1 (MP+G) and MG tumors with DNp53 (MG+P). Whereas expression of Gfi1 had no effect on the growth of MP tumors (Extended Data 1h-i), expression of DNp53 had a dramatic effect on MG tumors, allowing them to grow in immunocompetent (aB6) mice (Figure 2a-b). These studies suggested that inactivation of p53 renders tumor cells resistant to rejection by T cells.

To understand the mechanisms underlying this resistance, we analyzed MP and MG tumors for expression of molecules known to regulate immune responses. We observed no differences in expression of molecules that have been reported to regulate T cell responses, including cytotoxic T-lymphocyte associated protein 4 (Ctla4), Arginase 1 (Arg1), inducible nitric oxide synthase (iNOS, Nos2), indoleamine 2,3-dioxygenase (Ido1), transforming growth factor beta (TGF β), or interleukin-10 (IL-10) (Extended Data 1j). Although p53 has been shown to regulate programmed cell death ligand 1 (PD-L1, CD274) in other tumor types^{16,17}, we saw no differences in expression of this molecule in MP vs. MG tumors. Molecules that regulate activation of T-cells and dendritic cells, including OX40 ligand (OX40L, CD252, Tnfsf4), CD137 ligand (CD137L, 4-1BB-L, Tnfsf9), CD40, glucocorticoid-induced TNF-related ligand (GITRL, Tnfsf18), CD25 (IL2RA), CD62 ligand (CD62L, Sell), B Lymphocyte activation antigen B7-1 (CD80) and B lymphocyte activation antigen B7-2 (CD86) were also not differentially expressed (Extended Data 1k).

We did observe one striking difference between these tumors: whereas MG tumors expressed significant amounts of major histocompatibility complex I (MHC-I) on their surface, MP tumors expressed virtually none (Figure 2c). The lack of MHC-I was not a property of the stem cells from which these tumors were made, since cerebellar stem cells express substantial amounts of MHC-I (Extended Data 2a). Rather, the differences in MHC-I expression between MP and MG tumor cells were attributable to differences in p53: expression of DNp53 in MG tumors caused a marked downregulation of MHC-I (Figure 2d), while expression of WT p53 in MP tumors (generated using stem cells from p53 knockout mice, MPKO) increased levels of MHC-I (Figure 2e). The decrease in MHC-I was not specific to expression of DNp53, since it was also seen following shRNA-mediated knockdown of p53, both in MG tumor cells (Extended Data 2b and c) and in the human Group 3 MB cell line HD-MB03¹⁸ (Extended Data 2d and e). Thus, reduction of p53 expression or function results in loss of MHC-I.

To determine if p53 regulation of MHC-I expression is restricted to Myc-driven medulloblastoma, we examined tumors from conditional Patched-1 knockout (Math1-CreER^{T2}; Ptch1^{flox/flox}) mice, a model for SHH-driven medulloblastoma¹⁹. Like MG tumors, conditional Ptch1 knockout tumors are unable to grow following transplantation into

immunocompetent mice, but their growth can be restored by overexpression of DNp53 (Extended Data 2f), which causes downregulation of MHC-I expression (Extended Data 2g). Medulloblastoma patient-derived xenografts (PDXs) with p53 mutations, including both SHH and Group 3 tumors, also showed markedly lower levels of MHC-I (called HLA-I in humans) than PDXs expressing WT p53. Finally, murine and human diffuse intrinsic pontine glioma (DIPG) cells with WT p53 expressed significant amounts of MHC-I, whereas p53 mutant DIPG cells had little or no MHC-I on their surface (Extended Data 3k-n). Together, these data suggest that p53 function is necessary for expression of cell surface MHC-I. Although a link between p53 and MHC-I has been described in carcinoma and sarcoma cell lines^{20,21}, it has not been shown in brain tumors, and its significance for immune evasion has not been investigated *in vivo*.

p53 regulates expression of Erap1 and Tap1

To assess whether DNp53-mediated downregulation of MHC-I was responsible for the ability of MP tumor cells to evade T cell mediated-rejection, we examined whether loss of MHC-I would be sufficient to allow MG tumors to grow in immunocompetent mice. To this end, we generated MG tumors using neural stem cells from MHC-I knockout mice. As shown in Figure 2f, MHC-I knockout MG tumors were able to grow in aB6 mice. Thus, loss of MHC-I is sufficient to allow tumors to escape immune attack, providing an explanation for the ability of MP tumors to grow in immunocompetent mice.

Whereas MP tumors had markedly decreased cell surface MHC-I, they showed no significant differences in the levels of MHC-I mRNA (Figure 2g) or total cellular MHC-I protein (Figure 2h-i). This suggested that p53 was not regulating MHC-I expression, but rather, its cell surface localization. MHC-I can only traffic to the cell surface if it is properly loaded with peptides, a process that depends on cleavage of intracellular proteins by the proteasome, transport of the resultant peptides into the endoplasmic reticulum, and trimming of peptides so they fit properly into the peptide-binding pocket of the MHC-I complex²². Notably, at least two of the proteins required for this process – the peptide transporter Tap1 (transporter 1, ATP binding cassette subfamily B member) and the aminopeptidase Erap1 (endoplasmic reticulum aminopeptidase 1) – have been reported to be targets of p53^{20,21}. Thus, we speculated that the lack of MHC-I in MP tumors might be due to reduced expression of these proteins. As shown in Figure 3a-b, MG tumors express levels of Erap1 and Tap1 comparable to those seen in normal CD133+ neural stem cells. In contrast, MP tumors express significantly less Erap1 and Tap1 at both the mRNA (Figure 3a-d) and protein levels (Figure 3e). Moreover, transduction of MG tumors with DNp53 resulted in a dramatic downregulation of Tap1 and Erap1 (Figure 3c-e), whereas transduction of MP tumors with Gfi1 had no effect. Consistent with this, we found that human medulloblastoma samples with *TP53* mutations have significantly lower levels of *ERAP1* and *TAP1* than those with wild type *TP53* (Figure 3f-g). The correlation between p53 status and *ERAP1* expression is also seen in other tumor types, including acute myeloid leukemia (AML), head and neck cancer, lung cancer and stomach cancer (Extended Data 4a-d, Supplemental Table 1), and in cell lines derived from AML and DIPG (Extended Data 4e-h). Notably, no differences were observed in expression of the close homologs Tap2 and Erap2 in p53-

mutant vs. WT cells (Extended Data 4i-j). These results suggest that p53 regulates MHC-I localization by controlling expression of *Erap1* and *Tap1*.

To investigate whether p53 regulates expression of *Erap1* and *Tap1* directly, we performed chromatin immunoprecipitation (ChIP) using antibodies specific for p53, and PCR to detect the portions of the *Erap1* and *Tap1* promoters containing putative p53 binding sites. As shown in Figure 3h-j, p53 is bound to the *Erap1* and *Tap1* promoters in MG tumors, but not in MP tumors. Similarly, human medulloblastoma PDX cells with WT p53 (ICb1299, RCBM40) show binding of p53 to the *ERAP1* and *TAP1* promoters, whereas PDX cells with mutations in p53 (RCMB18, ICb984, BT-084) do not (Figure 3k-m). These results indicate that p53 directly regulates expression of *Erap1* and *Tap1*.

Erap1 and Tap1 are necessary for cell surface MHC-I expression and T cell mediated rejection

To determine if the resistance of MP tumor cells to T cell attack was caused by loss of *Tap1* or *Erap1*, we first tested whether downregulation of these proteins could phenocopy the effects of loss of p53. shRNA-mediated knockdown of *Erap1* in MG tumor cells caused a significant decrease in surface MHC-I expression (Figure 4a-b) and allowed these cells to grow in syngeneic mice (Figure 4c-d). Knockdown of *Tap1* also decreased MHC-I expression (Extended Data 5a-b), and allowed some tumors to grow in syngeneic mice (Extended Data 5c-d).

We also tested whether overexpression of *Erap1* and *Tap1* could rescue the defects in MHC-I expression in MP tumor cells. Overexpression of *Erap1* caused a marked upregulation of MHC-I in MP tumors (Extended Data 5e), and slowed the growth of MP tumors *in vivo* (Figure 4e,h-i). Overexpression of *Tap1* had a more modest effect on MHC-I expression (Extended Data 5f) and tumor growth (Figure 4f,h-i). However, co-expression of both *Erap1* and *Tap1* caused a significant increase in MHC-I expression (Figure 4g) and a significant delay in growth of MP tumors *in vivo* (Figure 4h-i). Thus, *Erap1* and *Tap1* both contribute to the resistance of *p53*-mutant tumors to immune rejection.

Agonists of TNFR2 and LTβR rescue MHC-I expression in p53-mutant tumor cells

Since the lack of surface MHC-I on *p53*-mutant medulloblastoma cells made them insensitive to T cell killing, we reasoned that restoring MHC-I expression could enhance rejection. Interferon alpha (IFNα)²³, interferon-gamma (IFNγ)²⁴ and Interleukin 12 (IL-12)²⁵ have all been reported to increase MHC-I expression in a variety of cell types^{24,26,27}, we tested the effects of these cytokines on MHC-I expression in our tumor models. IFNα and IL-12 had minimal effects on MHC-I expression in MG or MP tumors (Figure 5a-b and Extended Data 6c-d). IFNγ increased MHC-I expression on MG tumors, which already express MHC-I (Figure 5c), but had no effect on MP tumors, which lack MHC-I (Figure 5d). In contrast, tumor necrosis factor (TNF), which has been reported to enhance MHC-I expression in some cell types^{26,28,29}, caused a marked increase in MHC-I expression on both MG and MP tumors (Figure 5e-f). TNF also induced expression of MHC-I on human MB PDXs and primary MB patient samples (Extended Data 6e-j), and in murine and human DIPG cell lines (Extended Data 6k-l). MHC-I expression was also

induced by an antibody that functions as an agonist of lymphotoxin-beta receptor (LT β R), another member of the TNF receptor superfamily (Figure 5g-h). TNF and lymphotoxin-beta receptor agonist (LT β Rag), but not IFN γ , induced expression of *Erap1* and *Tap1* in MP tumor cells (Figures 5i-m). These studies suggest that TNF receptor signaling can overcome the effects of loss of p53, restoring *Erap1* and *Tap1* expression and rescuing surface MHC-I expression in p53 mutant tumor cells.

One way TNF and LT β Rag could induce MHC-I expression is by restoring expression or function of p53. To test whether this was the case, we generated MP tumors from p53 knockout mice (instead of using DNp53 to suppress p53 function). As shown in (Extended Data 6k-l) TNF and LT β Rag were still able to induce MHC-I expression even in the absence of p53. These data suggest that p53 is not required for rescue of MHC-I expression by TNF or LT β Rag.

TNFR signaling activates the canonical NF κ B pathway

To understand the mechanisms by which TNF and LT β Rag are acting, we first focused on the receptors that mediate their effects. TNF can bind to two distinct receptors – TNF receptor 1 (TNFR1) and TNF receptor 2 (TNFR2), while LT β Rag activates intra-cellular signaling by binding to LT β receptor (LT β R)³⁰. MP tumor cells express all three of these receptors (Extended Data 7a-c). However, treatment with an agonist antibody specific for TNFR1³¹ had no effect on MHC-I expression (Figure 6a), whereas a ligand specific for TNFR2³¹ potently induced MHC-I (Figure 6b). We also tested the effects of TNF and LT β Rag on MP tumors generated from TNFR1, TNFR2 or LT β R knockout mice. TNF was unable to induce MHC-I expression in tumors from TNFR2 knockout mice (Figure 6c), but was still able to do so in tumors from TNFR1 and LT β R knockout mice (Extended Data 7d-f). Conversely, LT β Rag failed to induce MHC-I in LT β R-knockout tumors, but could do so in TNFR1 and TNFR2 knockout tumors (Extended Data 7g-i). These data suggest that TNF and LT β Rag induce MHC-I expression through TNFR2 and LT β R respectively.

We also investigated the transcription factors that mediate MHC-I induction by TNF and LT β Rag. TNF receptor signaling often culminates in translocation of nuclear factor kappa B (NF κ B) family transcription factors – including RelA/p50 heterodimers (canonical pathway) and RelB/p52 heterodimers (non-canonical pathway) – from the cytoplasm to the nucleus³², where they bind to promoters and regulate transcription. As shown in Figure 6d-f, treatment of MP tumor cells with TNF resulted in translocation of RelA and RelB into the nucleus. To determine whether these transcription factors interact with the *Erap1* and *Tap1* promoters, we performed ChIP using antibodies specific for RelA and p50, and PCR to detect the portions of the *Erap1* and *Tap1* promoters containing putative RelA binding sites (no RelB sites were found in the *Erap1* or *Tap1* promoters). These studies demonstrated binding of RelA (Figure 6g-h) and p50 (Figure 6i-j) to the *Erap1* and *Tap1* promoters in TNF- or LT β Rag-treated cells, but not in control (untreated) cells. Moreover, knockdown of RelA or p50 in MP tumors abrogated the effects of TNF and LT β Rag MHC-I expression, whereas knockdown of RelB had no effect (Figure 6k, Extended Data 7j-l). These data indicate that TNF receptor signaling induces *Erap1*, *Tap1* and MHC-I expression by activating the canonical NF κ B pathway.

TNF enhances responses to immune checkpoint inhibitors *in vivo*

The fact that TNF could increase MHC-I expression in tumor cells raised the possibility that it could be used to sensitize p53-mutant tumor cells to immunotherapy. Previous studies have suggested that TNF can cause acute toxicity when administered to animals and people; this toxicity is seen at doses of 1000 µg/kg or higher^{33,34}. We therefore tested doses far below this to determine if they could induce MHC-I expression in MP tumor cells. As shown in Figure 7a-c, intracranial tumor-bearing mice showed a dose-dependent upregulation of MHC-I expression 24 hours after treatment with 0.1 µg/kg, 0.5 µg/kg and 1 µg/kg TNF, doses at which no toxicity is seen even after daily dosing for several weeks (Extended Data 8a). This increase in MHC-I expression was associated with an increase in the number of T cells within the tumor (Extended Data 8b-c). TNF also increased MHC-I expression in murine DIPG tumors 24 hours after treatment (Figure 7d). These studies suggest that low doses of TNF can be administered safely, can accumulate in brain tumor tissue, and can increase expression of MHC-I in tumor cells.

To determine whether these doses of TNF can be used to sensitize p53 mutant tumor cells to T cell killing, we transplanted MP tumor cells into a cohort of aB6 mice and treated them with vehicle, with the immune checkpoint inhibitor anti-PD-1, with low-dose TNF (0.1 - 1 µg/kg) or with the combination of anti-PD-1 and TNF (Figure 7e-f). Mice treated with vehicle had a median survival time of 20 days. Anti-PD-1 alone had little effect on tumor growth or survival (median survival 25 days). TNF alone had a modest effect on tumor growth (median survival 32, 37 and 38 days for 0.1, 0.5 and 1 µg/kg respectively). The combination of anti-PD-1 + 0.1 µg/kg TNF markedly inhibited tumor growth, leading to a 3.1-fold increase in median survival (median survival 52 days, Extended Data 8d-e). The combination of anti-PD-1 + 0.5 µg/kg TNF resulted in long-term tumor-free survival in 52% of mice (Extended Data 8f-g). Finally, the combination of anti-PD-1 + 1 µg/kg TNF led to complete rejection of tumors in all animals (Figure 7e-f). Importantly, these effects were dependent on induction of MHC-I, since no survival benefit was conferred by anti-PD-1 or TNF in tumors generated from MHC-I knockout neural stem cells (Extended Data 8h-i). Similar results were observed in a murine DIPG model (Figure 7g-h) and in conditional *Ptch1* knock-out tumor cells lacking p53 function (Extended Data 8j-k). These results suggest that TNF can be used to increase MHC-I expression and sensitize tumor cells to checkpoint inhibitor immunotherapy.

DISCUSSION

Immunotherapies, including dendritic cell vaccination and immune checkpoint inhibitors, are already being tested in patients with medulloblastoma³⁵. Our observation that p53-mutant tumors lack MHC-I expression and are resistant to immune attack *in vivo* suggests that before enrolling these patients in immunotherapy trials, it may be important to screen them for p53 pathway mutations, or to look directly at ERAP1, TAP1 and MHC-I expression. Given the importance of MHC-I loss in resistance to immunotherapy^{36,37}, we believe this kind of testing is critical for the success of immunotherapy trials.

In light of the prevalence of p53 mutations in human cancer, it will also be important to determine whether the relationship between p53 and MHC-I holds true in other

malignancies. Beyond our findings in medulloblastoma and DIPG, we also observed reduced expression of MHC-I in *TP53*-mutant AML cell lines and in p53-mutant mouse models of pancreatic cancer (data not shown). It is striking that in addition to its well-established roles in proliferation, survival and genomic stability, p53 also acts as a fundamental regulator of the adaptive immune system. Thus, p53 serves as a critical mechanism coordinating cellular failsafe mechanisms to protect genome homeostasis and elimination of malignant cells.

Our studies also demonstrate that treatment with TNF and LT β Rag can rescue MHC-I expression independently of p53, by inducing canonical NF κ B pathway activation and thus rendering tumors more sensitive to T cell-mediated immunotherapy. TNF has been in clinical trials as a single agent, but at the doses that were used, significant toxicity was observed^{33,38}. In that context, it is important to note that in our studies, even low doses of TNF are capable of inducing MHC-I expression. Indeed, the dose of TNF we have used to induce MHC-I in mice (1 μ g/kg, which is equivalent to a human pediatric dose of 2 μ g/m²)³⁹, is at least 100-fold lower than the maximum tolerated dose established in Phase I studies of TNF in children (200-300 μ g/m²)^{40,41}. Our studies suggest that low doses of TNF and related activators can be administered safely, and could be used as a “co-immunotherapy” along with immune checkpoint inhibitors, dendritic cell vaccines or viral oncolytic therapies.

One group of patients who might benefit significantly from this type of therapy is the subset of SHH MB patients who have *TP53* mutations. These patients have a dismal prognosis, and almost all die from their disease. Interestingly, it was recently discovered that a large proportion of these patients also harbor mutations in U1 small nuclear RNAs (snRNAs), which cause altered splicing of the RNAs encoding oncogenes and tumor suppressors⁴². Since aberrant splicing may lead to production of neoantigens, these tumors could be particularly sensitive to immunotherapy. However, if the *TP53* mutations in these tumors result in decreased cell surface MHC-I, these neoantigens cannot be presented to T cells. Rescuing MHC-I expression using TNF might allow these antigens to be presented, and might enhance the responses of these patients to immunotherapy.

It is important to note that tumors may exhibit multiple barriers to immunotherapy – low mutational burden, poor homing of T cells, immunosuppressive factors in the microenvironment – and simply restoring MHC-I by treating with TNF may not be sufficient to allow an effective immune response. However, it is also clear that without MHC-I expression, tumors cannot be recognized and killed by CD8+ T cells; thus, restoration of MHC-I may be one important component of effective T cell based immunotherapy. Given the low response rate in most immunotherapy trials, such co-immunotherapies could markedly increase the number of patients who benefit from immunotherapy in the clinic. In summary, our studies point to p53 mutation as a potential basis for resistance to immunotherapy, and identify TNF and LT β Rag as novel agents for overcoming this resistance and enhancing immune responses in patients.

ACCESSION CODES

The RNA sequencing data from human medulloblastoma samples are deposited in the European Genome-Phenome Archive (EGA) as EGAD00001001899, and EGAD00001004958.

Materials and Methods

Animals

C57BL/6J mice, used as a source of cerebellar stem cells, were obtained from the Sanford Burnham Prebys (SBP) Medical Discovery Institute Animal Facility. NOD-SCID IL2R-gamma null mice (NSG, JAX Stock # 005557) and B6(Cg)-*Tyr^{c-2J}*/J mice (Albino B6, JAX Stock # 000058), used as hosts for intracranial tumor transplantation, were purchased from Jackson Labs. Other mice used as sources of cerebellar stem cells include NSG-(K^bD^b)^{null} (NOD.Cg-*Prkdc^{scid}* *H2-K1^{tm1Bpe}* *H2-D1^{tm1Bpe}* *Il2rg^{tm1Wjl}*/SzJ) mice (MHC-I-null, JAX Stock # 023848), C57BL/6-*Tnfrsf1a^{tm1lmx}*/J mice (TNFR1 Knockout, JAX Stock #003242), B6.129S2-*Tnfrsf1b^{tm1Mwm}*/J mice (TNFR2 Knockout, JAX Stock #002620), B6.129S-*Tnfrsf1a^{tm1lmx}*-*Tnfrsf1b^{tm1lmx}*/J mice (TNFR1/2 double-knockout, JAX Stock # 003243) and B6.129S2-*Trp53^{tm1Tyj}*/J mice (p53 Knockout, JAX Stock # 002101), all from Jackson Labs. Mice were maintained in the animal facility at the Sanford Consortium for Regenerative Medicine. LTβR-deficient (LTβR^{-/-}) mice, provided by C.F.W., are described in ^{43,44}. All experiments were performed in accordance with national guidelines and regulations, and with the approval of the animal care and use committees at SBP and at the University of California San Diego (UCSD).

Retroviruses and Lentiviruses

pMSCV-IRES-GFP, pMSCV-MycT58A-IRES-Luciferase (Luc), pMSCV-DNp53-IRES-GFP, pMSCV-MycT58A-IRES-CD2, pMSCV-DNp53-IRES-Luc and pMSCV-GFI-IRES-GFP were described previously ^{11,14}. MP tumors for most experiments were generated by infecting cerebellar stem cells with viruses produced from pMSCV-MycT58A-IRES-Luc and pMSCV-DNp53-IRES-GFP. Exceptions include tumors used for p53 rescue experiments, in which cerebellar stem cells from p53-knockout mice were infected with pMSCV-MycT58A-IRES-Luc viruses, but not with pMSCV-DNp53-IRES-GFP; and tumors used for overexpression (GOF) of Erp1 and Ta1, which were generated using pMSCV-MycT58A-IRES-CD2 and pMSCV-DNp53-IRES-Luc. MG tumors for all experiments were generated by using pMSCV-MycT58A-IRES-Luc and pMSCV-GFI1-IRES-GFP, except those used for loss of function (LOF) of ERAP1 and TAP1, in which pMSCV-MycT58A-IRES-CD2 and pMSCV-GFI1-IRES-Luc were used instead.

For LOF of Erp1 and Tap1, lentiviral pLKO-GFP vectors encoding scrambled shRNA or shRNA targeting Erp1 (NM_030711, clones TRCN0000324260 for shERAP1#1, TRCN0000324289 for shERAP1#2), or Tap1 (NM_013683, clones TRCN0000066352 for shTAP1#1, TRCN0000066351 for shTAP1#2) were obtained from Sigma-Aldrich. For LOF of NFκB members in MP tumor cells, lentiviral pLKO-GFP vectors encoding scrambled shRNA or shRNA targeting RelA (NM_009045, clones TRCN0000235832 for shRelA#832,

TRCN0000235833 for shRelA#833), p50 (NM_0086, clones TRCN0000235484 for shp50#484, TRCN0000235485 for shp50#485) or RelB (NM_009046, clones TRCN0000233394 for shRelB#494, TRCN0000233392 for shRelB#495) were obtained from Sigma-Aldrich. For LOF of p53, pMSCV-DNp53-IRES-GFP or retroviral vector pMSCV-GFP encoding scrambled shRNA (pJSB42) or p53 shRNA (pJSB109, short hairpin sequence:

TGCTGTTGACAGTGAGCGACCCTGTCATCTTTTGTCCCTTTAGTGAAGCCACAGAT GTAAAGGGACAAAAGATGACAGGGGTGCCTACTGCCTCGGA and pJSB110, short hairpin sequence:

TGCTGTTGACAGTGAGCGCCCACTACAAGTACATGTGTAATAGTGAAGCCACAGAT GTATTACACATGTACTTGTAGTGGATGCCTACTGCCTCGGA). All pJSB constructs were provided by James Scott-Browne (Rao Laboratory, La Jolla Institute for Allergy and Immunology).

For GOF experiments, the *Erap1* construct (pFB-CT10HF-LIC-ERAP1, Addgene, plasmid #39174), a gift from Nicola Burgess-Brown and the *Tap1* construct (MGC-TAP1 from Dharmacon) were used to clone *Erap1* and *Tap1* respectively into pMSCV-IRES-GFP. The p53-encoding construct was bought from Addgene, a gift from Shinya Yamanaka (pMXs-p53, Addgene, plasmid # 22725) ⁴⁵.

Tissue Culture

MP and MG Tumor Generation and Tumor Cell Preparation—Cerebellar stem/progenitor cells (Prominin1+ cells) were purified by fluorescence activated cell sorting (FACS) from the cerebella of postnatal day 5–7 (P5–P7) C57BL/6J pups or MHC-I, TNFR1, TNFR2, TNFR1/2, LTβR or p53 knockout pups as previously described ^{11,14}. To generate MP or MG tumors, cells were infected with viruses encoding Myc and either DNp53 (for MP tumors) or Gfi1 (for MG tumors). After overnight infection, cells were washed and 10,000 cells were stereotactically injected into the cerebellum of 6- to 8-week-old NSG or Albino B6 (aB6) mice. Animals were monitored weekly using in vivo bioluminescence imaging and euthanized when they showed signs of MB. Tumors were then dissociated and resuspended in NeuroCult medium with proliferation supplement (STEMCELL Technologies) for subsequent experiments.

Tumor cells from conditional *Ptch1* knockout (Math1-CreERT²; *Ptch1*^{flox/flox}) mice, a model for SHH-driven medulloblastoma, were generated as previously described ¹⁹. 0.5 to 1 million cells from frozen tumor stocks were stereotactically implanted into the cerebellum of NSG mice. When animals showed signs of MB, tumors were harvested, resuspended in NeuroCult media with proliferation supplement (STEMCELL Technologies), and used for further experiments.

Human MB cell lines, Patient-Derived Xenografts (PDXs) and primary patient samples—The medulloblastoma cell line HD-MB03 (obtained from Till Milde, Hopp Children's Cancer Center, Heidelberg) ¹⁸, was cultured in RPMI-1640 medium (Gibco-ThermoFisher), supplemented with 10% fetal bovine serum (Seradigm), 100 units/mL

penicillin, and 100 µg/mL streptomycin (Invitrogen), 0.1mM non-essential amino acids (Invitrogen).

Primary MB samples and PDX lines used for this study are listed in Supplemental Table 2 and in the Life Sciences Reporting Summary. Primary MB samples were collected during surgery at Rady Children's Hospital (San Diego, CA). PDX lines were generated by implanting patient cells directly into the cerebellum of NSG mice, and propagating them from mouse to mouse without in vitro passaging; the identity and subgroup of each line was validated by gene expression and/or methylation analysis. For all experiments, cells were isolated from tumor-bearing mice, resuspended in NeuroCult media with proliferation supplement (STEMCELL Technologies), and assayed.

Human Acute Myeloid Lymphoma (AML) cell lines—MV4-11⁴⁶, MOLM13⁴⁶, HL60⁴⁶, THP1⁴⁷, U937⁴⁸, ML-2⁴⁹, P31⁵⁰ cells were cultured in suspension in RPMI 1640 medium (Gibco-ThermoFisher), supplemented with 10% fetal bovine serum (Seradigm), 100 units/mL penicillin, and 100 µg/mL streptomycin (Invitrogen).

Human and mouse DIPG cell lines—p53 WT (PDGF-B; H3.3K27M; PTEN deficient⁵¹) and p53-deficient (PDGF-B; H3.3K27M; p53 loss (Cre), or NS-PKC)⁵² murine DIPG lines (created in the laboratory of O.J.B.) were cultured in NeuroCult medium with proliferation supplement (STEMCELL Technologies), 100 units/mL penicillin, and 100 µg/mL streptomycin (Invitrogen), 2ug/mL Heparin (Sigma-Aldrich), 20ng/mL FGF (Peprotech) and 10ng/mL EGF (Peprotech). The human p53 deficient DIPG cell line, SF8628⁵³ (generated by R. Hashizume, Northwestern University) was cultured in DMEM medium (Gibco-ThermoFisher), supplemented with 10% fetal bovine serum (Seradigm), 100 units/mL penicillin, and 100 µg/mL streptomycin (Invitrogen). The human p53 WT DIPG cell line, SU-DIPG⁵⁴ (generated by M. Monje, Stanford University) was cultured in Neurobasal media (-A) (Invitrogen), B27 (-A) (Invitrogen), 10ng/mL Heparin (Sigma-Aldrich), 20ng/mL FGF (Peprotech) and 10ng/mL EGF (Peprotech).

Flow Cytometry and Cell Sorting

Cells were suspended in PBS + 5% FBS (PBS-FBS), and then stained with fluorescent dye-conjugated antibodies (anti-mouse MHC Class I (H2 Kb/Db), anti-human HLA A/B/C, anti-mouse CD3, anti-mouse CD8, anti-mouse B220, anti-mouse CD11b, anti-mouse NK1.1 or isotype control (see Antibody List in Supplemental Table 3 and in the Life Sciences Reporting Summary)) for 1h at 4°C. Samples were washed 3 times with PBS-FBS, and then run on a LSRFortessa flow cytometer (BD Biosciences) and analyzed using FlowJo (TreeStar). For all samples, data from 10,000 viable cells were acquired. Gating strategy is presented in Extended Data 9. For cell sorting, infected cells were resuspended in PBS-FBS for sorting using a BD FACSAria II (BD Bioscience), based on GFP expression.

In vitro and in vivo treatments

Cytokines used *in vitro* were purchased from Peprotech and resuspended in DMSO. Cells were treated with 50 pg/ml TNF, 20 ng/ml IFNα, 20 ng/ml IFNγ or 1 µg/mL IL-12. LTβR agonist (produced in the Ware lab at SBP)⁴⁴ was used at 1.6 µg/ml. TNFR1 and TNFR2

agonists (from the Wajant lab ³¹) were used at 0.5µg/mL and 1.67nM respectively. The same amount of DMSO was added to control cells. MHC-I expression was assessed 48h after treatment.

TNF for *in vivo* use was purchased from Biolegend and resuspended in DMSO at a concentration of 200 µg/ml. Before treatment, TNF-DMSO stock was diluted in PBS to a final concentration of 0.1 ng/µl. Mice for *in vivo* experiments are transplanted at 6-8 weeks old. Treatments were started when bioluminescent signal reached approximately 5×10^6 . Mice were randomized based on bioluminescent signal. Equal number of males and females were used. Blinding was not used for the study because quantitative bioluminescent imaging and standard software analysis were used. No subjective analysis of tumor burden was necessary. Tumor bearing mice were injected retro-orbitally with 0.1 µg/kg, 0.5 µg/kg or 1 µg/kg TNF under general anesthesia (2.5% isoflurane, 2% O₂). For PD-1 blockade experiments, anti-PD-1 (clone RMP1.14, BioXCell) was administered intra-peritoneally (i.p.) at a dose of 10 mg/kg. For combinations of TNF and anti-PD-1, each cycle of treatment consisted of a dose of TNF on day 1, a dose of anti-PD-1 on day 2, and no treatment on day 3. Mice received up to 20 cycles of treatment. Control mice were treated with TNF alone, anti-PD-1 alone or PBS. Mice were monitored for signs of morbidity or toxicity (>20% weight loss), and were euthanized if these were observed.

For T-cell depletion experiments, mice were injected i.p. with anti-Thy-1 (clone T24/31, BioXcell) or anti-CD8 (clone 2.43, BioXCell) or rat IgG2b isotype control (BioXCell) at a dose of 20 mg/kg, 2 days before tumor transplantation, and again on the day of transplantation. Treatment was continued at a dose of 10 mg/kg every 4 days until the end of the experiment.

For *in vivo* experiments involving murine DIPG cells, 100,000 cells were stereotactically injected into the brains of 3 day-old Albino B6 (aB6) mice. Animals were monitored by weekly bioluminescence imaging starting at 21 days of age, and euthanized when they showed signs of tumors.

***In Vivo* Bioluminescence Imaging**

Mice injected with cells from MP or MG tumors were subjected to weekly bioluminescence imaging. Mice were given i.p. injections of 150 ng/g D-Luciferin (Caliper Life Sciences, cat# 12279) and anesthetized with 2.5% isoflurane in an induction chamber. Five minutes after injection, animals were imaged using the Xenogen Spectrum (IVIS-200) imaging system. Signal quantification (number of photons emitted/second) was performed using Living Image software.

Quantitative Reverse Transcriptase Polymerase Chain Reaction (qRT-PCR)

Cells were washed twice with cold PBS and lysed for total RNA extraction using a Qiagen RNeasy Kit. Extracted RNAs were then used to generate cDNA using the iScript cDNA synthesis kit (Bio-Rad). Real-time PCR was carried out in triplicate using SYBR Green PCR Master Mix (Bio-Rad) on a Bio-Rad CFX384 Real-Time System. The primers used in this study are described in Supplemental Table 4.

Chromatin ImmunoPrecipitation (ChIP)

15×10⁶ tumor cells were cross-linked with 1% formaldehyde for 10 min at 37°C and lysed in buffer containing 1% SDS, 10 mM EDTA and 50 mM Tris pH 8.1. Sonicated chromatin was immunoprecipitated overnight with p53, RelA or p50 antibodies or with IgG as a negative control, and protein A/G plus agarose magnetic beads (Thermo Fisher Scientific).

Antibodies used in this study are described in Supplemental Table 3. Immunoprecipitated chromatin was eluted in a buffer containing 1% SDS with 0.1M NaHCO₃. The cross-links were reversed by the addition of NaCl (200 mM final) for 4h at 65°C. DNA was extracted using phenol-chloroform. Real-time quantitative PCR was performed using SYBR Green PCR Master Mix (Bio-Rad) with primers specific for the promoter or 3'UTR (negative control) of genes. Putative transcription factor binding sites in the promoters of the *Erap1* and *Tap1* genes were identified using ConTra V3 software⁵⁵ (<http://bioit2.irc.ugent.be/contrav3/#/step/1>). Analysis of an input sample was used to normalize each sample (DCt (normalized CHIP) = (Ct(chip)-(Ctinput-log2(input dilution factor))), and fold induction was calculated by comparing p53, RelA or p50 antibody precipitates with their corresponding IgG antibody precipitates. The primers used in this study are described in Supplemental Table 4.

Cell Lysis and Western Blotting

Cells were washed twice with cold PBS and lysed in RIPA buffer with protease and phosphatase inhibitor cocktail (Cell Signaling, Cat# 5872) for 30 min on ice. Cell lysates were centrifuged at 13,000 rpm for 15 min at 4°C. Protein concentrations were measured using a Pierce BCA Protein Assay Kit (Thermo Scientific #23225). Nuclear/cytoplasmic protein extracts were prepared according to manufacturer's guidelines (Thermo Fisher Scientific, Cat#78833). Equal amounts of protein were separated by 10% SDS-PAGE and transferred to nitrocellulose membranes (Invitrogen, Cat# LC2006). The membranes were blocked with 5% nonfat dried milk (Bio-Rad) for 1 hr at room temperature and incubated in primary antibodies diluted in 5% nonfat dried milk overnight at 4°C with gentle shaking. Membranes were washed 3 times with Tris-buffered saline with 0.1% Tween-20 (TBS-T), and incubated with corresponding secondary antibodies for 45 min at room temperature. Membranes were washed again as described above, imaged, and quantified using Image Lab 5.0 (Bio-Rad). Antibodies used in this study are described in Supplemental Table 3.

RNA sequencing of human medulloblastoma samples

Sequencing reads were mapped by STAR version 2.5.1b on FASTA which includes the human reference genome "hs37d5" by 1000 Genomes Project Phase II, spike-in sequences of profile C1_2 ERCC spike-in concentrations used for C1 fluidigm and Caltech profile 3 spike-ins by ENCODE with the option '--outFilterMultimapNmax 20 --alignSJoverhangMin 8 --alignMatesGapMax 200000 --alignIntronMax 200000 --alignSJDBoverhangMin 10 --alignSJstitchMismatchNmax 5 -1 5 5 --outSAMmultNmax 20 --twopassMode Basic'⁴²

Fragments Per Kilobase per Million mapped reads (FPKM) was calculated using edgeR version 3.14.0 followed by counting reads using GENCODE v19 gtf file and htseq version 0.6.0 with the setting "--stranded reverse -m union".

TCGA analysis

Fold change in mRNA levels was assessed by subtracting the average log median normalized values of mRNA for each gene in *TP53* mutant tumors from *TP53* wild type tumors. A negative value indicates decreased fold change, and a positive value, increased. Data for analysis was downloaded from cBio portal and frozen on 09/18/2018. For each cancer site, the dataset from the most recent TCGA publication was used. In all cases, mRNA levels originated from RNAseq analyses. P-values for differences in fold change across cancer sites were downloaded from cBio data portal. P-values for differences in mRNA expression in boxplots were assessed using non-parametric Wilcoxon Rank Sum tests.

Statistics and Reproducibility

Statistical analysis was performed using GraphPad Prism software. All data are presented as means $\pm$ SD unless otherwise stated. Based on experience with studies of this type, we performed power analysis, and determined that with 8 mice per group we will have a power of 90% to detect an average survival difference of 14 days with a significance of $p=0.05$; thus survival experiments included at least 8 mice/group and were repeated at least 3 times. Replicates represent tumor samples harvested from independent mice. Data distribution was assumed to be normal but this was not formally tested. No animals or data points were excluded from any experiments.

Comparisons between different groups were made using two-sided Student's *t* test or two-way ANOVA as appropriate. The statistical significance of Kaplan-Meier survival curves was assessed using a two-sided log-rank (Mantel-Cox) test. For all experiments, confidence intervals were 95%, and *p* values of 0.05 or lower were considered statistically significant. Statistical details can be found in the figure legends and in Supplemental Table 1 for the TCGA analysis shown in Extended Data 4.

Data and materials availability

All data are available in the main paper or the supplementary materials. Raw data for all figures will be available as Source Data files. The human medulloblastoma sequencing data used in Figure 3f and 3g are deposited in the European Genome-Phenome Archive (EGA) as EGAD00001001899, and EGAD00001004958.

Extended Data

Extended Data Fig. 1. Immune microenvironment in p53-mutant and wild-type murine medulloblastoma.

(a-d) Intracranially transplanted MP tumors (black) and MG tumors (red) labelled with GFP were harvested 2, 5 and 7 days after transplant (n=3 mice per tumor type per day). Percentages of B-cells (a, B220 positive), Macrophages (b, CD11b positive), Natural killer (NK) cells (c, NK1.1 positive) and cytotoxic T-cells (d, CD8 positive) were determined by FACS analysis. The average curve and single data points are shown for each condition. p-values were determined by two-sided unpaired t-test. In (d), * p=0.0014. (e-g) Depletion of

T-cells (α CD8, red line), or no T-cell depletion (Control, black line), was performed in aB6 MG tumor-bearing mice. (e) Bioluminescence imaging of representative mice. (f) Quantification of the average bioluminescence signal (dark lines) and the signals for individual mice (light lines) for each group. (i) Survival curves for control mice ($n = 6$); and α CD8-treated mice ($n=6$). p-value for survival curve was determined by a two-sided log-rank (Mantel-Cox) test. All error bars represent the mean $\pm$ SD. (h-i) MP tumor cells or MP tumor cells overexpressing GFI1 (MP+G) were transplanted into NSG or aB6 mice. Bioluminescence imaging of representative mice (h) and survival curves ($n=10$) (i) are shown. p-values for the difference in survival between MP and MP+G transplanted into NSG or aB6 were determined by two-sided log-rank (Mantel-Cox) test. In (i), NS (not significant), $p=0.608$ (NSG) and $p=0.597$ (aB6). (j-k). Expression of T-cell suppression molecules (j) and Dendritic cell and T-cell activation molecules (k) in MP tumors ($n=3$) and MG tumors ($n=3$) were measured by qRT-PCR. Quantification of the relative expression of the genes for three independent tumors is shown; data points represent values for individual tumors. Bars represent means $\pm$ SD. p-values were determined by two-sided unpaired t-test. None of the differences between MP and MG were significant.

Extended Data Fig. 2. Knockdown of p53 decreases MHC-I expression in murine and human medulloblastoma cells.

(a) FACS histogram of MHC-I expression in normal neural stem cells (NSCs). Isotype control is shown in gray and MHC-I staining in black. Quantification of the mean fluorescence intensity (MFI) values for three independent NSC samples (isolated from three separate litters of mice) is shown below the histogram; data points represent MFIs for individual NSC samples. p-value, determined by two-sided unpaired t-test, is indicated on bar graph. (b and d) FACS histograms of MHC-I or HLA-I expression in murine MG tumor cells (b) and human HD-MB03 tumor cells (d) transduced with control shRNA (black histograms) or shRNAs that target sequences conserved in mouse and human *p53* (shp53

#109, blue histograms; shp53 #110, pink histograms). Quantification of the MFI values for three independent tumors are shown below each histogram; data points represent MFIs for individual tumor samples. p-values were determined by two-sided unpaired t-test. p-values are indicated on each bar graph. (c and e) Protein levels of p53 and β -Actin were determined by western blotting in MG tumor cells (c) and HD-MB03 cells (e). Western blots are cropped at the molecular weights for p53 (53 kDa) and β -actin (42 kDa); original blots are available in Source Data. (f) Murine Ptch-KO MB tumor cells overexpressing dominant negative p53 (Ptch-KO MB-DNP53) were transplanted into NSG or aB6 mice. Survival curves (n=8) are shown. p-values for the difference in survival between Ptch-KO MB and Ptch-KO MB-DNP53 were determined by two-sided log-rank (Mantel-Cox) test. NS (not significant), p=0.6939 (NSG) and ** p=0.0021 (aB6). (g) FACS histograms of MHC-I expression in Ptch-KO MB (black) and Ptch-KO MB-DNP53 (blue) tumor cells. Quantification of MFI values for three independent tumor samples is shown below the histogram; data points represent MFIs for each tumor sample. p-value, determined by two-sided unpaired t-test, is indicated on bar graph.

Extended Data Fig. 3. p53 status determines MHC-I expression in medulloblastoma patient-derived xenografts and DIPG cells.

(a-j) FACS histograms of HLA-I expression in human medulloblastoma PDXs expressing mutant *p53* (upper panel, a-e) or wild-type *p53* (lower panel, f-j). Isotype controls are shown in black and HLA-I staining in purple. (e, j) Quantification of average mean fluorescence intensity (MFI) for triplicate samples of each human medulloblastoma PDX. Two-sided t-tests were used to evaluate the significance of the difference between HLA-I and isotype control staining for *p53*-mutant tumors (e, NS, not significant $p=0.3793$) and for *p53*-WT tumors (j, * $p=0.0112$). (k-n) FACS histograms of MHC-I expression in *p53* wild-type (k and m) or *p53* deficient (l and n) murine DIPG lines (k-l) and human DIPG lines (m-n). Isotype controls are shown in black and murine MHC-I and human HLA-I staining in

purple. Quantification of the MFI values for three independent tumor samples are shown below each histogram; data points represent MFIs for individual tumor samples. p-values were determined by two-sided unpaired t-test and are indicated on each bar graph.

Extended Data Fig. 4. p53 mutations are associated with decreased ERAP1 expression in other cancers.

(a-b) Expression of *ERAP1* and *TAP1* in *p53*-mutant relative to *p53* wild type tumors in TCGA datasets. Cancer sites are indicated along the X-axis. Each column represents the fold change in expression in *p53* mutant relative to *p53* WT tumors. Positive values indicate increased expression of *ERAP1* (a) or *TAP1* (b) mRNA levels in *p53* mutant tumors

compared to *p53* wild-type (WT) tumors; negative values indicate decreased expression of *ERAP1* or *TAP1* in *p53* mutant tumors compared to WT tumors; 0 indicates no change in expression. Error bars indicate standard error of the mean. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; ns, not significant, based on two-sided Wilcoxon Rank Sum test. Precise p -/q-values and sample sizes are provided in Supplemental Table 1. (c-d) Box plots describing mRNA expression of *ERAP1* (c) and *TAP1* (d) in patient tumors from TCGA datasets (AML, Head and Neck, lung and stomach cancer) with *p53*-mutant samples shown in red and *p53*-wild type (WT) samples shown in blue. Box plot center lines show median, box limits indicate the 25th and 75th percentiles, lower and upper whiskers extend 1.5 times the interquartile range from the 25th and 75th percentiles, respectively. p -values represent significance, based on two-sided Wilcoxon Rank Sum test. (e, f) *Erap1* (e) and *Tap1* (f) expression in *p53*-WT (blue) and mutant (red) human AML cell lines. (g, h) *Erap1* (g) and *Tap1* (h) expression in *p53*-WT (blue) and mutant (red) murine and human DIPG cell lines. *Erap1* and *Tap1* expression were measured by qRT-PCR. Quantification of 3 independent experiments is shown. Error bars represent means $\pm$ SD. p -values were determined by two-sided unpaired t-test comparing *p53* WT and mutant lines. In (e) * $p = 0.0457$; in (f) NS, not significant, $p = 0.7724$; in (g) ** $p = 0.007$ (murine) and * $p = 0.033$ (human); in (h) ** $p = 0.0009$ (murine) and * $p = 0.028$ (human). (i-j) *Erap2* (i) and *Tap2* (j) expression in MP tumors ($n = 3$), MG tumors ($n = 3$), MG tumors overexpressing DNp53 (MG+P) ($n = 3$) were measured by qRT-PCR. Error bars represent means $\pm$ SD. p -values were determined by two-sided unpaired t-test. In (j) NS, not significant, $p = 0.21$ (MG vs MP), $p = 0.394$ (MG vs MG+P); In (i) NS, $p = 0.112$ (MG vs MP) and $p = 0.149$ (MG vs MG+P).

Extended Data Fig. 5. Perturbation of Tap1 alters expression of cell surface MHC-I.

(a-d) MG tumor cells were transduced with control shRNA (shCtl) or shRNAs targeting *Tap1* (shTap1#1, shTap1#2). Knockdown efficiency was determined by western blotting (a). Western blots are cropped at the molecular weights for Tap1 (68 kDa) and β-actin (42 kDa); original blots are available in Source Data. MHC-I expression was determined by FACS in control cells (shCtl, black) and *Tap1* knockdown cells (shTap1, red) (b). Quantification of the mean fluorescence intensity (MFI) values for three independent tumors is shown below each histogram; data points represent MFIs for individual tumor samples. p-values, determined by two-sided unpaired t-test, are indicated on bar graph. (c-d) *Tap1* knockdown cells were transplanted into aB6 mice. Bioluminescence imaging of representative mice (c)

and survival curves (n=6) (d) are shown. p-values for the difference in survival between shTap1 and shCtl were determined using the two-sided log-rank (Mantel-Cox) test. *p=0.0198 (shTap1#1) and *p=0.0246 (shTap1#2). (e-f) MP tumor cells were transduced with empty vector (vect) or vectors encoding *Erap1* or *Tap1*. Analysis of MHC-I expression by FACS in control cells (vect, black), *Erap1*-overexpressing cells (e) or *Tap1*-overexpressing cells (f) (blue histograms in e and f). Quantification of the MFI for three independent experiments is shown below the histogram; data points represent MFIs for individual tumor samples. The p-values were determined by two-sided unpaired t-test and are indicated on each bar graph.

Extended Data Fig. 6. TNF increases MHC-I expression in human medulloblastoma and DIPG. MG (a) and MP (b) tumor cells were treated *in vitro* for 48h with interleukin-12 (IL-12, 1 $\mu\text{g/mL}$). MHC-I expression in untreated cells (Ctl, black histograms) and cells treated with IL-12 (pink) were analyzed by FACS. (c-h) Fresh patient samples (c, d) and PDXs (e-h) from *p53*-mutant MB patients were treated *in vitro* for 48h with TNF (50 pg/mL). HLA-I expression in untreated cells (Ctl, black histograms) and cells treated with TNF (green histograms) were analyzed by FACS. In (c) and (d), no error bars are shown because no replicates were possible from primary patient samples. (i, j) cells from *p53*-mutant murine (i) and human (j) DIPG cell lines were treated with TNF (50 pg/mL) for 48h. Murine MHC-I and human HLA-I expression was analyzed by FACS in untreated cells (Ctl, black histograms) and cells treated with TNF (green histograms). (k-l) MP tumor cells generated from *p53* knockout (*p53* KO) mice were treated for 48h with TNF (50 pg/mL) or $\text{Lt}\beta\text{Rag}$ (1.6 $\mu\text{g/mL}$). MHC-I expression was analyzed by FACS in untreated cells (Ctl, black histograms) and cells treated with TNF (green histograms) or $\text{Lt}\beta\text{Rag}$ (blue histograms). For each FACS analysis (except for fresh MB patient samples in c and d), quantification of the mean fluorescence intensity (MFI) for three independent experiments is shown below each histogram; data points represent MFIs for individual tumor samples. p-values were determined by two-sided unpaired t-test. p-values are indicated on each bar graphs.

Extended Data Fig. 7. TNFR2 and LTβR are necessary for induction of MHC-I by TNF and LTβRag.

(a-c) TNFR1 (a, green), TNFR2 (b, magenta) and LTβR (c, blue) expression on MP tumor cells were analyzed by FACS. Isotype control antibody is shown in gray. (d-f) MP tumor cells generated from *TNFR1* knockout mice (*TNFR1* KO, d), *TNFR1/2* double knockout mice (*TNFR1/2* KO, e) or *LTβR* knockout mice (*LTβR* KO, f) were treated for 48h with TNF (50 pg/mL). MHC-I expression in untreated cells (Ctl, black histograms) and cells treated with TNF (green histograms) were analyzed by FACS. (g-i) MP tumor cells generated from *TNFR1* knockout mice (*TNFR1* KO, g), *TNFR2* knockout mice (*TNFR2* KO, h) or *LTβR* knockout mice (*LTβR* KO, i) were treated for 48h with LTβRag (1.6 μg/

mL). MHC-I expression in untreated cells (Ctl, black histograms) and cells treated with LT β Rag (blue histograms) were analyzed by FACS. For each FACS analysis, quantification of the mean fluorescence intensity (MFI) for three independent tumor samples is shown below each histogram; data points represent MFIs for individual tumor samples. p-values were determined by two-sided unpaired t-test, and are indicated on the corresponding graphs. (j-l) MP tumor cells were transduced with control shRNA (shCtl) or shRNAs targeting *RelA* (shRelA#832, shRelA#833), *p50* (shp50#484, shp50#485) or *RelB* (shRelB#494, shRelB#495). Knockdown efficiency of RelA (j), p50 (k) and RelB (l) was determined by western blot. Western blots are cropped at the molecular weights for RelA (65 kDa), p50 (50 kDa), RelB (68 kDa) and GAPDH (37 kDa); original blots are available in Source Data.

Extended Data Fig. 8. TNF increases sensitivity to immune checkpoint inhibitors in a dose-dependent manner.

(a) TNF toxicity was evaluated in aB6 mice after daily dosing with TNF at 0.5 $\mu\text{g/kg}$ (red line), 1 $\mu\text{g/kg}$, 10 $\mu\text{g/kg}$ and 20 $\mu\text{g/kg}$ ($n=3$ per dose of TNF). Body weight was measured every 5 days. (b-c) MP tumor cells were transplanted into aB6 mice, and after 10 days, animals were treated with vehicle (Ctl) or TNF (0.5 $\mu\text{g/kg}$) for 9 days, and tumor tissue was harvested and dissociated. Percentages of isolated cells representing tumor cells (GFP+) and CD3+ T-cells (anti-CD3-APC) were determined by FACS analysis. (b) Representative dot plots (isotype control antibody, black dots; anti-CD3 antibody, green dots) for control mice (left panel) and TNF-treated mice (right panel) are shown, with percentages of cells in each

quadrant indicated on the plot. (c) Bar graph indicating the average percentage of CD3+ T-cells in tumors from control mice (n=3, black) and TNF-treated mice (n=3, green). p-value was determined by two-sided unpaired t-test. ** p=0.0051). (d-k) MP tumor cells (d-g), *MHC-I* knock-out (KO) MP tumor cells (h, i) or DNp53-expressing Ptch-KO (j, k) tumor cells were transplanted into aB6 mice. 15 days later, mice were treated with vehicle (Ctl, black), anti-PD-1 (10 mg/kg, blue), TNF (0.1 – 1 µg/kg, as indicated in panels d, f, h and j, green) or anti-PD-1 and TNF (orange). For combinations of TNF and anti-PD-1, each cycle of treatment consisted of a dose of TNF on day 1, a dose of anti-PD-1 on day 2, and no treatment on day 3. Mice received up to 20 cycles of treatment. (d, f, h, j) Bioluminescence imaging of representative mice. (e, g, i, k) Survival curves (in (e) : n=10 ; in (g) : n=25 ; in (i) : n=16; in (k): n=10 per group). p-values for the difference in survival between Ctl mice and treated mice were determined by a two-sided log-rank (Mantel-Cox) test. (e) *p=0.019 (anti-PD-1), **p=0.0007 (TNF), ***p=9.3E-06 (TNF + anti-PD-1); (g) *** p=8.1E-06 (anti-PD-1), p=2.6E-10 (TNF), p=5.3E-12 (TNF + anti-PD-1); (i) NS, Not Significant p=0.29 (anti-PD-1), p=0.151 (TNF), p=0.094 (TNF + anti-PD-1) ; (k) *** p=8.1E-05 (anti-PD-1), p=3.5E-06 (TNF), p=3.5E-06 (TNF + anti-PD-1).

Extended Data Fig. 9. Representative gating strategy used for analysis of HLA A/B/C in human cells or MHC-I expression in murine cells.

(a-d) Gating strategy for unlabeled human tumor cells (data for RCMB18 are shown). (a) Dot plot representing FSC-A and SSC-A profile of the entire cell population, with box indicating intact cells (as opposed to debris and clumps) selected for analysis. (b) Dot plot showing the FSC-H and FSC-A profile of the population selected in (a), with a box indicating the single cell (as opposed to doublet) population selected for analysis. (c) Dot plot showing 7-AAD staining vs. FSC-A of the population in (b), with a box indicating the viable (7-AAD-negative) cell population selected for analysis. (d) Analysis of HLA-I staining of the cell population in (c). (e-i) Gating strategy for GFP-labeled murine (MG)

tumor cells. (e) Dot plot representing FSC-A and SSC-A profile of the entire cell population, with box indicating intact cells (as opposed to debris and clumps) selected for analysis. (f) Dot plot showing the FSC-H and FSC-A profile of the population selected in (e), with a box indicating the single cell (as opposed to doublet) population selected for analysis. (g) Dot plot showing 7-AAD staining vs. FSC-A of the population in (f), with a box indicating the viable (7-AAD-negative) cell population selected for analysis. (h) Dot plot showing GFP expression vs. FSC-A in the population selected in (g), with a box indicating the tumor cells (GFP+) selected for analysis. (i) Analysis of MHC-I staining of the population in (h).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the Animal Facilities at UCSD and SBP for help with animal care and husbandry; Silvia Tacheva Grigorova for assistance generating conditional *Ptch1* knockout mice, Y. Altman of the Flow Cytometry Shared Resource at SBP and Jesus Olvera and Cody Fine of the Stem Cell Core at UCSD for assistance with FACS sorting and analysis. We are also indebted to Till Milde at the Hopp Children's Cancer Center, Heidelberg for providing HD-MB03 and BT-084 cells and to Michelle Monje at Stanford University for sharing human DIPG cell lines, and to James Scott-Browne and Anjana Rao at the La Jolla Institute for Allergy and Immunology for providing p53 shRNA constructs. We thank Suzanne Baker and Laura Hover at St Jude Children's Research Hospital, Laura Attardi at Stanford University, Olivier Ayrault and Antoine Forget at Institut Curie, and Jeremy Rich, Brianna Prager and Leo Kim from UCSD for help with analysis of gene and protein expression in p53 mutant and WT tumors. M.D.T. is supported by the NIH (R01CA148699 and R01CA159859), The Pediatric Brain Tumour Foundation, The Terry Fox Research Institute, The Canadian Institutes of Health Research, The Cure Search Foundation, b.r.a.i.n. child, Meagan's Walk, SWIFTY Foundation, The Brain Tumour Charity, Genome Canada, Genome BC, Genome Quebec, the Ontario Research Fund, Worldwide Cancer Research, V-Foundation for Cancer Research, and the Ontario Institute for Cancer Research through funding provided by the Government of Ontario. M.D.T. is also supported by a Canadian Cancer Society Research Institute Impact grant, a Cancer Research UK Brain Tumour Award, and by a Stand Up To Cancer (SU2C) St. Baldrick's Pediatric Dream Team Translational Research Grant (SU2C-AACR-DT1113) and SU2C Canada Cancer Stem Cell Dream Team Research Funding (SU2C-AACR-DT-19-15) provided by the Government of Canada through Genome Canada and the Canadian Institutes of Health Research, with supplementary support from the Ontario Institute for Cancer Research through funding provided by the Government of Ontario. M.D.T. is also supported by the Garron Family Chair in Childhood Cancer Research at the Hospital for Sick Children and the University of Toronto. This work was supported by funding from the National Cancer Institute (2R01 CA159859-06 and P30 CA30199 (R.J.W.-R.)), P01 CA177322 (C.F.W.), K22 CDA CA229613 (S.H.), NIH (R01 AI048073) (C.F.W.), Forschungsgemeinschaft (DFG, German Research Foundation) (324392634-TRR221 and WA 1025/31-1) (H.W.), the V Foundation, the Pediatric Brain Tumor Foundation, Accelerate Brain Cancer Cure, Alex's Lemonade Stand Foundation (R.J.W.-R. and O.J.B.), William's Superhero Fund (R.J.W.-R.), the McDowell Charity Trust (R.J.W.-R.), Madox's Warriors (O.J.B.), Cristian Rivera Foundation (O.J.B.), Fly a kite Foundation (O.J.B.), and the California Institute for Regenerative Medicine (R.J.W.-R.). This work is dedicated to the memory of Errol McDowell.

REFERENCES

1. Tabori U, et al. Universal poor survival in children with medulloblastoma harboring somatic TP53 mutations. *J Clin Oncol* 28, 1345–1350 (2010). [PubMed: 20142599]
2. Hill RM, et al. Combined MYC and P53 defects emerge at medulloblastoma relapse and define rapidly progressive, therapeutically targetable disease. *Cancer cell* 27, 72–84 (2015). [PubMed: 25533335]
3. Morrissy AS, et al. Divergent clonal selection dominates medulloblastoma at recurrence. *Nature* 529, 351–357 (2016). [PubMed: 26760213]
4. Kantoff PW, et al. Sipuleucel-T immunotherapy for castration-resistant prostate cancer. *N Engl J Med* 363, 411–422 (2010). [PubMed: 20818862]
5. Miller JF & Sadelain M The journey from discoveries in fundamental immunology to cancer immunotherapy. *Cancer cell* 27, 439–449 (2015). [PubMed: 25858803]

6. Rosenberg SA & Restifo NP Adoptive cell transfer as personalized immunotherapy for human cancer. *Science* 348, 62–68 (2015). [PubMed: 25838374]
7. Topalian SL, Drake CG & Pardoll DM Immune checkpoint blockade: a common denominator approach to cancer therapy. *Cancer cell* 27, 450–461 (2015). [PubMed: 25858804]
8. Braun DA, Burke KP & Van Allen EM Genomic Approaches to Understanding Response and Resistance to Immunotherapy. *Clinical cancer research : an official journal of the American Association for Cancer Research* 22, 5642–5650 (2016). [PubMed: 27698000]
9. [ClinicalTrials.gov. NCT03173950](#): Immune Checkpoint Inhibitor Nivolumab in People With Select Rare CNS Cancers.
10. [ClinicalTrials.gov. NCT01326104](#): Vaccine Immunotherapy for Recurrent Medulloblastoma and Primitive Neuroectodermal Tumor.
11. Pei Y, et al. HDAC and PI3K Antagonists Cooperate to Inhibit Growth of MYC-Driven Medulloblastoma. *Cancer cell* 29, 311–323 (2016). [PubMed: 26977882]
12. Pei Y, et al. An animal model of MYC-driven medulloblastoma. *Cancer cell* 21, 155–167 (2012). [PubMed: 22340590]
13. Bowman T, et al. Tissue-specific inactivation of p53 tumor suppression in the mouse. *Genes Dev* 10, 826–835 (1996). [PubMed: 8846919]
14. Northcott PA, et al. Enhancer hijacking activates GFI1 family oncogenes in medulloblastoma. *Nature* 511, 428–434 (2014). [PubMed: 25043047]
15. Zurita E, et al. Genetic polymorphisms among C57BL/6 mouse inbred strains. *Transgenic Res* 20, 481–489 (2011). [PubMed: 20506040]
16. Cortez MA, et al. PDL1 Regulation by p53 via miR-34. *J Natl Cancer Inst* 108(2016).
17. Wang X, et al. Tumor suppressor miR-34a targets PD-L1 and functions as a potential immunotherapeutic target in acute myeloid leukemia. *Cell Signal* 27, 443–452 (2015). [PubMed: 25499621]
18. Milde T, et al. HD-MB03 is a novel Group 3 medulloblastoma model demonstrating sensitivity to histone deacetylase inhibitor treatment. *Journal of neuro-oncology* 110, 335–348 (2012). [PubMed: 23054560]
19. Yang ZJ, et al. Medulloblastoma can be initiated by deletion of Patched in lineage-restricted progenitors or stem cells. *Cancer cell* 14, 135–145 (2008). [PubMed: 18691548]
20. Zhu K, et al. p53 induces TAP1 and enhances the transport of MHC class I peptides. *Oncogene* 18, 7740–7747 (1999). [PubMed: 10618714]
21. Wang B, Niu D, Lai L & Ren EC p53 increases MHC class I expression by upregulating the endoplasmic reticulum aminopeptidase ERAP1. *Nat Commun* 4, 2359 (2013). [PubMed: 23965983]
22. Leone P, et al. MHC class I antigen processing and presenting machinery: organization, function, and defects in tumor cells. *J Natl Cancer Inst* 105, 1172–1187 (2013). [PubMed: 23852952]
23. Palmer KJ, Harries M, Gore ME & Collins MK Interferon-alpha (IFN-alpha) stimulates anti-melanoma cytotoxic T lymphocyte (CTL) generation in mixed lymphocyte tumour cultures (MLTC). *Clin Exp Immunol* 119, 412–418 (2000). [PubMed: 10691911]
24. Neumann H, Schmidt H, Cavalie A, Jenne D & Wekerle H Major histocompatibility complex (MHC) class I gene expression in single neurons of the central nervous system: differential regulation by interferon (IFN)-gamma and tumor necrosis factor (TNF)-alpha. *J Exp Med* 185, 305–316 (1997). [PubMed: 9016879]
25. Slatter TL, et al. Antitumor cytotoxicity induced by bone-marrow-derived antigen-presenting cells is facilitated by the tumor suppressor protein p53 via regulation of IL-12. *Oncoimmunology* 5, e1112941 (2016). [PubMed: 27141366]
26. Campbell IL, Oxbrow L, West J & Harrison LC Regulation of MHC protein expression in pancreatic beta-cells by interferon-gamma and tumor necrosis factor-alpha. *Mol Endocrinol* 2, 101–107 (1988). [PubMed: 3135484]
27. Mattsson R, et al. Placental MHC class I antigen expression is induced in mice following in vivo treatment with recombinant interferon-gamma. *J Reprod Immunol* 19, 115–129 (1991). [PubMed: 1901087]

28. Lavi E, et al. Tumor necrosis factor induces expression of MHC class I antigens on mouse astrocytes. *J Neuroimmunol* 18, 245–253 (1988). [PubMed: 2452831]
29. Andrews JS, Berger AE & Ware CF Characterization of the receptor for tumor necrosis factor (TNF) and lymphotoxin (LT) on human T lymphocytes. TNF and LT differ in their receptor binding properties and the induction of MHC class I proteins on a human CD4+ T cell hybridoma. *J Immunol* 144, 2582–2591 (1990). [PubMed: 1969453]
30. Locksley RM, Killeen N & Lenardo MJ The TNF and TNF receptor superfamilies: integrating mammalian biology. *Cell* 104, 487–501 (2001). [PubMed: 11239407]
31. Chopra M, et al. Exogenous TNFR2 activation protects from acute GvHD via host T reg cell expansion. *J Exp Med* 213, 1881–1900 (2016). [PubMed: 27526711]
32. Lawrence T The nuclear factor NF-kappaB pathway in inflammation. *Cold Spring Harb Perspect Biol* 1, a001651 (2009). [PubMed: 20457564]
33. Ciancio MJ, Hunt J, Jones SB & Filkins JP Comparative and interactive in vivo effects of tumor necrosis factor alpha and endotoxin. *Circ Shock* 33, 108–120 (1991). [PubMed: 2049814]
34. Biesmans S, et al. Peripheral Administration of Tumor Necrosis Factor-Alpha Induces Neuroinflammation and Sickness but Not Depressive-Like Behavior in Mice. *Biomed Res Int* 2015, 716920 (2015). [PubMed: 26290874]
35. Sonabend AM, et al. Medulloblastoma: challenges for effective immunotherapy. *J Neurooncol* 108, 1–10 (2012). [PubMed: 22173741]
36. Sade-Feldman M, et al. Resistance to checkpoint blockade therapy through inactivation of antigen presentation. *Nat Commun* 8, 1136 (2017). [PubMed: 29070816]
37. Paulson KG, et al. Acquired cancer resistance to combination immunotherapy from transcriptional loss of class I HLA. *Nat Commun* 9, 3868 (2018). [PubMed: 30250229]
38. Roberts NJ, Zhou S, Diaz LA Jr. & Holdhoff M Systemic use of tumor necrosis factor alpha as an anticancer agent. *Oncotarget* 2, 739–751 (2011). [PubMed: 22036896]
39. Reagan-Shaw S, Nihal M & Ahmad N Dose translation from animal to human studies revisited. *FASEB J* 22, 659–661 (2008). [PubMed: 17942826]
40. Furman WL, et al. Phase I clinical trial of recombinant human tumor necrosis factor in children with refractory solid tumors: a Pediatric Oncology Group study. *J Clin Oncol* 11, 2205–2210 (1993). [PubMed: 8229135]
41. Seibel NL, et al. Phase I study of tumor necrosis factor-alpha and actinomycin D in pediatric patients with cancer: a Children's Cancer Group study. *J Immunother Emphasis Tumor Immunol* 16, 125–131 (1994). [PubMed: 7804527]
42. Suzuki H, et al. Recurrent non-coding U1-snRNA mutations drive cryptic splicing in Shh medulloblastoma. *Nature* (2019).

REFERENCES FOR METHODS

43. Futterer A, Mink K, Luz A, Kosco-Vilbois MH & Pfeffer K The lymphotoxin beta receptor controls organogenesis and affinity maturation in peripheral lymphoid tissues. *Immunity* 9, 59–70 (1998). [PubMed: 9697836]
44. Banks TA, et al. A lymphotoxin-IFN-beta axis essential for lymphocyte survival revealed during cytomegalovirus infection. *J Immunol* 174, 7217–7225 (2005). [PubMed: 15905567]
45. Hong H, et al. Suppression of induced pluripotent stem cell generation by the p53-p21 pathway. *Nature* 460, 1132–1135 (2009). [PubMed: 19668191]
46. Kojima K, et al. p53 activation of mesenchymal stromal cells partially abrogates microenvironment-mediated resistance to FLT3 inhibition in AML through HIF-1alpha-mediated down-regulation of CXCL12. *Blood* 118, 4431–4439 (2011). [PubMed: 21868571]
47. Tsuchiya S, et al. Establishment and characterization of a human acute monocytic leukemia cell line (THP-1). *Int J Cancer* 26, 171–176 (1980). [PubMed: 6970727]
48. Sundstrom C & Nilsson K Establishment and characterization of a human histiocytic lymphoma cell line (U-937). *Int J Cancer* 17, 565–577 (1976). [PubMed: 178611]

49. Smardova J, Pavlova S, Svitakova M, Grochova D & Ravcukova B Analysis of p53 status in human cell lines using a functional assay in yeast: detection of new non-sense p53 mutation in codon 124. *Oncol Rep* 14, 901–907 (2005). [PubMed: 16142349]
50. Weisberg E, et al. Inhibition of Wild-Type p53-Expressing AML by the Novel Small Molecule HDM2 Inhibitor CGM097. *Mol Cancer Ther* 14, 2249–2259 (2015). [PubMed: 26206331]
51. Misuraca KL, et al. Pax3 expression enhances PDGF-B-induced brainstem gliomagenesis and characterizes a subset of brainstem glioma. *Acta Neuropathol Commun* 2, 134 (2014). [PubMed: 25330836]
52. Lewis PW, et al. Inhibition of PRC2 activity by a gain-of-function H3 mutation found in pediatric glioblastoma. *Science* 340, 857–861 (2013). [PubMed: 23539183]
53. Hashizume R, et al. Characterization of a diffuse intrinsic pontine glioma cell line: implications for future investigations and treatment. *J Neurooncol* 110, 305–313 (2012). [PubMed: 22983601]
54. Caretti V, et al. Human pontine glioma cells can induce murine tumors. *Acta Neuropathol* 127, 897–909 (2014). [PubMed: 24777482]
55. Kreft L, et al. ConTra v3: a tool to identify transcription factor binding sites across species, update 2017. *Nucleic Acids Res* 45, W490–W494 (2017). [PubMed: 28472390]

Figure 1: p53-mutant medulloblastomas are resistant to T cell-mediated rejection. MP tumor cells (a and b) or MG tumor cells (c and d) were transplanted into the cerebellum of NSG (black line) or Albino B6 (aB6, red line) mice. Bioluminescence imaging of representative mice (a and c) and survival curves (b and d) are shown (MP NSG, $n = 12$; MP aB6, $n = 12$; MG NSG, $n = 12$ and MG aB6, $n = 45$). (e-f) Intracranially transplanted MP tumors (black) and MG tumors (red) labelled with GFP were harvested 2, 5 and 7 days after transplant ($n=3$ mice per tumor type per day). Percentages of GFP⁺ tumor cells (e) and percentages of CD3⁺ T cells (f) were determined by FACS analysis. The average curve and single data points are shown for each condition. p-values were determined by two-sided unpaired t-test. In (e), * $p=0.009$; ** $p=0.0004$. In (f), ** $p=0.00017$. (g-i) Depletion of T-cells (α Thy1, blue line), or no T cell depletion (Control, black line), were performed in aB6 MG tumor-bearing mice. (g) Bioluminescence imaging of representative mice. (h) Quantification of the average bioluminescence signal (dark lines) and the signals for individual mice (light lines) for each group. (i) Survival curves for control mice ($n = 7$); and α Thy1-treated mice ($n=7$). p-value for survival curve was determined by a two-sided log-rank (Mantel-Cox) test. All error bars represent the mean $\pm$ SD.

Figure 2: Inactivation of p53 results in loss of cell surface MHC-I.

(a-b) MG tumor cells or MG tumor cells overexpressing DNp53 (MG+P) were transplanted into NSG or aB6 mice. Bioluminescence imaging of representative mice (a) and survival curves (n=15) (b) are shown. p-values for the difference in survival between MG and MG+P transplanted into NSG or aB6 were determined by two-sided log-rank (Mantel-Cox) test. In (b), NS (not significant), p=0.1932 (NSG) and ** p=0.0011 (aB6). (c) FACS histograms of MHC-I expression in MP (black) and MG (red) tumor cells. (d) MHC-I expression in MG tumor cells (red) and MG tumor cells overexpressing DNp53 (MG+P, brown), (e) MHC-I expression in MP-p53 knockout (MPKO) tumor cells (black) and MPKO tumor cells overexpressing p53 (MPKO+p53, blue). Isotype controls are shown in gray. Quantification

of the mean fluorescence intensity (MFI) values for three independent tumors are shown below each histogram; data points represent MFIs for individual tumor samples. p-values were determined by two-sided unpaired t-test. p-values are indicated on each bar graph. (f) MG tumor cells generated from MHC-I knockout (MHC-I KO) mice were transplanted into NSG or aB6 mice. Survival curves are shown (n=15 for each group). p-values for the difference in survival between MG (aB6) and MG MHC-I KO (aB6) were determined by a two-sided log-rank (Mantel-Cox) test. $p=1.9\text{E-}12$ (NSG versus aB6) and $p=5.7\text{E-}07$ (MHC-I KO versus MHC-I WT). (g) Expression of MHC class I K(b) and D(b) mRNA was analyzed by qRT-PCR. p-values were determined by two-way ANOVA test. NS: $p=0.448$ (MG) ; $p=0.099$ (MG+P). (h-i) Protein levels of MHC class I and β -Actin were determined by western blotting and quantification of 3 independent experiments is shown (i). Western blots are cropped at the molecular weights for MHC-I (45 kDa) and β -actin (42 kDa); original blots are available in Source Data. p-values were determined by two-way ANOVA test, NS: $p=0.267$ (MG) ; $p=0.614$ (MG+P). Individual data points represent expression in independent tumor samples (n=3). All error bars represent means $\pm$ SD.

Figure 3: p53 regulates expression of Erap1 and Tap1.

Erap1 (a, c, e) and Tap1 (b, d, e) expression in normal neural stem cells (NSCs, n=3), MP tumors (n=3), MG tumors (n=3), MG tumors overexpressing DNp53 (MG+P) (n=3) and MP tumors overexpressing GFI1 (MP+G) (n=3) were measured by qRT-PCR (a-d) and by western blotting (e); quantification of 3 independent experiments is shown below the western blot. Error bars represent means $\pm$ SD. Western blots are cropped at the molecular weights for Erap1 (120 kDa), Tap1 (68kDa) and β -actin (42 kDa); original blots are available in Source Data. p-values were determined by two-sided unpaired t-test. In (a), NS: p=0.21 (MG vs NSC), ** p=0.0019 (MG vs MP). In (b) NS, p=0.33 (MG vs NSC); *** p=1.9E-05. In (c), *** p=1E-04 (MP vs MG) and p=8E-05 (MG vs MG+P). In (d), ***

p=2.3E-05 (MP vs MG), ** p=0.003 (MG vs MG+P, Tap1). In (e), * refers to p=0.002 (MP vs MG, Erap1), p=0.0052 (MG vs MG+P, Erap1), p=0.0097 (MP vs MG, Tap1), p=0.0049 (MG vs MG+P, Tap1). (f-g) Analysis of *ERAP1* (f) and *TAP1* (g) mRNA levels in human Sonic Hedgehog-associated MB with mutant (red) or wild-type (WT, blue) TP53. Fragments Per Kilobase per Million mapped reads (FPKM) of *ERAP1* and *TAP1* are shown. p-values were determined by two-sided Wilcoxon rank sum test. Boxplot center lines show median, box limits indicate the 25th and 75th percentiles, lower and upper whiskers extend 1.5 times the interquartile range (IQR) from the 25th and 75th percentiles, respectively. (h) Putative p53 binding sites in the mouse *Erap1* and *Tap1* promoters. (i-j) Chromatin Immunoprecipitation (ChIP) and PCR analysis of p53 binding site-containing regions in the *Erap1* (i) and *Tap1* (j) promoters. qPCR results for anti-p53 ChIP (IP p53) or isotype control ChIP (IP Ctl) in MG (n=3) and MP (n=3) tumor cells. p-values were determined by two-sided unpaired t-test * p=0.0046 (Erap1) and ** p=0.0011 (Tap1). (k) Putative p53 binding sites in the human *ERAP1* and *TAP1* promoters. (l-m) ChIP and PCR analysis of the p53 binding site-containing regions in the *ERAP1* (l) and *TAP1* (m) promoters. qPCR results for anti-p53 ChIP (IP p53) or isotype control ChIP (IP Ctl) in p53 mutant PDXs (RCMB18, Icb984, BT084) and p53 wild-type PDXs (Icb1299, RCMB40) (n=3 for each tumor type). p-values were determined by two-sided unpaired t-test. * p=0.026 (Icb1299, ERAP1), p=0.01 (Icb1299, TAP1), p=0.002 (RCMB40, TAP1) ; ** p=0.001 (RCMB40, ERAP1).

Figure 4: Erap1 and Tap1 are necessary for cell surface MHC-I expression and T cell mediated rejection.

(a-d) MG tumor cells were transduced with control shRNA (shCtl) or shRNAs targeting Erap1 (shErap1#1, shErap1#2). Knockdown efficiency was determined by western blotting in three independent tumor samples (a). MHC-I expression was determined by FACS in control cells (shCtl, black) and Erap1 knockdown cells (shErap1, red). Western blots are cropped at the molecular weights for Erap1 (120 kDa) and β -actin (42 kDa); original blots are available in Source Data. (b). Quantification of the mean fluorescence intensity (MFI) values for three independent tumors is shown below each histogram; data points represent MFIs for individual tumor samples. p-values, determined by two-sided unpaired t-test, are

indicated on each bar graph. (c-d) Erap1 knockdown cells were transplanted into aB6 mice. Bioluminescence imaging of representative mice (c) and survival curves (n=6) (d) are shown. p-values for the difference in survival between shErap1 and shCtl were determined using the two-sided log-rank (Mantel-Cox) test. * p=0.024; ** p= 0.0075. (e-i) MP tumor cells were transduced with empty vector (vect) or vectors encoding Erap1, Tap1 or both. Erap1 (e) and Tap1 (f) expression levels were assessed by western blotting in three independent tumor samples. Western blots are cropped at the molecular weights for Erap1 (120 kDa), Tap1 (68kDa) and β -actin (42 kDa); original blots are available in Source Data. (g) Analysis of MHC-I expression by FACS in control cells (vect, black) and Erap1 + Tap1 overexpressing cells (pink). Quantification of the mean fluorescence intensity (MFI) for three independent experiments is shown below the histogram; data points represent MFIs for individual tumor samples. The p-value was determined by two-sided unpaired t-test. (h, i) MP tumor cells transduced with control vector (vect, black), Tap1 (green), Erap1 (blue) or Erap1 + Tap1 (pink) were orthotopically transplanted into aB6 mice. Bioluminescence imaging of representative mice (h) and survival curves (n=6 per group) (i) are shown. p-values were determined by the two-sided log-rank (Mantel-Cox) test. NS (Ctl vs. Tap1), p=0.79; * (Ctl vs. Erap1) p=0.0047; ** (Ctl vs. Erap1+Tap1) p=0.0005. Mice carrying MP tumors expressing Erap1 + Tap1 survive significantly longer than mice carrying tumors expressing Erap1 alone (p=0.0016 for Erap1 vs. Erap1 + Tap1) or Tap1 alone (p = 0.0005 for Tap1 vs. Erap1 + Tap1).

*** $p=0.0004$. In (j), NS, $p=0.453$, * $p=0.0086$, ** $p=0.003$. (k) Tap1 and Erap1 protein levels were assayed by western blotting. Western blots are cropped at the molecular weights for Erap1 (120 kDa), Tap1 (68kDa) and β -actin (42 kDa); original blots are available in Source Data. Relative protein levels of Erap1 (l) and Tap1 (m) normalized to actin are shown for three independent samples. Error bars represent mean $\pm$ SD; data points represent expression values for individual tumor samples. p-values were determined by two-sided unpaired t-test. In (l), NS $p=0.133$, * $p=0.0025$, ** $p=0.0004$; In (m), NS = 0.378, * $p=0.029$, ** $p=0.0116$.

Figure 6: TNF induction of MHC-I depends on TNFR2 signaling through NFκB.

(a, b) FACS analysis of MHC-I expression in MP tumor cells that were untreated (Ctl, black histograms) or treated for 48h with TNFR1 agonist (0.5μg/mL, purple histogram, a) or TNFR2 agonist (1.67nM, red histogram, b). (c) MP tumor cells generated from TNFR2 knockout (TNFR2 KO) mice were treated for 48h with no stimulus (Ctl, black histogram) or with TNF (50 pg/mL, green histogram) and then analyzed by FACS. Quantification of the mean fluorescence intensity (MFI) for three independent experiments is shown below each histogram; data points represent MFIs for individual tumor samples. p-values were determined by two-sided unpaired t-test. p-values are indicated on the corresponding graphs. (d-f) MP tumor cells were untreated (NT) or treated with TNF for 15', 30' or 1h. Expression

of RelA and RelB protein were assessed by western blotting in nuclear extracts (d) and in total cellular protein extracts (e). Histone H3 and GAPDH were used as controls for nuclear and total extracts respectively. Western blots are cropped at the molecular weights for RelA (65 kDa), RelB (68kDa), Histone H3 (17kDa) and GAPDH (37 kDa); original blots are available in Source Data. (f) Quantification of Western blots for RelA (black) and RelB (blue) protein levels in nucleus relative to levels in total extract at 0', 15', 30' and 60' for three independent experiments. p-values were determined by two-sided unpaired t-test. For RelA, * p=0.049 (15') ; ***p=0.0032 (30') ; **p=0.0043 (1h); for RelB: NS, p=0.088 (15') ; *p=0.037 (30') ; NS p=0.073 (1h). (g, i) Putative binding sites for the RelA-p50 heterodimer in the mouse *Erap1* and *Tap1* promoters. (h, j) MP tumor cells were treated for 1h with no stimulus (ctl), TNF or LtβRag. Chromatin Immunoprecipitation (ChIP) was performed using antibodies specific for RelA (h) or p50 (j), and precipitates were analyzed by qPCR for the presence of the RelA-p50 binding site-containing regions of the *Erap1* and *Tap1* promoters. Data represent the mean fold-induction ± SD. p values were determined by two-sided unpaired t test. In (h), * p=0.0034, **p=0.003, ***p=2.6E-06 (TNF, *Erap1*), p=1E-04 (TNF, *Tap1*). (k) MP tumor cells were transduced with control shRNA (shCtl) or shRNAs targeting RelA (shRelA#832, shRelA#833), p50 (shp50#484, shp50#485) or RelB (shRelB#494, shRelB#495). MHC-I expression was determined by FACS in control cells (shCtl) and RelA, p50 or RelB knockdown cells treated in vitro for 48h with vehicle (DMSO, ctl), TNF or LtβRag. Data shown represent quantification of mean fluorescence intensity (MFI) for three independent experiments; data points represent MFIs for individual tumor samples. p values were determined by two-sided unpaired t test. shCtl : ** p=0.0019, *** p=1.1E-05; shRelA (#832): NS, Not significant p=0.063 (TNF) and p=0.098 (LtβRag); shRelA (#833): *p=0.035, **p=0.009; shp50 (#484): NS, Not significant p=0.341 (TNF), p=0.125 (LtβRag); shp50 (#485): NS, p=0.601, *p=0.049 ; shRelB (#494) *** p=1.8E-05 (TNF), p=2.3E-04 (LtβRag); shRelB (#495): *** p=9E-06 (TNF), p=1.6E-05 (LtβRag).

Figure 7: TNF enhances responses to immune checkpoint inhibitors in vivo.

(a-c) MP tumor cells were transplanted into aB6 mice, and 10 days later mice were treated with $0.1 \mu\text{g/kg}$ (a), $0.5 \mu\text{g/kg}$ (b), $1 \mu\text{g/kg}$ (c) TNF (green histograms) or vehicle (Ctl, black histograms). 24h after treatment tumors were harvested and MHC-I expression was determined by FACS. (d) Murine p53 knockout DIPG tumor cells were transplanted into aB6 mice, and 21 days later mice were treated with $1 \mu\text{g/kg}$ TNF (green) or vehicle (Ctl, black). 24h after treatment MHC-I expression was determined by FACS. Quantification of the mean fluorescence intensity (MFI) for three independent mice per condition is shown below each histogram; data points represent MFIs for individual tumor samples. p-values were determined by two-sided unpaired t-test. p-values are indicated on the corresponding

graphs. (e-h) MP tumor cells (e-f) or murine DIPG tumor cells (g-h) were transplanted into aB6 mice. 15 days later (for MP) or 21 days later (for DIPG) mice were treated with vehicle (Ctl, black), anti-PD-1 (10 mg/kg, blue), TNF (1 µg/kg, green) or anti-PD-1 and TNF (orange). For combinations of TNF and anti-PD-1, each cycle of treatment consisted of a dose of TNF on day 1, a dose of anti-PD-1 on day 2, and no treatment on day 3. Mice received up to 20 cycles of treatment. (e, g) Bioluminescence imaging of representative mice. (f, h) Survival curves (n=10 per group for MP and n=15 per group for DIPG). p-values for the difference in survival between Ctl mice and TNF + anti-PD-1 treated mice were determined by a two-sided log-rank (Mantel-Cox) test. (f) * p=0.019, ***p=2.5E-06 (TNF), p=2.4E-07 (TNF + anti-PD-1); (h) * p=0.028, *** p=3.6E-09 (TNF), p=1E-12 (TNF + anti-PD-1).