Aortic root remodeling in bicuspid and tricuspid aortic valves—long-term results

Shunsuke Miyahara¹ · Irem Karliova¹ · Christian Giebels¹ · Ulrich Schneider¹ · Shunsuke Matsushima¹ · Hans-Joachim Schäfers¹

Received: 4 December 2018 / Revised: 14 January 2019 / Accepted: 16 January 2019 / Published online: 15 February 2019
© Indian Association of Cardiovascular-Thoracic Surgeons 2019

Abstract

Background Root remodeling is one form of valve-preserving root replacement to treat patients with aortic regurgitation (AR) and root aneurysm. We have consistently used it for patients with and without connective tissue disease and different aortic valve morphologies. The objective of this retrospective study was to review 23 years of experience with root remodeling.

Methods and patients Between 10/95 and 9/2018, 1004 of 1038 root remodeling procedures were performed in patients with tricuspid ($n = 589$) or bicuspid ($n = 414$) anatomy of the aortic valve in our institution. Aortic aneurysm was present in 932 cases, 73 procedures were performed for acute aortic dissection type A. The severity of aortic regurgitation ranged from grade I to grade IV (mean 2.5 ± 0.8).

Results All patients underwent root remodeling, concomitant operations were performed in 433, and cusp repair in 883 instances. Hospital mortality was 2%. Overall freedom from reoperation was 92% at 10 years and 89% at 15 years. It was 94% for tricuspid valves at 10 and 15 years, and 88% for bicuspid aortic valves at 10 years and 80% at 15 years ($p = 0.003$).

Conclusion In conclusion, root remodeling is a viable option in valve-preserving root replacement. If combined with careful assessment and, if necessary, correction of aortic valve form reproducible restoration of aortic valve function can be achieved.

Keywords Aortic root remodeling · Long term · Bicuspid valve

Introduction

Root remodeling was proposed by Yacoub as a valve-preserving operation to treat patients with AR and root aneurysm [1]. The goal was to normalize valve form and function by replacing all dilated aortic root tissue and restoring normal geometry.

The early results were positive, but the technique did not always result in adequate valve competence [2, 3]. In particular, the late results of the original series were suboptimal, in that a relevant proportion of patients developed recurrent and relevant AR [4]. The inferior stability has been related to the lack of annular stabilization in root remodeling [5].

Based on our own positive results and the better preservation of cusp motion, we routinely employed root remodeling for root

aneurysm, initially for patients without marked annular dilatation [6]. We observed that root aneurysm not infrequently coexisted with cusp prolapse [7], and we later found that the probability of prolapse seemed to increase with increasing degree of AR [8]. In addition, we realized that cusp prolapse could be induced or aggravated through reduction of intercommissural distance in root replacement [9]. We found that the combination of root remodeling and cusp repair was safe and led to improved aortic valve function postoperatively [10].

The finding of postoperative prolapse stimulated research on the normal configuration of aortic valve function. We proposed the effective height concept with the possibility of intraoperative measurement of cusp configuration; this helped in achieving normalized postoperative valve function irrespective of the preoperative degree of AR [11, 12]. Based on decreased durability of isolated aortic valve repair we introduced the concept of suture annuloplasty also as addition to root remodeling [13]. Its use resulted in higher proportion of competent aortic valves [14].

After initial careful exploration of root remodeling [15], we were positively impressed by the reproducibility of achieving

✉ Hans-Joachim Schäfers
h-j.schaefers@uks.eu

¹ Department of Thoracic and Cardiovascular Surgery, Saarland University Medical Center, Homburg, Saar, Germany

good aortic valve function. We extended its application to bicuspid aortic valves [16]. We hypothesized that normalized aortic valve configuration should lead to low early failure rate, and the near-physiologic cusp motion also to good valve durability.

The objective of this retrospective study was to review 23 years of experience with root remodeling, both for tricuspid and also bicuspid aortic valves.

Methods and patients

Between 10/95 and 9/2018, root remodeling was performed in 1038 patients in our institution. The aortic valve anatomy was tricuspid in 589, bicuspid in 414, and unicuspid in 34 individuals. The individuals with unicuspid valves were excluded from this analysis. The mean age of the remaining 1004 patients ranged from 3 to 86 years (mean 54 ± 15 years), 833 were male.

Aortic aneurysm was present in 932 cases; 73 procedures were performed for acute aortic dissection type A. Partial aortic arch involvement was present in 264 instances with aneurysm; in 44 cases, the aneurysm involved the total arch.

The primary indication for surgery was severe AR ($n = 600$), acute dissection ($n = 73$), or aortic aneurysm ($n = 312$). The severity of aortic regurgitation ranged from grade I to grade IV (mean 2.5 ± 0.8 median 2.5).

Cardiac comorbidity included coronary artery disease ($n = 132$), mitral regurgitation ($n = 43$), tricuspid regurgitation ($n = 37$), and persistent atrial fibrillation ($n = 53$). Previous aortic or cardiac surgery had been performed in 22 patients.

The operative technique has been described in detail previously and has undergone only minimal changes [17]. The chest was opened by a median sternotomy, and aortic and right atrial cannulation was used; only in the presence of acute dissection the right axillary artery was used for arterial inflow. Following root mobilization and excision of sinus wall a tubular graft was tailored to accommodate the configuration of the aortic root. For tricuspid valves, a 120° configuration was used; in most instances of bicuspid valves, two symmetric tongues with a 180° orientation were created. The graft was sutured to the cusp insertion lines and the cusps were inspected for prolapse. If present, prolapse was corrected by central plication sutures on the cusp margin [18], if necessary extending it into the belly of the cusp.

Originally remodeling was primarily used for patients with a basal diameter of less than 30 mm, and a graft size chosen 1 to 2 mm smaller than the basal diameter. With increasing experience, all patients were included, and graft size was chosen according to the body surface area of the patient (24 mm for $< 1.8\text{m}^2$, 26 mm for 1.9 to 2.2m^2 , and 28 mm for 2.3m^2 and larger). Since 2004, visual inspection for prolapse was replaced by measurement of effective height [11]; any

effective height of less than 9 mm was considered as prolapse and corrected. Since 2009, a suture annuloplasty was added in the last 503 patients; braided polyester was utilized in 66 and PTFE in 437 [13]. Postoperative anticoagulation consisted of aspirin (100 mg/day) given for 2 months unless persistent atrial fibrillation had been present preoperatively; in this instance, coumadin was used.

All patients were followed; follow-up was complete in 1000 patients (99.5%). Cumulative follow-up was 7607 patient years, mean 7.6 ± 5.2 years.

Statistical methods

Data are expressed as means $\pm$ standard deviation and categorical data are expressed frequencies or ratios of patients. Freedom from reoperation was analyzed using the Kaplan-Meier method. Statistical differences in Kaplan-Meier survival estimates were determined using the log-rank test. Values of $p < 0.05$ were considered to indicate statistical significance. Data were analyzed using JMP 9.0.2 for Windows (SAS Institute, Cary, NC).

Results

All patients underwent root remodeling with replacement of the ascending aorta. Concomitant procedures were performed in 433 patients; these included partial ($n = 264$) or total arch replacement ($n = 44$), coronary artery bypass grafting ($n = 132$), mitral repair ($n = 43$), tricuspid repair ($n = 37$), or left atrial ablation ($n = 53$).

Cusp repair was performed in 883 patients, with plication for prolapse correction being the most frequent maneuver ($n = 848$; 84%). In 71 patients, partial cusp replacement or augmentation with pericardium was performed, mostly in the setting of bicuspid anatomy. In tricuspid aortic valves, 1 cusp was plicated in 188, 2 cusps in 194, and 3 cusps in 98 cases. In bicuspid valves, 1 cusp was plicated in 81 and both in 275 instances.

The time of extracorporeal circulation varied from 51 to 464 min (mean 103 ± 33), myocardial ischemic time ranged from 42 to 222 min (mean 75 ± 19).

Sixteen patients died in hospital for a mortality of 1.6%. The causes of death were cardiac ($n = 5$), multi-organ failure ($n = 6$), sepsis ($n = 2$), cerebral ($n = 2$), and pulmonary embolism ($n = 1$). No pacemaker implantation was required for atrioventricular conduction disturbance. In 1 instance, a pacemaker was implanted for sinus node dysfunction following atrial ablation. Actuarial survival was 94.3% at 5, 89.6% at 10 years, and 86.1% at 15 years (Fig. 1).

A total of 54 patients underwent valve-related reoperation between 1 month and 15 years postoperatively. The aortic

Fig. 1 Kaplan-Meier analysis of overall survival following root remodeling. Dotted lines indicate 95% confidence interval

valve morphology was tricuspid in 23 and bicuspid in 31 cases. Reasons for reoperation were secondary AR ($n = 39$), endocarditis ($n = 7$), and aortic stenosis ($n = 6$).

Overall freedom from reoperation was 91.8% at 10 years and 88.5% at 15 years (Fig. 2a). It was 94.2% at 10 and 93.6% at 15 years for tricuspid valves, and 87.5% at 10 years and 79.5% at 15 years for bicuspid aortic valves. Freedom from reoperation was significantly better in tricuspid versus bicuspid anatomy ($p = 0.0030$; Fig. 2b). Actual freedom from AR $\geq 2+$ in all study cohort was 89.1% and 86.5% at 10 and 15 years, respectively (Fig. 3). In this analysis, there was no significant difference between tricuspid valve and bicuspid valve ($p = 0.76$). Patients with dilated aortic annulus (≥ 30 mm) had no inferiority in terms of freedom from reoperation ($p = 0.73$) (Fig. 4), also in patients with tricuspid valve ($p = 0.16$) and bicuspid valve ($p = 0.14$).

Valve-related complications were rare. Early endocarditis (< 3 months) occurred in 1 instance and led to reoperation in the acute stage of the disease. Late endocarditis was observed in 6 patients between 16 and 141 months postoperatively. They were all treated surgically.

Discussion

Root remodeling was designed as a technique of valve-sparing aortic replacement. Despite positive initial results, subsequently suboptimal durability was reported, both in root aneurysm and acute dissection [19]. In particular, patients with connective tissue disease seemed to have inferior valve stability after root remodeling, even though 10-year freedom from reoperation was identical to reimplantation [20]. Most authors

have related the limitations to the lack of annular stabilization in root remodeling [20]; exact analyses of the valve failures, however, have not been published.

Stimulated by the early results of Yacoub, we have also used root remodeling, initially in patients with lesser degree of annular dilatation [17]. We found remodeling to be the shorter procedure, saving an average of 30 min of myocardial ischemic time compared to valve reimplantation. With time, the difference has become even more apparent, as our average ischemic time for isolated remodeling is now in the range of 50 to 60 min compared to approximately 2 h for reimplantation in the hands of experienced surgeons [21, 22]. In addition, remodeling is not limited in its applicability by muscle in the sinuses, i.e., discrepancy between aorto-ventricular junction and basal ring [23].

Later, we found out that reduction of root dimensions through aortic replacement (of any form) could lead to induction of cusp prolapse through reduction of intercommissural distance [9]. Since prolapse was found to be safely corrected by cusp plication [7], we combined these techniques in order to generate near-normal aortic valve form and function. This was facilitated by the use of the effective height concept [11], which had not been utilized by us in the early experience; it was later also used by other groups [24, 25, 30].

Using this concept, we have continued to apply root remodeling for an increasing proportion of patients with root aneurysm. In fact, remodeling has become our primary approach to all individuals with root enlargement and a non-calcified aortic valve irrespective of the degree of preoperative AR. We have been able to show that annular diameter is already reduced through remodeling even in the absence of an annuloplasty [26]. Also in root dilatation with acute

Fig. 2 **a** Kaplan-Meier analysis of freedom from reoperation following root remodeling. Dotted lines indicate 95% confidence interval. **b** Kaplan-Meier analysis of freedom from reoperation following root remodeling for different aortic valve morphologies. Stability is significantly superior for tricuspid aortic valve anatomy compared to bicuspid. TAV tricuspid aortic valve, BAV bicuspid aortic valve

dissection, we have seen the best stability with remodeling [27]. One may argue that reduction of annular size improves coaptation height [28] and thus compensates inaccuracies of cusp repair. If the surgeon feels that an annuloplasty may improve the durability of his repair (as yet unproven), a simple suture annuloplasty [29] may be added. Another form of annuloplasty may be used; it has been associated, however, with prolonged myocardial ischemic times [30].

The current results confirm the hypothesis that good early and stable late valve function can be achieved with this approach. Our current results have to be seen in the light that the majority of patients in our series had severe AR preoperatively, while this was less in other series [5, 12, 19, 20, 31, 32]. In retrospect, the suboptimal results with root remodeling in other groups can probably be best explained by the limitations in

detection of cusp prolapse use its correction. Cusp configuration in those series was assessed by judgment only; even though most operations were done by experienced surgeons, symmetric cusp prolapse may have evaded intraoperative inspection. In fact, persistent or recurrent prolapse (defined by suboptimal effective height) was the strongest predictor of failure in our earlier experience; one has to consider that most of the procedures with this finding were performed before the introduction of measuring effective height intraoperatively.

Prolapse can also explain the observed discrepancies between root remodeling and valve reimplantation [2, 20, 32], since generally smaller grafts are used for root remodeling than for valve reimplantation. Reproducible detection of generalized prolapse requires a specific approach, such as the measurement of effective height [11]. This consideration would

Fig. 3 Kaplan-Meier analysis of freedom from AR $\geq 2+$ following root remodeling. Dotted lines indicate 95% confidence interval. AR aortic regurgitation

also explain our previous observation that larger root size and larger annular size were associated with lesser valve durability [33], most likely due to unrecognized induction of prolapse before the use of the effective height concept [9]. If this prolapse is corrected systematically—as we have done in the past 14 years—the results should at least be comparable. Indeed, our results in patients with tricuspid aortic valves equal those that have been obtained by others with valve reimplantation.

We have also observed that remodeling accommodates the specifics of root anatomy seen in bicuspid valve [16] anatomy. We were positively impressed by the early results [16], and initially the data in the setting of bicuspid anatomy were even superior to those with tricuspid valves [6]. Based on our experience with bicuspid valves [34], we have slightly modified the approach in bicuspid valves. In most BAVs with

configuration of the commissures $> 140^\circ$, we now create complete symmetry of the root by creating two equal tongues. In very asymmetric BAVs with a commissural orientation closer to 120° , we now employ a concept similar to tricuspid valves. This has improved cusp mobility, even though we have not yet seen an effect on durability.

Despite good valve stability, we have explored whether valve function and durability could be improved further by adding an annuloplasty [14]. So far, we have not seen a reduction in the early and mid-term need for reoperation. Nonetheless, as we reported previously for repair of bicuspid valves, the use of annuloplasty has increased the proportion of valves completely competent [35]. It remains to be seen whether annuloplasty will have an additional positive effect on long-term durability.

Fig. 4 Kaplan-Meier analysis of freedom from reoperation following root remodeling in patients with and without dilation of AVJ. AVJ aorto-ventricular junction

The limitations of the current study are primarily related to its retrospective nature, the stepwise adaptations to the learning process, and the lack of a control group. On the other hand, the basic principles of the operation have remained largely constant over time. The results may not be transferable to a setting in which eye-balling only is used for assessment of valve configuration, since intraoperative measurement of effective height has been employed in the vast majority of our operations. Finally, longer follow-up will be needed to determine the very long-term results of this procedure.

In conclusion, root remodeling continues to be a valid option in valve-preserving root replacement. If combined with careful assessment and, if necessary, correction of aortic cusp prolapse, reproducible restoration of aortic valve function can be achieved. The concept of root remodeling can be applied to tricuspid as well as bicuspid aortic valves, even though long-term durability is inferior with non-tricuspid anatomy.

Compliance with ethical standards

This study was approved by the local ethics committee. Informed consent was waived for data analysis under anonymized form.

Conflict of interest The authors declare that they have no conflicts of interest.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Sarsam MA, Yacoub M. Remodeling of the aortic valve anulus. *J Thorac Cardiovasc Surg.* 1993;105:435–8.
- Luciani GB, Casali G, Tomezzoli A, Mazzucco A. Recurrence of aortic insufficiency after aortic root remodeling with valve preservation. *Ann Thorac Surg.* 1999;67:1849–52.
- Leyh RG, Fischer S, Kallenbach K, et al. High failure rate after valve-sparing aortic root replacement using the "remodeling technique" in acute type A aortic dissection. *Circulation.* 2002;106:1229–33.
- Yacoub MH, Gehle P, Chandrasekaran V, Birks EJ, Child A, Radley-Smith R. Late results of a valve-preserving operation in patients with aneurysms of the ascending aorta and root. *J Thorac Cardiovasc Surg.* 1998;115:1080–90.
- Erasmí AW, Sievers HH, Bechtel JF, Hanke T, Stierle U, Misfeld M. Remodeling or reimplantation for valve-sparing aortic root surgery? *Ann Thorac Surg.* 2007;83:S752–6.
- Aicher D, Langer F, Lausberg H, Bierbach B, Schäfers HJ. Aortic root remodeling: ten-year experience with 274 patients. *J Thorac Cardiovasc Surg.* 2007;134:909–15.
- Langer F, Graeter T, Nikoloudakis N, Aicher D, Wendler O, Schäfers HJ. Valve-preserving aortic replacement: does the additional repair of leaflet prolapse adversely affect the results? *J Thorac Cardiovasc Surg.* 2001;122:270–7.
- Asano M, Kuniyama T, Aicher D, El Beyrouti H, Rodionychewa S, Schäfers HJ. Mid-term results after sinotubular junction remodeling with aortic cusp repair. *Eur J Cardiothorac Surg.* 2012;42:1010–5.
- Schäfers HJ, Aicher D, Langer F, Lausberg HF. Preservation of the bicuspid aortic valve. *Ann Thorac Surg.* 2007;83:S740–5.
- Langer F, Aicher D, Kissinger A, et al. Aortic valve repair using a differentiated surgical strategy. *Circulation.* 2004;110:1167–73.
- Schäfers HJ, Bierbach B, Aicher D. A new approach to the assessment of aortic cusp geometry. *J Thorac Cardiovasc Surg.* 2006;132:436–8.
- Lansac E, Di Cerna I, Sleilaty G, et al. An aortic ring: from physiologic reconstruction of the root to a standardized approach for aortic valve repair. *J Thorac Cardiovasc Surg.* 2010;140:S28–35.
- Schäfers HJ, Aicher D. Root remodeling for aortic root dilatation. *Ann Cardiothorac Surg.* 2013;2:113–6.
- Aicher D, Schneider U, Schmied W, Kuniyama T, Tochii M, Schäfers HJ. Early results with annular support in reconstruction of the bicuspid aortic valve. *J Thorac Cardiovasc Surg.* 2013;145:S30–4.
- Graeter TP, Aicher D, Langer F, Wendler O, Schäfers HJ. Mid-term results of aortic valve preservation: remodelling vs. reimplantation. *Thorac Cardiovasc Surg.* 2002;50:21–4.
- Aicher D, Langer F, Kissinger A, Lausberg H, Fries R, Schäfers HJ. Valvesparing aortic root replacement in bicuspid aortic valves: a reasonable option? *J Thorac Cardiovasc Surg.* 2004;128:662–8.
- Schäfers HJ, Fries R, Langer F, Nikoloudakis N, Graeter T, Grundmann U. Valve-preserving replacement of the ascending aorta: remodeling versus reimplantation. *J Thorac Cardiovasc Surg.* 1998;116:990–6.
- Aicher D, Langer F, Adam O, Tscholl D, Lausberg H, Schäfers HJ. Cusp repair in aortic valve reconstruction: does the technique affect stability? *J Thorac Cardiovasc Surg.* 2007;134:1533–8.
- David TE, Feindel CM, Webb GD, Colman JM, Armstrong S, Maganti M. Long-term results of aortic valve-sparing operations for aortic root aneurysm. *J Thorac Cardiovasc Surg.* 2006;132:347–54.
- David TE, Armstrong S, Maganti M, Colman J, Bradley TJ. Long-term results of aortic valve-sparing operations in patients with Marfan syndrome. *J Thorac Cardiovasc Surg.* 2009;138:859–64.
- David TE, Feindel CM, David CM, Manlhiot C. A quarter of a century of experience with aortic valve-sparing operations. *J Thorac Cardiovasc Surg.* 2014;148:872–99.
- Boodhwani M, de Kerchove L, Watremez C, et al. Assessment and repair of aortic valve cusp prolapse: implications for valve-sparing procedures. *J Thorac Cardiovasc Surg.* 2011;141:917–25.
- Anderson RH. Clinical anatomy of the aortic root. *Heart.* 2000;84:670–3.
- Miyahara S, Omura A, Sakamoto T, et al. Impact of postoperative cusp configuration on midterm durability after aortic root reimplantation. *J Heart Valve Dis.* 2013;22:509–16.
- Rankin JS, Gaca JG. Techniques of aortic valve repair. *Innovations (Phila).* 2011;6:348–54.
- Kuniyama T, Arimura S, Sata F, Giebels C, Schneider U, Schäfers HJ. Aortic annulus does not dilate over time after aortic root remodeling with or without annuloplasty. *J Thorac Cardiovasc Surg.* 2018;155:885–94 e3.
- Kuniyama T, Neumann N, Kriechbaum SD, Nomura R, Schäfers HJ. Long-term outcome of aortic root remodeling for patients with and without acute aortic dissection. *Circ J.* 2017;81:1824–31.
- Marom G, Halevi R, Haj-Ali R, Rosenfeld M, Schäfers HJ, Raanani E. Numerical model of the aortic root and valve: optimization of graft size and sinotubular junction to annulus ratio. *J Thorac Cardiovasc Surg.* 2013;146:1227–31.
- Schneider U, Aicher D, Miura Y, Schäfers HJ. Suture annuloplasty in aortic valve repair. *Ann Thorac Surg.* 2016;101:783–5.

30. Lansac E, Di Cerna I, Sleilaty G, et al. Remodeling root repair with an external aortic ring annuloplasty. *J Thorac Cardiovasc Surg.* 2017;153:1033–42.
31. De Paulis R, Scaffa R, Nardella S, et al. Use of the Valsalva graft and longterm follow-up. *J Thorac Cardiovasc Surg.* 2010;140:S23–7.
32. Bethea BT, Fitton TP, Alejo DE, et al. Results of aortic valve-sparing operations: experience with remodeling and reimplantation procedures in 65 patients. *Ann Thorac Surg.* 2004;78:767–72.
33. Kuniyara T, Aicher D, Rodionicheva S, et al. Preoperative aortic root geometry and postoperative cusp configuration primarily determine long-term outcome after valve-preserving aortic root repair. *J Thorac Cardiovasc Surg.* 2012;143:1389–95.
34. Aicher D, Kuniyara T, Abou Issa O, Brittner B, Graber S, Schäfers HJ. Valve configuration determines long-term results after repair of the bicuspid aortic valve. *Circulation.* 2011;123:178–85.
35. Schneider U, Hofmann C, Aicher D, Takahashi H, Miura Y, Schäfers HJ. Suture annuloplasty significantly improves the durability of bicuspid aortic valve repair. *Ann Thorac Surg.* 2017;103:504–10.