affect case fatality (6). These findings are in line with a previous trial that demonstrated reduced intubation rates in the most hypoxemic patients (3) and that mortality is not affected by HFNC put forward the complexity of SARS-CoV-2 infection, in which the underlying lungs do not hold typical features of acute respiratory distress syndrome (7, 8). Instead, acute fibrinous and organizing pneumonia with organizing intraalveolar fibrin associated with notorious endothelial injury can be found in postmortem biopsies (7, 8). Moreover, the proportion of pulmonary embolism and acute kidney and myocardial injury are reported in much higher proportions than in typical acute respiratory distress syndrome (9). Finally, this study highlights that HFNC was as safe as standard oxygen in a large cohort of patients with COVID-19. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Alexandre Demoule, M.D., Ph.D.*
Assistance Publique-Hôpitaux de Paris Sorbonne Université
Paris, France

Antoine Vieillard Baron, M.D., Ph.D.
University of Versailles Saint-Quentin en Yvelines
Villejuif, France

Michael Darmon, M.D., Ph.D.
Université de Paris
Paris, France

Alexandra Beurton, M.D., Ph.D.
Assistance Publique-Hôpitaux de Paris Sorbonne Université
Paris, France

Guillaume Géri, M.D., Ph.D.
University of Versailles Saint-Quentin en Yvelines
Villejuif, France

Guillaume Voiriot, M.D., Ph.D.
Assistance Publique-Hôpitaux de Paris Sorbonne Université
Paris, France

Thibault Dupont, M.D.
Lara Zafrani, M.D., Ph.D.
Université de Paris
Paris, France

Lola Girodias, M.D.
University of Versailles Saint-Quentin en Yvelines
Villejuif, France

Vincent Labbé, M.D.
Martin Dres, M.D., Ph.D.
Muriel Fartoukh, M.D., Ph.D.
Assistance Publique-Hôpitaux de Paris Sorbonne Université
Paris, France

Elie Azoulay, M.D., Ph.D.
Université de Paris
Paris, France

*Corresponding author (e-mail: alexandre.demoule@aphp.fr).

References

1. Mauri T, Turrini C, Eronia N, Grasselli G, Volta CA, Bellani G, *et al*. Physiologic effects of high-flow nasal cannula in acute

hypoxemic respiratory failure. *Am J Respir Crit Care Med* 2017;195:1207–1215.

2. Mauri T, Alban L, Turrini C, Cambiaghi B, Carlesso E, Taccone P, *et al*. Optimum support by high-flow nasal cannula in acute hypoxemic respiratory failure: effects of increasing flow rates. *Intensive Care Med* 2017;43:1453–1463.
3. Frat J-P, Thille AW, Mercat A, Girault C, Ragot S, Perbet S, *et al*; FLORALI Study Group; REVA Network. High-flow oxygen through nasal cannula in acute hypoxemic respiratory failure. *N Engl J Med* 2015;372:2185–2196.
4. Rochwerf B, Granton D, Wang DX, Helviz Y, Einav S, Frat JP, *et al*. High flow nasal cannula compared with conventional oxygen therapy for acute hypoxemic respiratory failure: a systematic review and meta-analysis. *Intensive Care Med* 2019;45:563–572.
5. Alhazzani W, Möller MH, Arabi YM, Loeb M, Gong MN, Fan E, *et al*. Surviving Sepsis Campaign: guidelines on the management of critically ill adults with coronavirus disease 2019 (COVID-19). *Intensive Care Med* 2020;46:854–887.
6. Azoulay E, Lemiale V, Mokart D, Nseir S, Argaud L, Pène F, *et al*. Effect of high-flow nasal oxygen vs standard oxygen on 28-day mortality in immunocompromised patients with acute respiratory failure: the HIGH randomized clinical trial. *JAMA* 2018;320:2099–2107.
7. Copin M-C, Parmentier E, Duburcq T, Poissy J, Mathieu D; Lille COVID-19 ICU and Anatomopathology Group. Time to consider histologic pattern of lung injury to treat critically ill patients with COVID-19 infection. *Intensive Care Med* 2020;46:1124–1126.
8. Lax SF, Skok K, Zechner P, Kessler HH, Kaufmann N, Koelblinger C, *et al*. Pulmonary arterial thrombosis in COVID-19 with fatal outcome: results from a prospective, single-center, clinicopathologic case series. *Ann Intern Med* [online ahead of print] 14 May 2020; DOI: 10.7326/M20-2566.
9. Puelles VG, Lütgehetmann M, Lindenmeyer MT, Sperhake JP, Wong MN, Allweiss L, *et al*. Multiorgan and renal tropism of SARS-CoV-2. *N Engl J Med* 2020;383:590–592.

Copyright © 2020 by the American Thoracic Society

Sex Differences in Right Ventricular–Pulmonary Arterial Coupling in Pulmonary Arterial Hypertension

To the Editor:

Right ventricular (RV) adaptation is a major determinant of survival in pulmonary arterial hypertension (PAH) (1). The right ventricle initially compensates for the PAH-induced increase in afterload through hypertrophy and increased contractile function, maintaining

Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) project number 268,555,672; SFB 1213, Project B08; NIH 1R01HL144727–01A1; and VA Merit Review Award BX002042–07.

Author Contributions: K.T., M.J.R., A.Y., H.A.G., R.N., R.V., H.G., R.J.T., W.S., and T.L.: study design. K.T., M.J.R., H.A.G., H.G., and W.S.: patient recruitment. K.T., M.J.R., A.Y., H.A.G., R.N., R.V., H.G., R.J.T., and W.S.: data collection and analysis. K.T., M.J.R., and H.G.: statistical analyses. K.T., M.J.R., A.Y., H.A.G., R.N., R.V., H.G., R.J.T., W.S., and T.L.: drafting of the manuscript. K.T., M.J.R., A.Y., H.A.G., R.N., R.V., H.G., R.J.T., W.S., and T.L.: critical revision of the manuscript for important intellectual content. K.T. was the principal investigator, had access to all the data in the study, and takes full responsibility for the integrity and accuracy of the data analysis.

This letter has a related editorial.

Originally Published in Press as DOI: 10.1164/rccm.202003-0807LE on June 5, 2020

Table 1. Clinical Characteristics

	Female	Male	P Value
<i>n</i>	33	24	—
Age, yr	53 ± 14	53 ± 12	0.98
PAH subtype, <i>n</i> (%)			0.40
Idiopathic PAH	24 (72.7)	22 (91.7)	—
Associated PAH	9 (27.3)	2 (8.3)	—
6-min-walk distance, m (<i>n</i> = 42)	401 ± 141	343 ± 146	0.20
B-type natriuretic peptide, pg/ml (<i>n</i> = 54)	91 (38–246)	99 (46–295)	0.91
TAPSE/PASP ratio, mm/mm Hg (<i>n</i> = 52)	0.31 (0.21–0.40)	0.33 (0.19–0.38)	0.85
Prior PAH therapy, <i>n</i> (%)			
Dual combination therapy	9 (27.3)	9 (37.5)	0.41
Triple combination therapy	8 (24.2)	9 (37.5)	0.28
Right heart catheterization			
Mean pulmonary artery pressure, mm Hg	42 (36–52)	44 (34–55)	0.84
Pulmonary vascular resistance, Wood units	7.3 (5.6–9.8)	6.3 (4.3–11.3)	0.38
Cardiac index, L/min/m ²	2.8 ± 0.7	2.9 ± 0.6	0.66
Pulmonary artery wedge pressure, mm Hg	7 (6–11)	9 (8–11)	0.18
Right atrial pressure, mm Hg	6 (4–10)	7 (5–9)	0.66
RV pressure–volume loop measurements			
Single-beat			
Ea, mm Hg/ml	0.80 (0.56–0.96)	0.72 (0.46–1.10)	0.97
Ees, mm Hg/ml	0.68 (0.38–0.91)	0.48 (0.29–0.63)	0.03
Ees/Ea ratio	0.92 (0.58–1.24)	0.59 (0.35–0.99)	0.03
Eed, mm Hg/ml (<i>n</i> = 51)	0.14 (0.05–0.23)	0.16 (0.06–0.24)	0.84
Tau, ms	33 ± 8	41 ± 9	<0.01
RVSWI, ml · mm Hg/m ²	1,777 (1,233–2,997)	1,541 (1,035–2,670)	0.54
Multibeat (<i>n</i> = 37, 21 female, 16 male)			
Ees, mm Hg/ml	0.70 ± 0.30	0.46 ± 0.23	0.01
Ees/Ea ratio	0.85 (0.56–1.41)	0.52 (0.29–0.74)	0.01
Cardiac magnetic resonance (<i>n</i> = 55)			
RV mass index, g/m ²	34 ± 14	46 ± 21	0.01
RV end-diastolic volume index, ml/m ²	96 (80–121)	105 (89–162)	0.20
RV mass/volume ratio	0.34 (0.27–0.38)	0.33 (0.28–0.48)	0.40
RV ESV index, ml/m ²	59 (45–92)	69 (59–107)	0.17
RV ejection fraction, %	40 (35–48)	35 (27–41)	0.33
SV/ESV	0.78 ± 0.32	0.57 ± 0.29	0.02

Definition of abbreviations: Ea = arterial elastance; Eed = end-diastolic elastance; Ees = end-systolic elastance; ESV = end-systolic volume; PAH = pulmonary arterial hypertension; PASP = pulmonary artery systolic pressure; RV = right ventricular; RVSWI = right ventricular stroke work index; SV = stroke volume; TAPSE = tricuspid annular plane systolic excursion. Normally and nonnormally distributed data are presented as means ± SD and medians (interquartile ranges), respectively. Between-group differences were analyzed by Student's *t* tests (for normally distributed variables), Pearson chi-square tests (for categorical variables), or Mann-Whitney *U* tests (for nonnormally distributed variables).

RV–pulmonary arterial (PA) coupling. However, as PAH worsens, these compensatory mechanisms fail, leading to RV–PA uncoupling and eventual RV failure (1). In a cardiac magnetic resonance imaging (cMRI) study, female patients with PAH exhibited better RV ejection fraction than their male counterparts (2). RV ejection fraction is load dependent, and pressure–volume (PV) analysis is considered the gold standard to characterize RV–PA coupling (3). To date, no study has directly evaluated whether the superior RV systolic function in female versus male patients with PAH translates into better RV–PA coupling.

We investigated RV–PA coupling using high-fidelity measurements of PV relationships in 57 patients with PAH as part of a prospective study of RV function in PAH and chronic thromboembolic pulmonary hypertension (ClinicalTrials.gov: NCT03403868). The diagnosis of PAH conformed to updated recommendations (4). The study population included a proportion of previously published

patients (3, 5, 6). Real-time PV loop assessment was performed as previously described (3), using the Inca PV Loop System (CD Leycom) with a 4F PV catheter (CA-Nr 41063; CD Leycom) inserted in the RV apex. Measurements were repeated during inferior vena cava occlusion for at least 10 seconds at expiratory breath hold for multibeat determinations. Multibeat end-systolic elastance (Ees) was determined by a tangent fitted on the end-systolic portions of the PV loops at decreasing venous return. We calculated arterial elastance (Ea) as end-systolic pressure/stroke volume (SV). RV–PA coupling was defined as the Ees/Ea ratio. Tau, a load-dependent measure of early diastolic relaxation (7), was calculated from the reciprocal of the natural logarithm of the early maximal fall in ventricular pressure during diastole. We calculated end-diastolic elastance (which is load independent) from the relationship $dP/dV = \alpha\beta \times e^{\beta \times \text{end-diastolic volume}}$ at calculated end-diastolic volumes, where dP/dV is delta

Figure 1. Sex stratification of pressure–volume loop parameters and cardiac magnetic resonance–derived RV mass index and SV/ESV in patients with PAH. Compared with male patients, female patients had significantly improved (A) RV contractility (multibeat Ees; all PAH, $n = 37$; idiopathic PAH, $n = 31$), (B) RV–pulmonary arterial coupling (multibeat Ees/Ea; all PAH, $n = 37$; idiopathic PAH, $n = 31$), (C) early diastolic relaxation (Tau; all PAH, $n = 57$), (D) RV mass index (all PAH, $n = 55$), and (E) SV/ESV (all PAH, $n = 55$). Data are presented as (A, C, and E) means and SD for normally distributed variables or (B and D) medians and interquartile ranges for nonnormally distributed variables. Between-group differences were analyzed by Student's t tests or Mann-Whitney U tests (SPSS, version 25.0; IBM). BSA = body surface area; Ea = arterial elastance; Ees = end-systolic elastance; ESV = end-systolic volume; MWU = Mann-Whitney U test; PAH = pulmonary arterial hypertension; RV = right ventricular; SV = stroke volume.

pressure/delta volume (1). RV mass index and volume measures were determined by cMRI. Data distributions were tested for normality using Kolmogorov–Smirnov tests and visual assessment of histograms. All studies were approved by the local ethics committee. All patients gave written informed consent.

Of the 57 patients with PAH, 37 had multibeam data (Table 1). The study initially included single-beat data only. After establishment of vena cava occlusion, the multibeam method was also used. Therefore, not all patients had multibeam data. Female patients demonstrated significantly higher Ees and Ees/Ea than male patients (Figures 1A and 1B), suggesting higher contractility and better RV-PA coupling despite similar afterload (Ea). Tau and RV mass index were lower in female patients (Figures 1C and 1D), suggesting better diastolic adaptation. However, the mass/volume relationship (which has been associated with clinical worsening in PAH [8]) and end-diastolic elastance showed no significant difference. Although the SV/end-systolic volume ratio was higher in female patients (Figure 1E), we observed no female-to-male differences in other key hemodynamic, echocardiographic, and cMRI parameters (Table 1). Exclusion of patients with associated PAH revealed similar findings (Figure 1). We observed no differences in female patients after stratification for age.

Previously, only indirect evidence for a sex-based difference in RV-PA coupling was available; in a large retrospective study of veterans with pre- and/or postcapillary pulmonary hypertension, women exhibited lower right atrial pressure despite having higher pulmonary vascular resistance than men, suggesting better RV adaptation to the increased afterload (9). However, most of those patients did not have PAH, and factors such as volume status may have affected right atrial pressure.

The present results show that female patients with PAH have higher contractility and better RV-PA coupling than male patients. In fact, the median Ees/Ea ratio was above the RV-PA uncoupling threshold of 0.8 (3) in female patients but below that threshold in male patients. This might represent a significant contributor to the survival advantage noted in female patients with PAH (2). We observed a consistent pattern using the volume-only method to assess RV-PA coupling (Ees and Ea have a common pressure term, so Ees/Ea can be simplified as a ratio of volumes [SV/end-systolic volume]). However, the tricuspid annular plane systolic excursion/systolic pulmonary artery pressure ratio (previously validated as a noninvasive surrogate for Ees/Ea [5]) was not significantly different in female and male patients. This may be a type II error because of the small sample size and insufficient precision of the ratio. Jacobs and colleagues demonstrated that 40% of the survival advantage in female versus male patients can be explained by a higher RV ejection fraction in female patients after initiation of treatment (2). RV ejection fraction was not significantly higher in women versus men in the present study, but this was also probably a type II error related to the small sample size. However, we assessed several potential confounders, including cMRI data, which did not reveal many sex differences (Table 1), whereas RV-PA coupling did. Especially in light of the relatively small study population, parameters of RV-PA coupling are obviously more sensitive to detect these changes.

Mechanisms underlying the superior RV adaptation in female patients with PAH are not yet known. We hypothesize that the lesser extent of RV hypertrophy in female versus male patients with PAH may make RV cardiomyocytes less prone to substrate depletion and ischemia. We have previously demonstrated in a rodent model that the female sex

steroid 17 β -estradiol attenuates RV proapoptotic and proinflammatory signaling, mitochondrial abnormalities, and oxidative stress and increases RV abundance of the proangiogenic and procontractile peptide apelin (10). Other sex hormones as well as nonhormonal factors may also contribute. However, estradiol and other sex hormone levels and menopausal status were not defined in the current analysis. Further investigations into mechanisms underlying the superior RV adaptation in female patients are ongoing in our laboratory.

In summary, our data demonstrate superior RV-PA coupling, better diastolic relaxation, and lower RV mass index in female versus male patients with PAH. These differences may underlie the superior survival of female patients. Our observations provide a rationale and basis for further evaluations of sex-based RV structural differences in PAH. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Acknowledgment: Editorial assistance was provided by Claire Mulligan, Ph.D. (Beacon Medical Communications Ltd.), funded by the University of Giessen.

Khodr Tello, M.D.*[†]
Manuel J. Richter, M.D.*
Athithan Yogeswaran
Universities of Giessen and Marburg Lung Center
Giessen, Germany

Hossein A. Ghofrani, M.D.
Universities of Giessen and Marburg Lung Center
Giessen, Germany
Kerckhoff Heart, Rheuma and Thoracic Center
Bad Nauheim, Germany
and
Imperial College London
London, United Kingdom

Robert Naeije, M.D.
Erasme University Hospital
Brussels, Belgium

Rebecca Vanderpool, Ph.D.
University of Arizona
Tucson, Arizona

Henning Gall, M.D., Ph.D.
Universities of Giessen and Marburg Lung Center
Giessen, Germany

Ryan J. Tedford, M.D.
Medical University of South Carolina
Charleston, South Carolina

Werner Seeger, M.D.
Universities of Giessen and Marburg Lung Center
Giessen, Germany

Tim Lahm, M.D.
Indiana University School of Medicine
Indianapolis, Indiana
and
Richard L. Roudebush VA Medical Center
Indianapolis, Indiana

*These authors contributed equally to this work.

[†]Corresponding author (e-mail: khodr.tello@innere.med.uni-giessen.de).

References

1. Vonk Noordegraaf A, Chin KM, Haddad F, Hassoun PM, Hemnes AR, Hopkins SR, *et al.* Pathophysiology of the right ventricle and of the pulmonary circulation in pulmonary hypertension: an update. *Eur Respir J* 2019;53:1801900.
2. Jacobs W, van de Veerdonk MC, Trip P, de Man F, Heymans MW, Marcus JT, *et al.* The right ventricle explains sex differences in survival in idiopathic pulmonary arterial hypertension. *Chest* 2014;145:1230–1236.
3. Tello K, Dalmer A, Axmann J, Vanderpool R, Ghofrani HA, Naeije R, *et al.* Reserve of right ventricular-arterial coupling in the setting of chronic overload. *Circ Heart Fail* 2019;12:e005512.
4. Simonneau G, Montani D, Celermajer DS, Denton CP, Gatzoulis MA, Krowka M, *et al.* Haemodynamic definitions and updated clinical classification of pulmonary hypertension. *Eur Respir J* 2019;53:1801913.
5. Richter MJ, Peters D, Ghofrani HA, Naeije R, Roller F, Sommer N, *et al.* Evaluation and prognostic relevance of right ventricular-arterial coupling in pulmonary hypertension. *Am J Respir Crit Care Med* 2020;201:116–119.
6. Tello K, Dalmer A, Vanderpool R, Ghofrani HA, Naeije R, Roller F, *et al.* Impaired right ventricular lusitropy is associated with ventilatory inefficiency in pulmonary arterial hypertension. *Eur Respir J* 2019;54:1900342.
7. Sys SU, Brutsaert DL. Diagnostic significance of impaired LV systolic relaxation in heart failure. *Circulation* 1995;92:3377–3380.
8. Badagliacca R, Poscia R, Pezzuto B, Nocioni M, Mezzapesa M, Francone M, *et al.* Right ventricular remodeling in idiopathic pulmonary arterial hypertension: adaptive versus maladaptive morphology. *J Heart Lung Transplant* 2015;34:395–403.
9. Ventetuolo CE, Hess E, Austin ED, Barón AE, Klinger JR, Lahm T, *et al.* Sex-based differences in veterans with pulmonary hypertension: results from the veterans affairs-clinical assessment reporting and tracking database. *PLoS One* 2017;12:e0187734.
10. Frump AL, Goss KN, Vayl A, Albrecht M, Fisher A, Tursunova R, *et al.* Estradiol improves right ventricular function in rats with severe angioproliferative pulmonary hypertension: effects of endogenous and exogenous sex hormones. *Am J Physiol Lung Cell Mol Physiol* 2015;308:L873–L890.

Copyright © 2020 by the American Thoracic Society

Preprocurement *In Situ* Donor Lung Tissue Gene Expression Classifies Primary Graft Dysfunction Risk

To the Editor:

Primary graft dysfunction (PGD) is a form of acute lung injury following lung transplantation that is responsible for 50% of all 30-day mortality experienced and has an attributable risk of death of 23% at 1 year (1). PGD is the primary driver of morbidity after transplant and predisposes patients to death from other causes (2). Given that donors are often assessed over a period of days prior to transplantation, prediction of recipient PGD risk based on donor characteristics is an attractive concept. Thus far, reliable clinical PGD risk classification has

been elusive. We previously showed that upregulation of innate immune pathways in recipient blood and lung lavage fluid occurs soon after transplantation (3, 4). We therefore hypothesized that innate immune transcripts from donor lung tissue obtained prior to transplantation could be used to predict recipient PGD risk.

We conducted a prospective cohort study of matched donor/recipient pairs transplanted at the University of Pennsylvania between October 2011 and December 2017. Institutional review board approval and patient written informed consent were collected prior to enrollment. Those not included in the analysis had insufficient sample, did not consent for participation, or failed to be collected by the procurement surgeon owing to circumstance (e.g., donation after circulatory death donor or time limitations). A 1 × 1 cm lung tissue biopsy was collected *in situ* in the donor prior to cross clamp and stored immediately in RNAlater. Gene expression profiles were obtained using the Affymetrix Human ST 2.1 (ThermoFisher Scientific) oligonucleotide microarray. Raw expression data were corrected for background, quantile normalized using the robust multiarray average method, and summarized at 23,819 gene levels (5).

We focused on TLR (Toll-like receptor) and NLR (Nod-like receptor) signaling pathways from the Kyoto Encyclopedia of Genes and Genomes Pathway Database because our prior work demonstrated association with PGD in bronchial wash and blood (3, 4). We applied a feed-forward deep learning method for PGD classification (6). Recipient body mass index, diagnosis, pulmonary artery pressure, and donor smoking status were considered to be included in the model as previously identified (7). A set of covariates, including body mass index, cytomegalovirus status, cardiopulmonary bypass use, transplant type, and total ischemic time, were subsequently selected to optimize Akaike's information criterion in a logistic regression model. We used cross entropy as the objective loss function, the hyperbolic tangent function as the activation function, two hidden layers with sizes 200 and 100, a learning rate of 0.001, and an epoch of 300. We used five-fold cross-validation to avoid overfitting. We report the receiver operating characteristic curve, the area under the receiver operating characteristic curve, and prediction accuracy at a cutoff probability optimized for sensitivity and specificity. The results were also compared with a deep learning model that used the five covariates as the only predictors.

We enrolled 113 subjects, of whom 28 (24.8% [95% confidence interval, 17.1–33.8]) developed grade 3 PGD defined at 48–72 hours. Differences in donor and recipient characteristics or operative factors between subjects with and without PGD appear in Table 1. Table 2 summarizes the predictive classifier performance between clinical variables alone or in combination with our identified pathways. The TLR signaling pathway demonstrated superior discriminant ability to covariates alone ($P=0.014$) in contrast to the NLR signaling pathway ($P=0.296$). The TLR-based classifier improved PGD prediction by 0.20 over the model using clinical factors alone. With respect to diagnostic test performance, TLR signaling demonstrated superior performance to covariates alone and not NLR (Table 2).

The clinical implications of this TLR classifier depend on PGD prevalence and context of use. For enrichment purposes using the values from Table 2, the predicted PGD probability could be enriched two-fold over baseline (20% PGD pretest probability [odds of 0.25]; 2.673 likelihood ratio [+]; the *post hoc* test odds would be 0.67 [0.25 × 2.673], corresponding to a *post hoc* test probability of 40% [0.67/(1 + 0.67)]). Conversely, for efforts focused on PGD risk reduction, this classifier could reduce the predicted probability of PGD by more than 60% (*post*

Supported by NIH grants HL087115, HL081619, HL096845, HL115354, HL116656, HL135227, and HL121406 and by Robert Wood Johnson grant AMFDP70640.

Author Contributions: Conception and design: E.C., J.D.C., and R.F. Data acquisition, analysis, and interpretation: E.C., M.Y., Y.S., T.B., V.G., N.M., C.A.B., J.M.D., J.D.C., and R.F. Drafting the manuscript for important intellectual content, final approval, and accountability for the work presented: E.C., M.Y., Y.S., T.B., V.G., N.M., C.A.B., J.M.D., J.D.C., and R.F.

Originally Published in Press as DOI: 10.1164/rccm.201912-2436LE on May 28, 2020