Published in final edited form as:

J Am Chem Soc. 2020 September 23; 142(38): 16205–16210. doi:10.1021/jacs.0c08035.

P(III)/P(V)-Catalyzed Methylation of Arylboronic Acids and Esters: Reductive C–N Coupling with Nitromethane as a Methylamine Surrogate

Gen Li[†], Ziyang Qin^{†,‡}, Alexander T. Radosevich^{†,*}

[†]Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States.

Abstract

The direct reductive *N*-arylation of nitromethane by organophosphorus-catalyzed reductive C–N coupling with arylboronic acid derivatives is reported. This method operates by the action of a small ring organophosphorus-based catalyst (1,2,2,3,4,4-hexamethylphosphetane *P*-oxide) together with a mild terminal reductant hydrosilane to drive the selective installation of the methylamino group to (hetero)aromatic boronic acids and esters. This method also provides for a unified synthetic approach to isotopically-labeled *N*-methylanilines from various stable isotopologues of nitromethane (i.e. CD₃NO₂, CH₃¹⁵NO₂ and ¹³CH₃NO₂), revealing this easy-to-handle compound as a versatile precursor for the direct installation of the methylamino group.

Graphical Abstract

Nitromethane ($\text{H}_3\text{C--NO}_2$) is an industrially important commodity chemical, primarily used as a solvent, stabilizer, and fuel additive.¹ Also, $\text{H}_3\text{C--NO}_2$ is a common and useful carbon pronucleophile in synthesis (Figure 1A, left),^{2–5} but it is comparatively less developed as an amination reagent. Recently, Jiao and coworkers reported $\text{H}_3\text{C--NO}_2$ as a precursor to " $\text{H}_2\text{N--X}$ " equivalents by reductive Nef-like decomposition,⁶ but an alternative reductive *N*-functionalization of $\text{H}_3\text{C--NO}_2$ that retains the " $\text{H}_3\text{C--N}$ " substructure would trace a role for

*Corresponding Author radosevich@mit.edu.

[‡]Current address: Department of Chemistry, University of Science and Technology of China, 96 Jinzhai Road, Hefei, Anhui 230026, China

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

General methods and synthetic procedures (.pdf)

¹H, ²H, ¹³C, ¹⁵N, ¹⁹F and ³¹P NMR spectra (.pdf).

The authors declare no competing financial interests.

this inexpensive and easily-handled liquid as a potential surrogate for methylamine—a gaseous List 1 controlled precursor⁷—especially in catalytic coupling chemistry (Figure 1A, right).^{8–11} At present, reductive *N*-functionalization of H₃C–NO₂ is limited to two isolated examples: Niggemann has reported reductive *N*-benzylation of nitromethane mediated by stoichiometric B₂pin₂ in the presence of excess BnZnBr,¹² and Suarez-Pantiga and Sanz have reported a triphenylphosphine-mediated reductive *N*-phenylation of nitromethane catalyzed by an oxomolybdenum(VI) compound under microwave irradiation.¹³ We report here a general catalytic method for the methylation of arylboronic acids and esters with H₃C–NO₂ by reductive C–N coupling driven by P^{III}/P^V=O redox cycling (Figure 1B). These results expand the scope of reductive C–N coupling, generalize the use of H₃C–NO₂ as a methylamine surrogate in reductive cross coupling, and provide a unique pathway for introducing stable isotopes (¹⁵N, ¹³C, ²H) widely-used for metabolic tracing.¹⁴

Prior work has established P^{III}/P^V=O catalysis^{15–17} as a viable approach to reductive *N*-functionalization of nitroarene (Ar–NO₂) substrates,^{18–20} but the translation of this technique to reductive *N*-functionalization of H₃C–NO₂ requires that the desired C–N coupling sequence (Figure 1B) outcompete numerous unproductive but well-known decomposition pathways, both for H₃C–NO₂ (e.g. Nef reaction) and potential deoxygenation intermediates (e.g. tautomerization of H₃C–NO→H₂C=NOH and H₃C–N:→H₂C=NH, *inter alia*). In principle, the base-free conditions of the P^{III}/P^V=O catalyzed nitro functionalization, which depends on a rate-limiting (3+1) cheletropic addition of phosphine and nitro substrate, might suppress these processes. However, the nitro moiety of H₃C–NO₂ is inherently less reactive than Ar–NO₂ as cycloaddition partner due to a larger local frontier orbital energy gap ($E = 1.0$ eV, Figure 1C),²¹ resulting in less favorable pairwise orbital interactions that therefore impose a stringent demand of the biphilic character of the phosphetane catalyst.

Investigation of the reductive *N*-functionalization of H₃C–NO₂ was initiated with these considerations in mind. Empirical observations (Table 1) indicate that a catalytic system comprising the organophosphorus O-atom transfer catalyst **1**•[O]²² and a terminal hydrosilane reductant indeed successfully achieve a methylation of arylboronic acids by P^{III}/P^V=O-catalyzed intermolecular C–N coupling. With 4-fluorophenylboronic acid (**3**) as a representative substrate, the reductive functionalization of 1.0 equiv of commercial reagent grade H₃C–NO₂ (**2**) by 10 mol% of **1**•[O] and 2 equiv of PhSiH₃ gave a promising 35% yield of *N*-methyl-4-fluoroaniline (**4**) (Table 1, entry 1), but simply increasing equivalencies of inexpensive H₃C–NO₂ (3.0 equiv, entry 2) resulted in excellent yield of **4** (95% NMR yield, 90% isolated yield). A high yield was similarly obtained with phosphetane **1** as a precatalyst (entry 3)—consistent with P^{III}/P^V=O redox cycling—but omission of either the precatalyst **1**•[O] or phenylsilane failed to provide methylation product **4** (entries 4–5). Inexpensive and bench-stable polymethylhydrosiloxane (PMHS) is also a viable terminal reductant for the reaction, albeit with somewhat lower efficiency (85%, entry 6). *In situ* spectral monitoring of the reductive C–N coupling showed that isotopically enriched nitromethane-*d*₃ (δ 4.32 ppm) is cleanly converted to the product *N*-(methyl-*d*₃)-4-bromoaniline (δ 2.79 ppm), and no side products or long-lived intermediates were observed (Figure S2). The relatively high concentration of the P^{III} catalyst resting state **1** presumably

drives the methylation forward along the productive reaction pathway in preference to possible side pathways, including the favorable isomerization of nitrosomethane to formaldoxime.^{23,24}

As depicted in Figure 2A, the $P^{III}/P^V=O$ -catalyzed methylation method shows excellent functional group tolerance, several facets of which provide for chemoselectivities that complement transition-metal catalyzed C–N coupling methods. For example, halogen substituents are well-tolerated (**5a**, **8–10**, **16**) in the $P^{III}/P^V=O$ -catalyzed methylation and intermolecular C–N coupling occurs exclusively at the boronic acid position. Also, the orthogonal reactivity to the nitro group with respect to other nitrogen-containing functional groups permits selective aryl methylation even in the presence of free $-NH_2$ groups (**16**) without explicit protection. Carbonyl functionalities such as esters (**13**) and ketones (**17**, **19**) that might be susceptible to acylation or condensation reaction with H_3C-NH_2 are retained when H_3C-NO_2 is used as a surrogate. Furthermore, substrates with other functional groups including benzyloxy (**7**), tetrahydropyranyl ether (**14**), thiomethyl (**15**), and trifluoromethyl (**18**) groups all yielded the corresponding *N*-methylaniline products in good yields. The method is also applicable to compounds with known bioactive core structures, such as derivatives of celecoxib (**18**), estrone (**19**), and Pittsburgh compound B²⁵ (**20**).

The ability to use of H_3C-NO_2 as a simple synthetic precursor for the MeHN– group presents the opportunity for the preparation of diverse stable-isotope labeled products—useful tools in both laboratory²⁶ and pharmaceutical²⁷ research—through a unified synthetic strategy. As illustrated in Figure 2B, a suite of isotopically labeled *N*-methylaniline products **5b–5d** are accessible with programmed labeling as dictated by the initial isotopic composition of the nitromethane isotopologue (viz. $^2H_3C-NO_2$ (**2b**), $H_3^{13}C-NO_2$ (**2c**), $H_3C-^{15}NO_2$ (**2d**)). The relative ease with which the isotopologues of nitromethane are accessed make this an attractive approach to potentially valuable stable isotope labeled compounds.

Initial attempts to directly transfer the foregoing $P^{III}/P^V=O$ -catalyzed conditions to methylation of 3-quinolylboronic acid resulted in a disappointing 19% yield of the target compound **21** (Figure 3A, yield in parenthesis). Studies showed that protodeboronation of the heteroarylboronic acid was a primary pathway limiting the productive C–N coupling in this case.²⁸ Correspondingly, $P^{III}/P^V=O$ -catalyzed methylation of 3-quinolylboronic acid 1,3-propanediolate ester showed a significant improvement in C–N coupling yield (93%), in line with the relative stability of boronic esters to protodeboronation as compared to their parent boronic acids.²⁷ This phenomenon is general; *B*-heteroaryl-1,3,2-dioxaborinanes including a variety of five- and six-membered heterocyclic derivatives, such as isoquinoline (**22**), indazole (**23**) pyridine (**24**, **27**), thiophene (**25**) and pyrimidine (**26**), were successfully transformed to the corresponding *N*-methylamine derivatives with improved yields relative to their arylboronic acid congeners. *B*-Aryl-1,3,2-dioxaborinanes containing electron-donating (**28**) or electron-withdrawing group (**29**) transform smoothly to the desired products, showing the generality of utilizing boronate esters for methylation reactions. Other common boronate ester residues such as a neopentyl glycol ester (e.g. Ar-Bnep **30**, Figure 3B) and pinacol esters (e.g. Ar-Bpin

31–33, Figure 3C) gave similar efficiency in this methylation process. Notably, subjection of benzene-1,4-diboronic acid bispinacol ester to the standard reaction conditions afforded single methylation product **31** with excellent yield; evidently the second (pinacolato)boryl moiety becomes electronically deactivated for amination following an initial reductive C–N coupling.

Competition experiments evidence a differential reactivity of $\text{H}_3\text{C–NO}_2$ as compared to Ar–NO_2 substrates in $\text{P}^{\text{III}}/\text{P}^{\text{V}}=\text{O}$ -catalyzed reductive C–N coupling. As illustrated in Figure 4A, an equimolar mixture of four components—namely $\text{H}_3\text{C–NO}_2$ (**2**), 4-NC- $\text{C}_6\text{H}_4\text{–NO}_2$ (**34**), 3,5-(F_3C) $_2\text{–C}_6\text{H}_3\text{–Bpin}$ (**35**) and 4- $\text{H}_3\text{CO–C}_6\text{H}_4\text{–B(OH)}_2$ (**36**)—was subjected to standard reductive C–N coupling conditions catalyzed by **1**•[O] in a one pot manner. In the event, only two of the possible four products—specifically methylaniline **29** and diarylamine **37**—were observed in greater than 5% yield. Consistent with the FMO rationale in Figure 1,¹⁵ⁱ this observation implies that nitroarene **34** reacts more quickly than nitromethane **2** via (3+1) cheletropic addition with **1**, and goes on to couple selectively with the more reactive boronic acid partner **36**. In a subsequent event, $\text{H}_3\text{C–NO}_2$ then reacts to give **29** by methylation of the more inert boronic ester **35**, proceeding without diminished yield for these reaction partners.²⁹ This kinetically-controlled chemoselectivity was further demonstrated by the modular synthesis of *N,N'*-difunctionalized phenylenediamine (Figure 4B, **42**). The three-component coupling of $\text{H}_3\text{C–NO}_2$, Ph–B(OH)_2 (**40**), and 4-nitrophenylboronic ester (**41**) produced **42** via sequential amination reactions with almost complete suppression of undesired methylaminated product (**6**).

A synthesis of the anxiolytic triflubazam³⁰ (**45**, Figure 5) contextualizes the reductive C–N coupling of nitromethane within the growing portfolio of $\text{P}^{\text{III}}/\text{P}^{\text{V}}$ -catalyzed methods promoted by phosphetane catalyst **1**•[O]. Specifically, $\text{P}^{\text{III}}/\text{P}^{\text{V}}$ -catalyzed reductive C–N coupling of phenylboronic acid and 4,4,5,5-tetramethyl-2-(2-nitro-5-(trifluoromethyl)phenyl)-1,3,2-dioxaborolane (**43**), followed in situ by $\text{P}^{\text{III}}/\text{P}^{\text{V}}$ -catalyzed amidation^{15h,31} with monoethyl malonate gives diarylamide **44** in a one-pot sequence. Subsequent $\text{P}^{\text{III}}/\text{P}^{\text{V}}$ -catalyzed methylation by C–N coupling with $\text{H}_3\text{C–NO}_2$ installs the MeHN– moiety and induces intramolecular cyclization to close the diazepinedione ring, furnishing the medicinal target **45** in an overall 51% yield for the two-pot, three-step, all- $\text{P}^{\text{III}}/\text{P}^{\text{V}}$ -catalyzed sequence.

In summary, we have demonstrated that nitromethane is an inexpensive and easy-to-handle synthetic equivalent for installation of the MeHN– fragment via $\text{P}^{\text{III}}/\text{P}^{\text{V}}=\text{O}$ catalysis. Readily-available boronic acids and esters are selectively methylaminated in the presence of various functional groups and heteroaromatics. The method serves as a robust complementary tactic to transition metal catalyzed C–N coupling techniques relying on the use of MeNH_2 or related surrogates. With respect to our efforts to develop $\text{P}^{\text{III}}/\text{P}^{\text{V}}=\text{O}$ -catalyzed reductive C–N coupling reactions as modular approach to complex amine synthesis, this study represents our first success with nitroalkane substrates and portends future developments in the deoxygenative *N*-functionalization of higher nitroalkane homologues to access more elaborate *N*-alkyl products

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

Financial support was provided by NIH NIGMS (GM114547). The authors thank Prof. A. Wendlandt (MIT) and Drs. J.M. Lipshultz, T.V. Nykaza and J.C. Cooper for helpful discussion.

REFERENCE

1. Markofsky SB Nitro Compounds, Aliphatic In Ullmann's Encyclopedia of Industrial Chemistry, 7th edition; Ley C, Ed.; Wiley-VCH: Weinheim, 2012; Vol. 24, pp 291–300.
2. (a)Ono N The Nitro-aldol (Henry) Reaction In The Nitro Group in Organic Synthesis; Feuer H, Ed.; Wiley-VCH: New York, 2001, pp 30–69;(b)Luzzio FA The Henry Reaction: Recent Examples. Tetrahedron 2001, 57, 915–945.
3. Noble A; Anderson JC Nitro-Mannich Reaction. Chem. Rev 2013, 113, 2887–2939. [PubMed: 23461586]
4. (a)Ballini R; Bosica G; Fiorini D; Palmieri A; Petrini M Conjugate Additions of Nitroalkanes to Electron-Poor Alkenes: Recent Results. Chem. Rev 2005, 105, 933–971. [PubMed: 15755081] (b)Manzano R; Andrés JM; Álvarez R; Muruzábal MD; De Lera ÁR; Pedrosa R Enantioselective Conjugate Addition of Nitro Compounds to α,β -Unsaturated Ketones: An Experimental and Computational Study. Chem. Eur. J 2011, 17, 5931–5938. [PubMed: 21500295] (c)Ballini R; Palmieri A Formation of Carbon-Carbon Double Bonds: Recent Developments via Nitrous Acid Elimination (NAE) from Aliphatic Nitro Compounds. Adv. Synth. Catal 2019, 361, 5070–5097.
5. (a)Li CJ; Li Z Green Chemistry: The Development of Cross-Dehydrogenative Coupling (CDC) for Chemical Synthesis. Pure Appl. Chem 2006, 78, 935–945.(b)Li CJ Cross-Dehydrogenative Coupling (CDC): Exploring C–C Bond Formations beyond Functional Group Transformations. Acc. Chem. Res 2009, 42, 335–344. [PubMed: 19220064] (c)Girard SA; Knauber T; Li CJ The Cross-Dehydrogenative Coupling of Csp³-H Bonds: A Versatile Strategy for C–C Bond Formations. Angew. Chem. Int. Ed 2014, 53, 74–100.(d)Varun BV; Dhineshkumar J; Bettadapur KR; Siddaraju Y; Alagiri K; Prabhu KR Recent Advancements in Dehydrogenative Cross Coupling Reactions for C–C Bond Formation. Tetrahedron Lett 2017, 58, 803–824.
6. Liu J; Zhang C; Zhang Z; Wen X; Dou X; Wei J; Qiu X; Song S; Jiao N Nitromethane as a Nitrogen Donor in Schmidt-Type Formation of Amides and Nitriles. Science 2020, 367, 281–285.
7. Records and Reports of Listed Chemicals and Certain Machines; Importation and Exportation of Certain Machines. Code of Federal Regulations Part 1310, Title 21, (2019).
8. Hughes EC; Veatch F; Eilersich V N-Methylaniline From Chlorobenzene and Methylamine. Ind. Eng. Chem 1950, 42, 787–790.
9. Coupling with methylamine solutions:(a)Fors BP; Watson DA; Biscoe MR; Buchwald SL A Highly Active Catalyst for Pd-Catalyzed Amination Reactions: Cross-Coupling Reactions Using Aryl Mesylates and the Highly Selective Monoarylation of Primary Amines Using Aryl Chlorides. J. Am. Chem. Soc 2008, 130, 13552–13554. [PubMed: 18798626] (b)Henderson JL; Buchwald SL Efficient Pd-Catalyzed Amination Reactions for Heterocycle Functionalization. Org. Lett 2010, 12, 4442–4445. [PubMed: 20860403] (c)Fors BP; Buchwald SL A Multiligand Based Pd Catalyst for C–N Cross-Coupling Reactions. J. Am. Chem. Soc 2010, 132, 15914–15917. [PubMed: 20979367] (d)Crawford SM; Lavery CB; Stradiotto M BippyPhos: A Single Ligand with Unprecedented Scope in the Buchwald-Hartwig Amination of (Hetero)Aryl Chlorides. Chem. Eur. J 2013, 19, 16760–16771. [PubMed: 24281816] Ullman C–N Cross Coupling:(e)Jiao J; Zhang XR; Chang NH; Wang J; Wei JF; Shi XY; Chen ZG A Facile and Practical Copper Powder-Catalyzed, Organic Solvent-and Ligand-Free Ullmann Amination of Aryl Halides. J. Org. Chem 2011, 76, 1180–1183. [PubMed: 21261263] (f)Siddegowda MS; Yathirajan HS; Ramakrishna RA A Ligand-Free and Base-Free Copper Catalyzed Reaction: Arylation of Ammonia and Primary Amines as Their Acetate Salts. Tetrahedron Lett 2012, 53, 5219–5222.(g)Kumar AS; Ramani T; Sreedhar B Magnetically Separable CuFe₂O₄ Nanoparticles in PEG: A Recyclable Catalytic System for the Amination of

Aryl Iodides. *Synlett* 2013, 24, 938–942.(h)Wang D; Kuang D; Zhang F; Yang C; Zhu X Room-Temperature Copper-Catalyzed Arylation of Dimethylamine and Methylamine in Neat Water. *Adv. Synth. Catal* 2015, 357, 714–718.

10. Coupling with methylammonium salts:(a)Green RA; Hartwig JF Palladium-Catalyzed Amination of Aryl Chlorides and Bromides with Ammonium Salts. *Org. Lett* 2014, 16, 4388–4391. [PubMed: 25133675] (b)Green RA; Hartwig JF Nickel-Catalyzed Amination of Aryl Chlorides with Ammonia or Ammonium Salts. *Angew. Chem. Int. Ed* 2015, 54, 3768–3772.(c)Zeng L; Fu H; Qiao R; Jiang Y; Zhao Y Efficient Copper-Catalyzed Synthesis of *N*-Alkylanthranilic Acids via an Ortho-Substituent Effect of the Carboxyl Group of 2-Halobenzoic Acids at Room Temperature. *Adv. Synth. Catal* 2009, 351, 1671–1676.
11. Coupling with protected methylamine surrogates:(a)Trabanco AA; Vega JA; Fernandez MA Fluorous-Tagged Carbamates for the Pd-Catalyzed Amination of Aryl Halides. *J. Org. Chem* 2007, 72, 8146–8148. [PubMed: 17877407] (b)Bernardi P; Dembech P; Fabbri G; Ricci A; Seconi G A General and Convenient Procedure for the Synthesis of *N*- Alkylarylamines and *N*- Alkylheteroarylamines by Electrophilic Amination of Cuprates with *N*-Alkylhydroxylamines. *J. Org. Chem* 1999, 64, 641–643.
12. Rauser M; Ascheberg C; Niggemann M Direct Reductive N-Functionalization of Aliphatic Nitro Compounds. *Chem. Eur. J* 2018, 24, 3970–3974. [PubMed: 29378085]
13. Suárez-Pantiga S; Hernández-Ruiz R; Virumbrales C; Pedrosa MR; Sanz R Reductive Molybdenum-Catalyzed Direct Amination of Boronic Acids with Nitro Compounds. *Angew. Chem. Int. Ed* 2019, 58, 2129–2133.
14. (a)Harbeson SL; Tung RD Deuterium in Drug Discovery and Development. *Annu. Rep. Med. Chem* 2011, 46, 403–417.For examples highlighting the importance of isotope labeling:(b)Atzrodt J; Derdau V Pd- and Pt- catalyzed H/D exchange methods and their application for internal MS standard preparation from a Sanofi-Aventis perspective *J. Labelled Compd. Radiopharm* 2010, 53, 674–685.(c)Gant TGJ Using Deuterium in Drug Discovery: Leaving the Label in the Drug. *J. Med. Chem* 2014, 57, 3595–3611. [PubMed: 24294889] (d)Atzrodt J; Derdau V; Kerr WJ; Reid M Deuterium- and Tritium-Labelled Compounds: Applications in the Life Sciences. *Angew. Chem. Int. Ed* 2018, 57, 1758–1784.(e)Zachleder V; Vítová M; Hlavová M; íková Š; Mojzeš P; Heumann H; Becher JR; Bišová K Stable isotope compounds - production, detection, and application. *Biotechnol. Adv* 2018, 36, 784–797. [PubMed: 29355599]
15. (a)Reichl KD; Dunn NL; Fastuca NJ; Radosevich AT Biphilic Organophosphorus Catalysis: Regioselective Reductive Transposition of Allylic Bromides via P^{III}/P^V Redox Cycling. *J. Am. Chem. Soc* 2015, 137, 5292–5295. [PubMed: 25874950] (b)Zhao W; Yan PK; Radosevich AT A Phosphetane Catalyzes Deoxygenative Condensation of α -Keto Esters and Carboxylic Acids via $P^{III}/P^V=O$ Redox Cycling. *J. Am. Chem. Soc* 2015, 137, 616–619. [PubMed: 25564133] (c)Lin Y-C; Hatzakis E; McCarthy SM; Reichl KD; Lai T-Y; Yennawar HP; Radosevich AT P–N Cooperative Borane Activation and Catalytic Hydroboration by a Distorted Phosphorous Triamide Platform. *J. Am. Chem. Soc* 2017, 139, 6008–6016. [PubMed: 28398750] (d)Nykaza TV; Harrison TS; Ghosh A; Putnik RA; Radosevich AT A Biphilic Phosphetane Catalyzes N–N Bond-Forming Cadogan Heterocyclization via $P^{III}/P^V=O$ Redox Cycling. *J. Am. Chem. Soc* 2017, 139, 6839–6842. [PubMed: 28489354] (e)Nykaza TV; Ramirez A; Harrison TS; Luzung MR; Radosevich AT Biphilic Organophosphorus-Catalyzed Intramolecular Csp^2 –H Amination: Evidence for a Nitrenoid in Catalytic Cadogan Cyclizations. *J. Am. Chem. Soc* 2018, 140, 3103–3113. [PubMed: 29389114] (f)Nykaza TV; Cooper JC; Li G; Mahieu N; Ramirez A; Luzung MR; Radosevich AT Intermolecular Reductive C–N Cross Coupling of Nitroarenes and Boronic Acids by $P^{III}/P^V=O$ Catalysis. *J. Am. Chem. Soc* 2018, 140, 15200–15205. [PubMed: 30372615] (g)Ghosh A; Lecomte M; Kim-Lee S-H; Radosevich AT Organophosphorus-Catalyzed Deoxygenation of Sulfonyl Chlorides: Electrophilic (Fluoroalkyl)Sulfonylation by $P^{III}/P^V=O$ Redox Cycling. *Angew. Chem. Int. Ed* 2019, 58, 2864–2869.(h)Lecomte M; Lipshultz JM; Kim-Lee S-H; Li G; Radosevich AT Driving Recursive Dehydration by P^{III}/P^V Catalysis: Annulation of Amines and Carboxylic Acids by Sequential C–N and C–C Bond Formation. *J. Am. Chem. Soc* 2019 141, 12507–12512. [PubMed: 31345031] (i)Li G; Nykaza TV; Cooper JC; Ramirez A; Luzung MR; Radosevich AT An Improved $P^{III}/P^V=O$ -Catalyzed Reductive C–N Coupling of Nitroaromatics and Boronic Acids by Mechanistic Differentiation of Rate- and Product-Determining Steps. *J. Am. Chem. Soc* 2020, 142, 6786. [PubMed: 32178514] (j)Nykaza TV; Li G; Yang J; Luzung MR;

- Radosevich AT $P^{III}/P^V=O$ -Catalyzed Cascade Synthesis of *N*-Functionalized Azaheterocycles. *Angew. Chem. Int. Ed* 2020, 59, 4505–4510.
16. For a review of $P^{III}/P^V=O$ redox cycling, see: (a) Marsden SP Catalytic Variants of Phosphine Oxide-Mediated Organic Transformations in Sustainable Catalysis; Dunn PJ, Hii KK, Krische MJ, Williams MT, Eds.; John Wiley & Sons, Inc.: New York, 2013; pp 339–361. (b) Guo H; Fan YC; Sun Z; Wu Y; Kwon O Phosphine Organocatalysis. *Chem. Rev* 2018, 118, 10049–10293. [PubMed: 30260217]
 17. For recent examples of $P^{III}/P^V=O$ redox cycling, see: (a) O'Brien CJ; Tellez JL; Nixon ZS; Kang LJ; Carter AL; Kunkel SR; Przeworski KC; Chass GA Recycling the Waste: The Development of a Catalytic Wittig Reaction *Angew. Chem. Int. Ed* 2009, 48, 6836–6839. (b) O'Brien CJ; Lavigne F; Coyle EE; Holohan AJ; Doonan BJ Breaking the Ring through a Room Temperature Catalytic Wittig Reaction *Chem. Eur. J* 2013, 19, 5854–5858. [PubMed: 23526683] (c) O'Brien CJ; Nixon ZS; Holohan AJ; Kunkel SR; Tellez JL; Doonan BJ; Coyle EE; Lavigne F; Kang LJ; Przeworski KC Part I: The Development of the Catalytic Wittig Reaction *Chem. Eur. J* 2013, 19, 15281–15289. [PubMed: 24115040] (d) Coyle EE; Doonan BJ; Holohan AJ; Walsh KA; Lavigne F; Krenske EH; O'Brien CJ Catalytic Wittig Reactions of Semi- and Nonstabilized Ylides Enabled by Ylide Tuning *Angew. Chem. Int. Ed* 2014, 53, 12907–12911. (e) van Kalker HA; Leenders SHAM; Hommersom CRA; Rutjes FPJT; van Delft FL In Situ Phosphine Oxide Reduction: A Catalytic Appel Reaction *Chem. Eur. J* 2011, 17, 11290–11295. [PubMed: 21882274] (f) van Kalker HA; Bruins JJ; Rutjes FPJT; van Delft FL Organophosphorus-Catalysed Staudinger Reduction *Adv. Synth. Catal* 2012, 354, 1417–1421. (g) Lee C; Chang T; Yu J; Reddy GM; Hsiao M; Lin W Synthesis of functionalized furans via chemoselective reduction/Wittig reaction using catalytic triethylamine and phosphine. *Org. Lett* 2016, 18, 3758–3761. [PubMed: 27434727] (h) Saleh N; Voituriez A Synthesis of 9H-pyrrolo[1,2-*a*]indole and 3H-pyrrolizine derivatives via a phosphine-catalyzed umpolung addition/intramolecular Wittig reaction. *J. Org. Chem* 2016, 81, 4371–4377. [PubMed: 27080174] (i) Saleh N; Blanchard F; Voituriez A Synthesis of nitrogen containing heterocycles and cyclopentenone derivatives via phosphine catalyzed Michael addition/intramolecular Wittig reaction. *Adv. Synth. Catal* 2017, 359, 2304–2315. (j) Zhang K; Cai L; Yang Z; Houk KN; Kwon O Bridged [2.2.1] bicyclic phosphine oxide facilitates catalytic γ -umpolung addition–Wittig olefination. *Chem. Sci* 2018, 9, 1867–1872. [PubMed: 29732112] (k) Cai L; Zhang K; Chen S; Lepage RJ; Houk KN; Krenske EH; Kwon O Catalytic Asymmetric Staudinger–Aza-Wittig Reaction for the Synthesis of Heterocyclic Amines. *J. Am. Chem. Soc* 2019, 141, 9537–9542. [PubMed: 31184143] (l) Lorton C; Castanheiro T; Voituriez A Catalytic and Asymmetric Process via $P^{III}/P^V=O$ Redox Cycling: Access to (Trifluoromethyl)Cyclobutenes via a Michael Addition/Wittig Olefination Reaction. *J. Am. Chem. Soc* 2019, 141, 10142–10147. [PubMed: 31194912]
 18. (a) Sapountzis I; Knochel P A New General Preparation of Polyfunctional Diarylamines by the Addition of Functionalized Arylmagnesium Compounds to Nitroarenes. *J. Am. Chem. Soc* 2002, 124, 9390–9391. [PubMed: 12167031] (b) Doyle W; Staubitz A; Knochel P Mild Synthesis of Polyfunctional Benzimidazoles and Indoles by the Reduction of Functionalized Nitroarenes with Phenylmagnesium Chloride. *Chem. Eur. J* 2003, 9, 5323–5331. [PubMed: 14613142] (c) Kopp F; Sapountzis I; Knochel P Preparation of Polyfunctionalized Amines by the Addition of Functionalized Organomagnesium Reagents to Nitrosoarenes *Synlett*, 2003, 885–887. (d) Sapountzis I; Knochel P A New Method for the Selective Amination of 1,3- and 1,4-Dinitrobenzenes and Protected Nitroanilines Leading to Polyfunctional 1,3- and 1,4-Disubstituted Anilines. *Synlett* 2004, 955–958. (e) Dhayalan V; Saemann C; Knochel P Synthesis of polyfunctional secondary amines by the addition of functionalized zinc reagents to nitrosoarenes. *Chem. Commun* 2015, 51, 3239–3242. (f) Gao H; Xu Q-L; Ess DH; Kürti L Transition-Metal-Free, Low-Temperature Intramolecular Amination of Aromatic C-H Bonds: Rapid Synthesis of Fused Heterocycles. *Angew. Chem. Int. Ed* 2014, 53, 2701–2705. (g) Rauser M; Ascheberg C; Niggemann M Electrophilic Amination with Nitroarenes. *Angew. Chem. Int. Ed* 2017, 56, 11570–11574.
 19. (a) Gui J; Pan C-M; Jin Y; Qin T; Lo JC; Lee BJ; Spergel SH; Mertzman ME; Pitts WJ; La Cruz TE; Schmidt MA; Darvatkar N; Natarajan SR; Baran PS Practical olefin hydroamination with nitroarenes. *Science* 2015, 348, 886–891. [PubMed: 25999503] (b) Cheung CW; Hu X Amine synthesis via iron-catalysed reductive coupling of nitroarenes with alkyl halides. *Nat. Commun* 2016, 7, 12494. [PubMed: 27515391] (c) Cheung CW; Hu X Nickel-Catalyzed Reductive Transamidation of Secondary Amides with Nitroarenes. *ACS Catal* 2017, 7, 7092–7096.

20. Xiao J; He Y; Ye F; Zhu S Remote Sp³ C–H Amination of Alkenes with Nitroarenes. *Chem* 2018, 4, 1645–1657.
21. See SI for Kohn-Sham frontier orbital eigenvalues for H₃C–NO₂ and H₅C₆–NO₂ at the ωB97XD/6–311++G(2d,2p) level of theory.
22. For a preparation of **1**•[O], see: Nykaza TV; Cooper JC; Radosevich AT *anti*-1,2,2,3,4,4-Hexamethylphosphetane 1-Oxide. *Org. Synth* 2019, 96, 418–435. [PubMed: 31902967]
23. (a)Adeney PD; Bouma WJ; Radom L; Rodwell WR Nitrosomethane and Its Nitron and Oxime Isomers. A Theoretical Study of 1,2- and 1,3-Intramolecular Hydrogen Shifts. *J. Am. Chem. Soc* 1980, 102, 4069–4074.(b)Long JA; Harris NJ; Lammertsma K Formaldehyde Oxime ⇌ Nitrosomethane Tautomerism. *J. Org. Chem* 2001, 66, 6762–6767. [PubMed: 11578232]
24. Indeed, when formaldoxime trimer was used instead of nitromethane under otherwise optimal conditions, no methylation product **4** is formed. See SI.
25. (a)Klunk WE; Engler H; Nordberg A; Wang Y; Blomqvist G; Holt DP; Bergström M; Savitcheva I; Huang GF; Estrada S; Ausen B; Debnath ML; Barletta J; Price JC; Sandell J; Lopresti BJ; Wall A; Koivisto P; Antoni G; Mathis CA; Langstrom B Imaging Brain Amyloid in Alzheimer's Disease with Pittsburgh Compound-B. *Ann. Neurol* 2004, 55, 306–319. [PubMed: 14991808] (b)Rowe CC; Ellis KA; Rimajova M; Bourgeat P; Pike KE; Jones G; Frapp J; Tochon-Danguy H; Morandau L; O'Keefe G; Price R; Raniga P; Robins P; Acosta O; Lenzo N; Szeoke C; Salvado O; Head R; Martins R; Masters C; Ames D; Villemagne VL Amyloid Imaging Results from the Australian Imaging, Biomarkers and Lifestyle (AIBL) Study of Aging. *Neurobiol. Aging* 2010, 31, 1275–1283. [PubMed: 20472326]
26. Lloyd-Jones GC; Muñoz MP Isotopic Labelling in the Study of Organic and Organometallic Mechanism and Structure: An Account. *J. Label. Compd. Radiopharm* 2007, 50, 1072–1087.
27. (a)Mutlib AE Application of Stable Isotope-Labeled Compounds in Metabolism and in Metabolism-Mediated Toxicity Studies. *Chem. Res. Toxicol* 2008, 21, 1672–1689. [PubMed: 18702535] (b)Chokkathukalam A; Kim DH; Barrett MP; Breitling R; Creek DJ Stable Isotope-Labeling Studies in Metabolomics: New Insights into Structure and Dynamics of Metabolic Networks. *Bioanalysis* 2014, 6, 511–524. [PubMed: 24568354] (c)Rinkel J; Dickschat JS Recent Highlights in Biosynthesis Research Using Stable Isotopes. *Beilstein J. Org. Chem* 2015, 11, 2493–2508. [PubMed: 26734097] (d)Pons G; Rey E Stable Isotopes Labeling of Drugs in Pediatric Clinical Pharmacology. *Pediatrics* 1999, 104, 633–639. [PubMed: 10469806] (e)Kerfah R; Plevin MJ; Sounier R; Gans P; Boisbouvier J Methyl-Specific Isotopic Labeling: A Molecular Tool Box for Solution NMR Studies of Large Proteins. *Curr. Opin. Struct. Biol* 2015, 32, 113–122. [PubMed: 25881211] (f)Gant TG Using Deuterium in Drug Discovery: Leaving the Label in the Drug. *J. Med. Chem* 2014, 57, 3595–3611. [PubMed: 24294889]
28. (a)Lennox AJJ; Lloyd-Jones GC Selection of Boron Reagents for Suzuki-Miyaura Coupling. *Chem. Soc. Rev* 2014, 43, 412–443. [PubMed: 24091429] (b)Cox PA; Leach AG; Campbell AD; Lloyd-Jones GC Protodeboronation of Heteroaromatic, Vinyl, and Cyclopropyl Boronic Acids: PH-Rate Profiles, Autocatalysis, and Disproportionation. *J. Am. Chem. Soc* 2016, 138, 9145–9157. [PubMed: 27355973] (c)Chen L; Sanchez DR; Zhang B; Carrow BP “cationic” Suzuki-Miyaura Coupling with Acutely Base-Sensitive Boronic Acids. *J. Am. Chem. Soc* 2017, 139, 12418–12421. [PubMed: 28862445]
29. A competition reaction involving reductive coupling of nitromethane under the standard condition in the presence of both 4-fluorophenylboronic acid and 4-chlorophenylboronic acid pinacol ester showed that boronic acid is more reactive than boronic acid pinacol ester (Page S37). Evidently, there is no inherent preference of the nitroalkane for the boronic ester.
30. (a)Itil TM; Akpinar S; Fink M Controlled clinical and quantitative EEG studies of triflubazam (ORF 8063) in patients with anxiety syndrome. *Curr. Ther. Res* 1976; 19, 307. [PubMed: 5248] (b)Csanlosi I; Pereira-Oran J; Case G; Werblowsky J; Rickels K Triflubazam (ORF 8063), a new benzodiazepine in anxiety neurosis. *Curr. Ther. Res* 1977, 22, 166.(c)Nicholson AN; Stone BM; Clarke CH Effect of the 1,5-benzodiazepines, clobazam and triflubazam, on sleep in man. *Br. J. Clin. Pharmacol* 1977, 4, 567. [PubMed: 20917]
31. Lenstra DC; Rutjes FPJT; Mecinovi J Triphenylphosphine-Catalysed Amide Bond Formation Between Carboxylic Acids and Amines. *Chem. Commun* 2014, 50, 5763–5766.

A. C- vs N-functionalization of nitromethane

B. Envisioned P^{III}/P^V reaction sequence

Do neutral P^{III}/P^V conditions enable reductive N-functionalization over competing pathways arising from lability of α -C-H positions?

C. Electronic considerations concerning biphilic reactivity of P

Does phosphetane biphilicity permit FMO-controlled pericyclic (3+1) addition even for less reactive nitro substrates like $\text{H}_3\text{C}-\text{NO}_2$?

Figure 1.

A) Uses of nitromethane in organic synthesis. B) Reaction sequence leading to formation of N-methylanilines via P^{III}/P^V=O-catalyzed reductive C–N coupling of nitromethane with boronic acids (esters); C) Electronic challenge presented by high frontier orbital gap of $\text{H}_3\text{C}-\text{NO}_2$. FMO=frontier molecular orbital.

Figure 2.

Synthetic scope of P^{III}/P^V=O-catalyzed C–N coupling of nitromethane and arylboronic acids. See SI for full experimental details and conditions. Yields are reported for material obtained following purification and isolation.

Figure 3. Synthetic scope of $\text{P}^{\text{III}}/\text{P}^{\text{V}}=\text{O}$ -catalyzed C–N coupling of nitromethane and (hetero)arylboronic esters. See SI for full experimental details and conditions. Yields are reported for material obtained following purification and isolation. ^a Yields for reactions with boronic acids were determined by ^1H NMR with dibromomethane as the internal standard.

Figure 4.

A) Reaction of a four component mixture illustrating differential reactivity of H_3CNO_2 vs ArNO_2 and ArB(OH)_2 vs ArB(OR)_2 . (B) Selective three-component coupling via sequential C–N coupling reactions.

triflubazam
51% yield over three steps

Figure 5.

Two-pot, three step synthesis of triflubazam by an all- $\text{P}^{\text{III}}/\text{P}^{\text{V}}$ -catalyzed sequence. Reaction conditions: (a) **43** (1.0 equiv), PhB(OH)_2 (1.1 equiv), Ph_2SiH_2 (4.0 equiv), **1•[O]** (30 mol%), CPME, 120 °C ; then add monoethyl malonate (1.2 equiv), diethylbromomalonate (1.5 equiv), 40 °C; (b) **44** (1.0 equiv), $\text{H}_3\text{C-NO}_2$ (3.0 equiv), Ph_2SiH_2 (3.0 equiv), **1•[O]** (15 mol %), CPME, 120 °C.

Table 1.Discovery and Control Experiments for Organophosphorus-Catalyzed *N*-methylaniline Synthesis.^a

Entry	silane	R ₃ P=O	Yield (%) ^b
1 ^c	PhSiH ₃	1•[O]	35
2	PhSiH ₃	1•[O]	95 (90) ^d
3	PhSiH ₃	1	94
4	PhSiH ₃	None	0
5	None	1•[O]	0
6 ^e	PMHS	1•[O]	85

^aUnless otherwise noted, reactions were carried out with **1•[O]** (10 mol %), phenylsilane (0.5 mmol, 2.0 equiv), 4-fluorophenylboronic acid (0.25 mmol) and nitromethane (0.75 mmol, 3.0 equiv) in CPME (0.5 mL) at 120 °C for 18 h.

^bYields were determined by ¹⁹F NMR integration with the aid of an internal standard.

^c1.0 equiv of **2** was used.

^dIsolated yield.

^e24 h reaction time.