Understanding the molecular basis on the biological suppression of bacterial leaf blight of anthurium exerted by *Bacillus subtilis* (BIO3) through proteomic approach

S. Nakkeeran¹ · M. Suganyadevi¹ · S. Rajamanickam¹

Received: 23 July 2020 / Accepted: 26 September 2020 / Published online: 10 October 2020
© King Abdulaziz City for Science and Technology 2020

Abstract

We attempted to study the antibacterial activity of rhizospheric *Bacillus* spp., to curb the bacterial blight of anthurium caused by *Xanthomonas axonopodis* pv. *dieffenbachiae* (Xad). Twenty-eight bacterial isolates from rhizospheric regions were identified as different *Bacillus* spp. and *Ochrobactrum* sp. using 16S rRNA gene sequencing. *B. subtilis* BIO3 effectively inhibited the growth of Xad up to 1450.7 mm², and extracted volatile organic metabolites from the isolate BIO3 inhibited the growth of Xad up to 1024 mm². Tritrophic interaction of anthurium leaves bacterized with *B. subtilis* BIO3 and challenged with Xad resulted in the expression of 12 unique proteins compared to untreated control. Mascot Peptide Mass Fingerprint-based identification indicated that one was glutathione peroxidase, involved in defence mechanism, other six proteins were identified as leghemoglobin II, CTP synthase-like, predicted protein (*Physcomitrella patens*), centromere-associated protein E, grain size protein, and five proteins were hypothetical proteins. Foliar application with 1% liquid formulations (10⁸ CFU/ml) of *B. subtilis* BIO3 significantly suppressed the bacterial leaf blight of anthurium up to 78% over untreated control and also increased the stem length and flower yield.

Keywords Anthurium bacterial blight · *Bacillus* spp · 2D-PAGE analysis · Expressed protein · Disease suppression

Introduction

Anthurium (*Anthurium andreanum* Linden) is the second largest cut flower cultivated in tropical humid climate. In India, it is commercially grown in Tamil Nadu, Kerala, Madhya Pradesh, Karnataka, Goa and Maharashtra. The flowers of anthurium are popular among flower arrangers because of their attractive colours, increased vase life, bold effect and long-lasting qualities (Teixeira da Silva et al. 2015; Prakasha et al. 2017). The greatest advantage of anthurium is it produces flowers throughout the year around. Though anthurium has commercial potential, its cultivation is hindered due to the bacterial leaf blight caused by *Xanthomonas*

axonopodis pv. *dieffenbachiae* (Xad). The disease was first reported from Brazil in 1960 (Nishijima 1988) and is prevalent in almost all anthurium growing regions of the world. The disease was reported in Tamil Nadu by Sathyanarayana et al. (1998) and in the state of Kerala by Dhanya et al. (2016). Symptoms of bacterial blight appeared on the leaves were water-soaked spots on the abaxial surface coupled with faint chlorosis, yellow zone along the leaf margins, necrotization on flower stalk and drying of the entire flower stalk (Aysan and Sahin 2003; Balan et al. 2014; Suganyadevi et al. 2015). A larger number of chemicals have been deployed for the management of bacterial disease. Unfortunately, the application of bactericides including antibiotics might result in the development of resistance among the bacterial pathogen. However, the inherent disadvantages of continuous usage of chemicals in ecosystem have warned for the search of alternatives. In attempt to aiming towards the reduction of synthetic chemical, extensive research on exploitation of rhizosphere and endophytic PGPR towards management is being taken as they are safe for environment and human beings.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s13205-020-02456-2>) contains supplementary material, which is available to authorized users.

✉ S. Nakkeeran
nakkeeranayya@tnau.ac.in

¹ Department of Plant Pathology, Tamil Nadu Agricultural University, Coimbatore 641 003, India

Several *Bacillus* strains were demonstrated to manage the plants against many fungal, bacterial and viral diseases (Srinivasan and Mathivanan 2011; Dheepa et al. 2016). The plant growth-promoting rhizobacteria (PGPR) showed much attention because of their advantage over other microbial agents, which include broad-spectrum antimicrobial activity and growth-promoting capacity (Lian et al. 2011). Previous findings indicated that *Bacillus* strains have a greater capacity to produce compounds such as lipopeptide, which can be cyclic or linear, mainly consisting of 7 to 11 amino acid residues linked to β -amino or β -hydroxy fatty acids (Athukorala et al. 2009). The *Bacillus* isolates have been reported with antimicrobial peptide genes responsible for the biosynthesis of various antibiotics including iturin, surfactin, fengycin and bacillomycin D (Isabel et al. 2011; Vinodkumar et al. 2017). Application of PGPR challenged by the plant pathogens triggers defence proteins and suppress the ingress of plant pathogens. Proteomics has been widely used in the field of understanding stress responses and constitutive differences between developmental stages or genotypes. Proteomics is an ideal tool for the dissection of plant microbe interactions (Saveetha et al. 2009; Prabhukarthikeyan et al. 2017). The research behind the tritrophic interaction between anthurium (leaf), biocontrol agent and *Xad* is only at the stage of infancy. With this background, we attempted to unveil the mechanism of action through metabolomic and proteomic approaches to suppress bacterial blight and to manage bacterial blight of anthurium using *Bacillus* spp. under protected cultivation.

Materials and methods

Isolation of bacterial pathogen

The bacterial blight pathogen *X. axonopodis* pv. *diefenbachiae* (*Xad*) was isolated from infected anthurium variety Rosette expressing the typical symptoms of

bacterial blight on nutrient agar medium as described (Balan et al. 2014). The purified pathogenic bacterium was prepared by suspending the bacterial cells in 50 ml of sterile nutrient broth from 24-h-old bacterial culture and adjusted to approximately 10^6 cells/ml. To prove the Koch postulate, fully expanded leaves were spray inoculated thrice at weekly intervals without pin prick method. The plants were covered with transparent polythene sheet of 150 gauge thickness and maintained in mist chamber at 28 ± 2 °C and at relative humidity of 90–95%. Three replications were maintained with 5 plants per replication. Likewise, uninoculated healthy control was also maintained separately in the humid chambers. The virulence of the isolate was assessed by the expression of symptoms.

PCR detection of bacterial pathogen using 16S rRNA and housekeeping genes

The genomic DNA from *Xad* was isolated using CTAB method described (Knapp and Chandless 1996). Amplification of 16S rDNA gene was performed using universal primers, 799F (5' AACMGGATTAGATACCKG 3') and 1193R (5' ACGTCATCCCCACCTTCC 3') (Weisburg et al. 1991). Apart from it, the pathogen was also confirmed with gene-specific primers corresponding to ATP synthase subunit beta gene (*atpD*), molecular chaperone (*dnaK*), elongation factor P (*Efp*), ton B-dependent receptor (*fyuA*), glutamine synthetase gene (*glnA*) and DNA gyrase subunit B gene (*gyrB*) (Table 1) (Nadia et al. 2013). Amplification was performed with a total reaction volume of 50 μ l in Eppendorf Mastercycler, Germany. The PCR products were resolved on 1.2% agarose stained with ethidium bromide (0.5 μ g/ml) and photographed. After electrophoresis, the gel was sliced along with PCR fragment and purified with QIAquick Gel Purification Kit (QIAGEN, California) according to the standard manufacturer's protocol and sequenced at Chromous Biotech Pvt. Ltd, Bangalore.

Table 1 The gene-specific primers used in the present study

S. no.	Gene name	Primer sequence (5'–3')
1	ATP synthase subunit beta gene (<i>atpD</i>)	F-GGGCAAGATCGTTCAGAT R-GCTCTTGGTCGAGGTGAT
2	Molecular chaperone (<i>dnaK</i>),	F-GGTATTGACCTCGGCACCAC R-ACCTTCGGCATACTGGGTCT
3	Elongation factor P (<i>Efp</i>)	F-TCATCACCGAGACCGAATA R-TCCTGGTTGACGAACAGC
4	Ton B-dependent receptor (<i>fyuA</i>)	F-ACCATCGACARGGACTGGACC R-GTCGCCGAACAGGTTACAC
5	Glutamine synthetase gene (<i>glnA</i>)	F-ATCAAGGACAACAAGGTGCG R-GCGGTGAAGTTCAGGTAG
6	DNA gyrase subunit B gene (<i>gyrB</i>)	F-TGCGCGGCAAGATCCTCAAC R-GCGTTGCTCTCGATGAAGTC

Isolation and identification of rhizospheric bacteria

Twenty-eight *Bacillus* spp. were isolated from rhizoplane of anthurium germplasm accessions, viz. Rosette, Glamour, Temptation, Red dragon, Hawaiian red and Colorado red maintained at TNAU, Coimbatore, through serial dilution technique. The isolates of *Bacillus* spp. were identified according to the description given in Bergey's manual for Systematic Bacteriology and cryopreserved at -80°C in 60 per cent glycerol for further use. The isolated bacterial cultures were inoculated into nutrient broth, and the genomic DNA was extracted using CTAB method as described (Knapp and Chandlee 1996). To confirm the isolates as *Bacillus* spp., 16S rRNA intervening sequence-specific BCF1 (5' CGGGAGGCAGCAGTAGGGAAT 3') and BCR2 (5' CTCCCCAGGCGGAGTGCTTAAT 3') primers were used (Cano et al. 1994). Amplification was performed with a total reaction volume of 50 μl in Eppendorf Mastercycler, Germany. The PCR products were resolved on 1.2% agarose at 50 V, stained with ethidium bromide (0.5 $\mu\text{g}/\text{ml}$) and photographed using gel documentation system (Alpha Innotech Corporation, San Leandro, California). The nucleotide sequences of genomic product were aligned using CLUSTAL W program, and BLAST search program was used to identify the *Bacillus* isolates based on the similarity index (Larkin et al. 2007).

In vitro screening of bacterial antagonist

The antibacterial activities of different bacterial antagonists were tested against *Xad* by paper disc assay in nutrient agar medium (Bauer et al. 1996; Rajamanickam et al. 2018). The bacterial antagonists were inoculated in nutrient broth and incubated for 72 h at $28 \pm 2^{\circ}\text{C}$. One ml of 72-h-old *X. axonopodis* pv. *dieffenbachiae* maintained in nutrient broth consisting of 10^6 cells/ml was inoculated into nutrient agar medium. The NA medium seeded with the pathogen was poured into Petri plates (90 mm). The sterile paper disc dipped in bacterial antagonistic suspension was placed on *Xad* seeded medium apart from the periphery of the Petri plates at four directions. Three replications were maintained for each treatment and without antagonist was maintained as a control. The efficacy of bacterial antagonists against *Xad* was assessed by measuring the area of inhibition zone in mm^2 using the graph sheet.

Inhibitory activities of secondary metabolites against *Xad*

The secondary metabolites were extracted from ten effective *Bacillus* spp. identified as per the results of in vitro screening using the standard protocol (Prapagdee et al.

2012). The antibacterial activities of secondary metabolites isolated from effective antagonist were tested against *Xad* using agar well diffusion assay. Seventy-two-hour-old bacterial pathogen *Xad* (10^6 cells/ml) maintained in nutrient broth was added into the sterile molten NA medium at the rate of 1 ml/100 ml while pouring on to the Petri plates. Nutrient agar was poured into the sterile Petri plates and allowed to solidify. A well (5 mm in diameter) was made by punching the nutrient agar with a sterile cork borer on the corner of the plate in four directions by leaving a distance of 1 cm away from the periphery of the plates. Each well was poured with 50 μl of secondary metabolite separately in single plates to assess the antibacterial activity. The efficacy was assessed by measuring the area of inhibition zone (mm^2) using the graph sheet after 48 h of incubation at $28 \pm 2^{\circ}\text{C}$.

Bio-efficacy of *Bacillus* spp. under protected cultivation

The glasshouse experiment was conducted to assess the bio-efficacy of liquid formulation of *Bacillus* spp. against bacterial blight of anthurium under protected cultivation. Two-year-old anthurium var. Rosette grown in cocopeat was used throughout the experiment. As per the previous screening experiments, seven most effective isolates, viz. *B. cereus* (BTW-R23 and BTS-R32), *B. amyloliquefaciens* (BS4, BS2 and BIO3) and *B. subtilis* (BIO3 and BTW-R12), were cultured on nutrient agar medium. A loop full of culture from stock was inoculated into 1000 ml of nutrient broth and incubated in an orbital shaker at 150 rpm at $28 \pm 2^{\circ}\text{C}$ for 48 h. Later, the liquid biomass along with bacterial cells was mixed with 1% glycerol (10 ml), 1% tween 20 (10 ml) and 1% polyvinyl pyrrolidone 40000 MW (10 g) each separately (Vinodkumar et al. 2017). The resultant mixture was incubated in orbital shaker at 200 rpm for 5 min to ensure uniform mixing of the bacterial cells. One ml of the formulation consists of 10^8 cfu/ml. To assess the efficacy of antagonistic *Bacillus* spp., for the management of bacterial blight of anthurium, six foliar applications were given at weekly intervals using knapsack sprayer, after the initiation of the disease. *Pseudomonas fluorescens* pf1 (10^8 cfu/ml) obtained from the Department of Plant Pathology, TNAU, Coimbatore, was maintained as standard check to compare the efficacy of *Bacillus* spp. The incidence of bacterial blight, number of leaves, length of stem, number of flowers and height of the plants were observed after 20 days of last spray. Three replications were maintained for each treatment with 5 plants per replication. The trial was repeated twice, and pooled mean analysis for the two seasons has been carried out for validating the results.

Proteomic analysis for identification of defence response

In order to find the change in defence response under tri-trophic interactions against pathogen, effective isolate *B. subtilis* BIO3-treated anthurium plants were sampled to elucidate the proteomic analysis (Prabhukarthikeyan et al. 2017). The bulbs of the anthurium variety Rosette were surface sterilized with 2% sodium hypochlorite for 30 s and rinsed with sterile distilled water. Bulbs were individually placed in 30-cm-diameter pots with 30-cm spacing between the pots. Plants were grown under 70% UV-stabilized shade net with natural light for 8 weeks before inoculation. The temperature of the shade net was 30 °C during the day and 22 °C at night. Sixty-day-old plants were challenged with pathogen *Xad* and biocontrol agent *B. subtilis* BIO3. The application of liquid formulation of *B. subtilis* BIO3 was carried out 24 h prior to *Xad* inoculation; later on, seedlings were challenged with the *Xad*. After the inoculation, plants were covered with polythene cover and placed in a mist chamber for twenty days and was maintained ~100% humidity. The observations were made on disease development after twenty days of inoculation. The treatment details are T1-healthy control (absolute control); T2-pathogen inoculated (absolute control + *Xad*); T3-interaction (absolute control + *Xad* + foliar application of *B. subtilis* BIO3); and T4-biocontrol (absolute control + foliar application of *B. subtilis* BIO3). Three replications were maintained for each treatment with 5 plants per replications. To understand the induction of defence proteins in tritrophic interaction, leaves from the bacterized anthurium plants were collected, transferred to liquid nitrogen and processed for 2D-PAGE analysis.

Protein extraction

Frozen leaf tissues were ground in a mortar with liquid nitrogen and suspended in 10% trichloroacetic acid (TCA) in acetone with 0.07% dithiothreitol (DTT) at –20 °C for 1 h, followed by centrifugation for 20 min at 12,000 rpm. The pellets were washed with ice-cold acetone containing 0.07% DTT at –20 °C for 1 h and centrifuged again for 20 min at 12,000 rpm. The washing step was repeated three to four times until the supernatant was free of chlorophyll. The precipitated pellets were placed for one hour in a –80 °C freezer until frozen and then placed in a lyophilizer. The freeze-dry procedure was carried out under vacuum overnight until the pellet turns into a dry powder. A total of 10 mg of the dried powder was dissolved in 250 µl of sample buffer containing 7 M urea, 2 M thiourea, 4% 3-[(3-cholamidopropyl) dimethylammonia]-1-propanesulfonate (CHAPS), 0.5% ampholytes (Bio-Rad) and 0.7% DTT. The powder was dissolved in the extraction buffer and was gently shaken for

1 h and centrifuged for 20 min at 12,000 rpm at 28 ± 2 °C. The supernatant was distributed in 100 µl aliquots and maintained at –80 °C before 2D-PAGE analysis.

2D-PAGE

For analytical gels, the 17 cm IPG (Immobiline pH Gradient) strips (pH 4–7) were rehydrated overnight with 350 µl of rehydration buffer (8 M urea, 2% CHAPS, DTT (7 mg per 2.5 ml of rehydration buffer) and 0.5% (v/v) IPG buffer pH 4–7) containing the required quantity of proteins (150 µg) in a reswelling tray (Pharmacia Biotech., USA) at room temperature. The strips were allowed for about 12–24 h for rehydration and subjected to the first-dimension separation. Isoelectric focusing (IEF) was carried out at 20 °C with a Pharmacia Multiphore II kit. The running conditions were as follows: 500 V for 1 h followed by 1000 V for 1 h and finally 3000 V for 16 h. The focused strips were equilibrated twice for 15 min in 10 ml equilibration solution. The first equilibration was performed in a solution containing 6 M urea, 30% (w/v) glycerol, 2% (w/v) SDS, 1% (w/v) DTT and 50 mM Tris–HCl buffer. The second equilibration was performed in a solution modified by the replacement of DTT by 4% (w/v) iodoacetamide. After polymerization of the gel, gels were electrophoresed at 15 mA till the dye front reaches the bottom of the gel. Gels were removed from their gel cassettes and fixed in the fixatives (4:1:5 of methanol/acetic acid/water). The gels were stained by silver staining method as described (Blum et al. 1987). The reaction was stopped using a stopper reagent and rinsed again with deionized water.

Analysis of 2D-PAGE

Selected protein spots were excised manually from the 2D gel, and the protein on the gels was digested as per the protocol (Shevchenko et al. 1996). Pooled extracts of proteins were dried using a speed vacuum and the extracts were re-dissolved in 1 µl of extraction buffer and 1 µl of matrix solution and targeted onto the matrix-assisted laser desorption/ionization (MALDI) plate. The samples were completely dried onto the targeting plate and analysed by MALDI-TOF and GC–MS. MALDI-TOF experiments were performed on a Voyager-DE STR mass spectrometer (Applied Biosystems, Framingham, MA, USA) equipped with delay ion extraction. Mass spectra were obtained over a mass range of 800–3,000 Da. The proteins were identified by Mascot Peptide Mass Fingerprint (www.matrixscience.com) database research. The following parameters were used for database searches: taxonomy, viridiplantae, cleavage specificity, trypsin with 1 missed cleavage allowed peptide tolerance of 100–300 ppm for the fragment ions, allowed modifications, Cys carbamidomethyl (fixed) oxidation of Met (variable).

The peptide mass fingerprinting of the proteins was scored with the Mowse score.

Statistical analysis

Mean differences of the treatment were evaluated with ANOVA using Duncan's multiple-range test at 5% significance (Gomez and Gomez 1984). All the data were statistically analysed with IRRISTAT (version. 3/93, Biometrics unit, International Rice Research Institute) and consequently interpreted.

Results

Bacterial pathogen and pathogenicity

The bacterial pathogen was isolated from infected anthurium plants expressing the typical symptom of bacterial blight. The morphology of bacterial colony showed the presence of raised projection at the centre of the colony, and the margins were smooth, concave and bright yellow in colour on the nutrient agar medium. The bacterial pathogen was purified by streak plate method and stored at 4 °C under refrigeration. The pathogenicity assay proved that the bacterial pathogen expressed the typical symptoms on leaves, spathe and spadix of anthurium plants after 20 days of last spray inoculation. The symptoms associated with bacteria expressed the presence of water-soaked lesions on leaf margin and subsequently turned to necrotic brown.

Genotypic characterization of *Xad*

PCR to identify the bacterial pathogen associated with bacterial blight revealed the presence of genomic product with the fragment size of approximately 1500 bp corresponding to 16S rRNA gene. The amplified products were separated on 1.2% agarose gel and sequence determined. The rDNA homology searches using BLAST program though NCBI database confirmed the association of *X. axonopodis* pv. *dieffenbachiae* (*Xad*) causing bacterial blight of anthurium. The sequence had 99 per cent similarity to *Xad* available in NCBI database. The sequence of our isolate was submitted in NCBI database, bearing the accession number KU753854. Subsequently, the pathogen was also confirmed by multi-locus sequencing with housekeeping genes of *Xad*. The results revealed the amplicons of 868 bp, 1034 bp, 445 bp, 963 bp, 1094 bp and 1051 bp corresponding to ATP synthase subunit beta (*atpD*), Molecular chaperone *dnak* (*Dnak*), elongation factor (*efp*), ton B-dependent receptor (*fyuA*), glutamine synthetase type 1 (*glnA*) and DNA gyrase subunit B (*gyrB*) genes, respectively (Figure S1). The PCR products were

sequenced, and nucleotide sequence analysis using BLAST programme revealed that the sequences had 95–100 per cent similarity to the respective genes. The sequences of our isolate were submitted in NCBI database, bearing the accession no. KX033359, KX033362, KX033364, KX033370, KX033367 and KX033369, respectively.

Isolation and identification of bacterial antagonists

Twenty-eight isolates of bacterial antagonists were isolated from rhizoplane of anthurium and identified using 16S rRNA gene. All the isolates were amplified with a product size of 546 bp corresponding to 16S rRNA gene and sequence determined. The sequence homology search using NCBI BLAST revealed that out of 28 isolates 5 were *Bacillus amyloliquefaciens* (accession numbers KX363487, KU753858, KU753864, KU753857 and KU753861), 9 were *Ochrobactrum* spp., (accession numbers KU753868, KU753882, KU753883, KU753856, KU753859, KU753862, KU753872, KU753873 and KU753877), 4 were *B. subtilis* (accession numbers KX363486, KU753871, KU753863 and KU753866), 6 antagonists were *B. cereus* (accession numbers KU753881, KU753884, KU753874, KU753875, KU753876 and KX363485), and each one was from *Bacillus* sp., *Bacillus thuringiensis* (accession number KX363484), *Brevibacillus lateroporus* (accession number KU753879) and *B. licheniformis* (accession number KU753869), respectively (Table S1).

Screening of bacterial antagonists and secondary metabolites for the antibacterial activity against *Xad*

In vitro screening of bacterial antagonists against *Xad* indicated that *B. subtilis* BIO3 inhibited the growth of *Xad* over an area of 1450.7 mm² and was followed by *B. subtilis* BTWR12 and *B. amyloliquefaciens* BS2, which inhibited the growth of *Xad* over an area of 1419.7 and 1353.3 mm², respectively. However, the other assayed bacterial antagonists also inhibited the growth of *Xad* to a tune of 19.0 to 1324.3 mm² (Fig. 1). However, the least area of inhibition with 15.3 mm² was observed in *Ochrobactrum* sp. BTS-R20. Further, the secondary metabolites extracted from ten different effective *Bacillus* spp. were tested for their antibacterial activity against *Xad* in vitro. The maximum area of inhibition was observed with the metabolites extracted from *B. subtilis* BIO3 by accounting for the maximum area of inhibition of 1024 mm² (Fig. 2). It was followed by *B. subtilis* BTW-R12 (1019 mm²) and *B. amyloliquefaciens* BS2 (1015 mm²).

Fig. 1 Inhibition of growth of *X. axonopodis* pv. *dieffenbachiae* by antagonistic bacteria. Error bars indicate standard deviation obtained from three replicates per treatment. Different letters indicate significant differences between treatments

Fig. 2 Inhibition of growth of *X. axonopodis* pv. *dieffenbachiae* by secondary metabolites of effective *Bacillus* spp. Error bars indicate standard deviation obtained from three replicates per treatment. Different letters indicate significant differences between treatments

Effect of bacterial antagonists against bacterial blight and growth promotion under protected condition

The seven selected effective isolates of *Bacillus* spp., namely *B. subtilis* (BIO3 and BTW-R12), *B. amyloliquefaciens* (BS2, BS4 and BIO4) and *B. cereus* (BTW-R23 and BTS-R32), were tested for their efficacy on bacterial blight of anthurium and plant growth promotion activities under protected conditions. The result revealed that foliar application of *B. subtilis* BIO3 at weekly intervals for six times reduced the incidence of bacterial blight in susceptible rosette variety, which had the minimum disease index of 8.55 PDI as against 39.28 PDI in untreated control, which was 78.23% reduction over control. It was followed by *B. subtilis* BTW-R12 and *B. amyloliquefaciens* BS2 which suppressed the bacterial blight up to 67.92 and 63.37 per cent reduction over control, respectively. The *Pseudomonas fluorescens* pf1 used as a comparative check was less effective in reducing

the bacterial blight in comparison with *Bacillus* spp. (Figure 3). The results on plant growth promotion revealed that the application of *B. subtilis* BIO3 with six sprays at weekly intervals had the maximum numbers of leaves, stem, plant height (cm) and flowers, which were accounted for 11.12 (Nos.), 13.25 (Nos.), 29.60 cm and 7.23 (Nos.), respectively. This was followed by foliar application of *B. subtilis* BTW-R12 and *B. amyloliquefaciens* BS2 (Fig. 4).

Expressed proteins during the tripartite interaction

The protein spots were reproducibly resolved in all gels which resulted in similar protein spot locations across all the replications. The results revealed that around 400 clear and consistent protein spots were observed in expressed protein map. Out of 400 proteins observed, 12 protein spots were associated with pathogen inoculated and challenged with biocontrol agent *B. subtilis* BIO3. Among the 12 proteins, 8 proteins were up-regulated in the plants

Fig. 3 Efficacy of bacterial antagonists against bacterial blight of anthurium under protected cultivation. Error bars indicate standard deviation obtained from three replicates per treatment. Different letters indicate significant differences between treatments

Fig. 4 Plant growth promotion of anthurium by *Bacillus* spp. under protected cultivation. Error bars indicate standard deviation obtained from three replicates per treatment. Different letters indicate significant differences between treatments

bacterized with *B. subtilis* BIO3 and challenged with the *Xad* pathogen (P2, P4, P5, P6, P8, P10, P11 and P12) (Figs. 5, 6). However, the up-regulation of the eight different proteins was maximum in the plants bacterized with *B. subtilis* BIO3 and challenged with the pathogen. The up-regulated proteins, viz. P2, P4, P5, P6, P8, P10, P11, P12, were identified as leghemoglobin II in *Lupinus luteus*, hypothetical protein associated with *Phaseolus vulgaris*, predicted protein in *Physcomitrella patens*, centromere-associated protein E in *Morus notabilis*, hypothetical protein from *Vitis vinifera*, hypothetical protein from *Zea mays*, hypothetical protein from *Chlorella variabilis* and grain size protein from *Triticum aestivum* (Table 2). However, since the work on proteomic approaches in anthurium has not been done earlier by other workers, the functions of the hypothetical proteins are not clearly known.

The proteins P1 and P3 were down-regulated in pathogen-inoculated treatments compared to that of uninoculated control (Fig. 7). However, there was no change with respect to same proteins in the plants bacterized with *B. subtilis* BIO3 challenged with *Xad*. The down-regulated proteins were identified as hypothetical protein associated with *Oryza sativa japonica* and hypothetical protein associated with *Phaseolus vulgaris*, respectively. Likewise, the protein P7 known as predicted protein from *Micromonas* spp. was down-regulated in pathogen inoculated control. However, the same protein was up-regulated in plants bacterized with biocontrol agent and plants challenged with *Xad*, indicating that this protein might play a vital role in inducing the defence reaction. The protein P9 was identified as glutathione peroxidase and was up-regulated in pathogen inoculated control (Table 2). However, there was no change

Fig. 5 Proteome analysis of anthurium treated with *B. subtilis* BIO3 and challenged with pathogen *Xad* by 2-D gel electrophoresis. **a** (Anthurium + *Xad*), **b** (Anthurium + *Bacillus*), **c** (Anthurium + *Xad* + *Bacillus*) and **d** Healthy anthurium

in regulation of protein compared to the control in the anthurium leaves bacterized with *B. subtilis* BIO3 followed by challenge inoculation with the pathogen.

Discussion

The present study confirmed the potential of *Bacillus subtilis* BIO3 as an effective biocontrol agent against *X. axonopodis* pv. *dieffenbachiae* causing bacterial blight of anthurium. *B. subtilis* BIO3 exhibited a synergistic mode of action against anthurium bacterial blight. The direct mode of actions of *Bacillus* is production of antimicrobial peptides and volatile compounds which exhibited the antibacterial activity against pathogen *Xad*. The indirect mode of action includes the induction of defence proteins and promotes the plant growth and yield. In the present study, isolated bacterial pathogen *X. axonopodis* pv. *dieffenbachiae* was amplified with the specific primers corresponding to 16S rRNA and housekeeping like ATP synthase subunit beta gene (*atpD*),

Molecular chaperone (*dnaK*), Elongation factor P (*Efp*), ton B-dependent receptor (*fyuA*), glutamine synthetase gene (*glnA*) and DNA gyrase subunit B gene (*gyrB*). These results are in agreement with the findings of Nadia et al. (2013), who identified the plant pathogenic bacteria *X. axonopodis* pv. *dieffenbachiae* was associated with the anthurium leaf blight and housekeeping genes are the molecular sieve in identification of bacterial pathogen.

To manage the leaf blight of anthurium, 28 isolates of rhizospheric *Bacillus* spp. were isolated from anthurium. They were identified as *B. thuringiensis*, *B. lateroporus*, *B. subtilis*, *B. amyloliquefaciens*, *B. licheniformis*, *B. cereus* and *Ochrobactrum* sp. using 16S rRNA. The inhibitory effect of biocontrol agents was documented based on area inhibition under in vitro assay, which was correlated with the management of bacterial blight of anthurium under protected cultivations. Undeniably, bacterial antagonist *B. subtilis* BIO3 was most effective in inhibiting the bacterial

Fig. 6 Proteins up regulated in pathogen (*Xad*) and interaction (*Xad* + *B. subtilis* BIO3) compared with healthy control. P2, P4, P5, P6, P8, P10, P11 and P12 were differentially expressed proteins

pathogen *Xad* compared to other bacterial antagonists. *Bacillus* species are capable of producing a wide variety of secondary metabolites that has antimicrobial activities

against many fungal and bacterial pathogens. The capability of synthesis of secondary metabolites determines the ability of these species to control plant diseases. In

Table 2 Diversity of differential proteins expressed during tritrophic interaction in anthurium leaf tissues in response to foliar application with *Bacillus subtilis* BIO3 against *X. axonopodis* pv. *dieffenbachiae*

Spot ID	Accession No.	Protein name	Experimental				
			pI	MW (kDa)	Score	Matched peptides	Seq. Cov.
Hypothetical protein							
P1	gil222622568	Hypothetical protein [<i>Oryza sativa</i> Japonica Group]	11.71	12	55	4	53
P4	gil593793458	Hypothetical protein [<i>Phaseolus vulgaris</i>]	5.93	27	42	5	25
P8	gil147769915	Hypothetical protein [<i>Vitis vinifera</i>]	8.72	27	54	5	16
P10	gil413953152	Hypothetical protein [<i>Zea mays</i>]	6.75	27	49	5	18
P11	gil552847179	Hypothetical protein [<i>Chlorella variabilis</i>]	8.17	45	50	7	16
Protein involved in defence mechanism							
P9	gil255537447	Glutathione peroxidase, putative [<i>Ricinus communis</i>]	6.58	18	54	5	22
Other proteins							
P2	gil441461	Leghemoglobin II [<i>Lupinus luteus</i>]	5.95	16	41	4	29
P3	gil357150353	CTP synthase-like [<i>Brachypodium distachyon</i>]	5.78	63	46	7	14
P5	gil168001717	Predicted protein [<i>Physcomitrella patens</i>]	4.75	15	52	6	51
P6	gil587938546	Centromere-associated protein E [<i>Morus notabilis</i>]	6.07	94	56	10	13
P7	gil255088942	Predicted protein [<i>Micromonas</i> sp. RCC299]	8.33	2	41	4	19
P12	gil566086681	Grain size protein [<i>Triticum aestivum</i>]	7.79	93	46	4	52

our study, the maximum inhibitory activities of secondary metabolites were documented in *B. subtilis* BIO3 than the other *Bacillus* spp. Similarly, Suganyadevi et al. (2016) screened the different *Bacillus* isolates against bacterial blight pathogen and found the antagonistic activity of *B. mojavensis* strain KA3 against *Xad* under in vitro. Rajamanickam et al. (2018) reported the antibacterial activity of secondary metabolites of *B. subtilis* (PP and CL3) against *Pectobacterium carotovorum* pv. *carotovorum* causing rhizome rot of banana. The cell suspension and cell free culture filtrate of *B. amyloliquefaciens* (B014) have been reported to inhibit *X. axonopodis* pv. *dieffenbachiae* causing anthurium leaf blight (Li et al. 2012). Nakkeeran et al. (2019) identified the several antimicrobial compounds such as benzenemethanol, *n*-hexadecanoic acid, benzenepropanoic acid pyrrole, phenol and benzenoacetic acid from the secondary metabolites of *B. subtilis* (AP). Hence, the presence of antimicrobial compounds in *Bacillus* spp. served as a potent inhibitor of bacterial pathogen, which could be the reason for the suppression of bacterial blight of anthurium.

The efficacy of formulation of *Bacillus* species has already been demonstrated in different agriculture and horticulture crops against soil-borne disease under field conditions (Li et al. 2012; Mohan Kumar et al. 2015; Cheng et al. 2016). In the present study, foliar application *B. subtilis* BIO3 at weekly intervals significantly reduced the disease severity compared to untreated control. Similarly, Kumsingkaew and Akarapisan (2016) found that application of *B. subtilis* EPB14 recorded 50 per cent reduction in disease severity of bacterial blight of anthurium under greenhouse condition. The foliar application of *B. subtilis* (S-12) decreased the disease incidence of citrus canker incited by *X. axonopodis* pv. *citri* (Das et al. 2014). Application of endophytic *Bacillus* strains reduced the disease incidence of bacterial panicle blight of rice caused by *Burkholderia glumae*, under field condition and increased the plant growth promotion activity (Shrestha et al. 2016). Apart from the antibacterial activity of secondary metabolites, we also investigated the molecular mechanism of disease suppression mediated by *B. subtilis* BIO3 against *Xad* in anthurium through proteomic approach. The differentially expressed

Fig. 7 Proteins down regulated in pathogen (*Xad*); down and up regulated in interaction (*Xad*+*B. subtilis* BIO3). P1, P3, P7 and P9 were differentially expressed proteins

proteins were identified during tripartite interactions and the proteins spots gave reproducible and consistent patterns. In the present investigation, 12 proteins were associated with pathogen inoculated and challenged with biocontrol agent *B. subtilis* BIO3. Among the 12 proteins, 10 proteins were up-regulated in plants bacterized with *B. subtilis* BIO3, challenged with the pathogen *Xad*. These proteins were hypothetical proteins characterized from *Physcomitrella patens*, *Morus notabilis*, *Vitis vinifera*, *Zea mays*, *Chlorella variabilis* and *Triticum aestivum*. This finding was supported by Prabhukarthikeyan et al. (2017) who emphasized that protein expression of *Curcuma longa* was up-regulated in interaction treatments and matched with uncharacterized proteins of *Cucumis melo* and *Phaseolus vulgaris*. Similarly, Shen et al. (2003) identified the rice leaf sheath proteome with 400 proteins resolved on 2D gel. These results indicated that the proteins induced were hypothetical proteins, whose function is not yet identified. Out of twelve differentially expressed proteins, protein (P7) was up-regulated in tripartite interaction in anthurium and was homologous to predicted protein from *Micromonus* spp., which was associated with stronger resistance to pathogens. Similarly, Saveetha et al. (2009) reported that sixteen proteins were up-regulated

and seven proteins were down-regulated in rice leaf sheath treated with *Pseudomonas fluorescens* strain KH-1. These proteins were involved in general metabolism, defence, ISR, biotic and abiotic stress. This clearly indicated that the differentially expressed unique proteins in anthurium treated with *B. subtilis* BIO3 and challenge inoculated with *Xad* would be a novel protein involved in plant defence. To our knowledge, the current study was the first application of a metabolomic and proteomic approach to elucidate the mechanism of resistance to bacterial blight of anthurium under protected cultivation. The bacterial antagonists *B. subtilis* BIO3 isolated from rhizospheric region produced wide variety of antibacterial compounds including small peptides and fatty acids which inhibited the growth of bacterial pathogen. The foliar application of *B. subtilis* BIO3 in anthurium plants has up-regulated hypothetical proteins identified through 2D-PAGE analysis. The foliar application of *B. subtilis* BIO3 was effective in inhibiting the disease incidence and enhanced the plant growth promotion activity under protected cultivation. This could be the effective management strategy for the biological control of bacterial blight of anthurium, which plays an important role in sustainable agriculture.

Acknowledgement The author acknowledges DST-FIST, Department of Science and Technology, Government of India, New Delhi, for the infrastructure facilities for conducting the experiments.

Author contributions NS and RS have performed the laboratory experiments, glasshouse experiment, statistical analysis, data recording and writing the original draft. SM has performed the proteomic study. All authors proofread and reviewed the manuscript.

Compliance with ethical standards

Conflict of interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

Human and animal rights This article does not contain any studies with human participants or animals performed by any of the authors.

Informed consent Informed consent was obtained from all individual participants included in the study.

References

- Athukorala SNP, Fernando WGD, Rashid KY (2009) Identification of antifungal antibiotics of *Bacillus* species isolated from different microhabitats using polymerase chain reaction and MALDI-TOF mass spectrometry. *Can J Microbiol* 55:1021–1032
- Aysan Y, Sahin F (2003) First report of bacterial blight of anthurium caused by *Xanthomonas axonopodis* pv. *dieffenbachiae* in Turkey. *New Dis Rep* 7:5
- Balan S, Sajeesh PK, Abraham K (2014) Morphological and biochemical characterization of *Xanthomonas axonopodis* pv.

- dieffenbachiae* (McCulloch & Pirone) causing bacterial blight of *anthurium* (Schott.). Trends in Biosciences 7(8):636–640
- Bauer AW, Kirby WM, Sherris JC, Turck M (1966) Antibiotic susceptibility testing by a standardized Single disk method. Am J Clin Pathol 45:493–496
- Blum H, Beier H, Gross HJ (1987) Improved silver staining of plant proteins, RNA and DNA in polyacrylamide gels. Electrophoresis 8:93–99
- Cano RJ, Borucki MK, Highby-Schweitzer M, Poinar HN, Poinar GO, Pollard KJ (1994) *Bacillus* DNA in fossil bees: an ancient symbiosis. Appl Environ Microbiol 60(6):2164–2167
- Cheng J, Jaiswal KS, Yang SH, Suh JW (2016) Endophytic *Bacillus subtilis* MJMP2 from Kimchi inhibits *Xanthomonas oryzae* pv. *oryzae*, the pathogen of rice bacterial blight disease. J Appl Biol Chem 59(2):149–154
- Das R, Mondal B, Mondal P, Khatua DC, Mukherjee N (2014) Biological management of citrus canker on acid lime through *Bacillus subtilis* (S-12) in West Bengal, India. J Biopest 7:38–41
- Dhanya S, Sreekumar J, Jisha MS (2016) Evaluation of antagonistic potential and induction of flowering in *Anthurium andreaeanum* L. by indigenous *Pseudomonas taiwanensis*. Int J Agric Innov Res 4(4):760–766
- Dheepa R, Vinodkumar S, Renukadevi P, Nakkeeran S (2016) Phenotypic and molecular characterization of chrysanthemum white rust pathogen *Puccinia horiana* (Henn) and the effect of liquid based formulation of *Bacillus* spp. for the management of chrysanthemum white rust under protected cultivation. Biol Control. <https://doi.org/10.1016/j.biocontrol.2016.09.006>
- Gomez KA, Gomez AA (1984) Statistical procedure for agricultural research. Wiley, New York
- Isabel M, Jordi C, Emilio M (2011) Antimicrobial peptide genes in *Bacillus* strains from plant environments. Int Microbiol 14:213–223
- Knapp JE, Chandless JM (1996) RNA/DNA mini-prep from a single sample of orchid tissue. Biotechniques 21:54–56
- Kumsingkaew S, Akarapisan A (2014) Efficiency of *Bacillus subtilis* EPB14 as Biocontrol to control bacterial leaf blight of anthurium. Int J Agric Technol 10(3):755–766
- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG (2007) Clustal W and Clustal X version 2.0. Bioinformatics 23(21):2947–2948
- Li SB, Fang M, Zhou RC, Huang J (2012) Characterization and evaluation of the endophyte *Bacillus* B014 as a potential biocontrol agent for the control of *Xanthomonas axonopodis* pv. *dieffenbachiae*-Induced blight of Anthurium. Biol Control 63(1):9–16
- Li SB, Xu SR, Zhang RN, Liu Y, Zhou RC (2015) The antibiosis action and rice induced resistance, mediated by a lipopeptide from *Bacillus amyloliquefaciens* B014, in controlling rice disease caused by *Xanthomonas oryzae* pv. *oryzae*. J Microbiol Biotechnol 26(4):748–756. <https://doi.org/10.4014/jmb.1510.10072>
- Lian L, Xie L, Zheng L, Lin Q (2011) Induction of systemic resistance in tobacco against *Tobacco mosaic virus* by *Bacillus* spp. Biocontrol Sci Technol 21:281–292
- Mohan Kumar SP, Chowdappa P, Krishna V (2015) Development of seed coating formulation using consortium of *Bacillus subtilis* OTPB1 and *Trichoderma harzianum* OTPB3 for plant growth promotion and induction of systemic resistance in field and horticultural crops. Indian Phytopath 68(1):25–31
- Nadia MH, Hajri A, Boureau T, Darrasse A, Durand K, Brin C, Fischer-LeSaux M, Manceau C, Poussier S, Pruvost O, Lemaire C, Marie-Agnes J (2013) Evolutionary history of the plant pathogenic bacterium *Xanthomonas axonopodis*. PLoS ONE 8(3):e58474
- Nakkeeran S, Surya T, Vinodkumar S (2019) Antifungal potential of plant growth promoting *Bacillus* species against blossom blight of rose. J Plant Growth Regul. <https://doi.org/10.1007/s00344-019-09966-1>
- Nishijima WT (1988) Anthurium blight: An overview. In: A. M. Alvarez (Eds) Proceedings of the first anthurium blight conference, Honolulu, USA, pp. 6–8
- Prabhukarthikeyan SR, Manikandan R, Durgadevi D, Keerthana U, Harish S, Karthikeyan G, Raguchander T (2017) Bio-suppression of turmeric rhizome rot disease and understanding the molecular basis of tripartite interaction among *Curcuma longa*, *Pythium aphanidermatum* and *Pseudomonas fluorescens*. Biol Control 111:23–31
- Prakash DP, Ramya G, Srinivasalu GB (2017) Morphogenetic responses in *Anthurium andreaeanum* (Hort) cultivars. Curr Agric Res J 5(1):135–140
- Prapagdee B, Tharasaithong L, Nanthaphot R, Paisitwiroj C (2012) Efficacy of crude extract of antifungal compounds produced from *Bacillus subtilis* on prevention of anthracnose disease in *Dendrobium* Orchid. Environ Asia 5(1):32–38
- Rajamanickam S, Karthikeyan G, Kavino M, Manoranjitham SK (2018) Biohardening of micro-propagated banana using endophytic bacteria induces plant growth promotion and restrain the rhizome rot disease caused by *Pectobacterium carotovorum* subsp. *carotovorum*. Sci Hortic 231:179–187
- Sathyanarayana N, Reddy OR, Latha S, Rajak RL (1998) Interception of *Xanthomonas campestris* pv *Dieffenbachiae* on Anthurium plants from the Netherlands. Plant Dis 82(2):262
- Saveetha K, Karthiba L, Raveendran M, Saravanakumar D, Suresh S, Raguchander T, Balasubramanian P, Samiyappan R (2009) Understanding the molecular basis of plant growth promotional effect of *Pseudomonas fluorescens* on rice through protein profiling. Proteome Sci 7:1–8
- Shen S, Jing Y, Kuang T (2003) Proteomics approach to identify wound response related proteins from rice leaf sheath. Proteomics 3:527–535
- Shevchenko A, Wilm M, Vorm O, Mann M (1996) Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels. Anal Chem 68:850–858
- Shrestha BK, Karki HS, Groth DE, Nootjarin JN, Ham JH (2016) Biological control activities of rice-associated *Bacillus* sp. strains against sheath blight and bacterial panicle blight of rice. PLoS ONE 11(1):e0146764
- Srinivasan K, Mathivanan N (2011) Plant growth promoting microbial consortia mediated classical biocontrol of sunflower necrosis virus disease. J Biopest 4(1):65–72
- Suganyadevi M, Nakkeeran S, Renukadevi P (2015) Molecular characterization of plant quarantine pathogen *Xanthomonas axonopodis* pv *dieffenbachiae* causing bacterial blight in anthurium and its management. J Mycol Plant Pathol 45(4):336–341
- Teixeira da Silva JA, Dobranszki J, Winarto B, Zeng S (2015) Anthurium in vitro: a review. Sci Hortic 186:266–298
- Vinodkumar S, Nakkeeran S, Renukadevi P, Malathi VG (2017) Biocontrol potentials of antimicrobial peptide producing *Bacillus* species: multifaceted antagonists for the management of stem rot of carnation caused by *Sclerotinia sclerotiorum*. Front Microbiol 8:1–13
- Weisburg WG, Barns SM, Pelletier DA, Lane DJ (1991) 16S ribosomal DNA amplification for phylogenetic study. J Bacteriol 173(2):697–703
- Suganyadevi M, Renukadevi P, Nakkeeran S (2016) Efficacy of biocontrol agents and bactericides for the management of bacterial blight incited by *Xanthomonas axonopodis* pv. *dieffenbachiae* in *Anthurium andreaeanum*. Int J Plant Protect 9(1):292–296