Published in final edited form as:

Psychopharmacology (Berl). 2020 November ; 237(11): 3383–3397. doi:10.1007/s00213-020-05619-4.

Acute effects of alcohol on error-elicited negative affect during a cognitive control task

Roberto U. Cofresí, Bruce D. Bartholow

Department of Psychological Sciences, University of Missouri

Abstract

Rationale: Alcohol intoxication can dampen negative affective reactions to stressors. Recently, it has been proposed that these acute anxiolytic effects of alcohol may extend to dampening of negative affective reactions to error commission during cognitive control tasks. Nonetheless, empirical verification of this claim is lacking.

Objectives: Test the acute effect of alcohol on negative affective reactions to errors during an effort-demanding cognitive control task.

Methods: Healthy, young adult social drinkers (N=96 [49 women], 21–36 years old) were randomly assigned to consume alcohol (0.80 g/kg; n = 33 [15 female]), active placebo (0.04 g/kg; n = 33 [18 women]), or a non-alcoholic control beverage (n = 30 [16 women]) before completing the Eriksen flanker task. Corrugator supercilii (Corr) activation, a psychophysiological index of negative affect, was tracked across the task. Two neurophysiological reactions to errors, the error-related negativity (ERN) and the error positivity (Pe), were also measured.

Results: Erroneous actions increased Corr activation in the control and (to a lesser extent) placebo groups, but not in the alcohol group. Error-induced Corr activation was coupled to ERN and Pe in the control, but not in the alcohol and placebo groups. Error-induced Corr activation was not coupled to post-error performance adjustments in any group.

Conclusions: The ability of alcohol to dampen error-related negative affect was verified. It was also shown that placebo alone can disrupt coupling of affective and (neuro)cognitive reactions to errors. Although its behavioral relevance remains to be demonstrated, more attention should be paid to the role of affect in action monitoring and cognitive control processes.

Terms of use and reuse: academic research for non-commercial purposes, see here for full terms. <http://www.springer.com/gb/open-access/authors-rights/aam-terms-v1>

Correspondence concerning this article should be addressed to Roberto Cofresí, University of Missouri, Department of Psychological Sciences, Columbia, MO, 65211. cofresir@missouri.edu Phone: 469-323-0181.

Publisher's Disclaimer: This Author Accepted Manuscript is a PDF file of a an unedited peer-reviewed manuscript that has been accepted for publication but has not been copyedited or corrected. The official version of record that is published in the journal is kept up to date and so may therefore differ from this version.

Compliance with Ethical Standards

All procedures were approved by the University of Missouri Institutional Review Board.

Conflicts of Interest

On behalf of all authors, the corresponding author states that there is no conflict of interest.

Keywords

action monitoring; alcohol; anterior cingulate cortex; blame attribution; cognitive control; corrugator supercilii; ERP; errors; negative affect; performance adjustments

Cognitive control refers to a set of cognitive processes that allow individuals to guide their behavior in accordance with internal goals (Alexander & Brown, 2010; Braver, 2012; Gratton et al., 2018). A core component of cognitive control is the ability to adjust behavior in response to varying situational demands (Botvinick et al., 2001). Considerable research points to a neural circuit centered on the dorsal anterior cingulate cortex (ACC) as critical to cognitive control, particularly for signaling when adjustments in control are needed, i.e., following errors (Carter et al., 1998; Davis et al., 2005; Hall et al., 2007; Smith et al., 2019; Van Veen et al., 2001; Wang et al., 2005; Yeung et al., 2004). ACC responses to errors can be observed in the amplitude of the scalp-recorded error-related negativity (ERN), a response-locked event-related potential (ERP) component that is much larger during incorrect versus correct responses (Gehring et al., 1993; Holroyd & Coles, 2002; Van Veen & Carter, 2002) and that has been localized to ACC and neighboring structures in medial prefrontal cortex (Debener et al., 2005; Dehaene et al., 1994; Herrmann et al., 2004).

Impaired cognitive control is one of the acute effects of alcohol (e.g., Casbon et al., 2003; Guillot et al., 2010). Supporting evidence comes in part from studies showing that alcohol reduces or eliminates adjustments in performance that typically occur following errors (Bailey et al., 2014; Bartholow et al., 2012; Ridderinkhof et al., 2002). Additional evidence comes from studies showing alcohol reduces error-related neurophysiological responses (Anderson et al., 2011; Marinkovic et al., 2012), including the ERN (Bailey et al., 2014; Bartholow et al., 2012; Easdon et al., 2005; Nelson et al., 2011; Ridderinkhof et al., 2002).

At least four explanations for alcohol-induced cognitive control impairment have been offered. The first is that impairment reflects alcohol-induced *global* deficits in executive functions (Giancola, 2000; Pihl et al., 2003). The second is that impairment reflects alcohol-induced deficits in visual stimulus processing (Yeung et al., 2007; Yeung & Cohen, 2006). Both of these explanations were challenged by Bailey et al. (2014), who showed no effect of alcohol on conflict adaptation in behavioral performance or neurophysiological correlates of conflict monitoring (reactive control) processes during sequences of correct response trials. Acute effects of alcohol on performance and neurophysiology were observed only after error commission. A third explanation is that impaired cognitive control reflects deficits in error detection/recognition under alcohol as evidenced by reduced ERN (Ridderinkhof et al., 2002; also suggested in Yeung et al., 2007). This explanation was challenged by Bartholow et al. (2012) and Bailey et al. (2014), who replicated the finding that the ERN is reduced under alcohol, but also showed that participants given alcohol were just as accurate in detecting/recognizing their mistakes as participants given control and placebo beverages. A fourth explanation, proposed by Bartholow et al. (2012), is that impaired cognitive control might reflect alcohol-induced reduction of affective reactions to failures of control.

Negative affective reactions to failures of control, such as error commission, are theorized to motivate post-error increases in attention and improvements in performance (K. Aarts et al.,

2013; Dignath et al., 2020; Inzlicht et al., 2015; Proudfit et al., 2013; Saunders et al., 2015). The idea that negative affective reactions may be especially important for cognitive control is rooted in a long history of research showing that the ACC is critically involved in the evaluation of distress and pain (e.g., Ballantine et al., 1967; Rainville et al., 1997; Talbot et al., 1991), that failures of control such as error commission are aversive (Hajcak et al., 2004; Hajcak & Foti, 2008), and that error-elicited activity in the ACC indexed by ERN amplitude covaries with the motivational significance of errors (Gehring et al., 1993; Gehring & Taylor, 2004; Hajcak et al., 2005). Viewed from this perspective, acute effects of alcohol on error processing in cognitive control tasks could reflect the drug's well-known anxiolytic properties.

Acute alcohol has well-documented anxiolytic effects in humans (for review, see: Greeley & Oei, 1999; Sayette, 1999, 2017). Specifically, alcohol dampens subjective negative affective reactions (Bradford et al., 2013; Bujarski & Ray, 2014; Levenson et al., 1980; Ray et al., 2009, 2013; Sher et al., 2007) as well as facial expressions of negative affect (Kushner et al., 1997; Sayette et al., 1992, 2012) in a dose-dependent manner. It similarly dampens negative affective reactions at the level of central nervous system activity (e.g., Curtin, Patrick, Lang, Cacioppo, & Birbaumer, 2001; Franken, Nijs, Muris, & Van Strien, 2007; Gorka, Fitzgerald, King, & Phan, 2013) and autonomic nervous system regulation (e.g., Bradford et al., 2013; Donohue et al., 2007; Sher et al., 1994, 2007; Udo et al., 2009; Vaschillo et al., 2008).

Yet, the degree to which negative affect elicited by erroneous actions is reduced by alcohol remains unclear. Indirect support for this idea comes from studies showing that both alcohol-induced ERN reduction and impairments of post-error performance adjustment can be mediated by alcohol-induced decreases in subjective negative affect (Bartholow et al., 2012), and from studies showing that anxiolytic medication (e.g., lorazepam) also reduces the ERN (De Bruijn et al., 2004). Direct support for this idea, however, requires an approach in which alcohol's effects on negative affective reactions can be precisely time-locked to error commission.

Decades of research point to reactivity in the corrugator supercilii, the facial muscles that furrow the brow, as an excellent measure for this purpose (Cacioppo et al., 1984, 1986, 1988; Cacioppo & Petty, 1981; Larsen et al., 2003; Tan et al., 2012; Tassinari et al., 1989; Vrana, 1993). Corrugator supercilii reactivity, measured via electromyography (EMG) (Fridlund & Cacioppo, 1986), has been advanced as an indicator of automatic (involuntary) expression of negative affect (Dimberg et al., 1998, 2000, 2002; Dimberg & Thunberg, 1998). Corrugator supercilii EMG (cEMG) activity increases during exposure to different kinds of aversive stimuli (e.g., electric shock, loud noise, disgusting odors, upsetting pictures and videoclips, angry and fearful faces, negative emotion words) (Cacioppo et al., 1986; Dimberg et al., 1998, 2000; Dimberg & Thunberg, 1998; Hermann et al., 2000; Larsen et al., 2003; Neumann et al., 2005; Rymarczyk et al., 2011; Sestito et al., 2013). Additionally, aversive stimulus-elicited cEMG activity may reflect neural activity in the ACC because the latter has projections to the brainstem facial nucleus, which innervates the corrugator (Cattaneo & Pavesi, 2014; Shackman et al., 2011). Consequently, error-elicited cEMG activity may reflect error-elicited activity in ACC. In keeping with this idea, response-locked cEMG waveforms show amplified activity shortly after error commission (Elkins-Brown et

al., 2016; Lindström et al., 2013), and this activity appears related to the magnitude of the error positivity (Pe), an ERP component that follows the ERN and has been linked to distinct aspects of error processing, such as conscious error recognition (Overbeek et al., 2005). Like the ERN, the Pe is thought to emanate from the ACC (Falkenstein et al., 2000; Herrmann et al., 2004).

Here, we revisited the experiment reported by Bailey et al. (2014), which aimed to characterize performance (accuracy, response time, confidence) as well as stimulus- and response-locked ERPs (N2, FSW, ERN) during a classic, effort-demanding cognitive control task, the Eriksen flanker task (B. A. Eriksen & Eriksen, 1974), among participants who had consumed an alcohol, placebo, or control beverage. The breath alcohol concentrations and placebo manipulation check for this experiment were also published in the Bailey et al. report. In the present study, we examined never-before-reported cEMG activity elicited by errors and correct responses during the flanker task. We predicted (1) that response-locked cEMG activity would be enhanced on erroneous versus correct response trials; (2) that cEMG activity elicited by errors would be dampened following alcohol consumption; (3) that cEMG activity elicited by errors would correlate with ACC activation, as reflected in the amplitude of the ERN and/or Pe components of the response-locked ERP; and (4) that cEMG activity elicited by errors would predict post-error performance adjustments.

Method

Participants

A total of 96 healthy young adult social drinkers (46% female; $M_{\text{age}} = 23.19$ yrs; $N = 96$) from Columbia, Missouri completed the single-session experimental study. Recruitment strategy and study eligibility criteria were previously reported (Bailey et al., 2014). Characteristics of the final analytic sample for this report ($N = 74$) are presented in Supplemental Table 1.

Materials

Beverage Administration—Details were previously published in Bailey et al. (2014). Briefly, participants were randomly assigned to consume one of three beverages during the experiment: an alcohol beverage (dose = 0.80 g/kg), an active alcohol placebo beverage (dose = 0.04 g/kg), or a control beverage. The alcohol beverage was 5:1, tonic to vodka (50% ABV). The placebo beverage was 5:1, tonic to diluted vodka (9:1 flattened tonic to vodka). Doses were calculated based on estimated total body water and the duration of the drinking period (15 min) using published formulae (Curtin & Fairchild, 2003). The control beverage was tonic. Beverages were divided into three drinks (each consumed over 5 min). After the third drink, participants sat idle for another 5 min to permit complete absorption. Alcohol and placebo groups were told the beverage contained “a moderate amount of alcohol.” The control group was told that the beverage contained no alcohol.

Breath Alcohol Concentration (BrAC)—The concentration of alcohol in exhaled breath was measured using a breathalyzer (Alco-Sensor IV; Intoximeters, Inc., St. Louis, MO, USA). BrAC was never shown to participants.

Cognitive Control Task—As previously reported in detail by Bailey et al. (2014), participants completed an arrows-version of the Eriksen flanker task (B. A. Eriksen & Eriksen, 1974; C. W. Eriksen & Hoffman, 1973; Gratton et al., 1992) adapted from Ridderinkhof et al. (2002). Stimulus arrays were presented for 100 ms, and contained a left- or right-facing central arrow plus two similarly- or oppositely-facing flanker arrows on each side. Participants identified the direction of the central arrow by pressing a left-hand or right-hand button on a ms-accurate button box. Left- and right-hand responses were equally frequent. On correct response-compatible trials, flanker arrows faced the same direction as the central arrow. On correct response-incompatible trials, flankers faced the direction opposite to the target. Correct response-compatible and -incompatible arrays were presented pseudo-randomly and occurred with equal probability. Following a button press on each trial, participants rated their confidence in the correctness of their response (see Hester et al., 2005; Nieuwenhuis et al., 2001; Payne et al., 2005). Three sec later, an inter-trial interval (randomly varying between 1,100 and 1,500 ms) occurred, after which the next trial began. Over the course of seven practice blocks (28 trials/block), participants were titrated to a speed-accuracy balance that produced approximately 10% errors. Participants making fewer errors were instructed to speed up; those making more errors were instructed to slow down. No feedback was given during the subsequent experimental trials (10 blocks total, 80 trials/block).

Electrophysiological Recording—The scalp electroencephalogram (EEG) was recorded continuously throughout the flanker task from 32 tin electrodes embedded in a stretch-lycra cap (ElectroCap, Eaton, OH). Electrode sites were prepared such that measured impedance of the skin was ≤ 5 k Ω . The EEG was referenced to the right mastoid during recording. The surface electromyogram (EMG) was recorded throughout the task from 0.25 cm Ag-AgCl electrodes in a bipolar recording configuration placed about 1 cm apart over the left corrugator supercilii, following EMG recording guidelines (Fridlund & Cacioppo, 1986). The EMG signal was grounded to the middle of the forehead near the hairline. Both EEG and EMG signals were amplified using a Synamps2 amplifier (Compumedics Neuroscan, Charlotte, NC) and sampled at 1000 Hz. The EEG signal was band-pass filtered online (0.05 to 40 Hz). A 20-500 Hz band-pass filter was applied offline to the EMG to attenuate all slow non-muscle potentials (De Luca et al., 2010; Fridlund & Cacioppo, 1986; Van Boxtel, 2001). After recording, the EEG was re-referenced to an average of the two mastoids. Ocular artifacts (blinks) were removed from EEG and EMG using regression-based procedures (Semlitsch et al., 1986).

EEG data were segmented into epochs of -200 to 1200 ms of post-response activity. Epochs containing artefactual deflections (exceeding ± 75 μ V; e.g., due to major muscle movement) were rejected (28.36% of trials/subject). Average voltage from 200 to 100 ms before response onset was subtracted from the rest of the waveform. Inspection of the grand average ERP waveforms indicated a negative-going voltage deflection peaking between 25 and 75 ms post-response that was maximal over fronto-central scalp, corresponding to the ERN and its correct response analog, the correct-related negativity (CRN), as shown in Figure 3A. There was also a later, broader, positive-going deflection peaking between 150 and 450 ms that was maximal over centra-pari eta I scalp, corresponding to the Pe and its

correct response analog, the correct positivity (Pc), as shown in Figure 4A. Consequently, the ERN/CRN in each retained epoch was quantified as the average voltage in the 25 – 75 ms post-response window on frontal and central electrode sides (Fz, F3, F4, FCz, FC3, FC4, Cz, C3, C4). The Pe/Pc was quantified in each retained epoch as the average voltage in the 150 – 450 ms post-response window on centro-parietal and parietal electrode sides (CPz, CP3, CP4, Pz, P3, P4).

cEMG data were segmented into two sets of epochs: 200 ms of pre-stimulus activity and -200 to 1200 ms post-response activity. The root mean square voltage was computed for each 100 ms bin. Epochs containing artefactual deflections (i.e., exceeding ± 3.5 standard deviations from the average activity in the epoch or from the average activity across all epochs for the subject) were rejected (1.98% of trials/subject for pre-stimulus epochs and 1.91% of trials/subject for response-locked epochs). Retained pre-stimulus epochs were averaged together (collapsing the two bins) to produce each subject's overall average pre-trial cEMG (presented in Online Supplemental Information). Retained response-locked cEMG epochs were baseline corrected by subtracting the average root mean square voltage in bin “-2” (which corresponded to the -200 to -100 ms post-response window) from the other bins in the epoch.

Procedure

Figure 1 summarizes the experimental procedure and shows the timing of different within-session events. Procedural details were previously published in Bailey et al. (2014).

Results

Of 96 participants, three (all control group) were excluded due to equipment malfunction (they also were excluded in [Bailey et al., 2014]). Stored data were lost for two participants (1 alcohol group, 1 placebo group). Data from four participants (1 alcohol group, 2 control group, 1 placebo group) were excluded due to unstable baseline cEMG.¹ Data from 11 participants (2 alcohol group, 3 control group, 6 placebo group) were excluded because fewer than six error trials with artifact-free cEMG and ERP data were available for analysis.² Thus, our final analytic sample consisted of 74 participants (21 control group, 28 alcohol group, 25 placebo group). Given that this sample differs substantially from that used by Bailey et al. (2014), it was necessary to re-analyze certain measures critical for understanding the current report. Re-analyses of BrAC, flanker task performance, and ERN data are presented in the main text. Re-analyses of placebo manipulation check and subjective intoxication data are presented in Online Supplemental Information. Readers also are referred to Online Supplemental Information for analyses of pre-trial cEMG and subjective affect data. Briefly, pre-trial cEMG levels were reduced in the alcohol relative

¹As indicated by mean pre-trial cEMG ± 2.5 SD from the grand mean. Such extreme values are most likely due to poor cEMG recording (e.g., poor skin preparation, poor electrode placement, sweating).

²Prior psychometric work on ERP indices of performance monitoring (Olvet & Hajcak, 2009; Rietdijk et al., 2014) indicates that ideally six or more artifact-free error trials are needed to measure error-related neurophysiological signals reliably within individuals. Initial psychometric work on response-locked cEMG has focused on within-person standardized activity, and suggests that ideally 14 or more artifact-free error trials are needed for reliable estimates of error-induced activity (Elkins-Brown et al., 2016, 2017). Since it is unclear that the latter guideline can be applied to non-standardized cEMG (given that standardization distorts the EMG waveform), we applied the “six or more artifact-free error trials” guideline from the performance monitoring ERP psychometrics literature.

placebo and control groups (but note that response-locked cEMG data were baseline-corrected). There were no beverage effects on subjective affect data. Manipulation check and subjective intoxication scores indicated a convincing placebo.

BrAC

Repeated measures ANOVA indicated a significant main effect of the repeated measure (5 levels of “assessment:” # 2-6), $F(4, 104) = 5.207, p < 0.001, \eta^2 = 0.003$. As shown in Figure 2, pairwise comparisons indicated an increase in BrAC from assessment 3 to 4 capturing the final ascent to peak BrAC, $t(27) = 5.340, p < 0.001, d = 2.094$, and successive decreases in BrAC from assessment 4 through 6 capturing initial descent from peak BrAC, $t(27) = 1.995, p = 0.057, d = 0.782$.

Response-locked ERPs

ERN/CRN—ANOVA³ on ERN/CRN mean amplitudes considered the between-subject factor of beverage group (3 levels: alcohol, placebo, control), the within-subject factor of current trial response accuracy (2 levels: correct, incorrect), and their interaction, controlling for the within-subject factor of electrode (9 levels: Fz, F3, F4, FCz, FC3, FC4, Cz, C3, C4). ANOVA detected a significant Beverage group x Response accuracy interaction, $F(2, 1315) = 5.520, p = .004, \eta^2 = 0.016$, evident in Figure 3B. We followed up on this interaction by testing a planned directional prediction based on what was found in the larger sample (Bailey et al., 2014). Specifically, we tested whether the ERN was significantly *less negative* for the alcohol group compared to the control and placebo groups using a 1-sided, independent samples *t*-test. This prediction was confirmed: $t(72) = 2.026, p = 0.023, d = 0.478$. In contrast, the ERN was similar between the control and placebo groups, $t(44) = 0.082, p = 0.247, d = 0.025$.

Pe/Pc—This response-locked ERP component was not considered by Bailey et al. (2014), but we were interested in it as a potential predictor of response-locked cEMG activity and thus, quantified and analyzed it here. ANOVA³ on Pe/Pc mean amplitudes considered the between-subject factor of beverage group (3 levels: alcohol, placebo, control), the within-subject factor of current trial response accuracy (2 levels: correct, incorrect), and their interaction, controlling for the within-subject factor of electrode (6 levels: CPz, CP3, CP4, Pz, P3, P4). Neither the main effect of beverage nor its interaction with response accuracy was significant, $F(2, 876) = 2.180, p = 0.114, \eta^2 = 0.007$, in keeping with Figure 4A. ANOVA detected only a significant main effect of response accuracy, $F(1, 876) = 611.320, p < 0.001, \eta^2 = 0.408$. This within-subject effect was due to a larger (*more positive*) mean amplitude following erroneous compared to correct responses, $t(73) = 12.732, 2\text{-sided } p < 0.001, d = 2.980$.

³Repeated measures ANOVA was used for response-locked ERP analyses (instead of linear mixed modeling [LMM], which was used for response-locked cEMG analyses) because the effect of beverage on ERN/CRN and Pe/Pc mean amplitudes is not of central interest in this report. Furthermore, similar results were obtained when we re-analyzed the ERN/CRN and Pe/Pc mean amplitudes at the trial-level using LMM, following Volpert-Esmond et al. (2018). For simplicity, only ANOVA results are presented.

Response-locked cEMG Activity

Trial-by-trial response-locked corrugator EMG waveforms were analyzed using linear mixed models (LMMs) in R version 3.6.0 using packages lme4 (Bates et al., 2015), lmerTest (Kuznetsova et al., 2017), and emmeans (Lenth, 2019) in order to best account for the nested structure of repeated psychophysiological measurements (E. Aarts et al., 2014; Page-Gould, 2019). Technical details are presented in Online Supplemental Information.

Hypothesis 1.—To test the prediction that response-locked cEMG activity would be enhanced on erroneous versus correct response trials, we considered the simple slopes of bin and bin² as a function of response accuracy. In keeping with our prediction, the simple slope of bin was more positive for erroneous compared to correct response trials, $b \pm SE = 0.028 \pm 0.006 \mu\text{V}/\text{bin}$, $z = 5.022$, $p < .001$, and the simple slope of bin² was more negative for erroneous compared to correct response, $b \pm SE = 0.0019 \pm 0.0005 \mu\text{V}/\text{bin}^2$, $z = 4.146$, $p < .001$.

Hypothesis 2.—To test the prediction that error-elicited cEMG activity would be dampened in the alcohol relative to control and placebo groups, we first considered the simple slopes of bin and bin² as a function of response accuracy within groups. We then considered whether the simple slopes of bin and bin² for erroneous response trials differed among groups. In keeping with our prediction, the simple slope of bin was more positive for erroneous compared to correct response trials in the control group, $b \pm SE = 0.052 \pm 0.011 \mu\text{V}/\text{bin}$, $z = 4.839$, $p < .001$, and in the placebo group, $b \pm SE = 0.029 \pm 0.010 \mu\text{V}/\text{bin}$, $z = 2.910$, $p = .004$, but not in the alcohol group, $b \pm SE = 0.004 \pm 0.008 \mu\text{V}/\text{bin}$, $z = 0.494$, $p = .6211$. Similarly, the simple slope of bin² was more negative for erroneous compared to correct response trials in the control group, $b \pm SE = 0.0037 \pm 0.0009 \mu\text{V}/\text{bin}^2$, $z = 4.101$, $p < .001$, and the placebo group, $b \pm SE = 0.0027 \pm 0.0008 \mu\text{V}/\text{bin}^2$, $z = 3.389$, $p < .001$, but not the alcohol group, $b \pm SE = -0.0006 \pm 0.0007 \mu\text{V}/\text{bin}^2$, $z = 0.863$, $p = .388$. Furthermore, the simple slopes of bin for erroneous response trials were significantly more positive for the control and placebo groups compared to the alcohol group, $b \pm SE = 0.042 \pm 0.013 \mu\text{V}/\text{bin}$, $z = 3.350$, $p < .001$, and the corresponding simple slopes of bin² were significantly more negative, $b \pm SE = -0.004 \pm 0.001 \mu\text{V}/\text{bin}^2$, $z = 3.398$, $p < .001$. The simple slopes of bin and bin² for erroneous response trials were not significantly different between the control and placebo groups, $z = 0.423$, $p = .672$.⁴ Group differences are evident in the average and LMM-estimated response-locked cEMG waveforms (Figure 5A–B).

Hypothesis 3.—To test the prediction that error-elicited cEMG activity would correlate with error-elicited neural activity in the ACC, as reflected in the amplitude of the ERN and/or Pe components of the response-locked ERP, we added new effects to the best LMM of the trial-by-trial cEMG waveform. Since this hypothesis concerns within-trial associations within-persons, we first isolated within-person trial-by-trial changes in ERN and Pe

⁴For completeness, we also considered whether the simple slopes of bin and bin² for correct response trials differed among groups. Ordering of the simple slope of bin was: placebo > alcohol > control, pairwise comparisons: $z = 2.337$, $p = .019$. Ordering of the simple slope of bin² was: control > alcohol ($z = 1.793$, $p = .073$) > placebo, pairwise comparisons: $z = 3.020$, $p = .005$. These patterns are consistent with highly specific cEMG reactivity to errors in the control group, and some specificity loss in the alcohol or placebo group, though most evident in the latter.

following previous work from our laboratory (Von Gunten et al., 2018). Technical details are presented in Online Supplemental Information.

In keeping with our prediction, in the control group, erroneous response-elicited cEMG activity tended to be greater overall when the same-trial ERN was large (i.e., more negative), $b \pm SE = -0.005 \pm 0.002 \mu V \text{ cEMG per } \mu V \text{ ERN}$, $z = -2.965$, $p = .003$. As shown in Figure 6, this association was reversed in the placebo group, $b \pm SE = 0.007 \pm 0.001 \mu V \text{ cEMG per } \mu V \text{ ERN}$, $z = 4.866$, $p < .001$, and nullified in the alcohol group, $b \pm SE = 0.0005 \pm 0.0011 \mu V \text{ cEMG per } \mu V \text{ ERN}$, $z = 0.471$, $p = .638$. Also, in keeping with our prediction, for the control group, the simple slope of bin in erroneous response trials became significantly more positive with increasing (i.e., more positive) Pe, $b \pm SE = 0.523 \pm 0.110 \mu V \text{ cEMG per bin per } \mu V \text{ Pe}$, $z = 4.746$, $p < .001$, and the corresponding simple slope of bin^2 became significantly more negative, $b \pm SE = -0.0585 \pm 0.0092 \mu V \text{ cEMG per bin}^2 \text{ per } \mu V \text{ Pe}$, $z = -6.385$, $p < .001$, effects evident in Figure 7. In contrast, for the placebo group, the simple slopes of bin and bin^2 in erroneous response trials were not significantly affected associated with Pe, $z = 1.174$, $p = .241$. For the alcohol group, the simple slope of bin in erroneous response trials became significantly more negative with increasing (i.e., more positive) Pe, $b \pm SE = -0.305 \pm 0.091 \mu V \text{ cEMG per bin per } \mu V \text{ Pe}$, $z = -3.352$, $p < .001$, and the corresponding simple slope of bin^2 became significantly more positive, $b \pm SE = 0.025 \pm 0.008 \mu V \text{ cEMG per bin}^2 \text{ per } \mu V \text{ Pe}$, $z = 3.332$, $p < .001$.

Hypothesis 4.—To test the prediction that post-error cognitive control adjustments would be informed by affective reactivity to error commission, we first evaluated evidence for post-error reduction of interference (PERI) in the flanker task, a post-error behavioral effect linked to cognitive control processes (Burle et al., 2002; Danielmeier & Ullsperger, 2011; King et al., 2010; Ridderinkhof et al., 2002). The flanker task interference effect manifests as lower accuracy (probability of correct response), and larger (slower) correct response times (RTs), on trials with correct response-incompatible relative to -compatible flanker stimuli. Consequently, we fit LMMs of trial-by-trial accuracy and RT as a function of beverage group, current trial flanker type (correct response-compatible vs. -incompatible) and previous trial response accuracy (correct vs. erroneous response) accounting for time on task. After determining the best LMMs of accuracy and RT, we added new effects corresponding to isolated within-person, trial-by-trial changes in previous trial response-locked cEMG activity, following Von Gunten et al. (2018). Technical details are presented in Online Supplemental Information. As shown in Figure 8, PERI was evident in accuracy, but not in RT, and did not differ by beverage group. Other effects evident in the best LMMs (e.g., beverage effects on the flanker task interference effect, beverage effects on other post-error adjustments) are presented in Online Supplemental Information. Critically, contrary to our prediction, there was no evidence for associations between error-elicited cEMG on one trial and performance (accuracy or RT) on the next trial.

Discussion

The current study examined negative affective reactions to error commission in the flanker task following consumption of an alcohol, placebo, or control beverage by measuring response-locked cEMG activity. Four hypotheses were tested: (1) cEMG activity would be

enhanced following erroneous versus correct responses; (2) error-elicited cEMG activity would be dampened in the alcohol group; (3) error-elicited cEMG activity and error-elicited neural activity in the ACC, as indexed by the ERN and/or Pe components of the response-locked ERP, would be correlated; and (4) error-elicited cEMG activity would be correlated with post-error adjustments in cognitive control task performance.

Hypothesis 1 was supported, replicating previous work on error-elicited cEMG activity (Berger et al., 2020; Dignath et al., 2019; Elkins-Brown et al., 2016, 2017; Lindström et al., 2013). Hypothesis 2 was also supported, providing empirical weight to the idea that alcohol consumption might decrease negative affective reactions to erroneous actions in effort-demanding cognitive control tasks, as first proposed by Bartholow et al. (2012). Although not hypothesized, we also found that pre-trial cEMG activity in the flanker task was selectively dampened in the alcohol group (see Online Supplemental information). Given that response-locked cEMG activity was corrected for pre-response activity, this effect suggests that there may be acute effects of alcohol on both tonic affective state and phasic error reactivity. Previous studies using between-subject designs and similar participants have failed to find acute effects of alcohol on stimulus-locked cEMG in the context of a passive picture-viewing task (Curtin et al., 1998; Glautier et al., 2001; Stritzke et al., 1995), which is consistent with the idea that certain effects of alcohol are restricted to active and self-relevant contexts (for review, see Sayette, 2017).

Hypothesis 3 was supported in the control group, providing *in principle* replication of findings reported by Elkins-Brown et al. (2016) and confirmation of ideas proposed by Lindström et al. (2013). This finding suggests that negative affective reactions to errors, at least as indexed by cEMG activity, are shaped by both early (ERN) and late (Pe) error processing-related neural activity in the ACC. Specifically, under normal circumstances, the magnitude of negative affect elicited by error commission may be set primarily by early ACC activity captured in the ERN, whereas how quickly that negative affective response is emitted relative to error commission and how quickly it dissipates appear to be tuned by later ACC activity captured in the Pe.

Hypothesis 3 was not supported in the alcohol or placebo group. Within-trial associations between ERN or Pe and error-elicited cEMG activity in these groups were inconsistent. ERN failed to predict overall error-elicited cEMG activity in the alcohol group, perhaps in keeping with selective dampening of the ERN in this group. In contrast, Pe predicted a dip below pre-response baseline for error-elicited cEMG activity in the alcohol group, although the Pe did not differ across groups. Nonetheless, error-elicited cEMG activity was largely abolished in the alcohol group. Consequently, it is possible that within-trial ERP-cEMG associations in the alcohol group are statistical artifacts. It is difficult to apply the same logic to explain within-trial ERP-cEMG associations in the placebo group. Erroneous actions still elicited cEMG activity in the placebo group, even if less robustly than in the control group. Yet, in the placebo group, ERN predicted dips below pre-response baseline in overall error-elicited cEMG activity, and Pe failed to predict error-elicited cEMG activity. There were no differences in ERN or Pe between the placebo and control groups.

We are left with one substantive factor to explain differential ERP-cEMG coupling in the control versus placebo groups, namely, activation of alcohol use-outcome expectancies in the latter group. To the extent that affective reactions to error commission are shaped by attributional processes, then the expectation of sub-optimal performance due to alcohol (i.e., attribution to an external cause) could explain why error-elicited cEMG was uncoupled from ERN and Pe in placebo-consuming participants (see Testa et al., 2006). It also could explain dampened error-elicited cEMG in placebo- vs. control beverage-consuming participants despite equivalent ERN and Pe. Consequently, one reason why errors failed to elicit cEMG in alcohol-consuming participants may have been that they too were liable to externalize blame for their mistakes. There is some precedent for such an account (e.g., Critchlow, 1987; Isleib et al., 1988). Future studies could test this idea by manipulating participants' beliefs about the locus of performance deficits prior to task completion (e.g., participants could be convinced that alcohol does *not* diminish performance on the task while covertly titrating task difficulty to promote error commission). Alternatively, participants' biases toward explicitly externalizing vs. internalizing success and failure could be measured (e.g., using false-feedback) while sober vs. intoxicated.

Hypothesis 4 was not supported. Performance data provided no support for the idea that error-elicited negative affect is coupled to post-error adjustments in performance, or our contention that alcohol-induced diminution of error-elicited negative affect has implications for adjustment of cognitive control. In hindsight, we propose two reasons for this apparent failure. First, the experiment was not optimally designed to test questions about the association between affective reactivity to errors and subsequent control adjustment *per se*. Rather, it was designed specifically to test alcohol's effect on the first part of this premise, i.e., that alcohol reduces negative affective reactions to errors, as indicated by ERN and error-elicited cEMG activity. This was our primary interest. The experiment was designed to provide an alternative explanation for alcohol-induced reduction of the ERN, the first demonstration of which was reported by Ridderinkhof et al. (2002), who argued that alcohol reduced the ERN by impairing participants' ability to recognize when they made mistakes. Our alternative hypothesis has been that alcohol *does not* impair the ability to recognize when errors are made—indeed, data from this experiment (as reported in Bailey et al., 2014) and a separate experiment (Bartholow et al., 2012) strongly support this idea—but rather that alcohol reduces the ERN because it mollifies negative affective reactions to control failures—a prediction confirmed by the response-locked cEMG data reported here. The desire to test this alternative hypothesis led us to modify Ridderinkhof et al.'s original paradigm by introducing response accuracy confidence judgments following each trial. This design feature had the unfortunate side-effect of introducing long inter-trial intervals (≈ 5 s) into the task, which make it difficult to test whether error-elicited negative affect on a given trial influences performance on the subsequent trial.

A second possible reason why we failed to find associations between error-elicited cEMG activity and subsequent-trial performance is that participants likely had little motivation to adjust their behavioral performance. There was no penalty for errors or slow responses (and no performance feedback during the experimental trials), and no real benefit for accurate or fast responses. Over the course of a nearly hour-long task, participants likely lost motivation for engaging control in a consistent manner. Thus, although the cEMG and ERN data

support the idea that errors in the task elicited some negative affect, and that alcohol reduced affective reactions to errors, there likely was little motivation on the part of participants to translate affective reactions into control adjustments (e.g., see Boksem et al., 2006).

Findings from the current study should be considered in light of its strengths and weaknesses. The current study improved upon previous work in two important ways. First, it joins two studies (Dignath et al., 2019; Elkins-Brown et al., 2017) in showing that error-elicited cEMG activity can be observed in the absence of explicit feedback about response accuracy as well as in the absence of a threat (risk) of erroneous response-contingent punishment, which has been shown to amplify overall (Curtin et al., 1998) and error-elicited cEMG activity (Lindström et al., 2013). Second, the current study used statistical methods (i.e., LMM) that are arguably more statistically powerful yet conservative, if not at least more appropriate, for modeling psychophysiological data (E. Aarts et al., 2014; Page-Gould, 2019). The use of this methodology allowed us to model the response-locked cEMG waveform, its sensitivity to within- and between-person factors, and change in the waveform across the task.

Despite these strengths, the current study was not without limitations. First, we tested a single alcohol dose in a between-subject design, yet effects of alcohol on affective state and reactivity can vary in extent, nature, and specificity with dose (Donohue et al., 2007; for review, see Sayette, 2017). Second, participants were predominantly Non-Hispanic White young adults. Third, we found no acute effect of alcohol on subjective affect, despite repeated assessment, including during the flanker task (see Online Supplemental Information). However, subjective affect assessment during the task was not time-locked to errors, unlike in (Spunt et al., 2012). Fourth, the probability of errors across the task was low, limiting the number of error trials available for analysis. Accuracy was titrated to be 90% correct, by design, to ensure that beverage effects on cognitive control task-related ERPs were not confounded by major differences in the frequency of correct versus incorrect responses. The relatively low frequency of errors in this study and others like it (Berger et al., 2020; Dignath et al., 2019; Elkins-Brown et al., 2016, 2017; Lindström et al., 2013) imposes psychometric constraints on measured responses to error commission independently of and/or in addition to the constraints imposed by the type of measurement (i.e., subjective, neuro/psycho-physiological). Thus, future studies should parametrically manipulate the frequency of error commission.

Keeping in mind its strengths and weaknesses, the current study has implications for understanding the acute effects of alcohol on cognition. Specifically, its findings suggest that the acute effects of alcohol on canonical cognitive control tasks sometimes may reflect acute effects of alcohol expectancy and pharmacology on the affective underpinnings of action monitoring, and more broadly, cognitive control. Continued work on the affective underpinnings of cognitive control (Dignath et al., 2020; Inzlicht et al., 2015) stands to improve not only our basic understanding of decision-making and self-regulation, but also how the latter might be affected by the acute and/or chronic effects of psychoactive substances, including alcohol.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

BDB designed the project and procured its funding. BDB and RUC formulated the research questions and wrote the manuscript together. RUC processed and analyzed the data and prepared the figures and tables. The authors are grateful to the research coordinators, undergraduate research assistants, doctoral and post-doctoral students, and laboratory managers that helped collect and archive the data for this project while working in the Social Cognition and Addiction Neuroscience Laboratory (SCANlab) over the years. Funding for the project was provided by NIH grant R21 AA017282 (BDB). Preparation of the manuscript was supported by NIH grant R01 AA025451 (BDB). RUC thanks the University of Missouri Department of Psychological Sciences Mission Enhancement Post-Doctoral Fellowship Fund and the local NIH-funded T32 program (AA013526) for enabling his contribution.

The authors have no conflicts of interest to declare. Funding and support for this work was provided by NIH grants AA017282 (BDB), AA025451 (BDB), AA013526 (RUC), and the University of Missouri College of Arts & Science Mission Enhancement Fund (RUC).

References

- Aarts E, Verhage M, & Veenvliet J (2014). A solution to dependency: using multilevel analysis to accommodate nested data. *Nature Neuroscience*, 77(4), 491–496. 10.1038/nn.3648
- Aarts K, De Houwer J, & Pourtois G (2013). Erroneous and correct actions have a different affective valence: Evidence from ERPs. *Emotion*, 13(5), 960–973. 10.1037/a0032808 [PubMed: 23795590]
- Alexander WH, & Brown JW (2010). Computational Models of Performance Monitoring and Cognitive Control. *Topics in Cognitive Science*, 2(4), 658–677. 10.1111/j.1756-8765.2010.01085.x [PubMed: 21359126]
- Anderson BM, Stevens MC, Meda SA, Jordan K, Calhoun VD, & Pearlson GD (2011). Functional Imaging of Cognitive Control During Acute Alcohol Intoxication. *Alcoholism: Clinical and Experimental Research*, 35(1), 156–165. 10.1111/j.1530-0277.2010.01332.x
- Bailey K, Bartholow BD, Saults JS, & Lust SA (2014). Give me just a little more time: Effects of alcohol on the failure and recovery of cognitive control. *Journal of Abnormal Psychology*, 123(1), 152–167. 10.1037/a0035662 [PubMed: 24661167]
- Ballantine HT, Cassidy WL, Flanagan NB, & Marino R (1967). Stereotaxic anterior cingulotomy for neuropsychiatric illness and intractable pain. *Journal of Neurosurgery*, 26(5), 488–495. 10.3171/jns.1967.26.5.0488 [PubMed: 5337782]
- Bartholow BD, Henry EA, Lust SA, Saults JS, & Wood PK (2012). Alcohol effects on performance monitoring and adjustment: affect modulation and impairment of evaluative cognitive control. *Journal of Abnormal Psychology*, 121(1), 173–186. 10.1037/a0023664 [PubMed: 21604824]
- Bates D, Mächler M, Bolker BM, & Walker S (2015). Fitting Linear Mixed-Effects Models Using {lme4}. *Journal of Statistical Software*, 67(1), 1–48. 10.18637/jss.v067.i01
- Berger A, Mitschke V, Dignath D, Eder A, & van Steenbergen H (2020). The face of control: Corrugator supercilii tracks aversive conflict signals in the service of adaptive cognitive control. *Psychophysiology*, 57(1), 1–13. 10.1111/psyp.13524
- Boksem MAS, Meijman TF, & Lorist MM (2006). Mental fatigue, motivation and action monitoring. *Biological Psychology*, 72(2), 123–132. 10.1016/j.biopsycho.2005.08.007 [PubMed: 16288951]
- Botvinick MM, Braver TS, Barch DM, Carter CS, & Cohen JD (2001). Conflict monitoring and cognitive control. *Psychological Review*, 108(3), 624–652. [PubMed: 11488380]
- Bradford DE, Shapiro BL, & Curtin JJ (2013). How bad could it be? Alcohol dampens stress responses to threat of uncertain intensity. *Psychological Science*, 24(12), 2541–2549. 10.1177/0956797613499923 [PubMed: 24145332]
- Braver TS (2012). The variable nature of cognitive control: A dual mechanisms framework. *Trends in Cognitive Sciences*, 16(2), 106–113. 10.1016/j.tics.2011.12.010 [PubMed: 22245618]

- Bujarski S, & Ray LA (2014). Subjective response to alcohol and associated craving in heavy drinkers vs. alcohol dependents: an examination of Koob's allostatic model in humans. *Drug and Alcohol Dependence*, 140, 161–167. 10.1016/j.drugalcdep.2014.04.015 [PubMed: 24837580]
- Burle B, Possamai CA, Vidal F, Bonnet M, & Hasbroucq T (2002). Executive control in the Simon effect: An electromyographic and distributional analysis. *Psychological Research*, 66(4), 324–336. 10.1007/s00426-002-0105-6 [PubMed: 12466929]
- Cacioppo JT, Martzke JS, Petty RE, & Tassinary LG (1988). Specific Forms of Facial EMG Response Index Emotions During an Interview: From Darwin to the Continuous Flow Hypothesis of Affect-Laden Information Processing. *Journal of Personality and Social Psychology*, 54(4), 592–604. 10.1037/0022-3514.54.4.592 [PubMed: 3367281]
- Cacioppo JT, & Petty RE (1981). Electromyograms as measures of extent and affectivity of information processing. *American Psychologist*, 36(5), 441–456. 10.1037/0003-066X.36.5.441 [PubMed: 7247103]
- Cacioppo JT, Petty RE, Losch ME, & Kim HS (1986). Electromyographic Activity Over Facial Muscle Regions Can Differentiate the Valence and Intensity of Affective Reactions. *Journal of Personality and Social Psychology*, 50(2), 260–268. 10.1037/0022-3514.50.2.260 [PubMed: 3701577]
- Cacioppo JT, Petty RE, & Marshall-Goodell B (1984). Electromyographic specificity during simple physical and attitudinal tasks: location and topographical features of integrated EMG responses. *Biological Psychology*, 18, 85–121. [PubMed: 6733194]
- Carter CS, Braver TS, Barch DM, Botvinick MM, Noll D, & Cohen JD (1998). Anterior cingulate cortex, error detection, and the online monitoring of performance. *Science*, 280(5364), 747–749. 10.1126/science.280.5364.747 [PubMed: 9563953]
- Casbon TS, Curtin JJ, Lang AR, & Patrick CJ (2003). Deleterious effects of alcohol intoxication: Diminished cognitive control and its behavioral consequences. *Journal of Abnormal Psychology*, 112(3), 476–487. 10.1037/0021-843X.112.3.476 [PubMed: 12943026]
- Cattaneo L, & Pavesi G (2014). The facial motor system. *Neuroscience and Biobehavioral Reviews*, 38, 135. 10.1016/j.neubiorev.2013.11.002 [PubMed: 24239732]
- Critchlow B (1987). Blaming the booze: the attribution of responsibility for drunken behavior. *Personality and Individual Differences*, 9(3), 451–473.
- Curtin JJ, & Fairchild BA (2003). Alcohol and cognitive control: Implications for regulation of behavior during response conflict. *Journal of Abnormal Psychology*, 112(3), 424–436. 10.1037/0021-843X.112.3.424 [PubMed: 12943021]
- Curtin JJ, Lang AR, Patrick CJ, & Stritzke WGK (1998). Alcohol and fear-potentiated startle: The role of competing cognitive demands in the stress-reducing effects of intoxication. *Journal of Abnormal Psychology*, 107(A), 547–557. 10.1037//0021-843X.107.4.547 [PubMed: 9830242]
- Curtin JJ, Patrick CJ, Lang AR, Cacioppo JT, & Birbaumer N (2001). Alcohol Affects Emotion Through Cognition. *Psychological Science*, 12(6), 527–531.
- Danielmeier C, & Ullsperger M (2011). Post-error adjustments. *Frontiers in Psychology*, 2(SEP), 1–10. 10.3389/fpsyg.2011.00233 [PubMed: 21713130]
- Davis KD, Taylor KS, Hutchinson WD, Dostrovsky JO, McAndrews MP, Richter EO, & Lozano AM (2005). Human Anterior Cingulate Cortex Neurons Encode Cognitive and Emotional Demands. *Journal of Neuroscience*, 25(37), 8402–8406. 10.1523/JNEUROSCI.2315-05.2005 [PubMed: 16162922]
- De Bruijn ERA, Hulstijn W, Verkes RJ, Ruigt GSF, & Sabbe BGC (2004). Drug-induced stimulation and suppression of action monitoring in healthy volunteers. *Psychopharmacology*, 177(1–2), 151–160. 10.1007/s00213-004-1915-6 [PubMed: 15578258]
- De Luca CJ, Donald Gilmore L, Kuznetsov M, & Roy SH (2010). Filtering the surface EMG signal: Movement artifact and baseline noise contamination. *Journal of Biomechanics*, 43(8), 1573–1579. 10.1016/j.jbiomech.2010.01.027 [PubMed: 20206934]
- Debener S, Ullsperger M, Siegel M, Fiehler K, von Cramon DY, & Engel AK (2005). Trial-by-Trial Coupling of Concurrent Electroencephalogram and Functional Magnetic Resonance Imaging Identifies the Dynamics of Performance Monitoring. *Journal of Neuroscience*, 25(50), 11730–11737. 10.1523/jneurosci.3286-05.2005 [PubMed: 16354931]

- Dehaene S, Posner MI, & Tucker DM (1994). Localization of a neural system for error detection and compensation. *Psychological Science*, 5(5), 303–305. 10.1111/j.1467-9280.1994.tb00630.x
- Dignath D, Berger A, Spruit IM, & van Steenbergen H (2019). Temporal dynamics of error-related corrugator supercilii and zygomaticus major activity: Evidence for implicit emotion regulation following errors. *International Journal of Psychophysiology*, 146(October), 101773 10.1016/j.ijpsycho.2019.10.003
- Dignath D, Eder AB, Steinhäuser M, & Kiesel A (2020). Conflict monitoring and the affective-signaling hypothesis—An integrative review. *Psychonomic Bulletin and Review*, 27(2), 193–216. 10.3758/s13423-019-01668-9 [PubMed: 31898269]
- Dimberg U, Hansson G, & Thunberg M (1998). Fear of snakes and facial reactions: A case of rapid emotional responding. *Scandinavian Journal of Psychology*, 39(2), 75–80. 10.1111/1467-9450.00059
- Dimberg U, & Thunberg M (1998). Rapid facial reactions to emotional facial expressions. *Scandinavian Journal of Psychology*, 39(1), 39–45. 10.1111/1467-9450.00054 [PubMed: 9619131]
- Dimberg U, Thunberg M, & Elmehed K (2000). Unconscious facial reactions to emotional facial expressions. *Psychological Science*, 11(1), 86–89. 10.1111/1467-9280.00221 [PubMed: 11228851]
- Dimberg U, Thunberg M, & Grunedal S (2002). Facial reactions to emotional stimuli: Automatically controlled emotional responses. *Cognition and Emotion*, 16(4), 449–471. 10.1080/02699930143000356
- Donohue KF, Curtin JJ, Patrick CJ, & Lang AR (2007). Intoxication level and emotional response. *Emotion*, 7(1), 103–112. 10.1037/1528-3542.7.1.103 [PubMed: 17352567]
- Easdon C, Izenberg A, Armilio ML, Yu H, & Alain C (2005). Alcohol consumption impairs stimulus- and error-related processing during a Go/No-Go Task. *Cognitive Brain Research*, 25(3), 873–883. 10.1016/j.cogbrainres.2005.09.009 [PubMed: 16256319]
- Elkins-Brown N, Saunders B, He F, & Inzlicht M (2017). Stability and reliability of error-related electromyography over the corrugator supercilii with increasing trials. *Psychophysiology*, 54(10), 1559–1573. 10.1111/psyp.12902 [PubMed: 28621433]
- Elkins-Brown N, Saunders B, & Inzlicht M (2016). Error-related electromyographic activity over the corrugator supercilii is associated with neural performance monitoring. *Psychophysiology*, 53(2), 159–170. 10.1111/psyp.12556 [PubMed: 26470645]
- Eriksen BA, & Eriksen CW (1974). Effects of noise letters upon the identification of a target letter in a nonsearch task. *Perception & Psychophysics*, 16(1), 143–149. 10.3758/BF03203267
- Eriksen CW, & Hoffman JE (1973). The extent of processing of noise elements during selective encoding from visual displays. *Perception & Psychophysics*, 14(1), 155–160. 10.3758/BF03198630
- Falkenstein M, Hoormann J, Christ S, & Hohnsbein J (2000). ERP components on reaction errors and their functional significance: A tutorial. *Biological Psychology*, 51(2–3), 87–107. 10.1016/S0301-0511(99)00031-9 [PubMed: 10686361]
- Franken IHA, Nijs IMT, Muris P, & Van Strien JW (2007). Alcohol selectively reduces brain activity during the affective processing of negative information. *Alcoholism: Clinical and Experimental Research*, 31(6), 919–927. 10.1111/j.1530-0277.2007.00424.x
- Fridlund AJ, & Cacioppo JT (1986). Guidelines of Human Electromyographic Research. *Psychophysiology*, 23(5), 567–589. [PubMed: 3809364]
- Gehring WJ, Goss B, Coles MGH, Meyer DE, & Donchin E (1993). A Neural System for Error Detection and Compensation. *Psychological Science*, 4(6), 385–390.
- Gehring WJ, & Taylor SF (2004). When the going gets tough, the cingulate gets going. *Nature Neuroscience*, 7(12), 1285–1287. 10.1038/nn1204-1285 [PubMed: 15643434]
- Giancola PR (2000). Executive functioning: A conceptual framework for alcohol-related aggression. *Experimental and Clinical Psychopharmacology*, 8(4), 576–597. 10.1037/1064-1297.8.4.576 [PubMed: 11127429]
- Glaetzer S, O'Brien J, & Dixon J (2001). Facial electromyographic (EMG) responses to emotionally significant visual images: Differences between light and heavy drinkers. *Drug and Alcohol Dependence*, 64(3), 337–345. 10.1016/S0376-8716(01)00136-3 [PubMed: 11672948]

- Gorka SM, Fitzgerald DA, King AC, & Phan KL (2013). Alcohol attenuates amygdala-frontal connectivity during processing social signals in heavy social drinkers: A preliminary pharmacofMRI study. *Psychopharmacology*, 229(1), 141–154. 10.1007/s00213-013-3090-0 [PubMed: 23584670]
- Gratton G, Coles MGH, & Donchin E (1992). Optimizing the Use of Information: Strategic Control of Activation of Responses. *Journal of Experimental Psychology: General*, 121(4), 480–506. 10.1063/1.1379753 [PubMed: 1431740]
- Gratton G, Cooper P, Fabiani M, Carter CS, & Karayanidis F (2018). Dynamics of cognitive control: Theoretical bases, paradigms, and a view for the future. *Psychophysiology*, 55(3), 1–29. 10.1111/psyp.13016
- Greeley J, & Oei T (1999). Alcohol and tension reduction In Leonard KE & Blane HT (Eds.), *Psychological Theories of Drinking and Alcoholism* (pp. 14–53). Guilford Press.
- Guillot CR, Fanning JR, Bullock JS, McCloskey MS, & Berman ME (2010). Effects of Alcohol on Tests of Executive Functioning in Men and Women: A Dose Response Examination. *Experimental and Clinical Psychopharmacology*, 18(5), 409–417. 10.1037/a0021053 [PubMed: 20939644]
- Hajcak G, & Foti D (2008). Errors are aversive: Defensive motivation and the error-related negativity. *Psychological Science*, 19(2), 103–108. 10.1111/j.1467-9280.2008.02053.x [PubMed: 18271855]
- Hajcak G, McDonald N, & Simons RF (2004). Error-related psychophysiology and negative affect. *Brain and Cognition*, 56(2 SPEC. ISS), 189–197. 10.1016/j.bandc.2003.11.001 [PubMed: 15518935]
- Hajcak G, Moser JS, Yeung N, & Simons RF (2005). On the ERN and the significance of errors. *Psychophysiology*, 42(2), 151–160. 10.1111/j.1469-8986.2005.00270.x [PubMed: 15787852]
- Hall JR, Bernat EM, & Patrick CJ (2007). Externalizing psychopathology and the error-related negativity. *Psychological Science*, 18(4), 326–333. 10.1111/j.1467-9280.2007.01899.x [PubMed: 17470258]
- Hermann C, Ziegler S, Birbaumer N, & Flor H (2000). Pavlovian aversive and appetitive odor conditioning in humans: subjective, peripheral, and electrocortical changes. *Experimental Brain Research*, 132(2), 203–215. 10.1007/s002210000343 [PubMed: 10853945]
- Herrmann MJ, Römmler J, Ehlis AC, Heidrich A, & Fallgatter AJ (2004). Source localization (LORETA) of the error-related-negativity (ERN/Ne) and positivity (Pe). *Cognitive Brain Research*, 20(2), 294–299. 10.1016/j.cogbrainres.2004.02.013 [PubMed: 15183400]
- Hester R, Foxe JJ, Molholm S, Shpaner M, & Garavan H (2005). Neural mechanisms involved in error processing: A comparison of errors made with and without awareness. *Neuroimage*, 27(3), 602–608. 10.1016/j.neuroimage.2005.04.035 [PubMed: 16024258]
- Holroyd CB, & Coles MGH (2002). The neural basis of human error processing: Reinforcement learning, dopamine, and the error-related negativity. *Psychological Review*, 109(4), 679–709. 10.1037/0033-295X.109.4.679 [PubMed: 12374324]
- Inzlicht M, Bartholow BD, & Hirsh JB (2015). Emotional foundations of cognitive control. *Trends in Cognitive Sciences*, 19(3), 126–132. 10.1016/j.tics.2015.01.004 [PubMed: 25659515]
- Isleib RA, Vuchinich RE, & Tucker JA (1988). Performance Attributions and Changes in Self-Esteem Following Self-Handicapping with Alcohol Consumption. *Journal of Social and Clinical Psychology*, 6(1), 88–103. 10.1521/jscp.1988.6.1.88
- King JA, Korb FM, Von Cramon DY, & Ullsperger M (2010). Post-error behavioral adjustments are facilitated by activation and suppression of task-relevant and task-irrelevant information processing. *Journal of Neuroscience*, 30(38), 12759–12769. 10.1523/JNEUROSCI.3274-10.2010 [PubMed: 20861380]
- Kushner MG, Massie ED, Gaskel S, Mackenzie T, Fiszdon J, & Anderson N (1997). Alcohol effects on the facial expressions of anxiety patients undergoing a panic provocation. *Addictive Behaviors*, 22(2), 275–280. 10.1016/S0306-4603(96)00009-3 [PubMed: 9113222]
- Kuznetsova A, Brockhoff PB, & Christensen RHB (2017). {lmerTest} Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, 82(13), 1–26. 10.18637/jss.v082.i13
- Larsen JT, Norris CJ, & Cacioppo JT (2003). Effects of positive and negative affect on electromyographic activity over zygomaticus major and corrugator supercilii. *Psychophysiology*, 40(5), 776–785. 10.1111/1469-8986.00078 [PubMed: 14696731]

- Lenth R (2019). emmeans: Estimated Marginal Means, aka Least-Squares Means. <https://cran.r-project.org/package=emmeans>
- Levenson RW, Sher KJ, Grossman LM, Newman J, & Newlin DB (1980). Alcohol and stress response dampening: Pharmacological effects, expectancy, and tension reduction. *Journal of Abnormal Psychology*, 89(4), 528–538. 10.1037/0021-843X.89.4.528 [PubMed: 7400453]
- Lindström BR, Mattsson-Mårn IB, Golkar A, & Olsson A (2013). In Your Face: Risk of Punishment Enhances Cognitive Control and Error-Related Activity in the Corrugator Supercilii Muscle. *PLoS ONE*, 8(6). 10.1371/journal.pone.0065692
- Marinkovic K, Rickenbacher E, Azma S, & Artsy E (2012). Acute alcohol intoxication impairs top-down regulation of stroop incongruity as revealed by blood oxygen level-dependent functional magnetic resonance imaging. *Human Brain Mapping*, 33(2), 319–333. 10.1002/hbm.21213 [PubMed: 21391268]
- Nelson LD, Patrick CJ, Collins P, Lang AR, & Bernat EM (2011). Alcohol impairs brain reactivity to explicit loss feedback. *Psychopharmacology*, 218(2), 419–428. 10.1007/s00213-011-2323-3 [PubMed: 21559803]
- Neumann R, Hess M, Schulz SM, & Alpers GW (2005). Automatic behavioural responses to valence: Evidence that facial action is facilitated by evaluative processing. *Cognition and Emotion*, 19(4), 499–513.
- Nieuwenhuis S, Richard Ridderinkhof K, Blom J, Band GPH, & Kok A (2001). Error-related brain potentials are differentially related to awareness of response errors: Evidence from an antisaccade task. *Psychophysiology*, 38(5), 752–760. 10.1017/S0048577201001111 [PubMed: 11577898]
- Olvet DM, & Hajcak G (2009). The stability of error-related brain activity with increasing trials. *Psychophysiology*, 46(5), 957–961. 10.1111/j.1469-8986.2009.00848.x [PubMed: 19558398]
- Overbeek TJM, Nieuwenhuis S, & Ridderinkhof KR (2005). Dissociable components of error processing: On the functional significance of the Pe vis-à-vis the ERN/Ne. *Journal of Psychophysiology*, 19(4), 319–329. 10.1027/0269-8803.19.4.319
- Page-Gould E (2019). Multilevel Modeling In Cacioppo JT, Tassinary LG, & Berntson GG (Eds.), *Handbook of Psychophysiology* (4th ed.). Cambridge University Press.
- Payne BK, Shimizu Y, & Jacoby LL (2005). Mental control and visual illusions: Toward explaining race-biased weapon misidentifications. *Journal of Experimental Social Psychology*, 41(1), 36–47. 10.1016/j.jesp.2004.05.001
- Pihl RO, Paylan SS, Gentes-Hawn A, & Hoaken PNS (2003). Alcohol affects executive cognitive functioning differentially on the ascending versus descending limb of the blood alcohol concentration curve. *Alcoholism: Clinical and Experimental Research*, 27(5), 773–779. 10.1097/01.ALC.0000065434.92204.A1
- Proudfit GH, Inzlicht M, & Mennin DS (2013). Anxiety and error monitoring: The importance of motivation and emotion. *Frontiers in Human Neuroscience*, 7(OCT), 8–11. 10.3389/fnhum.2013.00636 [PubMed: 23386818]
- Rainville P, Duncan GH, Price DD, Carrier B, & Bushnell MC (1997). Pain Affect Encoded in Human Anterior Cingulate But Not Somatosensory Cortex. *Science*, 277(5328), 968–971. 10.1126/science.277.5328.968 [PubMed: 9252330]
- Ray LA, Bujarski S, MacKillop J, Courtney KE, Monti PM, & Miotto K (2013). Subjective response to alcohol among alcohol-dependent individuals: effects of the μ -opioid receptor (OPRM1) gene and alcoholism severity. *Alcoholism: Clinical and Experimental Research*, 37 Suppl 1(1), E116–24. 10.1111/j.1530-0277.2012.01916.x
- Ray LA, MacKillop J, Leventhal AM, & Hutchison KE (2009). Catching the Alcohol Buzz: An Examination of the Latent Factor Structure of Subjective Intoxication. *Alcoholism: Clinical and Experimental Research*, 33(12), 2154–2161. 10.1111/j.1530-0277.2009.01053.x
- Ridderinkhof KR, De Vlugt Y, Bramlage A, Spaan M, Elton M, Snel J, & Band GPH (2002). Alcohol consumption impairs detection of performance errors in mediofrontal cortex. *Science*, 298(5601), 2209–2211. 10.1126/science.1076929 [PubMed: 12424384]
- Rietdijk WJR, Franken IHA, & Thurik AR (2014). Internal consistency of event-related potentials associated with cognitive control: N2/P3 and ERN/Pe. *PLoS ONE*, 9(7), 3–9. 10.1371/journal.pone.0102672

- Rymarczyk K, Biele C, Grabowska A, & Majczynski H (2011). EMG activity in response to static and dynamic facial expressions. *International Journal of Psychophysiology*, 79(2), 330–333. 10.1016/j.ijpsycho.2010.11.001 [PubMed: 21074582]
- Saunders B, Milyavskaya M, & Inzlicht M (2015). What does cognitive control feel like? Effective and ineffective cognitive control is associated with divergent phenomenology. *Psychophysiology*, 52(9), 1205–1217. 10.1111/psyp.12454 [PubMed: 26041054]
- Sayette MA (1999). Does Drinking Reduce Stress? *Alcohol Research and Health*, 23(4), 250–255. [PubMed: 10890821]
- Sayette MA (2017). The effects of alcohol on emotion in social drinkers. *Behaviour Research and Therapy*, 88, 76–89. 10.1016/j.brat.2016.06.005 [PubMed: 28110679]
- Sayette MA, Creswell KG, Dimoff JD, Fairbairn CE, Cohn JF, Heckman BW, Kirchner TR, Levine JM, & Moreland RL (2012). Alcohol and Group Formation: A Multimodal Investigation of the Effects of Alcohol on Emotion and Social Bonding. *Psychological Science*, 23(8), 869–878. 10.1177/0956797611435134 [PubMed: 22760882]
- Sayette MA, Smith DW, Breiner MJ, & Wilson GT (1992). The effect of alcohol on emotional response to a social stressor. *Journal of Studies on Alcohol*, 53(6), 541–545. 10.15288/jsa.1992.53.541 [PubMed: 1434629]
- Semlitsch HV, Anderer P, Schuster P, & Presslich O (1986). A Solution for Reliable and Valid Reduction of Ocular Artifacts, Applied to the P300 ERP. *Psychophysiology*, 23(6), 695–703. 10.1111/j.1469-8986.1986.tb00696.x [PubMed: 3823345]
- Sestito M, Umiltà MA, De Paola G, Fortunati R, Raballo A, Leuci E, Maffei S, Tonna M, Amore M, Maggini C, & Gallese V (2013). Facial reactions in response to dynamic emotional stimuli in different modalities in patients suffering from schizophrenia: a behavioral and EMG study. *Frontiers in Human Neuroscience*, 7(July 2013). 10.3389/fnhum.2013.00368
- Shackman AJ, Salomons TV, Slagter HA, Fox AS, Winter JJ, & Davidson RJ (2011). The integration of negative affect, pain and cognitive control in the cingulate cortex. *Nature Reviews Neuroscience*, 12(3), 154–167. 10.1038/nrn2994 [PubMed: 21331082]
- Sher KJ, Bartholow BD, Peuser K, Erickson DJ, & Wood MD (2007). Stress-response-dampening effects of alcohol: Attention as a mediator and moderator. *Journal of Abnormal Psychology*, 116(2), 362–377. 10.1037/0021-843X.116.2.362 [PubMed: 17516768]
- Sher KJ, Bylund DB, Walitzer KS, Hartmann J, & Ray-Prenger C (1994). Platelet Monoamine Oxidase (MAO) Activity: Personality, Substance Use, and the Stress-Response-Dampening Effect of Alcohol. *Experimental and Clinical Psychopharmacology*, 2(1), 53–81. 10.1037/1064-1297.2.1.53
- Smith EH, Horga G, Yates MJ, Mikell CB, Banks GP, Pathak YJ, Schevon CA, McKhann GM, Hayden BY, Botvinick MM, & Sheth SA (2019). Widespread temporal coding of cognitive control in the human prefrontal cortex. *Nature Neuroscience*, 22(November), 10.1038/s41593-019-0494-0
- Spunt RP, Lieberman MD, Cohen JR, & Eisenberger NI (2012). The phenomenology of error processing: The dorsal ACC response to stop-signal errors tracks reports of negative affect. *Journal of Cognitive Neuroscience*, 24(8), 1753–1765. 10.1162/jocn_a_00242 [PubMed: 22571460]
- Stritzke W GK, Patrick CJ, & Lang AR (1995). Alcohol and Human Emotion: A Multidimensional Analysis Incorporating Startle-Probe Methodology. *Journal of Abnormal Psychology*, 104(1), 114–122. 10.1037/0021-843X.104.1.114 [PubMed: 7897033]
- Talbot JD, Marrett S, Evans AC, Meyer E, Catherine M, Talbot JD, Marretr S, Evans AC, Meyer E, Bushnell MC, & Duncan GH (1991). Multiple Representations of Pain in Human Cerebral Cortex. *Science*, 251 (4999), 1355–1358. <http://www.jstor.org/stable/2875534> [PubMed: 2003220]
- Tan J-W, Walter S, Scheck A, Hrabal D, Hoffmann H, Kessler H, & Traue HC (2012). Repeatability of facial electromyography (EMG) activity over corrugator supercilii and zygomaticus major on differentiating various emotions. *Journal of Ambient Intelligence and Humanized Computing*, 3(1), 3–10. 10.1007/s12652-011-0084-9

- Tassinary LG, Cacioppo JT, & Geen TR (1989). A Psychometric Study of Surface Electrode Placements for Facial Electromyographic Recording: I. The Brow and Cheek Muscle Regions. *Psychophysiology*, 26(1), 1–16. 10.1111/j.1469-8986.1989.tb03125.x [PubMed: 2922450]
- Testa M, Fillmore MT, Norris J, Abbey A, Curtin JJ, Leonard KE, Mariano KA, Thomas MC, Nomensen KJ, George WH, VanZile-Tamsen C, Livingston JA, Saenz C, Buck PO, Zawacki T, Parkhill MR, Jacques AJ, & Hayman LW (2006). Understanding alcohol expectancy effects: Revisiting the placebo condition. *Alcoholism: Clinical and Experimental Research*, 30(2), 339–348. 10.1111/j.1530-0277.2006.00039.x
- Udo T, Bates ME, Mun E-Y, Vaschillo EG, Vaschillo B, Lehrer P, & Ray S (2009). Gender differences in acute alcohol effects on self-regulation of arousal in response to emotional and alcohol-related picture cues. *Psychology of Addictive Behaviors: Journal of the Society of Psychologists in Addictive Behaviors*, 23(2), 196–204. 10.1037/a0015015 [PubMed: 19586136]
- Van Boxtel A (2001). Optimal signal bandwidth for the recording of surface EMG activity of facial, jaw, oral, and neck muscles. *Psychophysiology*, 38(1), 22–34. 10.1017/S004857720199016X [PubMed: 11321618]
- Van Veen V, & Carter CS (2002). The timing of action monitoring processes in rostral and caudal anterior cingulate cortex. *Journal of Cognitive Neuroscience*, 14(4), 593–602. 10.1016/j.jeeh.2005.10.006 [PubMed: 12126500]
- Van Veen V, Cohen JD, Botvinick MM, Stenger VA, & Carter CS (2001). Anterior cingulate cortex, conflict monitoring, and levels of processing. *Neuroimage*, 14(6), 1302–1308. 10.1006/nimg.2001.0923 [PubMed: 11707086]
- Vaschillo EG, Bates ME, Vaschillo B, Lehrer P, Udo T, Mun E-Y, & Ray S (2008). Heart rate variability response to alcohol, placebo, and emotional picture cue challenges: Effects of 0.1-Hz stimulation. *Psychophysiology*, 45(5), 847–858. 10.1111/j.1469-8986.2008.00673.x [PubMed: 18513359]
- Volpert-Esmond HI, Merkle EC, Levens MP, Ito TA, & Bartholow BD (2018). Using trial-level data and multilevel modeling to investigate within-task change in event-related potentials. *Psychophysiology*, 55(5), 1–12. 10.1111/psyp.13044
- Von Gunten CD, Volpert-Esmond HI, & Bartholow BD (2018). Temporal dynamics of reactive cognitive control as revealed by event-related brain potentials. *Psychophysiology*, 55(3), e13007 10.1111/psyp.13007
- Vrana SR (1993). The psychophysiology of disgust: Differentiating negative emotional contexts with facial EMG. *Psychophysiology*, 30(3), 279–286. 10.1111/j.1469-8986.1993.tb03354.x [PubMed: 8497557]
- Wang C, Ulbert I, Schomer DL, Marinkovic K, & Halgren E (2005). Responses of Human Anterior Cingulate Cortex Microdomains to Error Detection, Conflict Monitoring, Stimulus-Response Mapping, Familiarity, and Orienting. *Journal of Neuroscience*, 25(3), 604–613. 10.1523/jneurosci.4151-04.2005 [PubMed: 15659596]
- Yeung N, Botvinick MM, & Cohen JD (2004). The neural basis of error detection: Conflict monitoring and the error-related negativity. *Psychological Review*, 111(4), 931–959. 10.1037/0033-295X.111.4.931 [PubMed: 15482068]
- Yeung N, & Cohen JD (2006). The Impact of Cognitive Deficits on Conflict Monitoring. *Psychological Science*, 17(2), 164–171. [PubMed: 16466425]
- Yeung N, Ralph J, & Nieuwenhuis S (2007). Drink alcohol and dim the lights: The impact of cognitive deficits on. *Cognitive, Affective, & Behavioral Neuroscience*, 7(4), 347–355.

Figure 1. Timeline of within-session events.

Numbers below the line indicate planned timing. **Arrival:** Participants provided informed consent, indicated compliance with pre-study protocols, and were randomly assigned to a beverage condition (alcohol, placebo, control). Participants then completed a questionnaire battery unrelated to the current report. **Dashed arrows:** At several times across the session, all participants completed brief questionnaires to assess subjective affect (see Online Supplemental Information). At these times, BrAC also was measured in alcohol and placebo group participants. At arrival, however, BrAC was measured in all participants to confirm sobriety (.000 g%). **Preparation (Prep.):** Experimenters escorted the participant to a recording chamber, measured height and weight, and then placed and tested electrodes for electrophysiological recording. **Practice:** Participants completed 5 blocks of practice trials for the cognitive control task before beverage administration, and 2 blocks of practice trials afterward. **Beverage Administration (Bev.):** Experimenters prepared beverages in front of participants, and participants then consumed the beverage (three drinks, 5 min/drink). **Task:** Participants completed 10 blocks of experimental trials of the cognitive control task with breaks occurring after blocks 3 and 7. **Post:** Experimenters detached electrodes and escorted participants to the restroom where the latter could wash recording gel from their face and hair. All groups then completed a post-experiment questionnaire. For the alcohol and placebo groups, the latter contained a placebo manipulation check (see Online Supplemental Information). All participants were then debriefed. **Exit:** Participants in the control and placebo groups were dismissed. Participants in the alcohol group were retained until BrAC .02 g%.

Figure 2. Breath Alcohol Concentration (BrAC) Across Post- Drinking Assessments.

Sample *M* and *SEM* are shown ($n = 28$). BrAC at assessment 1, which took place before drinking, was 0.000 g%. Assessment 2 took place 5 min after drinking and before starting the cognitive control task experimental trials. Assessments 3, 4, and 5 took place after cognitive control task experimental trial blocks 3, 7, and 10, respectively. Assessment 6 took place after electrodes were detached but before debriefing. Asterisk indicates $p < 0.05$.

Figure 3. ERN/CRN as a Function of Beverage Group and Response Accuracy.

A: Waveforms shown represent the average across frontal, fronto-central, and central electrodes elicited by correct (“Corr”) and erroneous (“Err”) responses. “R” on the x-axis denotes time of button press. Note that the y-axis is reversed flowing convention for ERPs. Window (25–75 ms) for component mean amplitude quantification denoted by the yellow rectangle. **B:** Mean amplitudes shown represent average across frontal, fronto-central, and central electrodes. Note that the y-axis is reversed because ERN/CRN are negative-going ERP components. Asterisk indicates $p < 0.05$. **A+B:** Sample $M \pm SEM$ shown for the no-alcohol control group ($n = 21$), placebo alcohol group ($n = 25$), and alcohol group ($n = 28$).

Figure 4. Pe/Pc as a Function of Beverage Group and Response Accuracy.

A: Waveforms shown represent the average across centro-parietal and parietal electrodes elicited by correct (“Corr”) and erroneous (“Err”) responses. “R” on the x-axis denotes time of button press. Note that the y- axis is reversed following convention for ERPs. Window (150–450 ms) or component mean amplitude quantification denoted by the yellow rectangle.

B: Mean amplitudes shown represent average across centro-parietal and parietal electrodes.

A+B: Sample $M \pm SEM$ shown for the no-alcohol control group ($n = 21$), placebo alcohol group ($n = 25$), and alcohol group ($n = 28$).

Figure 5. Response-Locked cEMG Activity as a Function of Beverage Group and Response Accuracy.

A: Sample M and SEM are shown. **B:** Estimated marginal population M and SE from the best conditional LMM are shown. **A-B:** Data represent $n = 21$ participants who consumed the no- alcohol control beverage, $n = 25$ who consumed the placebo alcohol beverage, and $n = 28$ who consumed the alcohol beverage. On the x-axes, “R” indicates button press. Y-axes differ between A-B. Some error bars are hidden underneath the point symbols.

Figure 6. Overall Response-Locked cEMG Activity as a Function of Beverage Group, Response Accuracy, and ERN/CRN.

Marginal population M and SE estimated from best conditional LMM are shown. Data represent $n = 21$ participants who consumed the no-alcohol control beverage, $n = 25$ who consumed the placebo alcohol beverage, and $n = 28$ who consumed the alcohol beverage.

Figure 7. Response-Locked cEMG Activity as a Function of Beverage Group, Response Accuracy, and Pe/Pc.

Marginal population M and SE estimated from best conditional LMM are shown. Data represent $n = 21$ participants who consumed the no-alcohol control beverage, $n = 25$ who consumed the placebo alcohol beverage, and $n = 28$ who consumed the alcohol beverage.

Figure 8. Correct Response Probability and Response Time (RT) as a Function of Current Trial Flanker Type and Previous Trial Response Accuracy.

A: Back-transformed marginal population M and SE estimated from best conditional generalized LMM (binomial, logit) of accuracy are shown. Post-error reduction of interference (PERI) was detected: difference in predicted probability of correct response when the current trial contains correct response-compatible vs. - incompatible flankers was significantly diminished when previous trial response was incorrect relative to correct. PERI effect was driven by a significant increase in the probability of correct response on correct response-incompatible trials after an incorrect relative to correct response. **B:** Marginal

population M and SE estimated from best conditional LMM of RT are shown. PERI was not detected: difference in RT due to flanker type was similar post-error vs. -correct trial. **A-B:** Data represent $n = 21$ participants who consumed the no-alcohol control beverage, $n = 25$ who consumed the placebo alcohol beverage, and $n = 28$ who consumed the alcohol beverage. Nevertheless, LMMs found no support for Beverage x Current Trial Type x Previous Trial Response Accuracy effects (these LMM-estimated marginal M s and SE s and corresponding sample M s and SEM s are presented in Supplemental Figure 5). Consequently, only the Current Trial Type x Previous Trial Response Accuracy effects are shown. Note that the Current Trial Type x Previous Trial Response Accuracy effect was not significant for RT, and was ultimately dropped from the best LMM of RT used to test effects of previous trial cEMG. Note also that all LMM-estimated M s and SE s shown control for other effects evident in the LMMs (e.g., Beverage x Current Trial Type, Beverage x Previous Trial Response Accuracy). Asterisk indicates $p < 0.05$.