Published in final edited form as:

Psychopharmacology (Berl). 2020 November ; 237(11): 3337–3355. doi:10.1007/s00213-020-05615-8.

Noradrenergic circuits in the forebrain control affective responses to novelty

Daniel Lustberg¹, Rachel P. Tillage¹, Yu Bai¹, Molly Pruitt², L. Cameron Liles¹, David Weinshenker^{1,*}

¹Emory University, Department of Human Genetics, Atlanta, GA 30322

²University of Maryland School of Medicine, Baltimore, MD 21201

Abstract

Rationale—In rodents, exposure to novel environments elicits initial anxiety-like behavior (neophobia) followed by intense exploration (neophilia) that gradually subsides as the environment becomes familiar. Thus, innate novelty-induced behaviors are useful indices of anxiety and motivation in animal models of psychiatric disease. Noradrenergic neurons are activated by novelty and implicated in exploratory and anxiety-like responses, but the role of norepinephrine (NE) in neophobia has not been clearly delineated.

Objective—We sought to define the role of central NE transmission in neophilic and neophobic behaviors.

Methods—We assessed dopamine β -hydroxylase knockout (*Dbh* $-/-$) mice lacking NE and their NE-competent (*Dbh* $+/-$) littermate controls in neophilic (novelty-induced locomotion; NIL) and neophobic (novelty-suppressed feeding; NSF) behavioral tests with subsequent quantification of brain-wide c-fos induction. We complimented the gene knockout approach with pharmacological interventions.

Results—*Dbh* $-/-$ mice exhibited blunted locomotor responses in the NIL task and completely lacked neophobia in the NSF test. Neophobia was rescued in *Dbh* $-/-$ mice by acute pharmacological restoration of central NE with the synthetic precursor L-3,4-dihydroxyphenylserine (DOPS), and attenuated in control mice by the inhibitory α 2-adrenergic autoreceptor agonist guanfacine. Following either NSF or NIL, *Dbh* $-/-$ mice demonstrated reduced c-fos in the anterior cingulate cortex, medial septum, ventral hippocampus, bed nucleus of the stria terminalis, and basolateral amygdala.

Conclusion—These findings indicate that central NE signaling is required for the expression of both neophilic and neophobic behaviors. Further, we describe a putative noradrenergic novelty network as a potential therapeutic target for treating anxiety and substance abuse disorders.

* Address correspondence to: David Weinshenker, Ph.D., Department of Human Genetics, Emory University School of Medicine, 615 Michael St., Whitehead 301, Atlanta, GA 30322, dweinsh@emory.edu.

Conflicts of Interest: The authors declare no conflicts of interest.

Keywords

norepinephrine; dopamine β -hydroxylase; neophobia; neophilia; novel environment; novelty suppressed feeding; guanfacine; c-fos; locus coeruleus; anxiety

Introduction

A novel environment may represent either an appetitive opportunity or a potential threat; the valence of the context is uncertain because it has never been encountered before (Kafkas and Montaldi 2018). In novel environments, it is adaptive for animals to exhibit initial anxiety-like behavior (neophobia) that gradually transitions to exploratory behavior (neophilia) once the environment has proven to be non-threatening (Barnett 1958; Dulawa and Hen 2005; Osorio-Gómez et al. 2018). Novelty-seeking and novelty-fearing personality traits correlate with substance abuse and anxiety disorders, respectively (Bardo et al. 1996; Dovey et al. 2008; Marcontell et al. 2003; Tulving et al. 1994; Weierich et al. 2010). In animal models of psychiatric disease, behavioral responses to novel environments are also strong predictors of drug-seeking and anxiety-like behavior (Dulawa 2009; Pawlak et al. 2008; Walker et al. 2009; Wingo et al. 2016).

Neophilic behavior can be assessed in mice with the novelty-induced locomotion (NIL) test, which measures exploration in a novel environment (Kabbaj et al. 2000; Walker et al. 2009). In the NIL test, high locomotor reactivity to novelty is interpreted as increased exploratory behavior, while low locomotor responses are interpreted as decreased exploratory behavior (Cubells et al. 2016; Schroeder et al. 2013; Stone et al. 1999). Neophobia can be assessed in mice with the novelty-suppressed feeding (NSF) test, a conflict-based model of anxiety-like behavior in which anxiety elicited by exposure to an unfamiliar environment suppresses feeding in hungry mice presented with food (Cryan and Sweeney 2011; Dulawa and Hen 2005). In the NSF test, longer latencies to begin feeding are interpreted as increases in neophobia.

Novel environments activate neural circuits involved in spatial memory (Takeuchi et al. 2016; Wagatsuma et al. 2018), attention (Aston-Jones et al. 2007; Gompf et al. 2010), motivation (Rinaldi et al. 2010; Struthers et al. 2005), arousal (Britton and Indyk 1990; Stone et al. 2005), and anxiety (Delini-Stula et al. 1984; File 2001; Sheth et al. 2008), which enables flexible responding to potential threats or rewards that may emerge under uncertain conditions (Aston-Jones et al. 1999; Kafkas and Montaldi 2018; Snyder et al. 2012). Although much is known about the biological mechanisms that drive conditioned responses to familiar contexts in animal models of anxiety and drug addiction (Koob and Simon 2009; Koob and Volkow 2016; Martin et al. 2009; Tovote et al. 2015), the basic neurochemistry and circuitry of the network that controls innate behavioral responses to novel environments remain largely unknown.

The locus coeruleus (LC) is the primary source of norepinephrine (NE) to the forebrain, projecting extensively within circuits that govern attention, arousal, emotion, and motivated behavior (Aston-Jones et al. 1999; Berridge et al. 1996; Sara and Bouret 2012; Uematsu et al. 2015). A longstanding theory is that the LC-NE system is sensitive to contextual novelty

(Gompf et al. 2010; Sara et al. 1995), and it is well established that the firing rate of LC neurons increases in novel environments and declines as the context becomes familiar (Aston-Jones and Bloom 1981; Vankov et al. 1995). Besides the LC, the brainstem contains smaller, scattered populations of non-cerulean adrenergic cell groups such as A2 neurons in the nucleus of the solitary tract (NTS), which also send axonal projections to the forebrain to regulate affective and motivated behavior (Olson et al. 2006; Ricardo and Koh 1978; Rinaman 2011; Robertson et al. 2016).

Hyperactivity of the LC-NE system has been implicated in the pathophysiology of anxiety and substance abuse disorders (Aston-Jones et al. 1994; Brady 1994; Goddard et al. 2010; Tanaka et al. 2000; Weinshenker and Schroeder 2007; West et al. 2009), and drugs that suppress the activity of the LC or reduce NE signaling have therapeutic efficacy for patients with these conditions (Aston-Jones and Kalivas 2008; Boehnlein and Kinzie 2007; Fox et al. 2012). Experimental disruptions of the LC-NE system also suppress innate behavioral responses to novel environments and stimuli (Archer et al. 1981; Delini-Stula et al. 1984; Harro et al. 1995; Neophytou et al. 2001; Sara et al. 1995; Stone et al. 1999).

The LC-NE system innervates a constellation of cortical and limbic regions implicated in spatial memory and novelty detection, including the anterior cingulate cortex (ACC), medial septum/diagonal band (MS/DB) complex, and hippocampus (Chandler 2016; Guiard et al. 2008; Uematsu et al. 2015). LC-NE signaling within the basolateral amygdala (BLA) elicits anxiety-like behavior (Llorca-Torralba et al. 2019; McCall et al. 2017; Siuda et al. 2016), and NE transmission within the bed nucleus of the stria terminalis (BNST) is associated with stress responses, innate fear, and reward seeking (Avery et al. 2016; Lebow and Chen 2016; Weinshenker and Schroeder 2007). LC-NE inputs to the anterior insula (AI) and medial orbitofrontal cortex (MO) modulate decision making, emotion, and motivated behavior (Mather et al. 2016; Petrides 2007; Rojas et al. 2015; Sadacca et al. 2017). Thus, the LC is well poised to orchestrate both exploratory and anxious behavior in unfamiliar environments (Berridge and Dunn 1989; McCall et al. 2015; Snyder et al. 2012).

In the present study, we investigated the consequences of genetic or pharmacological disruption of central NE signaling on neophilic and neophobic behavior in the NIL and NSF tests using NE-deficient (dopamine β -hydroxylase knockout; *Dbh* $-/-$) mice and NE-sufficient (*Dbh* $+/-$) controls (Thomas et al. 1995). We also determined the effects of pharmacological restoration of central NE synthesis on neophobia in *Dbh* $-/-$ mice and pharmacological reduction of NE transmission on neophobia in control mice (Mineur et al. 2015; Thomas et al. 1998). Finally, we measured how genetic NE deficiency alters c-fos induction in the LC and 15 of its target structures in the forebrain as a proxy measure of task-specific neuronal activity following NIL and NSF.

Methods

Subjects

Dbh $-/-$ mice were maintained on a mixed 129/SvEv and C57BL/6J background, as described (Thomas et al. 1998; Thomas et al. 1995). Pregnant *Dbh* $+/-$ dams were given drinking water supplemented with the β -adrenergic receptor (β AR) agonist isoproterenol

and α 1AR agonist phenylephrine (20 μ g/ml each) + vitamin C (2 mg/ml) from E9.5-E14.5, and L-3,4-dihydroxyphenylserine (DOPS; 2 mg/ml + vitamin C 2 mg/ml) from E14.5-parturition to prevent developmental lethality associated with homozygous *Dbh* deficiency (Mitchell et al. 2008; Thomas et al. 1998; Thomas et al. 1995). *Dbh* $-/-$ mice are easily identified by their ptosis phenotype, and genotypes were confirmed by PCR, as described (Mitchell et al. 2008; Thomas et al. 1995). *Dbh* $+/-$ mice behavior and NE levels are indistinguishable from wild-type (*Dbh* $+/+$) mice and were used as controls (Bourdlat-Parks et al. 2005; Mitchell et al. 2006).

All experiments included adult mice (3–8 months old) of both sexes. Because sex differences were not reported in the literature nor observed in pilot experiments, approximately equal numbers of male and female mice of the same *Dbh* genotype were included in all experiments, and data from both sexes were pooled. All animal procedures and protocols were congruent with the National Institutes of Health guidelines for the care and use of laboratory animals and were approved by the Emory University Animal Care and Use Committee. Mice were maintained on a 12 h light/12 h dark cycle with access to food and water *ad libitum*. For all experiments, behavioral testing was conducted during the light cycle.

Drugs

The following drugs were used for behavioral pharmacology experiments: the α 2AR agonist guanfacine hydrochloride (Sigma-Aldrich, St. Louis, MO), the peripheral aromatic acid decarboxylase inhibitor benserazide (Sigma-Aldrich), and the synthetic NE precursor l-3,4-dihydroxyphenylserine (DOPS; Lundbeck, Deerfield, IL). Guanfacine (0.3 mg/kg) was dissolved in sterile saline (0.9% NaCl) and injected i.p. at a volume of 10 ml/kg. Sterile saline vehicle was injected to control for any confounding effect of injection stress on behavior, and vehicle-treated animals were used for statistical comparison.

For the DOPS rescue experiment, DOPS was dissolved in distilled water with 2% HCl, 2% NaOH, and 2 mg/kg vitamin C, as described (Schank et al. 2008; Thomas et al. 1998). *Dbh* $-/-$ mice were injected s.c. with DOPS (1 g/kg) + benserazide (250 mg/kg), then tested 5 h later once NE levels peaked (Rommelfanger et al. 2007; Thomas et al. 1998). Benserazide was included to prevent the conversion of DOPS to NE in the periphery, thus restricting restoration of NE to the brain (Murchison et al. 2004). The vehicle control for the DOPS experiment was also administered 5 h before testing.

Novelty-induced locomotion (NIL)

Individual mice were removed from their home cages and placed into large novel cages (10" $\times$ 18" $\times$ 10") on locomotor activity testing racks (San Diego Instruments, San Diego, CA) (Schroeder et al. 2013; Weinshenker et al. 2002a). Novelty-induced ambulations (consecutive infrared beam breaks) were recorded for 1 h in 5 min bins. Testing occurred in a brightly lit room where the animals were housed. Test cages were covered with a lid and contained a thin layer of standard bedding substrate during NIL testing.

Novelty-suppressed feeding (NSF)

All chow was removed from the home cage of subject animals 24 h prior to behavioral testing. On the day of testing, mice were moved to the test room and allowed to habituate for at least 2 h before starting the test. For pharmacological experiments, mice were injected either 30 min (guanfacine) or 5 h (DOPS) prior to testing. Individual mice were removed from their home cages and placed into a large novel cage (10" × 18" × 10") with a single pellet of standard mouse chow in the center. The latency for the mouse to feed (grasp and bite the food pellet) was recorded using a stopwatch; mice that did not feed within 10 min were assigned a feeding latency score of 600 sec (Dulawa and Hen 2005; Tillage et al. 2020). Testing occurred in a dark room under red light to reduce the potentially confounding effect of bright light on anxiety-like behavior, and isolate behavioral responses induced purely by the novel environment. Test cages were uncovered and contained a thin layer of standard bedding substrate.

Feeding in the home cage and food neophobia

Mice were singly housed in clean standard mouse cages without chow for 24 h prior to behavioral testing to allow for cage familiarization. On the day of testing, mice were moved to the test room and allowed to habituate for at least 2 h before starting the test. For home cage feeding experiments, the cage lid was removed, and a standard food pellet was introduced into the cage. For food neophobia experiments, the cage lid was removed, and a standard food pellet coated in cinnamon (2 g cinnamon/100 g chow; Penzeys Spices, Wauwatosa, WI) was introduced into the cage (Modlinska and Stryjek 2016; Modlinska et al. 2015). The latency to feed was recorded for both tests using a stopwatch; mice that did not feed within 10 min were assigned a feeding latency score of 600 sec. As with NSF, testing occurred in a dark room under red light.

c-fos immunohistochemistry (IHC)

Dbh ^{-/-} and control mice were exposed to NIL or NSF tasks, after which they were left undisturbed in the test cage for 90 min (from the start of the task). The mice were then euthanized with an overdose of sodium pentobarbital (Fatal Plus, 150 mg/kg, i.p.; Med-Vet International, Mettawa, IL) for transcardial perfusion with cold 4% paraformaldehyde in 0.01 M PBS. Another group of *Dbh* ^{-/-} and control mice was selected for comparisons of brainwide c-fos induction in animals that were naïve to the behavioral tasks and were immediately perfused after removal from the home cage.

After extraction, brains were post-fixed for 24 h in 4% paraformaldehyde at 4°C, and then transferred to cryoprotectant 30% sucrose/PBS solution for 72 h at 4°C. Brains were embedded in OCT medium (Tissue-Tek; Sakura, Torrance, CA) and serially sectioned by cryostat (Leica) into 40-µm coronal slices spanning the entire brain between the LC and orbitofrontal cortex. Brain sections were stored in 0.01 M PBS (0.02% sodium azide) at 4°C before IHC.

For IHC, brain sections were blocked for 1 h at room temperature in 5% normal goat serum (NGS; Vector Laboratories, Burlingame, CA) diluted in 0.01 M PBS/0.1% Triton-X permeabilization buffer. Sections were then incubated for 48 h at 4°C in NGS blocking/

permeabilization buffer, including primary antibodies raised against c-fos (rabbit anti-c-fos, Millipore, Danvers, MA, ABE457; 1:5000) and the NE transporter (NET; chicken anti-NET, #260006, Synaptic System, Goettingen, Germany; 1:3000). After washing in 0.01 M PBS, sections were incubated for 2 h in blocking/permeabilization buffer with goat anti-rabbit AlexaFluor 488 and goat anti-chicken AlexaFluor 568 (Invitrogen, Carlsbad, CA; 1:500). After washing, the sections were mounted onto Superfrost Plus slides and coverslipped with Fluoromount-G plus DAPI (Southern Biotech, Birmingham, AL).

Viral tracing of LC terminal fields in forebrain targets

Adult mice expressing Cre recombinase under the *tyrosine hydroxylase* promoter (TH-Cre; Jax #008601) were anesthetized with isoflurane (for induction and maintenance; Patterson Veterinary Supply, Devens, MA), placed in a stereotaxic frame (David Kopf Instruments, Tujunga, CA), and administered meloxicam (Patterson Veterinary Supply; 5 mg/kg, s.c.) for analgesia prior to the start of surgery. Mice received bilateral infusions of AAV10-EF1a-DIO-hChR2-eYFP (#20298, Addgene; Watertown, MA) into the LC (AP:−5.4 mm, ML: ±1.2 mm, DV, −4.0 mm) with a 5 µL Hamilton syringe (VWR, Radnor, PA) and stereotaxic injector pump (Stoelting, Wood Dale, IL). A total volume of 0.6 µL virus diluted 1:1 with sterile artificial cerebrospinal fluid (Harvard Apparatus, Holliston, MA) was infused per side at a rate of 0.15 µL/min. The infusion needle was left in place for 5 min after each infusion to allow for viral diffusion.

Mice were given at least 3 weeks to recover from surgery and allow full viral expression before they were euthanized and perfused, as described above. Brains were processed for IHC to visualize LC cell bodies and axon terminals in target regions using antibodies raised against TH (rabbit anti-TH, P40101–0, Pel-Freez, Rogers, AR; 1:1000) and YFP (chicken anti-GFP, ab13970, abcam, Cambridge, MA; 1:1000). All steps for IHC were identical to those described above except primary antibody incubations occurred over 24 h and primary antibodies were detected with goat anti-chicken 488 and goat anti-rabbit 568 (Invitrogen; 1:500).

Fluorescent imaging and c-fos quantification

Fluorescent micrographs of immunostained sections were acquired on a Leica DM6000B epifluorescent upright microscope at 10× magnification with uniform exposure parameters for c-fos quantification. For the viral tracing experiment, micrographs were acquired at 20× for visualization of LC cell bodies and axon terminals. NET and TH antibodies were used to define LC cell bodies and identify noradrenergic axon terminals in target regions.

For c-fos quantification, we selected atlas-matched sections from each animal at the level of the LC and 15 of its forebrain targets. A standardized region of interest was drawn for all images to delineate the borders of discrete structures in all animals. Image processing and analysis were performed using ImageJ software. The analysis pipeline included background subtraction, intensity thresholding (Otsu method), and automated cell counting within defined regions of interest, guided by automated size and shape criteria for c-fos+ cells (size: 50–100 pixel², circularity: 0.6–1.0).

Statistical analysis

For NIL experiments, the effect of time on locomotion in *Dbh* $-/-$ and *Dbh* $+/-$ mice was compared using a two-way repeated measures ANOVA (genotype $\times$ time), with post hoc Bonferroni's tests for multiple comparisons. The within-trial habituation curves for NIL were fit with a simple linear regression for statistical comparison between *Dbh* genotypes. For NSF experiments, the effect of test cage familiarity on latency to feed after food deprivation was compared by two-way ANOVA (genotype $\times$ test environment), with post hoc Sidak's tests for multiple comparisons. Food neophobia between genotypes, NSF behavior between DOPS- and vehicle-treated *Dbh* $-/-$ mice, and NSF behavior between guanfacine- and vehicle-treated *Dbh* $+/-$ control mice were analyzed using unpaired t-tests.

For c-fos quantification, genotype differences were compared in the LC and 15 forebrain targets in naïve controls and after NIL or NSF. Comparisons were made within behavioral tasks and between genotypes by multiple t-tests using the Holm-Sidak correction for multiple comparisons. The threshold for adjusted significance was set at $p < 0.05$, and two-tailed variants of tests were used throughout. Analyses and graph design were performed using Prism v8 (GraphPad Software, San Diego, CA).

Results

Norepinephrine deficiency attenuates novelty-induced locomotion and promotes rapid habituation

To determine the contribution of NE to neophilic behavior, a cohort of age- and sex-matched *Dbh* $-/-$ and *Dbh* $+/-$ control mice were compared in the NIL test, with locomotor activity measured in 5 min bins across 1 h. A two-way repeated measures ANOVA (genotype $\times$ time) showed a main effect of genotype ($F(1,21) = 9.28$, $p < 0.01$), time ($F(11,231) = 30.16$, $p < 0.0001$), and a time $\times$ genotype interaction ($F(11,231) = 2.81$, $p = 0.02$). *Dbh* $-/-$ mice made fewer novelty-induced ambulations overall compared to control mice (1612 ± 582 vs 2405 ± 659), and post hoc analyses revealed that the *Dbh* $-/-$ mice were less active than control mice at the beginning of the task during the 5 min ($p = 0.01$), 10 min ($p < 0.01$), 15 min ($p = 0.03$), and 20 min ($p = 0.03$) time bins (Fig. 1a).

Mice of both genotypes showed functional novelty detection, exhibiting the most activity in time bins at the beginning of the task and within-trial habituation to the novel environment. However, simple linear regression analysis of ambulations over time revealed that variation in locomotor activity was predicted more strongly by variation in time in the control mice ($R^2 = 0.43$) than in *Dbh* $-/-$ mice ($R^2 = 0.23$). There was a significant difference in the slope of the fit lines for each *Dbh* genotype ($F(1,272) = 11.22$, $p < 0.001$). The flatter slope of the *Dbh* $-/-$ fit line suggests more rapid locomotor habituation to the novel environment in NE-deficient mice (Fig. 1b).

Central norepinephrine is necessary and sufficient for neophobia in the novelty-suppressed feeding test

To assess whether NE is also required for neophobia, the behavior of *Dbh* $-/-$ and control mice were compared in the NSF test. A two-way ANOVA (genotype $\times$ environment) showed

a main effect of genotype ($F(1,48) = 32.62$, $p < 0.0001$), environment ($F(1,48) = 30.38$, $p < 0.0001$), and a genotype $\times$ environment interaction ($F(1,48) = 25.39$, $p < 0.0001$). Post hoc comparisons revealed that in the novel test environment, *Dbh*^{-/-} mice ate much more rapidly than control mice (91.93 ± 44.43 vs 389.49 ± 162.81 s, $p < 0.0001$), but did not differ from control mice when assessed for feeding latency in the home cage (78.82 ± 44.79 vs 92.45 ± 47.43 sec, $p > 0.05$) (Fig. 2a). The amount of the food eaten over the course of 1 h following the NSF test also did not differ significantly between *Dbh*^{-/-} mice and controls (0.37 ± 0.05 vs 0.37 ± 0.05 g, $p > 0.05$). Thus, the genotype differences in feeding latency are likely to be related to reduced neophobia rather than increased hunger. As expected, *Dbh*^{+/-} mice ate significantly more rapidly in the home cage than they did in the novel cage ($t(48) = 7.57$, $p < 0.0001$). However, latency to feed did not differ between test environments in *Dbh*^{-/-} mice ($t(48) = 0.14$, $p > 0.05$), suggesting NE deficient mice exhibit behavioral indifference to context familiarity when hunger is a motivating factor.

We also measured the effects of NE deficiency in a variation of the NSF test with novel-scented food presented in a familiar environment. To this end, we coated standard mouse chow pellets with cinnamon, a novel olfactory stimulus that is neither innately appetizing nor aversive to rodents (Modlinska and Stryjek 2016; Modlinska et al. 2015; Scholtysik 1980). Similar to the novel environment version of the test, *Dbh*^{-/-} mice ate the novel-smelling pellet much more rapidly than control mice (115.3 ± 99.7 vs 423.6 ± 148.1 sec; $t(19) = 5.54$, $p < 0.0001$) (Fig. 2b).

To evaluate whether replacing central NE in *Dbh*^{-/-} mice would restore neophobic behavior in *Dbh*^{-/-} mice, we compared knockouts treated with vehicle or the synthetic NE precursor DOPS in the NSF test. DOPS treatment significantly increased the latency to feed in *Dbh*^{-/-} mice compared to vehicle treatment (366.4 ± 99.3 vs 88.8 ± 48.8 sec; $t(9) = 6.07$, $p < 0.001$), rescuing feeding latency to similar levels as control mice (Fig. 3a). To determine whether acutely disrupting NE transmission in normal mice confers resistance to neophobia, we also compared *Dbh*^{+/-} mice treated with vehicle or guanfacine (0.3 mg/kg) in the NSF test. Guanfacine is an agonist of α_2 a inhibitory autoreceptors and inhibits NE release (Fox et al. 2012). Guanfacine produced an anxiolytic effect in *Dbh*^{+/-} mice, significantly decreasing latency to feed in the novel environment (312.1 ± 56.1 vs 525.3 ± 76.4 sec; $t(16) = 6.58$, $p < 0.0001$) (Fig. 3b). Thus, neophobia can be restored in NE deficient mice by pharmacological rescue of brain NE synthesis, and attenuated in control mice by reducing NE release with guanfacine.

LC neurons project to forebrain regions implicated in responses to novelty

To determine whether LC neurons project to forebrain region implicated in novelty responses, we bilaterally injected a viral vector encoding Cre-dependent ChR2-eYFP into the LC of TH-Cre mice to label noradrenergic LC neurons and their axon terminals (Fig. 4a,b). Brainstem cell body expression of eYFP was histologically confirmed for all cases and was largely restricted to TH⁺ neurons in the LC, although moderate expression was also observed in the neighboring parabrachial nucleus (PBN) (Fig. 4b). TH-Cre activity in the parabrachial nucleus has been described previously (Lindeberg et al. 2004; Savitt et al. 2005), and was observed in neurons that were not immunoreactive for TH (Fig. 4b).

We next examined 15 forebrain regions previously implicated in novelty responses and reported to receive noradrenergic innervation from the LC (Robertson et al. 2016; Schwarz and Luo 2015; Uematsu et al. 2015; Zaborszky 1989), including the medial orbitofrontal cortex (mOFC), prelimbic (PrL) and infralimbic cortices (IL), anterior cingulate cortex (ACC), agranular insula (AI), claustrum (CL), medial septum/diagonal band (MS/DB), dorsal bed nucleus of the stria terminalis (dBNST), basolateral amygdala (BLA), dorsal dentate gyrus (dDG), dorsal hippocampal subfields CA1 (dCA1) and CA3 (dCA3), ventral hippocampus subfield CA1 (vCA1), lateral hypothalamus (LH), and the paraventricular nucleus of the thalamus (PVT) (Fig. 4a). All forebrain regions analyzed contained eYFP+ fibers, demonstrating direct inputs from the LC to novelty-sensitive forebrain targets (Fig. 4c). Because some YFP expression was observed in the PBN, which sends axonal projections to some of the same forebrain targets as the LC (Chiang et al. 2019), we confirmed that YFP+ terminals in target regions were noradrenergic (NET+) (Fig. 4d).

Norepinephrine deficiency is associated with blunted c-fos induction in several forebrain targets of the locus coeruleus following novelty exposure

To identify potential NE-sensitive novelty circuits from among the identified forebrain LC targets, *Dbh* $-/-$ and *Dbh* $+/-$ control mice were euthanized for quantification of c-fos+ cells in the LC and 15 of its forebrain targets (Fig. 4a) under naïve conditions and following NIL or NSF. c-fos expression was negligible in all regions examined in naïve mice, and there were no significant differences in c-fos+ cells between *Dbh* genotypes ($p > 0.05$ for all regions examined) (Lustberg et al. 2020) (Fig. S1a). Control and *Dbh* $-/-$ mice exposed to a novel environment demonstrated marked increases in c-fos induction in the LC and many of its forebrain targets (Fig. S1b) (Lustberg et al. 2020).

Exposure to the novel environment in either NSF or NIL tests resulted in intense c-fos induction in the LC regardless of *Dbh* genotype ($t(9) = 0.69$, $p > 0.05$ for NIL; $t(6) = 0.81$, $p > 0.05$ for NSF) (Fig. 5a,b), suggesting that NE deficiency does not impair LC activation in response to spatial novelty. There were also no genotype differences observed for the PVT, PrL, IL, dCA1, dCA3, and dDG ($p > 0.05$ for all comparisons) (Fig. 5c).

By contrast, profound genotype differences in c-fos induction emerged in several other forebrain targets of the LC during novelty exposure (Fig. 6a,b). Following either NIL or NSF, *Dbh* $-/-$ mice had significantly fewer c-fos+ cells than control mice in vCA1 ($t(9) = 5.02$, $p < 0.01$ for NIL; $t(9) = 5.13$, $p < 0.01$ for NSF), ACC ($t(9) = 4.86$, $p = 0.01$ for NIL; $t(9) = 5.20$, $p < 0.01$ for NSF), dBNST ($t(8) = 9.12$, $p < 0.001$ for NIL; $t(9) = 4.24$, $p = 0.02$ for NSF), MS/DB ($t(8) = 5.25$, $p < 0.01$ for NIL; $t(8) = 6.63$, $p < 0.01$ for NSF), and BLA ($t(8) = 5.92$, $p < 0.01$ for NIL; $t(9) = 5.63$, $p < 0.01$ for NSF) (Fig. 6a,c). After NSF, but not after NIL, NE-deficient mice also had fewer c-fos+ cells than controls in MO ($t(9) = 5.63$, $p < 0.01$), CL ($t(9) = 4.38$, $p = 0.02$), AI ($t(9) = 3.69$, $p = 0.04$), and LH ($t(9) = 5.69$, $p < 0.01$) (Fig. 6a,d).

Discussion

Norepinephrine is required for sustained locomotor activation in novel environments

In this study, we first compared neophilic behavior in NE-deficient and NE-competent littermate control mice in the NIL test. We found that over the course of 1 h, NE-deficient mice exhibited reduced locomotor activity and more rapid habituation in the novel test environment compared to control mice. These findings are congruent with previous reports from our lab and suggest that novelty may have reduced incentive value or intrinsic salience in NE-deficient animals (Cubells et al. 2016; Porter-Stransky et al. 2019; Weinshenker et al. 2002a).

Dbh $-/-$ mice show a modest deficit in motor learning in the rotarod test but exhibit normal locomotor activity in the open field test and home cage compared to littermate controls, and they can readily acquire new spatial memories (Marino et al. 2005; Thomas and Palmiter 1997). Importantly, novel environments elicit an initial increase in locomotion in *Dbh* $-/-$ mice that habituates over time, suggesting that spatial memory mechanisms that distinguish novel and familiar environments remain intact in the absence of NE. Previous work has also demonstrated functional single-trial learning of a novel environment in *Dbh* $-/-$ mice; total locomotion during the first exposure to the novel environment was significantly greater than total locomotion during re-exposure to the same environment 24 h later in both *Dbh* $-/-$ and control mice (Weinshenker et al. 2002a).

It is interesting to note that this particular type of “one-shot” contextual learning was recently shown to depend on LC transmission of dopamine (DA) to the dorsal hippocampus (Kempadoo et al. 2016; Takeuchi et al. 2016; Wagatsuma et al. 2018). Because DBH catalyzes the conversion of DA to NE, *Dbh* $-/-$ mice make DA instead of NE in LC neurons, and thus LC-DA transmission would theoretically be undisturbed (or even augmented) in *Dbh* $-/-$ mice (BourdlatParks et al. 2005; Schank et al. 2006; Weinshenker et al. 2002b).

Norepinephrine is necessary and sufficient for novelty-suppressed feeding

The present study also investigated the effect of NE deficiency on neophobia in the NSF test. NE-deficient mice demonstrated a total absence of neophobia in the novel test environment that could be rescued to control levels by restoring central NE synthesis with DOPS. In control mice, reducing NE release with guanfacine produced rapid anxiolytic effects in the NSF test. Together, these findings demonstrate that NE is both necessary and sufficient for species-typical neophobia in NSF. Importantly, feeding latencies and food consumption in NE-deficient mice did not differ from controls in the familiar home cage environment, ruling out the possibility that NE-deficient mice are simply hungrier after food deprivation than NE-competent mice (Cryan and Sweeney 2011; Dulawa and Hen 2005).

The profound lack of anxiety displayed by the *Dbh* $-/-$ mice in the NSF test contrasts sharply with the seemingly “normal” phenotype of these mice in canonical tests of anxiety, including the elevated plus and zero mazes (EPM/EZM), light/dark box (LDB), and open field test (OFT) (Lustberg et al. 2020; Marino et al. 2005; Schank et al. 2008). These conflict-based models of anxiety-like behavior are widely used in rodent studies, but they rely on the underlying assumption that mice are motivated to explore novel environments

(open arms of EPM/EZM, light compartment of LDB, center of OFT). The conflict in these canonical tasks is between innate fear (open arms, light, center of field) and the drive to explore the novel environment (Cryan and Sweeney 2011; Kalueff et al. 2007).

Given that *Dbh* $-/-$ mice exhibit blunted exploratory behavior in novel environments, it is possible that any anxiolytic phenotype would be occluded by a lack of neophilia. In other words, the behavioral measures of anxiolysis in these canonical anxiety tasks is exploratory behavior, which in NE-deficient mice is confounded by attenuated neophilia. In NSF, the conflict occurs between the fear of the novel environment and the motivation to eat. The behavioral measure of anxiolysis in NSF is latency to feed; thus, curiosity and neophilia are not a factor in this task.

The fact that a robust anxiolytic phenotype emerged in *Dbh* $-/-$ mice during NSF but not in more commonly employed tests of anxiety-like behavior should serve as a reminder that any behavioral model has underlying assumptions and associated confounds (Cryan and Sweeney 2011; Kalueff et al. 2007). We urge researchers investigating anxiety-like behavior to include NSF in the battery of behavioral testing in order to account for potentially confounding effects of neophilia in more canonical anxiety tests, which all fundamentally measure exploration.

NE-deficient mice show reduced c-fos induction in select targets of the LC following NIL and NSF

Following exposure to the novel environment in the NIL and NSF tests, robust c-fos induction was observed in both NE-deficient and control LC. This finding suggests that LC activation in response to novelty does not depend on NE signaling and may be induced by glutamatergic inputs to the LC, or an excitatory neuropeptide transmitter such as orexin or corticotropin-releasing factor (CRF) (Gompf et al. 2010; Soya et al. 2017; Valentino et al. 1993).

Despite demonstrating similar levels of LC activation compared to controls after NIL or NSF, *Dbh* $-/-$ mice had fewer c-fos+ cells in several forebrain targets of the LC, including vCA1, ACC, dBNST, MS/DB, and BLA. Because these regions were hypoactive in the absence of NE following novelty exposure in both tests (Fig. 7a), we propose that they comprise the nodes of a noradrenergic novelty network (NNN) that drives both neophilic and neophobic behavior (Fig. 7b). With outputs extending throughout the prefrontal cortex, striatum, hypothalamus, midbrain, and brainstem, the NNN is well positioned to coordinate affective and motor responses to unfamiliar environments (Lebow and Chen 2016; Li and Wang 2018; Padilla-Coreano et al. 2016; Stachniak et al. 2014; Tovote et al. 2015).

Recently, a population of spatially- and emotionally-tuned cells in vCA1 were shown to fire preferentially in anxiogenic environments (Jimenez et al. 2018). These “anxiety cells” are analogous to place cells in dCA1 which support spatial learning and memory, and exert their anxiogenic behavioral effects through excitatory glutamatergic projections to LH (Jimenez et al. 2018). The ventral hippocampus has been implicated in emotional learning and neophobia (Bannerman et al. 2002; Bannerman et al. 2004; Fanselow and Dong 2010;

Padilla-Coreano et al. 2016; Santarelli et al. 2003), but the effects of manipulating LC-NE signaling within vCA1 on neophobia have not been described.

Reciprocal connections between the LC and the ACC support vigilance in novel environments, and destruction of LC terminals in the ACC suppresses electrophysiological measures of arousal associated with exposure to contextual novelty (Gompf et al. 2010). Recently, our group reported hypoactivity in the ACC of NE-deficient mice following cage change stress or exposure to a novel environment containing marbles (Lustberg et al. 2020). Unlike control mice, NE-deficient mice demonstrated virtually no stress-induced repetitive behaviors following cage change or novelty exposure, suggesting that NE engagement of the ACC may be necessary for typical affective responses to contextual change (Egner 2011).

The dBNST is a region with a well-established role in innate fear, stress responses, and anxiety-like behavior (Avery et al. 2016; Lebow and Chen 2016). CRF-containing neurons in the dBNST are activated by NE and engage the hypothalamic-pituitary-adrenal axis to synchronize endocrine and behavioral responses to stress (Dabrowska et al. 2016; Egli et al. 2005; Vranjkovic et al. 2017). Activation of this region is also implicated in stress-induced reinstatement of drug-seeking behavior (Mantsch et al. 2016), an animal model of relapse, as well as threat-detection and stimulus evaluation under conditions of uncertainty (Lebow and Chen 2016).

Cholinergic neurons in the MS/DB of the basal forebrain are densely innervated by the LC (Bergado et al. 2007; Schwarz and Luo 2015; Schwarz et al. 2015), and control both exploratory and anxious behavior in novel environments (Bannerman et al. 2004; Carpenter et al. 2017; Myhrer 1989; Zhang et al. 2017). These cholinergic projection neurons likely modulate exploratory and avoidant behavior in novel environments via projections to the hippocampus and medial prefrontal cortex (Jiang et al. 2018; Tereshchenko et al. 2008). Although LC inputs to the MS/DB have not been well characterized, pharmacological studies suggest that the NE increases excitability of cholinergic neurons in this region (Berridge and Espana 2000; Berridge et al. 1996).

The BLA is critical for fear learning and anxiety-like behavior, but is also implicated in novelty responses (Balderston et al. 2011; Jhang et al. 2018). LC-NE signaling within the BLA elicits anxiety-like behavior in canonical anxiety tests as well measures of social anxiety (Llorca-Torralba et al. 2019; McCall et al. 2017; Siuda et al. 2016), but the role of the LC-NE → BLA circuit in neophobia has not been described.

NE-deficient mice show diminished c-fos induction in select targets of the LC following NSF

After NSF, but not after NIL, NE-deficient mice had fewer c-fos+ cells than controls in MO, AI, CL, and LH. The MO is a behavioral control center at the anterior pole of the brain that is bidirectionally connected to the LC (Aston-Jones and Cohen 2005; Rolls and Grabenhorst 2008; Sadacca et al. 2017). MO neurons participate in the evaluation of risk and reward, decision making, and executive control of emotional responses (Petrides 2007; Woon et al. 2019). The AI is a cortical region involved in processing gustatory and interoceptive information (Naqvi and Bechara 2010; Uddin et al. 2017), and it is implicated in panic and

anxiety disorders (Gehrlach et al. 2019; Klumpp et al. 2012; Poletti et al. 2015; Wittmann et al. 2014). NE signaling within the AI mediates food neophobia and is derived from the LC as well as non-cerulean brainstem groups (Robertson et al. 2016; Rojas et al. 2015). The CL is located between the AI and striatum, has connectivity with virtually all cortical regions, and is involved in contextual memory, novelty detection, allocation of attention, and action selection (Brown et al. 2017; Crick and Koch 2005; Qadir et al. 2018).

The LH is comprised of a heterogeneous population of neurons that are also bidirectionally connected to the LC and participate in feeding, anxiety, stress responses, and motivated behavior (Bonnavion et al. 2016; Tanaka et al. 2000). One possible reason that NE-deficient mice demonstrated hypoactivity in these regions after NSF but not after NIL may be the relative demands and parameters of these two tests. For instance, NSF requires the animals to be food-deprived before testing so that the mice are motivated to eat in the test. On test day, NSF also requires the animals to make a behavioral “choice” between avoidance and approach to the food pellet in the novel environment. Thus, NE transmission in NSF may modulate gustatory reward-evaluation and action selection circuits in MO, AI, and CL, as well as motivational and emotional circuits within the LH in the context of the NSF test. The NIL test does not require food deprivation or decision making, potentially masking genotype differences in NE-dependent c-fos induction in these regions.

NE-deficient mice show similar c-fos induction to controls in dorsal hippocampus and paraventricular thalamus after following NIL and NSF

By definition, novelty detection requires engagement of memory systems (Kafkas and Montaldi 2018). In order to determine that an environment is novel, it must be compared with existing representations of previously encountered environments that are stored in memory. Despite their striking absence of neophilia and neophobia, NE-deficient mice retain the ability to recognize an environment as novel or familiar (Weinshenker et al. 2002a). The dissociation between novelty detection and novelty responding in the NE-deficient animals suggests at least partially non-overlapping neural substrates for these operations (Bannerman et al. 2002; Harro et al. 1995; Kafkas and Montaldi 2014; Tereshchenko et al. 2008; Wingo et al. 2016).

Studies in rodents suggest that the hippocampus is functionally segregated along the dorsoventral axis; the dorsal hippocampus is required for spatial memory and navigation, while the ventral hippocampus is required for emotional memory and anxiety (Bannerman et al. 2002; Fanselow and Dong 2010). Although NE-deficient mice had reduced emotional responses and diminished neuronal activity in vCA1 after novelty exposure, they showed no impairment in the detection of spatial novelty and displayed neuronal activity similar to controls in dCA1, dCA3, and dDG after novelty exposure. *Dbh*^{-/-} mice also had similar levels of activity compared to controls in the PVT, a midline thalamic structure that integrates complex contextual signals from hypothalamus and brainstem and which becomes highly activated under conditions of uncertainty (Choi and McNally 2017; Kirouac 2015). As mentioned above, LC-DA transmission is theoretically intact in NE-deficient mice, and recent studies have shown that LC-DA transmission excites neurons within the dorsal hippocampus and PVT (Beas et al. 2018; Wagatsuma et al. 2018).

Limitations and future directions

A limitation of our study is that we cannot infer a causal role for LC-NE transmission to any of the identified forebrain regions in the expression of NIL and NSF behaviors. Other non-cerulean adrenergic cell groups in the brainstem, such as A2 neurons in the NTS, organize behaviors related to stress, reward, appetite, and affect via forebrain projections (Rinaman 2011). In addition to providing noradrenergic inputs to the LC itself, LC and A2 axon terminal fields overlap in prefrontal cortex (including PrL and IL), thalamus (including PVT), hypothalamus (including LH), and extended amygdala (including dBNST) (Ricardo and Koh 1978; van der Kooy et al. 1984). However, to our knowledge, the LC is the sole source of NE to the ACC, MS/DB, BLA, and hippocampus (including vCA1). The hypoactivity of these four nodes of the NNN, which receive NE exclusively from LC projections, following exposure to novelty suggests that the absence of LC-derived NE transmission at one or more these regions may underlie blunted affective responses to novelty in *Dbh* $-/-$ mice.

Another important consideration when interpreting these findings is that the LC is not a homogenous nucleus (Schwarz and Luo 2015; Uematsu et al. 2015), and NE transmission from distinct sets of LC cells can play functionally opposite roles on behavior (Buffalari and Grace 2007; Chandler et al. 2014). Moreover, elegant intersectional genetic techniques have led to even finer parcellations of adrenergic cell groups and interrogations of their functions (Robertson et al. 2013; Robertson et al. 2016). A recent and surprising example of the functional diversity of the central NE system is the finding that chemogenetic activation of NE neurons with a shared genetic history of *Hoxb1* actually reduced stress responses and anxiety-like behavior (Chen et al. 2019). These *Hoxb1* NE neurons are concentrated in the A2 nucleus and in the subcoeruleus regions below the LC, and send noradrenergic projections to the LC, hypothalamus, and extended amygdala (Chen et al. 2019). Thus, additional experiments using optogenetic and chemogenetic tools to bidirectionally manipulate distinct forebrain targets of the LC are necessary to functionally dissect the NNN and understand how NE organizes affective responses to novelty.

Although we have clearly demonstrated a critical role for central NE transmission in neophobia, the neurochemistry underlying innate anxiety is complex and likely to involve multiple neuromodulatory systems. Intriguingly, serotonin (5-HT) deficient mice (*Tph2* $-/-$) also exhibit dramatic reductions in neophobia in the NSF test (Angoa-Pérez et al. 2012; Mosienko et al. 2012). Moreover, these 5-HT deficient mice display increased aggressive and compulsive behaviors, which are absent in NE deficient mice (Angoa-Pérez et al. 2012; Lustberg et al. 2020; Marino et al. 2005; Mosienko et al. 2012). Indeed, the phenotypes of *Tph2* $-/-$ and *Dbh* $-/-$ mice are perfectly opposed with the notable exception of reduced neophobia in both mutants, suggesting that behavioral expression of neophobia may require both NE and 5-HT (Blier and El Mansari 2007).

Conclusions

In summary, these findings support an expanded role of central NE transmission in the expression of neophilic and neophobic behaviors. Guanfacine and related drugs have been used both on- and off-label for the treatment of anxiety and substance abuse disorders (Fox

et al. 2012; Hoehn-Saric et al. 1981), but this is the first study to demonstrate anxiolytic effects of guanfacine in the NSF test. We propose that guanfacine should be investigated for rapid anxiolytic effects in patients with context-specific anxiety symptoms, such as agoraphobia (Marazziti et al. 2012; Wittmann et al. 2014).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Lundbeck for providing the DOPS. This work was supported by the National Institutes of Health (AG061175, NS102306, and DA038453 to DW; GM8602–22 to DL; MH116622 to RPT).

References

- Angoa-Pérez M, Kane MJ, Briggs DI, Sykes CE, Shah MM, Francescutti DM, Rosenberg DR, Thomas DM, Kuhn DM. (2012). Genetic depletion of brain 5HT reveals a common molecular pathway mediating compulsivity and impulsivity. *Journal of neurochemistry* 121:974–984 [PubMed: 22443164]
- Archer T, Ogren S-O, Ross SB. (1981). N-2-chloroethyl-N-ethyl-2-bromobenzylamine hydrochloride (DSP4), a new selective noradrenaline neurotoxin, and taste neophobia in the rat. *Physiological Psychology* 9:197–202
- Aston-Jones G, Bloom F. (1981). Nonrepinephrine-containing locus coeruleus neurons in behaving rats exhibit pronounced responses to non-noxious environmental stimuli. *Journal of Neuroscience* 1:887–900 [PubMed: 7346593]
- Aston-Jones G, Cohen JD. (2005). An integrative theory of locus coeruleus-norepinephrine function: adaptive gain and optimal performance. *Annu. Rev. Neurosci.* 28:403–450 [PubMed: 16022602]
- Aston-Jones G, Iba M, Clayton E, Rajkowski J, Cohen J. (2007). The locus coeruleus and regulation of behavioral flexibility and attention: Clinical implications. *Brain norepinephrine: Neurobiology*:196–235
- Aston-Jones G, Kalivas PW. (2008). Brain norepinephrine rediscovered in addiction research. *Biological psychiatry* 63:1005 [PubMed: 18482610]
- Aston-Jones G, Rajkowski J, Cohen J. (1999). Role of locus coeruleus in attention and behavioral flexibility. *Biological Psychiatry* 46:1309–1320 [PubMed: 10560036]
- Aston-Jones G, Valentino RJ, Van Bockstaele EJ, Meyerson AT. (1994). Locus coeruleus, stress, and PTSD: neurobiological and clinical parallels.
- Avery S, Clauss J, Blackford J. (2016). The human BNST: functional role in anxiety and addiction. *Neuropsychopharmacology* 41:126 [PubMed: 26105138]
- Balderston NL, Schultz DH, Helmstetter FJ. (2011). The human amygdala plays a stimulus specific role in the detection of novelty. *Neuroimage* 55:1889–1898 [PubMed: 21256226]
- Bannerman D, Deacon R, Offen S, Friswell J, Grubb M, Rawlins J. (2002). Double dissociation of function within the hippocampus: spatial memory and hyponeophagia. *Behavioral neuroscience* 116:884 [PubMed: 12369808]
- Bannerman D, Matthews P, Deacon R, Rawlins J. (2004). Medial septal lesions mimic effects of both selective dorsal and ventral hippocampal lesions. *Behavioral neuroscience* 118:1033 [PubMed: 15506885]
- Bardo MT, Donohew RL, Harrington NG. (1996). Psychobiology of novelty seeking and drug seeking behavior. *Behavioural brain research* 77:23–43 [PubMed: 8762157]
- Barnett SA. (1958). Experiments on ‘neophobia’ in wild and laboratory rats. *British journal of psychology* 49:195–201 [PubMed: 13572791]
- Beas BS, Wright BJ, Skirzewski M, Leng Y, Hyun JH, Koita O, Ringelberg N, Kwon H-B, Buonanno A, Penzo MA. (2018). The locus coeruleus drives disinhibition in the midline thalamus via a

dopaminergic mechanism. *Nature neuroscience* 21:963–973 doi:10.1038/s41593-018-0167-4. [PubMed: 29915192]

Bergado JA, Frey S, López J, Almaguer-Melian W, Frey JU. (2007). Cholinergic afferents to the locus coeruleus and noradrenergic afferents to the medial septum mediate LTP-reinforcement in the dentate gyrus by stimulation of the amygdala. *Neurobiology of Learning and Memory* 88:331–341 doi:10.1016/j.nlm.2007.05.003. [PubMed: 17596976]

Berridge C, Espana R. (2000). Synergistic sedative effects of noradrenergic α 1- and β -receptor blockade on forebrain electroencephalographic and behavioral indices. *Neuroscience* 99:495–505 [PubMed: 11029541]

Berridge CW, Bolen SJ, Manley MS, Foote SL. (1996). Modulation of forebrain electroencephalographic activity in halothane-anesthetized rat via actions of noradrenergic β -receptors within the medial septal region. *Journal of Neuroscience* 16:7010–7020 [PubMed: 8824337]

Berridge CW, Dunn AJJoN. (1989). Restraint-stress-induced changes in exploratory behavior appear to be mediated by norepinephrine-stimulated release of CRF. 9:3513–3521

Blier P, El Mansari M. (2007). The importance of serotonin and noradrenaline in anxiety. *International journal of psychiatry in clinical practice* 11:16–23

Boehnlein JK, Kinzie DJJoPP. (2007). Pharmacologic reduction of CNS noradrenergic activity in PTSD: the case for clonidine and prazosin. 13:72–78

Bonnaivon P, Mickelsen LE, Fujita A, Lecea L, Jackson AC. (2016). Hubs and spokes of the lateral hypothalamus: cell types, circuits and behaviour. *The Journal of physiology* 594:6443–6462 [PubMed: 27302606]

Bourdélat-Parks BN, Anderson GM, Donaldson ZR, Weiss JM, Bonsall RW, Emery MS, Liles LC, Weinshenker D. (2005). Effects of dopamine β -hydroxylase genotype and disulfiram inhibition on catecholamine homeostasis in mice. *Psychopharmacology* 183:72–80 [PubMed: 16163519]

Brady LSJBrb. (1994). Stress, antidepressant drugs, and the locus coeruleus. 35:545–556

Britton DR, Indyk E. (1990). Central effects of corticotropin releasing factor (CRF): evidence for similar interactions with environmental novelty and with caffeine. *Psychopharmacology* 101:366–370 [PubMed: 1972997]

Brown SP, Mathur BN, Olsen SR, Luppi P-H, Bickford ME, Citri A. (2017). New Breakthroughs in Understanding the Role of Functional Interactions between the Neocortex and the Claustrum. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 37:10877–10881 doi:10.1523/JNEUROSCI.1837-17.2017. [PubMed: 29118217]

Buffalari DM, Grace AA. (2007). Noradrenergic modulation of basolateral amygdala neuronal activity: opposing influences of α -2 and β receptor activation. *Journal of Neuroscience* 27:12358–12366 [PubMed: 17989300]

Carpenter F, Burgess N, Barry C. (2017). Modulating medial septal cholinergic activity reduces medial entorhinal theta frequency without affecting speed or grid coding. *Scientific reports* 7:14573 [PubMed: 29109512]

Chandler DJ, Gao W-J, Waterhouse BD. (2014). Heterogeneous organization of the locus coeruleus projections to prefrontal and motor cortices. *Proceedings of the National Academy of Sciences* 111:6816–6821

Chandler DJBr. (2016). Evidence for a specialized role of the locus coeruleus noradrenergic system in cortical circuitries and behavioral operations. 1641:197–206

Chen Y-W, Das M, Oyarzabal EA, Cheng Q, Plummer NW, Smith KG, Jones GK, Malawsky D, Yakel JL, Shih Y-YI, Jensen P. (2019). Genetic identification of a population of noradrenergic neurons implicated in attenuation of stress-related responses. *Molecular Psychiatry* 24:710–725 doi:10.1038/s41380-018-0245-8. [PubMed: 30214043]

Chiang MC, Bowen A, Schier LA, Tupone D, Uddin O, Heinricher MM. (2019). Parabrachial Complex: A Hub for Pain and Aversion. *The Journal of Neuroscience* 39:8225 doi:10.1523/JNEUROSCI.1162-19.2019. [PubMed: 31619491]

Choi EA, McNally GP. (2017). Paraventricular thalamus balances danger and reward. *Journal of Neuroscience* 37:3018–3029 [PubMed: 28193686]

- Crick FC, Koch C. (2005). What is the function of the claustrum? Philosophical transactions of the Royal Society of London. Series B, Biological sciences 360:1271–1279 doi:10.1098/rstb.2005.1661. [PubMed: 16147522]
- Cryan JF, Sweeney FF. (2011). The age of anxiety: role of animal models of anxiolytic action in drug discovery. British journal of pharmacology 164:1129–1161 [PubMed: 21545412]
- Cubells JF, Schroeder JP, Barrie ES, Manvich DF, Sadee W, Berg T, Mercer K, Stowe TA, Liles LC, Squires KE, Mezher A, Curtin P, Perdomo DL, Szot P, Weinshenker D. (2016). Human Bacterial Artificial Chromosome (BAC) Transgenesis Fully Rescues Noradrenergic Function in Dopamine β -Hydroxylase Knockout Mice. PLOS ONE 11:e0154864 doi:10.1371/journal.pone.0154864. [PubMed: 27148966]
- Dabrowska J, Martinon D, Moaddab M, Rainnie DG. (2016). Targeting Corticotropin-Releasing Factor Projections from the Oval Nucleus of the Bed Nucleus of the Stria Terminalis Using Cell-Type Specific Neuronal Tracing Studies in Mouse and Rat Brain. Journal of neuroendocrinology 28:10.1111/jne.12442 doi:10.1111/jne.12442.
- Delini-Stula A, Mogilnicka E, Hunn C, Dooley D. (1984). Novelty-oriented behavior in the rat after selective damage of locus coeruleus projections by DSP-4, a new noradrenergic neurotoxin. Pharmacology Biochemistry Behavior 20:613–618
- Dovey TM, Staples PA, Gibson EL, Halford JCG. (2008). Food neophobia and 'picky/fussy' eating in children: a review. Appetite 50:181–193 [PubMed: 17997196]
- Dulawa SC (2009) Novelty-induced hypophagia In: Mood and Anxiety Related Phenotypes in Mice. Springer, pp 247–259
- Dulawa SC, Hen R. (2005). Recent advances in animal models of chronic antidepressant effects: the novelty-induced hypophagia test. Neuroscience Biobehavioral Reviews 29:771–783 [PubMed: 15890403]
- Egli RE, Kash TL, Choo K, Savchenko V, Matthews RT, Blakely RD, Winder DG. (2005). Norepinephrine modulates glutamatergic transmission in the bed nucleus of the stria terminalis. Neuropsychopharmacology 30:657 [PubMed: 15602500]
- Egner T (2011). Surprise! A unifying model of dorsal anterior cingulate function? Nature Neuroscience 14:1219 [PubMed: 21952261]
- Fanselow MS, Dong H-W. (2010). Are the dorsal and ventral hippocampus functionally distinct structures? Neuron 65:7–19 doi:10.1016/j.neuron.2009.11.031. [PubMed: 20152109]
- File SE. (2001). Factors controlling measures of anxiety and responses to novelty in the mouse. Behavioural brain research 125:151–157 [PubMed: 11682106]
- Fox HC, Seo D, Tuit K, Hansen J, Kimmerling A, Morgan PT, Sinha R. (2012). Guanfacine effects on stress, drug craving and prefrontal activation in cocaine dependent individuals: preliminary findings. Journal of psychopharmacology 26:958–972 [PubMed: 22234929]
- Gehrlach DA, Dolensek N, Klein AS, Roy Chowdhury R, Matthys A, Junghänel M, Gaitanos TN, Podgornik A, Black TD, Reddy Vaka N, Conzelmann K-K, Gogolla N. (2019). Aversive state processing in the posterior insular cortex. Nature neuroscience 22:1424–1437 doi:10.1038/s41593-019-0469-1. [PubMed: 31455886]
- Goddard AW, Ball SG, Martinez J, Robinson MJ, Yang CR, Russell JM, Shekhar A. (2010). Current perspectives of the roles of the central norepinephrine system in anxiety and depression. Depression and anxiety 27:339–350 [PubMed: 19960531]
- Gompf HS, Mathai C, Fuller PM, Wood DA, Pedersen NP, Saper CB, Lu J. (2010). Locus ceruleus and anterior cingulate cortex sustain wakefulness in a novel environment. Journal of Neuroscience 30:14543–14551 [PubMed: 20980612]
- Guiard BP, El Mansari M, Blier PJMp. (2008). Crosstalk between dopaminergic and noradrenergic systems in the rat ventral tegmental area, locus coeruleus, and dorsal hippocampus.
- Harro J, Orelund L, Vasar E, Bradwejn JJEN. (1995). Impaired exploratory behaviour after DSP-4 treatment in rats: implications for the increased anxiety after noradrenergic denervation. 5:447–455
- Hoehn-Saric R, Merchant AF, Keyser ML. (1981). Effects of clonidine on anxiety disorders. Archives of General Psychiatry 38:1278–1282 [PubMed: 7305609]

- Jhang J, Lee H, Kang MS, Lee H-S, Park H, Han J-H. (2018). Anterior cingulate cortex and its input to the basolateral amygdala control innate fear response. *Nature communications* 9:2744
- Jiang Y-Y, Zhang Y, Cui S, Liu F-Y, Yi M, Wan Y. (2018). Cholinergic neurons in medial septum maintain anxiety-like behaviors induced by chronic inflammatory pain. *Neuroscience Letters* 671:7–12 doi:10.1016/j.neulet.2018.01.041. [PubMed: 29410218]
- Jimenez JC, Su K, Goldberg AR, Luna VM, Biane JS, Ordek G, Zhou P, Ong SK, Wright MA, Zweifel L. (2018). Anxiety cells in a hippocampal-hypothalamic circuit. *Neuron* 97:670–683. e676 [PubMed: 29397273]
- Kabbaj M, Devine DP, Savage VR, Akil H. (2000). Neurobiological correlates of individual differences in novelty-seeking behavior in the rat: differential expression of stress-related molecules. *Journal of Neuroscience* 20:6983–6988 [PubMed: 10995843]
- Kafkas A, Montaldi D. (2014). Two separate, but interacting, neural systems for familiarity and novelty detection: A dual-route mechanism. *Hippocampus* 24:516–527 [PubMed: 24436072]
- Kafkas A, Montaldi D. (2018). How do memory systems detect and respond to novelty? *Neuroscience Letters* 680:60–68 doi:10.1016/j.neulet.2018.01.053. [PubMed: 29408218]
- Kalueff A, Wheaton M, Murphy D. (2007). What's wrong with my mouse model?: Advances and strategies in animal modeling of anxiety and depression. *Behavioural brain research* 179:1–18 [PubMed: 17306892]
- Kempadoo KA, Mosharov EV, Choi SJ, Sulzer D, Kandel ER. (2016). Dopamine release from the locus coeruleus to the dorsal hippocampus promotes spatial learning and memory. *Proceedings of the National Academy of Sciences* 113:14835–14840
- Kirouac GJ. (2015). Placing the paraventricular nucleus of the thalamus within the brain circuits that control behavior. *Neuroscience & Biobehavioral Reviews* 56:315–329 doi:10.1016/j.neubiorev.2015.08.005. [PubMed: 26255593]
- Klumpp H, Angstadt M, Phan KL. (2012). Insula reactivity and connectivity to anterior cingulate cortex when processing threat in generalized social anxiety disorder. *Biological Psychology* 89:273–276 [PubMed: 22027088]
- Koob GF, Simon EJ. (2009). The Neurobiology of Addiction: Where We Have Been and Where We Are Going. *Journal of drug issues* 39:115–132 doi:10.1177/002204260903900110. [PubMed: 20622969]
- Koob GF, Volkow ND. (2016). Neurobiology of addiction: a neurocircuitry analysis. *The lancet. Psychiatry* 3:760–773 doi:10.1016/S2215-0366(16)00104-8. [PubMed: 27475769]
- Lebow MA, Chen A. (2016). Overshadowed by the amygdala: the bed nucleus of the stria terminalis emerges as key to psychiatric disorders. *Molecular Psychiatry* 21:450–463 doi:10.1038/mp.2016.1. [PubMed: 26878891]
- Li L, Wang L. (2018). Modulation of Innate Defensive Responses by Locus Coeruleus-Superior Colliculus Circuit. *Journal of experimental neuroscience* 12:1179069518792035 [PubMed: 30127637]
- Lindeberg J, Usoskin D, Bengtsson H, Gustafsson A, Kylberg A, Söderström S, Ebendal T. (2004). Transgenic expression of Cre recombinase from the tyrosine hydroxylase locus. *Genesis* 40:67–73 doi:10.1002/gene.20065. [PubMed: 15452869]
- Llorca-Torralba M, Suárez-Pereira I, Bravo L, Camarena-Delgado C, Garcia-Partida JA, Mico JA, Berrocoso E. (2019). Chemogenetic silencing of the locus coeruleus–basolateral amygdala pathway abolishes pain-induced anxiety and enhanced aversive learning in rats. *Biological Psychiatry* 85:1021–1035 [PubMed: 30987747]
- Lustberg D, Iannitelli AF, Tillage RP, Pruitt M, Liles LC, Weinshenker D. (2020). Central norepinephrine transmission is required for stress-induced repetitive behavior in two rodent models of obsessive-compulsive disorder. *Psychopharmacology*:1–15 [PubMed: 31740993]
- Mantsch JR, Baker DA, Funk D, Lê AD, Shaham Y. (2016). Stress-Induced Reinstatement of Drug Seeking: 20 Years of Progress. *Neuropsychopharmacology* 41:335–356 doi:10.1038/npp.2015.142. [PubMed: 25976297]
- Marazziti D, Carlini M, Dell'Osso L. (2012). Treatment strategies of obsessive-compulsive disorder and panic disorder/agoraphobia. *Current topics in medicinal chemistry* 12:238–253 [PubMed: 22204483]

- Marcontell DK, Laster AE, Johnson J. (2003). Cognitive-behavioral treatment of food neophobia in adults. *Journal of anxiety disorders* 17:243–251 [PubMed: 12614666]
- Marino MD, Bourdélát-Parks BN, Liles LC, Weinshenker D. (2005). Genetic reduction of noradrenergic function alters social memory and reduces aggression in mice. *Behavioural brain research* 161:197–203 [PubMed: 15922045]
- Martin EI, Ressler KJ, Binder E, Nemeroff CB. (2009). The neurobiology of anxiety disorders: brain imaging, genetics, and psychoneuroendocrinology. *The Psychiatric clinics of North America* 32:549–575 doi:10.1016/j.psc.2009.05.004. [PubMed: 19716990]
- Mather M, Clewett D, Sakaki M, Harley CW. (2016). Norepinephrine ignites local hotspots of neuronal excitation: How arousal amplifies selectivity in perception and memory. *The Behavioral and brain sciences* 39:e200–e200 doi:10.1017/S0140525X15000667. [PubMed: 26126507]
- McCall JG, Al-Hasani R, Siuda ER, Hong DY, Norris AJ, Ford CP, Bruchas MR. (2015). CRH engagement of the locus coeruleus noradrenergic system mediates stress-induced anxiety. *Neuron* 87:605–620 [PubMed: 26212712]
- McCall JG, Siuda ER, Bhatti DL, Lawson LA, McElligott ZA, Stuber GD, Bruchas MR. (2017). Locus coeruleus to basolateral amygdala noradrenergic projections promote anxiety-like behavior. *eLife* 6:e18247 doi:10.7554/eLife.18247. [PubMed: 28708061]
- Mineur YS, Bentham MP, Zhou W-L, Plantenga ME, McKee SA, Picciotto MRJP. (2015). Antidepressant-like effects of guanfacine and sex-specific differences in effects on c-fos immunoreactivity and paired-pulse ratio in male and female mice. 232:3539–3549
- Mitchell HA, Ahern TH, Liles LC, Javors MA, Weinshenker D. (2006). The effects of norepinephrine transporter inactivation on locomotor activity in mice. *Biological psychiatry* 60:1046–1052 [PubMed: 16893531]
- Mitchell HA, Bogenpohl JW, Liles LC, Epstein MP, Bozyczko-Coyne D, Williams M, Weinshenker D. (2008). Behavioral responses of dopamine β -hydroxylase knockout mice to modafinil suggest a dual noradrenergic–dopaminergic mechanism of action. *Pharmacology Biochemistry Behavior* 91:217–222
- Modlinska K, Stryjek R. (2016). Food Neophobia in Wild Rats (*Rattus norvegicus*) Inhabiting a Changeable Environment-A Field Study. *PloS one* 11:e0156741–e0156741 doi:10.1371/journal.pone.0156741. [PubMed: 27254150]
- Modlinska K, Stryjek R, Pisula W. (2015). Food neophobia in wild and laboratory rats (multi-strain comparison). *Behavioural processes* 113:41–50 [PubMed: 25572457]
- Mosienko V, Bert B, Beis D, Matthes S, Fink H, Bader M, Alenina N. (2012). Exaggerated aggression and decreased anxiety in mice deficient in brain serotonin. *Translational psychiatry* 2:e122–e122 [PubMed: 22832966]
- Murchison CF, Zhang X-Y, Zhang W-P, Ouyang M, Lee A, Thomas SA. (2004). A distinct role for norepinephrine in memory retrieval. *Cell* 117:131–143 [PubMed: 15066288]
- Myhrer T (1989). Exploratory behavior and reaction to novelty in rats: effects of medial and lateral septal lesions. *Behavioral neuroscience* 103:1226 [PubMed: 2610915]
- Naqvi NH, Bechara A. (2010). The insula and drug addiction: an interoceptive view of pleasure, urges, and decision-making. *Brain Structure and Function* 214:435–450 [PubMed: 20512364]
- Neophytou SI, Aspley S, Butler S, Beckett S, Marsden CAJPIn-p, psychiatry b. (2001). Effects of lesioning noradrenergic neurones in the locus coeruleus on conditioned and unconditioned aversive behaviour in the rat. 25:1307–1321
- Olson VG, Heusner CL, Bland RJ, During MJ, Weinshenker D, Palmiter RD. (2006). Role of Noradrenergic Signaling by the Nucleus Tractus Solitarius in Mediating Opiate Reward. *Science* 311:1017 doi:10.1126/science.1119311. [PubMed: 16484499]
- Osorio-Gómez D, Guzmán-Ramos K, Bermúdez-Rattoni F (2018) Neurobiology of neophobia and its attenuation In: *Food Neophobia*. Elsevier, pp 111–128
- Padilla-Coreano N, Bolkan SS, Pierce GM, Blackman DR, Hardin WD, Garcia-Garcia AL, Spellman TJ, Gordon JA. (2016). Direct Ventral Hippocampal-Prefrontal Input Is Required for Anxiety-Related Neural Activity and Behavior. *Neuron* 89:857–866 doi:10.1016/j.neuron.2016.01.011. [PubMed: 26853301]

- Pawlak CR, Ho Y-J, Schwarting RKW. (2008). Animal models of human psychopathology based on individual differences in novelty-seeking and anxiety. *Neuroscience & Biobehavioral Reviews* 32:1544–1568 [PubMed: 18619487]
- Petrides M (2007). The orbitofrontal cortex: novelty, deviation from expectation, and memory. *Annals of the New York Academy of Sciences* 1121:33–53 [PubMed: 17872393]
- Poletti S, Radaelli D, Cucchi M, Ricci L, Vai B, Smeraldi E, Benedetti F. (2015). Neural correlates of anxiety sensitivity in panic disorder: a functional magnetic resonance imaging study. *Psychiatry Research: Neuroimaging* 233:95–101
- Porter-Stransky KA, Centanni SW, Karne SL, Odil LM, Fekir S, Wong JC, Jerome C, Mitchell HA, Escayg A, Pedersen NP. (2019). Noradrenergic Transmission at Alpha1-Adrenergic Receptors in the Ventral Periaqueductal Gray Modulates Arousal. *Biological Psychiatry* 85:237–247 [PubMed: 30269865]
- Qadir H, Krimmel SR, Mu C, Pouloupoulos A, Seminowicz DA, Mathur BN. (2018). Structural connectivity of the anterior cingulate cortex, claustrum, and the anterior insula of the mouse. *Frontiers in neuroanatomy* 12:100 [PubMed: 30534060]
- Ricardo JA, Koh ET. (1978). Anatomical evidence of direct projections from the nucleus of the solitary tract to the hypothalamus, amygdala, and other forebrain structures in the rat. *Brain research* 153:1–26 [PubMed: 679038]
- Rinaldi A, Romeo S, Agustin-Pavon C, Oliverio A, Mele A. (2010). Distinct patterns of Fos immunoreactivity in striatum and hippocampus induced by different kinds of novelty in mice. *Neurobiology of learning & memory* 94:373–381 [PubMed: 20736076]
- Rinaman L (2011). Hindbrain noradrenergic A2 neurons: diverse roles in autonomic, endocrine, cognitive, and behavioral functions. *American journal of physiology. Regulatory, integrative and comparative physiology* 300:R222–R235 doi:10.1152/ajpregu.00556.2010.
- Robertson SD, Plummer NW, de Marchena J, Jensen P. (2013). Developmental origins of central norepinephrine neuron diversity. *Nature Neuroscience* 16:1016–1023 doi:10.1038/nn.3458. [PubMed: 23852112]
- Robertson SD, Plummer NW, Jensen P. (2016). Uncovering diversity in the development of central noradrenergic neurons and their efferents. *Brain research* 1641:234–244 doi:10.1016/j.brainres.2015.11.023. [PubMed: 26612521]
- Rojas S, Diaz-Galarce R, Jerez-Baraona JM, Quintana-Donoso D, Moraga-Amaro R, Stehberg J. (2015). The insula modulates arousal-induced reluctance to try novel tastes through adrenergic transmission in the rat. *Frontiers in behavioral neuroscience* 9:164 [PubMed: 26175672]
- Rolls ET, Grabenhorst F. (2008). The orbitofrontal cortex and beyond: from affect to decision-making. *Progress in neurobiology* 86:216–244 [PubMed: 18824074]
- Rommelfanger K, Edwards G, Freeman K, Liles L, Miller G, Weinshenker D. (2007). Norepinephrine loss produces more profound motor deficits than MPTP treatment in mice. *Proceedings of the National Academy of Sciences* 104:13804–13809
- Sadacca BF, Wikenheiser AM, Schoenbaum G. (2017). Toward a theoretical role for tonic norepinephrine in the orbitofrontal cortex in facilitating flexible learning. *Neuroscience* 345:124–129 [PubMed: 27102419]
- Santarelli L, Saxe M, Gross C, Surget A, Battaglia F, Dulawa S, Weisstaub N, Lee J, Duman R, Arancio O. (2003). Requirement of hippocampal neurogenesis for the behavioral effects of antidepressants. *Science* 301:805–809 [PubMed: 12907793]
- Sara Susan J, Bouret S. (2012). Orienting and Reorienting: The Locus Coeruleus Mediates Cognition through Arousal. *Neuron* 76:130–141 doi:10.1016/j.neuron.2012.09.011. [PubMed: 23040811]
- Sara SJ, Dyon-Laurent C, Hervé A. (1995). Novelty seeking behavior in the rat is dependent upon the integrity of the noradrenergic system. *Cognitive Brain Research* 2:181–187 [PubMed: 7580400]
- Savitt JM, Jang SS, Mu W, Dawson VL, Dawson TM. (2005). Bcl-x is required for proper development of the mouse substantia nigra. *Journal of Neuroscience* 25:6721–6728 [PubMed: 16033881]
- Schank JR, Liles LC, Weinshenker D. (2008). Norepinephrine signaling through β -adrenergic receptors is critical for expression of cocaine-induced anxiety. *Biological psychiatry* 63:1007–1012 [PubMed: 18083142]

- Schank JR, Ventura R, Puglisi-Allegra S, Alcaro A, Cole CD, Liles LC, Seeman P, Weinshenker D. (2006). Dopamine β -hydroxylase knockout mice have alterations in dopamine signaling and are hypersensitive to cocaine. *Neuropsychopharmacology* 31:2221 [PubMed: 16395294]
- Scholtysik G (1980). Pharmacology of guanfacine. *British journal of clinical pharmacology* 10:21S–24S [PubMed: 6249333]
- Schroeder JP, Epps SA, Grice TW, Weinshenker D. (2013). The selective dopamine β -hydroxylase inhibitor nepicastat attenuates multiple aspects of cocaine-seeking behavior. *Neuropsychopharmacology* 38:1032 [PubMed: 23303068]
- Schwarz LA, Luo L. (2015). Organization of the locus coeruleus-norepinephrine system. *Current Biology* 25:R1051–R1056 [PubMed: 26528750]
- Schwarz LA, Miyamichi K, Gao XJ, Beier KT, Weissbourd B, DeLoach KE, Ren J, Ibanes S, Malenka RC, Kremer EJ, Luo L. (2015). Viral-genetic tracing of the input–output organization of a central noradrenaline circuit. *Nature* 524:88–92 doi:10.1038/nature14600. [PubMed: 26131933]
- Sheth A, Berretta S, Lange N, Eichenbaum H. (2008). The amygdala modulates neuronal activation in the hippocampus in response to spatial novelty. *Hippocampus* 18:169–181 [PubMed: 17960646]
- Siuda ER, Al-Hasani R, McCall JG, Bhatti DL, Bruchas MR. (2016). Chemogenetic and Optogenetic Activation of G α s Signaling in the Basolateral Amygdala Induces Acute and Social Anxiety-Like States. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* 41:2011–2023 doi:10.1038/npp.2015.371. [PubMed: 26725834]
- Snyder K, Wang W-W, Han R, McFadden K, Valentino RJ. (2012). Corticotropin-releasing factor in the norepinephrine nucleus, locus coeruleus, facilitates behavioral flexibility. *Neuropsychopharmacology* 37:520 [PubMed: 21993205]
- Soya S, Takahashi TM, McHugh TJ, Maejima T, Herlitze S, Abe M, Sakimura K, Sakurai T. (2017). Orexin modulates behavioral fear expression through the locus coeruleus. *Nature communications* 8:1606
- Stachniak TJ, Ghosh A, Sternson SM. (2014). Chemogenetic synaptic silencing of neural circuits localizes a hypothalamus→ midbrain pathway for feeding behavior. *Neuron* 82:797–808 [PubMed: 24768300]
- Stone E, Zhang Y, Rosengarten H, Yeretsian J, Quartermain D. (1999). Brain alpha 1-adrenergic neurotransmission is necessary for behavioral activation to environmental change in mice. *Neuroscience* 94:1245–1252 [PubMed: 10625064]
- Stone EA, Lin Y, Ahsan MR, Quartermain D. (2005). α 1-Adrenergic and α 2-adrenergic balance in the dorsal pons and gross behavioral activity of mice in a novel environment. *Psychopharmacology* 183:127 [PubMed: 16193333]
- Struthers WM, DuPriest A, Runyan J. (2005). Habituation reduces novelty-induced FOS expression in the striatum and cingulate cortex. *Experimental brain research* 167:136–140 [PubMed: 16041511]
- Takeuchi T, Duzsikiewicz AJ, Sonneborn A, Spooner PA, Yamasaki M, Watanabe M, Smith CC, Fernández G, Deisseroth K, Greene RW. (2016). Locus coeruleus and dopaminergic consolidation of everyday memory. *Nature* 537:357 [PubMed: 27602521]
- Tanaka M, Yoshida M, Emoto H, Ishii H. (2000). Noradrenaline systems in the hypothalamus, amygdala and locus coeruleus are involved in the provocation of anxiety: basic studies. *European journal of pharmacology* 405:397–406 [PubMed: 11033344]
- Tereshchenko Y, Brandewiede J, Schachner M, Irintchev A, Morellini F. (2008). Novelty-induced behavioral traits correlate with numbers of brainstem noradrenergic neurons and septal cholinergic neurons in C57BL/6J mice. *Behavioural brain research* 191:280–284
- Thomas SA, Marck BT, Palmiter RD, Matsumoto AM. (1998). Restoration of norepinephrine and reversal of phenotypes in mice lacking dopamine β -hydroxylase. *Journal of neurochemistry* 70:2468–2476 [PubMed: 9603211]
- Thomas SA, Matsumoto AM, Palmiter RD. (1995). Noradrenaline is essential for mouse fetal development. *Nature* 374:643 [PubMed: 7715704]
- Thomas SA, Palmiter RD. (1997). Disruption of the dopamine B-hydroxylase gene in mice suggests roles for norepinephrine in motor function, learning, and memory. *Behavioral neuroscience* 111:579 [PubMed: 9189272]

- Tillage RP, Sciolino NR, Plummer NW, Lustberg D, Liles LC, Hsiang M, Powell JM, Smith KG, Jensen P, Weinshenker D. (2020). Elimination of galanin synthesis in noradrenergic neurons reduces galanin in select brain areas and promotes active coping behaviors. *Brain Structure Function*:1–19 [PubMed: 31792694]
- Tovote P, Fadok JP, Lüthi A. (2015). Neuronal circuits for fear and anxiety. *Nature Reviews Neuroscience* 16:317 [PubMed: 25991441]
- Tulving E, Markowitsch HJ, Kapur S, Habib R, Houle S. (1994). Novelty encoding networks in the human brain: positron emission tomography data. *Neuroreport: An International Journal for the Rapid Communication of Research in Neuroscience*
- Uddin LQ, Nomi JS, Hebert-Seropian B, Ghaziri J, Boucher O. (2017). Structure and function of the human insula. *Journal of clinical neurophysiology: official publication of the American Electroencephalographic Society* 34:300 [PubMed: 28644199]
- Uematsu A, Tan BZ, Johansen JP. (2015). Projection specificity in heterogeneous locus coeruleus cell populations: implications for learning and memory. *Learning & memory (Cold Spring Harbor, N.Y.)* 22:444–451 doi:10.1101/lm.037283.114.
- Valentino RJ, Foote SL, Page ME. (1993). The Locus Coeruleus as a Site for Integrating Corticotropin-Releasing Factor and Noradrenergic Mediation of Stress Responses a. *Annals of the New York Academy of Sciences* 697:173–188 [PubMed: 7903030]
- van der Kooy D, Koda LY, McGinty JF, Gerfen CR, Bloom FE. (1984). The organization of projections from the cortex, amygdala, and hypothalamus to the nucleus of the solitary tract in rat. *Journal of Comparative Neurology* 224:1–24 [PubMed: 6715573]
- Vankov A, Hervé-Minvielle A, Sara SJ. (1995). Response to novelty and its rapid habituation in locus coeruleus neurons of the freely exploring rat. *European Journal of Neuroscience* 7:1180–1187 [PubMed: 7582091]
- Vranjkovic O, Pina M, Kash TL, Winder DG. (2017). The bed nucleus of the stria terminalis in drug-associated behavior and affect: a circuit-based perspective. *Neuropharmacology* 122:100–106 [PubMed: 28351600]
- Wagatsuma A, Okuyama T, Sun C, Smith LM, Abe K, Tonegawa S. (2018). Locus coeruleus input to hippocampal CA3 drives single-trial learning of a novel context. *Proceedings of the National Academy of Sciences* 115:E310–E316
- Walker QD, Schramm-Sapota NL, Caster JM, Waller ST, Brooks MP, Kuhn CM. (2009). Novelty-induced locomotion is positively associated with cocaine ingestion in adolescent rats; anxiety is correlated in adults. *Pharmacology Biochemistry and Behavior* 91:398–408
- Weierich MR, Wright CI, Negreira A, Dickerson BC, Barrett LF. (2010). Novelty as a dimension in the affective brain. *Neuroimage* 49:2871–2878 [PubMed: 19796697]
- Weinshenker D, Miller NS, Blizinsky K, Laughlin ML, Palmiter RD. (2002a). Mice with chronic norepinephrine deficiency resemble amphetamine-sensitized animals. *Proceedings of the National Academy of Sciences* 99:13873–13877
- Weinshenker D, Schroeder JP. (2007). There and back again: a tale of norepinephrine and drug addiction. *Neuropsychopharmacology* 32:1433 [PubMed: 17164822]
- Weinshenker D, White SS, Javors MA, Palmiter RD, Szot P. (2002b). Regulation of norepinephrine transporter abundance by catecholamines and desipramine in vivo. *Brain Research* 946:239–246 doi:10.1016/S0006-8993(02)02889-5. [PubMed: 12137927]
- West CHK, Ritchie JC, Boss-Williams KA, Weiss JM. (2009). Antidepressant drugs with differing pharmacological actions decrease activity of locus coeruleus neurons. *The international journal of neuropsychopharmacology* 12:627–641 doi:10.1017/S1461145708009474. [PubMed: 18950545]
- Wingo T, Nesil T, Choi J-S, Li MD. (2016). Novelty seeking and drug addiction in humans and animals: from behavior to molecules. *Journal of Neuroimmune Pharmacology* 11:456–470 [PubMed: 26481371]
- Wittmann A, Schlagenhauf F, Guhn A, Lueken U, Gaehlsdorf C, Stoy M, Bermpohl F, Fydrich T, Pfleiderer B, Bruhn H. (2014). Anticipating agoraphobic situations: the neural correlates of panic disorder with agoraphobia. *Psychological medicine* 44:2385–2396 [PubMed: 24398049]

- Woon EP, Sequeira MK, Barbee BR, Gourley SL. (2019). Involvement of the rodent prelimbic and medial orbitofrontal cortices in goal-directed action: A brief review. *Journal of Neuroscience Research* n/a doi:10.1002/jnr.24567.
- Zaborszky L (1989). Afferent connections of the forebrain cholinergic projection neurons, with special reference to monoaminergic and peptidergic fibers. *EXS* 57:12–32 doi:10.1007/978-3-0348-9138-7_2. [PubMed: 2533086]
- Zhang Y, Jiang Y-Y, Shao S, Zhang C, Liu F-Y, Wan Y, Yi M. (2017). Inhibiting medial septal cholinergic neurons with DREADD alleviated anxiety-like behaviors in mice. *Neuroscience Letters* 638:139–144 [PubMed: 27939976]

Fig. 1.

Assessment of neophilia in *Dbh* -/- and *Dbh* +/- mice. **a** Novelty-induced locomotor behavior was attenuated in *Dbh* -/- mice ($n = 11$) compared to controls ($n = 12$). Mice of both *Dbh* genotypes demonstrated the most activity at the start of the task followed by habituation, with activity diminished over time. However, in the four earliest time bins of the test, *Dbh* -/- mice were less active than controls. **b** The change in novelty-induced locomotion over time for both *Dbh* genotypes was assessed using a simple linear regression analysis. The slope of the dashed fit lines of within-trial habituation to the novel environment indicates a blunted initial response to novelty in *Dbh* -/- mice (represented by the Y-intercept) and a flatter slope, suggesting more rapid habituation to the novel environment. Error bars $\pm$ SEM, *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$.

Fig. 2.

Assessment of novelty-suppressed feeding and food neophobia in *Dbh* -/- and *Dbh* +/- mice. **a** Compared to *Dbh* +/- controls (n = 16), *Dbh* -/- (n = 14) ate familiar mouse chow much more rapidly in a novel environment. Unlike controls, *Dbh* -/- mice did not have a significant difference in feeding latency between the novel test environment and familiar home cage. The latency to feed did not differ between *Dbh* +/- controls (n = 11) and *Dbh* -/- mice (n = 11) when tested in the home cage. **b** Compared to *Dbh* +/- controls (n = 11), *Dbh* -/- mice (n = 10) ate the novel-scented cinnamon chow more rapidly in a familiar environment, indicating diminished food neophobia. Error bars $\pm$ SEM, **** p < 0.0001, n.s. = not significant.

Fig. 3.

Pharmacological restoration and attenuation of neophobia. **a** *Dbh* ^{-/-} mice in which central NE was restored with DOPS (n = 5) before NSF testing demonstrate significantly longer latencies to feed than *Dbh* ^{-/-} mice treated with vehicle (VEH; n = 6). **b** *Dbh* ^{+/-} control mice in which central NE transmission was reduced with guanfacine (n = 8) ate more rapidly than control mice treated with VEH (n = 10) in the NSF test. Error bars $\pm$ SEM, **** p < 0.0001, *** p < 0.001.

Fig. 4.

Viral tracing of locus coeruleus (LC) fibers in forebrain regions implicated in novelty. **a** Anatomical location of the LC and 15 target regions in the forebrain: medial orbitofrontal cortex (MO), prelimbic (PrL) and infralimbic cortices (IL), anterior cingulate cortex (ACC), agranular insula (AI), claustrum (CL), medial septum/diagonal band (MS/DB), dorsal bed nucleus of the stria terminalis (dBNST), basolateral amygdala (BLA), dorsal dentate gyrus (dDG), dorsal hippocampal subfields CA1 (dCA1) and CA3 (dCA3), ventral hippocampus subfield CA1 (vCA1), lateral hypothalamus (LH), and the paraventricular nucleus of the

thalamus (PVT). **b** TH-Cre mice received bilateral intra-LC infusions of a Cre-dependent viral vector encoding ChR2-eYFP, which becomes concentrated in axons and axon terminals and can be used as a tool for projection mapping. ChR2-DIO-YFP expression (green) substantially overlaps with immunostaining for TH (red) in the LC and is largely restricted to TH+ LC neurons, with some expression in TH- cells of the neighboring parabrachial nucleus (PNB). **c** Representative micrographs of YFP+ terminals from the LC in the dorsal bed nucleus of the stria terminals (dBNST), medial septum/diagonal band complex (MS/DB), basolateral amygdala (BLA), the CA1 subfield of the ventral hippocampus (vCA1), and the anterior cingulate cortex (ACC). **d** All YFP+ cell bodies in the LC and most YFP+ terminals in the ACC, dBNST, and MS/DB overlap with immunostaining for the norepinephrine transporter (NET), indicating that these axon terminals originate from LC neurons rather than the PBN. The scale bar denotes 100 μ M in all micrographs.

Fig. 5.

Assessing the effects of NE deficiency on neuronal activity in the locus coeruleus (LC) and forebrain target regions following exposure to a novel environment. **a** Quantification of c-fos + neurons in *Dbh* $-/-$ mice in the LC and forebrain targets after NSF (darker purple) and NIL (lighter purple), expressed as a percentage of the average number of c-fos+ cells in *Dbh* $+/-$ controls for each behavioral test and region. There were no significant differences between genotypes in the LC, dorsal hippocampal subfields CA1 (dCA1), CA3 (dCA3), and dentate gyrus (dDG), prelimbic cortex (PrL), infralimbic cortex (IL), or the paraventricular

nucleus of the thalamus (PVT). **b** Representative micrographs showing similar levels of c-fos (green) induction between *Dbh* genotypes after NSF in NET+ (red) LC neurons. Nuclei are counterstained with DAPI (blue). **c** *Dbh* genotype does not affect the number of c-fos+ cells observed following NSF in dorsal hippocampal subregions. Error bars denote group mean $\pm$ SEM. The scale bar denotes 100 μ M in all micrographs.

Fig. 6.

Visualization of NET+ noradrenergic fibers and c-fos induction in target regions after novelty-suppressed feeding (NSF). **a** c-fos induction in *Dbh* -/- mice in the LC and forebrain targets after NSF (darker purple) and NIL (lighter purple), expressed as a percentage of the average number of c-fos+ cells in *Dbh* +/- controls for each behavioral test and region. Dark purple stars indicate $p < 0.05$ for NSF, light purple stars indicate $p < 0.05$ for NIL. **b** Anatomical location of brain regions that were hypoactive in *Dbh* -/- mice after either NSF or NIL, including ventral hippocampus subfield CA1 (vCA1), anterior

cingulate cortex (ACC), dorsal bed nucleus of the stria terminalis (dBNST), medial septum/diagonal band (MS/DB), basolateral amygdala (BLA), medial orbitofrontal cortex (MO), claustrum (CL), agranular insula (AI), and lateral hypothalamus (LH). **c** Representative micrographs showing immunostaining for NET (red) and c-fos (green) between *Dbh* genotypes after NSF. Nuclei are counterstained with DAPI (blue). Following either NSF or NIL, *Dbh* $-/-$ mice showed fewer c-fos+ cells than controls in vCA1, ACC, dBNST, MS/DB, and BLA. **d** Following the NSF test, but not following NIL, *Dbh* $-/-$ mice showed fewer c-fos+ cells than controls in MO, CL, AI, and LH. Error bars denote group mean $\pm$ SEM. The scale bar denotes 100 μ M in all micrographs.

Fig. 7.

The noradrenergic novelty network (NNN). **a** Summary of effects of *Dbh* genotype on c-fos induction in the LC and forebrain targets following NSF (darker purple) or NIL (lighter purple). Downward arrows denote regional hypoactivity in *Dbh*^{-/-} mice and sideways green arrows denote no significant genotype differences following either test. Abbreviations: locus coeruleus (LC), medial orbitofrontal cortex (MO), prelimbic (PrL) and infralimbic cortices (IL), anterior cingulate cortex (ACC), agranular insula (AI), claustrum (CL), medial septum/diagonal band (MS/DB), dorsal bed nucleus of the stria terminalis (dBNST),

basolateral amygdala (BLA), dorsal dentate gyrus (dDG), dorsal hippocampal subfields CA1 (dCA1) and CA3 (dCA3), ventral hippocampus subfield CA1 (vCA1), lateral hypothalamus (LH), and the paraventricular nucleus of the thalamus (PVT). **b** LC-NE inputs to vCA1, ACC, dBNST, MS/DB, and BLA may support diverse and flexible behavioral responses in novel environments. The outputs of the various nodes of the NNN could “prime” either fear-related or exploratory-related circuits in unfamiliar contexts, thus controlling both neophobic and neophilic behavior. Dysregulation of the NNN could be a common feature in certain psychiatric conditions, including anxiety and substance abuse disorders.