Letter to the Editor

The Treatment of Gallstone Disease

by Carsten Gutt, Simon Schläfer, and Frank Lammert in issue 9/2020

Evidence Is Needed in Support of the Indication

The article provides a clear overview of the current consensus regarding the diagnostic evaluation and therapy of gallstone disease (1). As a general practitioner who has been using sonography for many years and has therefore been confronted with the very common gall bladder/gallstone diseases, I am somewhat surprised about the statement that cholecystectomy is indicated in asymptomatic gallstones >3 cm, polyps >1 cm, and porcelain gallbladder. Polyps >1 cm are rather rare, to date I have never come across any. Gallstones >3 cm I have seen, porcelain gallbladder only rarely. More than 40 years ago, before the use of internal medical abdominal sonography became widespread, the scientific consensus was that all patients carrying gallstones—which at that time usually were symptomatic—should undergo cholecystectomy because of an increased risk of cancer (of the gall bladder or common bile duct?). Subsequently this indication was no

longer mentioned in medical discussions or the literature, but it has now resurfaced for large gallstones, again without any evidence (such as cited references or the authors' own data). Why draw the limit at >3 cm?

The authors should provide evidence to support their indication for cholecystectomy primarily in gallstones >3cm but also in polyps >1 cm and porcelain gallbladder (all asymptomatic), because this statement means that more patients should be advised to undergo cholecystectomy.

DOI: 10.3238/arztebl.2020.0490a

Reference

1. Gutt C, Schläfer S, Lammert F: The treatment of gallstone disease. *Dtsch Arztebl Int* 2020; 117: 148–58.

Dr. med. Armin Lay

Praxis für Allgemeinmedizin
Dieblich
arminlay@web.de

In Reply:

Gallstones >3 cm are considered a risk factor for developing gallbladder cancer, because according to the literature the risk is tenfold. Gallstones >1.5 cm that have been present for a long time can also represent a risk (1).

Solitary gallbladder polyps >1 cm have a high probability of being of neoplastic origin. The cancer risk rises to 50% (2).

According to the literature, porcelain gallbladder is also a risk factor for developing gallbladder cancer. The risk is reported to be up to 62% (3).

For detailed information I refer readers to the updated S3 guideline from the German Society of Gastroenterology, Digestive and Metabolic Diseases (DGVS) and the German Society for General and Visceral Surgery (DGAV) for the prevention, diagnostic

evaluation, and treatment of gallstones (AWMF registry No. 021/008). DOI: 10.3238/arztebl.2020.0490b

References

1. Cariati A, Piromalli E, Cetta F: Gallbladder cancers: associated conditions, histological types, prognosis, and prevention. *Eur J Gastroenterol Hepatol* 2014; 26: 5562–9.
2. Okamoto M, Okamoto H, Kitakara F: Ultrasonic evidence of association of polyps and stones with gallbladder cancer. *Am J Gastroenterol* 1999; 94: 446–50.
3. Stephen AE, Berger DL: Carcinoma in the porcelain gallbladder: a relationship revisited. *Surgery* 2001; 129: 699–703.
4. Gutt C, Schläfer S, Lammert F: The treatment of gallstone disease. *Dtsch Arztebl Int* 2020; 117: 148–58.

On behalf of the authors

Prof. Dr. Dr. h.c. Carsten N. Gutt

Klinik für Allgemein-, Visceral-, Thorax- und Gefäßchirurgie,
Klinikum Memmingen
carsten.gutt@klinikum-memmingen.de

Conflict of interest statement

The authors of both contributions declare that no conflict of interest exists.