Letters to the Editor

Reducing the Risk of Preterm Birth by Ambulatory Risk Factor Management

by Prof. Dr. med. Richard Berger, Prof. Dr. med. Dr. h. c. Werner Rath, Prof. Dr. med. Harald Abele, PD Dr. med. Dr. Yves Garnier, PD Dr. med. Ruben-J. Kuon, and PD Dr. med. Holger Maul in issue 50/2019

Preterm Birth Rate in Germany—No Numbers Exist for This

The preterm birth numbers in Germany given in the article “Risk-related prevention of preterm birth in outpatient care” by Berger et al. (1), are wrong, if based on the usual use of [German] language. The authors refer to the Federal Evaluation of the Institute for Quality Assurance and Transparency in Health Care (*Institut für Qualitätssicherung und Transparenz im Gesundheitswesen*, IQTIG) on obstetrics. However, the IQTIG no longer publishes a volume with tables for all statistics and results relating to obstetrics (as was the case with the BQS reports on obstetrics), but rather only publishes the quality indicators for evaluation of obstetrics departments. For these, the institute does not count the number of preterm born children in relation to the number of pregnant women, but only the number of preterm births as number of born children. According to IQTIG, there were 761 176 births and 776 188 children born in Germany in 2017, with 66 730 children born preterm. This is 8.6% of all children born. This number is mentioned in the contribution by Berger et al. (1).

The same evaluation of the IQTIG shows that in 98.1% of all births were singletons, and 1.9% were multiples. The number of premature children that the IQTIG mentions includes about 15 000 children born as multiples, of approximately 7000 multiple deliveries. This would result in a total of around 59 000 preterm births, and a preterm delivery rate of 7.7%, and not 8.6%.

DOI: 10.3238/arztebl.2020.0509a

Reference

1. Berger R, Rath W, Abele H, Garnier Y, Kuon RJ, Maul H: Reducing the risk of preterm birth by ambulatory risk factor management. *Dtsch Arztebl Int* 2019; 116: 858–64.

Dr. med. Susanna Kramarz
Berlin, Germany
info@susanna-kramarz.de

Conflict of interest statement

The authors declare that no conflict of interest exists.

Notable Shortcomings

The review reports on risk factors and the prevention of preterm birth based on an international literature review from 2000 to 2019 (1). The following shortcomings can be noted:

- The development of preterm birth rates in Germany is shown based on a comparison of the years

2009 and 2017, such that a temporal trend cannot be recognized.

- This two-year comparison is incorrect, because preterm birth rates (2009) are compared with rates of babies born preterm (2017).

- The overview of the risk factors is incomplete. Among other things, the missing factors include body mass index, diabetes mellitus, gestational diabetes, iron deficiency without anemia, hypertension, thyroid dysfunction, psychological stress, the use of illicit drugs, weight gain during pregnancy, advanced maternal age, and ethnic origin (2).

- It is particularly critical that the article gives the risk measures (odds ratio [OR], relative risk [RR]) but not the prevalence of these risk factors among pregnant women in Germany, which is however critical for the preventive potential. Research on the prevalence would have revealed considerable deficits in the epidemiological data regarding the living conditions and behaviors of pregnant women as well as research needs. However, the dataset of the BabyCare program, with over 60 000 cases, could have been used as an approximation.

- The article states that rates of preterm births have not been reduced in many countries. Are the authors not aware of the drastic reduction in preterm birth rates in the United States since 2007 (3)?

As a conclusion, the authors note that a careful assessment of the risk factors should be carried out at the beginning of pregnancy. Practicing gynecologists have done this for decades. What is the new message?

DOI: 10.3238/arztebl.2020.0509b

References

1. Berger R, Rath W, Abele H, Garnier Y, Kuon RJ, Maul H: Reducing the risk of preterm birth by ambulatory risk factor management. *Dtsch Arztebl Int* 2019; 116: 858–64.
2. Department of Health, Australian Government: Clinical practice guidelines: pregnancy care. 2018. www.health.gov.au/resources/publications/pregnancy-care-guidelines (last accessed on 23 March 2020).
3. Schoen CN, Tabbah S, Iams JD: Why the United States preterm birth rate is declining. *Am J Obstet Gynecol*. 2015; 213: 175–80.

Dr. phil. Wolf Kirschner
Scientific Director BabyCare FBE
c/o Charité Frauenklinik Berlin, Germany
wkirschner@fbe.gmbh

Conflict of interest statement

The author's wife is the managing director of FBE GmbH. Dr. Kirschner has received reimbursement for travel expenses by FBE GmbH and receives a salary from FBE GmbH as scientific director.

Additional Information Necessary

I would like to thank the authors for their review on the prevention of preterm births—which in fact continues to pose an immense medical problem (1). It is surprising, however, that another important and proven risk factor for preterm births was not mentioned at all: previous abortions also lead to a higher rates of preterm births as well as to infants with a low birthweight in subsequent pregnancies (2, 3). This is plausible from a medical point of view, as iatrogenic trauma to the sensitive cervix can also permanently damage its function. In the article, only “prior medically indicated preterm birth” was included in the Table for risk factors, but this was more related to medically indicated cesarean section in previous pregnancies. Why wasn’t the proven risk factor “prior abortions” mentioned? As controversial as this topic is, isn’t this knowledge, along with the education of affected pregnant women and medical staff (who can only enable correct decision-making when they themselves have all the information for and against it), part of prevention of preterm births? Couldn’t lowering the rate of abortions (which is still very high) and encouraging a “yes” for the unborn child also reduce the rate of preterm births? Shouldn’t all available efforts be taken to do this—in terms of current and subsequent pregnancies? Perhaps the German study on the long term consequences and side effects of abortions intended by the German Federal Minister of Health, Spahn, will add further insights to this topic, in addition to the existing international studies.

DOI: 10.3238/arztebl.2020.0510a

References

- Berger R, Rath W, Abele H, Garnier Y, Kuon RJ, Maul H: Reducing the risk of preterm birth by ambulatory risk factor management. *Dtsch Arztebl Int* 2019; 116: 858–64.
- Thorp JM Jr, Hartmann KE, Shadigan E: Long term physical and psychological consequences of induced abortion: Review of the evidence. *Obst Gynecol Surv* 2002; 58: 67–79.
- Shah PS, Zao J, Knowledge Synthesis Group of Determinants of preterm/LBW births: Induced termination of pregnancy and low birthweight and preterm birth: a systematic review and meta-analyses. *BJOG* 2009; 116: 1425–42.

Dr. med. Michael Kiworr
Mannheim, Germany mkiworr@gmx.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Preconception Interventions Should Not Be Overlooked

Despite some positive results, this interesting review provides a sobering look at the limited possibilities of influencing preterm birth during pregnancy (1). Only three of the risk factors mentioned (short cervix in the second trimester, abstinence from nicotine, and avoidance of heavy physical work) provide evidence for the effectiveness of suitable interventions. However, many of the other risk factors listed are the result of missed preconception interventions, with reduced chances of a normal course of pregnancy right from the start. This also affects the risk of preterm pregnancy termination

for medical reasons, which now accounts for about a third of all preterm births. Common indications for iatrogenic preterm birth are hypertension and diabetic complications of pregnancy, which in turn are associated with pregravid maternal overweight and obesity (2). The chance of a spontaneous conception within a year is already reduced with a body mass index (BMI) >25 kg / m². It also is reduced with increasing age. The average age at first birth is now 30 years. The need for reproductive medicine measures increases with age, while success rates decrease (3). At the same time, there is an increase in the incidence of multiple pregnancies, which are associated with a high risk (relative risk [RR]: 7.7) of preterm birth. The preconception state of health is a key determinant for fertility and the course of pregnancy. As this state is affected by modifiable factors of influence that arise over relatively long periods of time, each woman should have sufficient competence in issues of sexual and reproductive health, and this should be promoted through preconception counseling (3, 4).

DOI: 10.3238/arztebl.2020.0510b

References

- Berger R, Rath W, Abele H, Garnier Y, Kuon RJ, Maul H: Reducing the risk of preterm birth by ambulatory risk factor management. *Dtsch Arztebl Int* 2019; 116: 858–64.
- Stubert J, Reister F, Hartmann S, Janni W: The risks associated with obesity in pregnancy. *Dtsch Arztebl Int* 2018; 115: 276–83.
- Röhl-Mathieu M: Nachhaltige Förderung der Frauengesundheit durch Preconception Counseling. *Gynäkologische praxis* 2016, Band 41/2: 215–31.
- Stephenson J, Heslehurst N, Hall J, et al.: Before the beginning: nutrition and lifestyle in the preconception period and its importance for future health. *Lancet* 2018; 391: 1830–41.

Dr. med. Marianne Röhl-Mathieu

Specialist physician for gynecology and obstetrics
German Professional Association of Gynecologists, Munich, Germany

Conflict of interest statement

The author declares that no conflict of interest exists.

As Always, the Devil Is in the Details

In addition to meta-analyses by the Cochrane Collaboration, there is another meta-analysis on the treatment of asymptomatic bacterial vaginosis (BV) in pregnancy, which includes five randomized controlled trials and demonstrates a significant reduction in late miscarriages and preterm births by therapy with clindamycin (1). As always, the devil is in the details: The effect depends on the week of gestation in which the pregnant women were examined, whether in addition to a BV, also an intermediate vaginal flora (Nugent score 4–6) was treated, on the exact details of the therapy (substance, dose, route of administration), whether the success of the treatment was monitored, and whether the therapy was repeated if necessary (note that there is a high recurrence rate of BV), among other factors. Meta-analyses that are not in line with Cochrane reviews should not be ignored.

The fact that progesterone therapy has proven useful for treating a short cervix is a step forward. But

wouldn't it be professionally much more consistent to prevent the cervix from shortening in the first place? Because cervical shortening is often caused by a BV or vaginal infection, it is important to recognize these as early as possible. For this reason, we have been recommending preventive measures for pregnant women since 1994, which they can carry out themselves early on in pregnancy and in short time intervals, as was not previously possible. The results are encouraging (2). Similar good results were reported by Hoyme et al. from Thuringia, Germany (3). These are not randomized controlled trials. However, as long as no such trials are available on this subject, the best available evidence comes from the results of the existing observational studies. It is not acceptable that one should remain inactive in this respect and wait until someone carries out a randomized trial.

DOI: 10.3238/arztebl.2020.0510c

References

1. Lamont RF, Nhan-Chang C-L, Sobel JD, et al.: Treatment of abnormal vaginal flora in early pregnancy with clindamycin for the prevention of spontaneous preterm birth: a systematic review and metaanalysis. *AJOG* 2011; 205: 177–190.
2. Saling E, Lütjke J, Schreiber M: Efficient and simple program including community-based activities for prevention of very early premature birth. In: Kurjak A, Chervenak FA, eds. *Textbook of Perinatal Medicine*. 3rd ed. Jaypee Brothers Medical Publishers; 2015: 1193–1200
3. Hoyme UB, Huebner J: Prevention of preterm birth is possible by vaginal pH screening, early diagnosis of bacterial vaginosis or abnormal vaginal flora and treatment. *Gynecol Obstet Invest* 2010; 70: 286–90.
4. Berger R, Rath W, Abele H, Garnier Y, Kuon RJ, Maul H: Reducing the risk of preterm birth by ambulatory risk factor management. *Dtsch Arztebl Int* 2019; 116: 858–64.

Prof. Dr. Erich Saling, MD
Jürgen Lütjke, MD

Erich Saling-Institute of Perinatal Medicine, Berlin; Germany
info@saling-institut.de

Conflict of interest statement

The authors declare that no conflict of interest exists.

Hematologic or Vascular Risks as Possible Causes

The standard risk factors for preterm birth are compiled in the article by Berger et al. (1). The authors note that, despite intensive preventive efforts, the risk rate for preterm births has remained unchanged in the past 10 years.

This may be due to the fact that when developing preventive strategies, hematologic and vascular risks, such as the von Willebrand disease or placental insufficiencies (2), have largely been ignored. Preeclampsia often occurs in late pregnancy and can be detected by the presence of a notch in detailed diagnostic examinations in the 22nd gestation week; nonetheless, the article does not mention that administration of either acetylsalicylic acid (ASA) before the 16th gestation week (3) or after the 20th gestation week (4), or low molecular weight heparin, leads to reduction of risk.

Additionnally, neither placental insufficiency or microcirculatory disorders as risk factors nor its treatments were mentioned in the article, although around 30% of pregnant women suffer from it to varying degrees. In these cases, administration of magnesium,

drinking a volume of 2 to 3 liters, and if necessary, administration of low molecular weight heparin is advisable, as these measures can significantly increase placental blood flow. A study is currently being conducted with pentaerythrityl tetranitrate to test its effectiveness in preventing fetal growth restriction (2).

Unfortunately, there was no mention of the HELLP syndrome, which, like preeclampsia, can be prevented by ASA and low molecular weight heparin in a subsequent pregnancy (1, 2).

DOI: 10.3238/arztebl.2020.0511a

References

1. Berger R, Rath W, Abele H, Garnier Y, Kuon RJ, Maul H: Reducing the risk of preterm birth by ambulatory risk factor management. *Dtsch Arztebl Int* 2019; 116: 858–64.
2. Groten T, Lehmann T, Schleußner E, PETN Study Group: Does Pentaerythrityltetranitrate reduce fetal growth restriction in pregnancies complicated by uterine mal-perfusion? *BMC Pregnancy childbirth* 2019; 19: 336.
3. Roberge S, Nicolaides K, Demers S, Hyett J, Chaillet N, Bujold E: The role of aspirin dose on the prevention of preeclampsia and fetal growth restriction: systematic review and meta-analysis. *Am J Obstet Gynecol* 2017; 216: 110–20.
4. Becker R, Keller T, Kiesewetter H, Fangerau H, Bittner U: Individual risk assessment of adverse pregnancy outcome by multivariate regression analysis may serve as basis for drug intervention studies: retrospective analysis of 426 high-risk patients including ethical aspects. *Arch Gynecol Obstet* 2013; 288: 41–8.

Prof. Dr. med. Dr.-Ing. Holger Kiesewetter

Hämostaseologikum MVZ GbR
Berliner Zentrum für Blutgerinnungsstörungen
und Humangenetik, Berlin, Germany
H.Kiesewetter@haemostaseologikum.com

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

We are grateful for the high level of interest and for the numerous Letters to the Editor regarding our article (1). This underscores the importance of this topic for daily practice.

In international literature, the term “preterm birth rate” is understood to mean the rate of babies born preterm. The percentages given in our work, 9.38% for 2009 and 8.60% for 2017, are the correct preterm birth rates for Germany, taken from the Cross-sectoral Quality in Healthcare (SQG, *Sektorenübergreifende Qualitätssicherung*) federal evaluation and the IQTIG federal evaluation. The preterm birth rates are not specified in the IQTIG federal evaluations. Thus, our information could therefore be misinterpreted.

In the United States, the preterm birth rates were 10.44% in 2007, and 10.02% in 2018 (2). One cannot speak of a drastic reduction here.

Due to the limited scope of our article in the *Deutsches Ärzteblatt* and the large number of risk factors, we had to carefully select the risk factors based on their clinical relevance in practice and their ability to be influenced. We are aware that this selection cannot satisfy all readers. Undisputedly, a prior curettage (for miscarriage, abortion), anemia, gestational diabetes, and pre-eclampsia are other risk factors for preterm birth. A recently published, randomized placebo-controlled trial showed that, the rate of spontaneous

preterm births can also be significantly reduced with low-dose aspirin as prophylaxis (3).

We thank you for pointing out that the risk factors should be given with respect to their prevalence in Germany. However, reliable data in this regard have so far been lacking.

Bacterial vaginosis is a significant risk factor for preterm birth. Nonetheless, a screening strategy (of vaginal pH measurement) only makes sense if evidence-based therapy is also available. Systemic or local administration of antibiotics is out of the question, as the PREMAVA trial recently showed (4). Antibiotics are highly potent drugs that can not only destroy the vaginal microbial flora, but also significantly affect the eubiosis of the maternal gut microbiome (for instance) and, as a result, can lead to diarrhea and abdominal complaints. Randomized controlled studies on other promising measures (especially displacement of pathogenic germs by promoting the physiological flora) in the prevention of preterm birth in bacterial vaginosis are not yet available.

The information on the possibility of lowering of the preterm birth rate through preconception measures could represent a worthwhile approach for future prospectively randomized prevention trials.

Our review was not about reducing iatrogenic preterm birth but rather about reducing spontaneous preterm birth.

Even if no significant reduction was observed for the preterm birth rate in Germany, it was important for us to demonstrate that the use of various preventive measures, such as progesterone, cerclage, and cervical pessary, can potentially reduce the risk of preterm birth (1).
DOI: 10.3238/arztebl.2020.0511b

References

1. Berger R, Rath W, Abele H, Garnier Y, Kuon RJ, Maul H: Reducing the risk of preterm birth by ambulatory risk factor management. *Dtsch Arztebl Int* 2019; 116: 858–64.
2. Statista: Percentage of preterm births in the United States from 1990 to 2018. www.statista.com/statistics/276075/us-preterm-birth-percentage/ (last accessed on 20 March 2020).
3. Hoffman MK, Goudar SS, Kodkany BS, et al.: Low-dose aspirin for the prevention of preterm delivery in nulliparous women with a singleton pregnancy (ASPIRIN): a randomised, double-blind, placebo-controlled trial. *Lancet* 2020; 395: 285–93.
4. Subtil D, Brabant G, Tilloy E, et al.: Early clindamycin for bacterial vaginosis in pregnancy (PREMEVA): a multicentre, double-blind, randomised controlled trial. *Lancet* 2018; 392: 2171–9.

On behalf of the authors

Prof. Dr. med. Richard Berger
Marienhaus Klinikum St. Elisabeth
Akademisches Lehrkrankenhaus
der Universitäten Mainz und Maastricht
Klinik für Gynäkologie und Geburtshilfe
Neuwied, Germany
richard.berger@marienhaus.de

Conflict of interest statement

Prof. Berger has been reimbursed for conference fees and travel expenses, and has received lecture honoraria, from Eickeler.