Published in final edited form as:

Neurobiol Aging. 2020 November ; 95: 168–175. doi:10.1016/j.neurobiolaging.2020.07.017.

Changes in the intracellular microenvironment in the aging human brain

Dinesh K. Deelchand, PhD^{1,*}, J. Riley McCarten, MD^{2,3}, Laura S. Hemmy, PhD^{2,4}, Edward J. Auerbach, PhD¹, Lynn E Eberly, PhD⁵, Małgorzata Marjańska, PhD¹

¹Center for Magnetic Resonance Research and Department of Radiology, University of Minnesota, Minneapolis, MN 55455, United States

²Geriatric Research, Education and Clinical Center, Veterans Affairs Health Care System, Minneapolis, MN 55417, United States

³Department of Neurology, University of Minnesota, 12-112 PWB, Minneapolis, MN 55455, United States

⁴Department of Psychiatry, University of Minnesota, F282/2A West, Minneapolis, MN 55454, United States

⁵Division of Biostatistics, University of Minnesota, Minneapolis, MN 55455, United States

Abstract

Normal brain aging is associated with changes occurring at all levels. This study investigates age-related differences in the brain intracellular microenvironment by comparing the apparent diffusion coefficients (ADC) and apparent transverse relaxation time-constants (T_2) of five neurochemicals (i.e., total *N*-acetyl-aspartate, total creatine, total choline, glutamate, and *myo*-inositol) between young and older adults. Thirty-two young healthy adults (18-22 years) and twenty-six older healthy adults (70-83 years) were recruited. Three brain regions were studied at 3T: prefrontal, posterior cingulate and occipital cortices. ADC and T_2 were measured using

*Corresponding author: Dinesh K. Deelchand, PhD, Center for Magnetic Resonance Research, University of Minnesota, 2021 6th St SE, Minneapolis, MN 55455, USA, Phone: (1) 612-625-8097, Fax: (1) 612-626-2004, deelc001@umn.edu.

Authors Contributions

Dinesh K. Deelchand: data acquisition, processing and analysis, writing and revising the manuscript.

J. Riley McCarten: conceptualization, clinical evaluation of older adults, revising the manuscript.

Laura S. Hemmy: conceptualization, cognitive evaluation of older adults, revising the manuscript.

Edward J. Auerbach: development of acquisition tools, revising the manuscript.

Lynn E Eberly: statistical analysis of the data, writing and revising the manuscript.

Małgorzata Marjańska: conceptualization, data acquisition, processing and analysis, writing and revising the manuscript.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosure statement

The authors disclose no conflicts of interest.

3. The data contained in this manuscript have not been previously published, have not been submitted elsewhere and will not be submitted elsewhere while under consideration at *Neurobiology of Aging*.

4. The study was approved by the Human Subjects' Protection Committee at the University of Minnesota. All recruited subjects provided written informed consent.

5. All authors have reviewed the contents of this manuscript, approved of its contents, and validate the accuracy of the data.

STEAM and LASER sequences, respectively. This study shows that the diffusivities of several neurochemicals are higher in older than in younger adults. In contrast, shorter apparent T_2 values for several metabolites were measured in older adults. Age-related difference in ADC and apparent T_2 of metabolites seem to be region-specific. Furthermore, this study shows that it is feasible to observe age-related differences in the cellular microenvironment of neurochemicals in the normal aging brain.

Keywords

ADC; diffusion; T_2 ; magnetic resonance spectroscopy; young

Introduction

Normal brain aging is associated with changes occurring at all levels from molecular to morphological, e.g., changes in blood flow and metabolism, decline in a number of cognitive functions, changes in the volume of the brain, the amount of white matter (WM), and an increase in the cerebral iron content (Peters, 2006). All of these changes occur to a different extent in the entire human brain.

Magnetic resonance imaging (MRI) is a non-invasive imaging modality, which has been widely used for volumetric and morphometric analysis of the brain. Using structural MRI, it was previously demonstrated that brain volume changes correlate with age (Walhovd et al., 2011). Interestingly, the prefrontal cortex (PFC) was found to be the most affected region while occipital cortex (OCC) and posterior cingulate cortex (PCC) were the least affected regions (Fjell and Walhovd 2010), consistent with the cognitive changes observed in the normal aging brain. In addition, diffusion-weighted imaging (DWI) assessing brain tissue microstructure showed that the apparent diffusion coefficients (ADC) of water gradually increased from young adulthood into late life (Klimas et al., 2013; Watanabe et al., 2013). On the other hand, fractional anisotropy, which is a measure of structural macroscopic anisotropy in the brain, was found to decrease with age in the frontal, parietal, and temporal lobes (Grieve et al., 2007).

Proton magnetic resonance spectroscopy (^1H MRS) is a technique which allows for non-invasive measurement of age-associated neurochemical concentrations *in vivo* in the brain (Marja ska et al., 2017). The most commonly measured neurochemicals are *N*-acetyl-aspartate (NAA), total creatine (tCr, creatine plus phosphocreatine), total choline (tCho, phosphorylcholine plus glycerophosphorylcholine), glutamate (Glu) and *myo*-inositol (mIns). These neurochemicals are preferentially concentrated in certain cell types: for example, NAA and Glu are predominantly located in neurons, tCr and tCho are found in both neuronal and glial cells, while mIns is thought to be localized exclusively in astrocytes (de Graaf, 2007; Ligneul et al., 2019). Age-related differences in neurochemical concentrations are also region-specific (Marja ska et al., 2017) where NAA and Glu were lower with age in OCC while mIns and tCr were higher with age in PCC.

The cellular microenvironment of neurochemicals can be examined using advanced MRS techniques such as diffusion-weighted (DW)-MRS and transverse (T_2) relaxometry. The

ADC value as measured by DW-MRS reflects the molecular displacement motion within the intracellular space (Ronen and Valette, 2015). By probing the diffusivity of metabolites, specific information on compartmentalization can be obtained at both cellular and subcellular levels (Palombo et al., 2018b; Valette et al., 2018). On the other hand, T_2 relaxation is a result of nuclear spin-spin interactions as reflected by MRS signal decay with time. T_2 is very sensitive to changes in molecular motion primarily through interaction of metabolites with structural or cystolic macromolecules (Öngür et al., 2010).

With DW-MRS, we recently reported that the diffusivity of five major metabolites (NAA, tCr, tCho, Glu and mIns) can be measured at 3 T (Deelchand et al., 2018b). Previously, it was demonstrated that the ADC of tNAA, tCr, and tCho were lower in older adults (Zheng et al., 2012). However, no difference in the ADC of water was found between the young and older cohorts, which is contradictory to previous DWI studies (Klimas et al., 2013; Watanabe et al., 2013). Several studies have reported the apparent T_2 relaxation time constants of metabolites with contradicting results in the older adult brains (Christiansen et al., 1993; Kirov et al., 2008; Kreis et al., 2005). We previously showed that apparent T_2 of NAA, tCr, and tCho can be measured reliably at high field strength, and these T_2 values were shorter in older adults by 10 to 23% in the OCC (Marjańska et al., 2013).

Therefore, the aims of this study were to compare trace/3 ADC values (i.e., averaged ADC along three orthogonal directions) and apparent T_2 relaxation time constants of the five major metabolites and water in the human brain between young and older adults in three brain regions: PCC, OCC, and PFC.

Methods

Subjects

Thirty-two young adults (9 males, age: 21 ± 1 years; range: 18-22 years) and twenty-six older adults (11 males, age: 74 ± 3 years, range: 70-83 years) were recruited after providing informed consent for the study approved by the Human Subjects Protection Committee at the University of Minnesota Institutional Review Board. All subjects completed a medical history questionnaire validated to recruit healthy older adults for cognitive research (Christensen et al., 1992) and controls for central nervous system (CNS) and cardiovascular disease, and were screened for cognitive impairment with the Montreal Cognitive Assessment (MoCA, (Nasreddine et al., 2005)). Study exclusions included MoCA scores < 24, neurological disorders (e.g., stroke, neurodegenerative conditions, epilepsy), a history of psychiatric diagnoses or current psychotropic medication use. Older adults were further evaluated with a neuropsychological battery to rule out any previously undetected cognitive deficits (Trail Making Test (Reitan, 1979), Digit Span Forward (Wechsler, 1997), Stoop (Golden, 1978), Boston Naming Test (Kaplan et al., 1983), verbal fluency (Benton et al., 1994), CVLT (Delis et al., 2000), Rey Complex Figure (Meyers and Meyers, 1995), Logical Memory (Wechsler, 1987), Block Design (Wechsler, 1997), Digit Symbol (Wechsler, 1997), Information (Wechsler, 1997), AmNART (Grober et al., 1991)). Young adults were evaluated with an abbreviated version of the older adult battery to rule out any gross cognitive deficits (Information, Digit Span, Block Design, Digit Symbol, verbal fluency, CVLT).

ADC and T_2 data were collected from three different brain regions in all subjects using the two MRS acquisition protocols in two randomized sessions. The order in which information from different brain regions was obtained was also randomized between subjects. Both sessions were carried out within five weeks. T_2 data were missing from one subject in each group since the two subjects did not come back for their measurements.

MR acquisition

All MR measurements were carried out on a whole-body 3 T Siemens Prisma^{fit} system (Siemens Medical Solutions, Erlangen, Germany). The standard body coil was used for radiofrequency (RF) excitation while the 32-channel receive-only head-coil was used for signal reception. ADC and T_2 data were acquired from three volumes-of-interest (VOIs): PCC ($2.5 \times 2.5 \times 2.5 \text{ cm}^3$), OCC ($3 \times 2.3 \times 2.3 \text{ cm}^3$), and PFC ($2.5 \times 2.5 \times 2.5 \text{ cm}^3$) (eFigure 1). For high reproducibility of voxel placement between acquisitions, the vendor-provided automatic voxel positioning technique called AutoAlign was utilized (van der Kouwe et al., 2005).

In each session, magnetization-prepared rapid gradient-echo (MPRAGE) images (1 mm^3 isotropic resolution, repetition time, $T_R = 2530 \text{ ms}$, echo time, $T_E = 3.65 \text{ ms}$, inversion time, $T_I = 1100 \text{ ms}$, flip angle = 7° , GRAPPA acceleration factor = 2) were acquired to position the VOIs. First- and second-order shims were automatically adjusted for each VOI using the system 3D gradient-echo shim, operated in the “Brain” mode. In addition, the B_1 field for the 90° pulse and the water suppression flip angles were calibrated for each VOI.

ADC data were measured using STEAM ($T_E/T_M/T_R = 21.22/105/3000 \text{ ms}$) as previously described (Deelchand et al., 2018b). Briefly, the water signal was suppressed using VAPOR interleaved with outer volume suppression (OVS) pulses. Diffusion weighting was applied using bipolar gradients in three orthogonal directions with positive gradients i.e., $[1, 1, -0.5]$, $[1, -0.5, 1]$, and $[-0.5, 1, 1]$ and negative gradients i.e., $[-1, -1, 0.5]$, $[-1, 0.5, -1]$, and $[0.5, -1, -1]$ to remove cross-term effects. Single shot metabolite spectra were acquired at two b -values: a null b -value (16 averages) and a high nominal b -value (48 averages, six diffusion-weighted directions) of 3172 s/mm^2 in all three regions. The diffusion gradient duration was 5.85 ms operating at 70 mT/m with a diffusion time of 118 ms . Water reference scans at both b -values were also acquired for eddy-current correction.

T_2 of metabolites and water were measured using LASER ($T_R = 3 \text{ s}$) as previously described (Deelchand et al., 2018a). Water suppression was also achieved with VAPOR, but no OVS pulses were used in the pulse sequence. Metabolite spectra were acquired at six different T_E s: 35, 140, 230, 290, 330, 400 ms with 8, 16, 32, 32, 64, and 128 averages, respectively. This nonuniform averaging scheme with more averages measured at longer TE was used to have signal-to-noise ratio (SNR) spectra at all T_E s. Water reference scans were also acquired for eddy-current correction and for the determination of the apparent T_2 of tissue water.

Individual single-shot spectra from both sessions were saved for further post-processing offline. All MRS data were acquired with 2048 complex data points using a spectral width of 6 kHz . The carrier frequency of the localization RF pulses was set to 3 ppm .

Tissue segmentation

MPRAGE images were segmented using the FMRIB Software Library (FSL) (Jenkinson et al., 2012). Fractions of gray matter (GM), white matter (WM), and cerebrospinal fluid (CSF) in each VOI were determined using an in-house written algorithm in MATLAB (MathWorks Inc., Natick, MA). For example, the GM fraction was estimated as the ratio of all voxels classified as GM to the total voxels in the VOI.

Spectral processing and quantification

All MR spectra were post-processed in MATLAB using the MRSpa package (<https://www.cmrr.umn.edu/downloads/mrspa/>). Eddy current effects were first corrected. For ADC data only, low SNR spectra were removed to avoid bias in the ADC values (Deelchand et al., 2018b). Single-shot frequency and phase corrections were then performed using cross-correlation and least-square algorithms. For the ADC data, spectra, at null b -value and for all six gradient diffusion directions, were summed separately. Similarly, spectra acquired at different T_E s were individually summed for the T_2 data.

All resulting spectra were analyzed with LCModel (Provencher, 1993) version 6.3-0G (Stephen Provencher Inc, Oakville, Ontario, Canada) without applying any baseline correction, zero-filling, or apodization functions to the *in vivo* data. All spectra were fitted between 0.5 and 4.1 ppm.

Basis sets for STEAM, and LASER at six different T_E s, were simulated using custom software in MATLAB based on density matrix formalism (Henry et al., 2006) using measured and published chemical shift and J -coupling values (Govind et al., 2015).

The basis set for both sequences consisted of nineteen metabolites: alanine, ascorbate, aspartate, creatine, γ -aminobutyric acid, glucose, glutamate, glutamine, glutathione, glycerophosphorylcholine, mIns, *scyllo*-inositol, lactate, NAA, NAAG, phosphocreatine, phosphorylcholine, phosphorylethanolamine, and taurine. Separate basis spectra were generated for the singlet and multiplet (i.e., CH_3 and CH_2 groups) resonances of NAA: denoted as sNAA and mNAA, respectively. In addition, the CH_3 and CH_2 groups of creatine and phosphocreatine were separated for the T_2 analysis. Due to differences in the macromolecular pattern recently observed between young and older adults (Marja ska et al., 2018), the macromolecular spectra were measured from both cohorts using the metabolite-nulled technique in STEAM (2 young subjects with 704 averages, 2 older subjects with 1280 averages) and LASER (2 young subjects with 512 averages, 2 older subjects with 1536 averages). These macromolecular spectra were included in the basis sets for each sequence. A singlet peak was also simulated at 1.45 ppm within LCModel to account for the falx cerebri lipid signal (McIntyre et al., 2007) observed in several of the older adults. Metabolites with Cramer-Rao lower bounds < 25% were selected for further analysis.

ADC values for tCr, tCho, tNAA, Glu and mIns were calculated as the mean ADC in all three gradient diffusion directions after determining the ADC value in each gradient diffusion direction using the geometric mean of the metabolite signals measured with both gradient polarities as previously reported (Deelchand et al., 2018b). Water ADC was also determined from the integral of the eddy current corrected water peak.

The apparent T_2 values for tCr, tCho, tNAA, Glu, and mIns were obtained by fitting the LCModel amplitudes of these metabolites with an exponential decay function (Mlynarik et al., 2001). Since water T_2 values correspond to different environments, the T_2 of tissue water and T_2 of water in CSF in each subject were determined by fitting the integrals of the eddy current corrected water peak at each T_E with a bi-exponential function (Whittall et al., 1997) using the known CSF content in each VOI from segmentation.

Statistics

Apparent T_2 , ADC, and tissue content data were summarized with means and standard deviations; no large outliers were seen. Young vs. older groups were compared using a linear mixed model, fit for each T_2 and each ADC measure separately; each model included group, region, and their interaction, and allowed for group and region specific variances. Models were adjusted for sex. Diagnostics indicated that model assumptions were well met. Tissue (WM, GM) and CSF contents were compared within region between young and older groups using one-way ANOVA. Bonferroni correction was used to control for multiple testing (6 neurochemicals in T_2 measurements and 5 neurochemicals in ADC measurements in the three brain regions), the threshold of significance was 0.0027 (0.05/18) and 0.0033 (0.05/15) for T_2 and ADC values of metabolites, respectively. The threshold of significance was 0.0083 (0.05/6) and 0.0167 for T_2 of water and ADC, respectively.

Results

ADC of metabolites and water

Typical diffusion-weighted STEAM spectra acquired from the PCC are shown in Figure 1 for one young and one older adult. Excellent high spectral quality, with minimal lipid contamination and flat baseline from efficient water suppression, was routinely obtained in this study. Consistent data quality was obtained in OCC and PCC as observed by the narrow water linewidths, with mean linewidths ranging from 6.0 to 8.5 Hz (eTable 1). However, the spectral linewidths were slightly broader in the PFC (mean linewidths ranging from 9.5 to 10.6 Hz) due to proximity of the frontal sinuses and nasal cavity, such that ADC data from 8 young and 14 older adults were excluded from further analysis.

At null b -value, similar spectral patterns with comparable SNR were observed between the young and older groups. At high b -value, even though the diffusion gradient strengths were identical in both cohorts, the spectral amplitudes were lower in older adults than young adults suggesting that ADC values of metabolites are higher in the older adult brain. The effects of cross-terms across the three diffusion directions are visible in the singlet peaks in the high b -value spectra (Figure 1).

The measured ADC values for five major metabolites and water for young and older adults are reported in Table 1. No gender differences were observed between the cohorts. The ADC values for tNAA, tCr, tCho and mIns were significantly higher ($P < 0.0033$) in older compared young adults in the PCC, and the ADC were higher between 8% to 19%. Similarly, in the OCC, only tCr had significantly higher ADC values (8%, $P = 0.001$) in

older adults. However, no significant difference in ADC values was observed for the five major metabolites in the PFC.

The ADC values of tissue water in all three brain regions were found to be 3.6 to 6.8% higher in older adults with the largest difference observed in the PCC (Table 1).

T_2 of metabolites and water

LASER spectra acquired from one young and one older subject at different T_{ES} in the PCC are shown in Figure 2. High quality spectra were consistently measured within and between subjects similar to the ADC measurements. Comparable water linewidths as observed in the ADC measurements were found in the PCC and the OCC, while in the PFC, the water linewidth was slightly broader. Based on the water linewidth > 10 Hz criterion, 9 young and 13 older T_2 datasets in the PFC region were excluded from further analysis.

Figure 2 shows a noticeable difference in the peak heights of tNAA and tCr, especially visible at long echo times (> 250 ms), suggesting that the apparent T_2 s of these metabolites are different between two cohorts.

Contrary to higher ADC values observed for some metabolites in the older adults, the apparent T_2 relaxation times of metabolites were found to be shorter in older adults (Table 2). No gender differences were observed between the cohorts. In the PCC, apparent T_2 of tNAA, tCr-CH₃, tCr-CH₂, tCho, and mIns were significantly lower ranging from 5% lower tCr-CH₃ to 11% lower NAA and tCr-CH₂. Similarly, in the OCC, significantly lower apparent T_2 values ($P < 0.0027$) were observed for tNAA, tCr-CH₃, tCr-CH₂, and mIns in the older adults' group, while Glu and tCho did not show any significant difference between the two groups.

The apparent T_2 of tissue water was significantly faster in the older cohort in all three regions studied (Table 2). On the other hand, the apparent T_2 of water in CSF was significantly faster in the young cohort in all three regions (Table 2).

Tissue content in VOIs

The VOI sizes for the regions studied were similar between the two age groups. Due to atrophy, the VOIs in the older adults contained a higher amount of CSF and a lower amount of GM (Figure 3). Since a larger volume of the VOI was occupied by CSF, in which there are no metabolites except glucose and lactate, the contribution of metabolites coming from GM and WM was different between the groups. In older adults, metabolites coming from WM contributed more signal than in younger adults since WM content was higher by 14% in PCC, 8% in OCC, and 13% in PFC in older adults.

Discussion

The present study shows that region-specific differences occur in the intracellular microenvironment in older vs. younger human brains. Mainly, the diffusivity of mIns, tCr and tissue water were significantly higher in the older healthy adults than in young healthy adults. The apparent T_2 relaxation time constants of neurochemicals and tissue water were

significantly shorter in the older cohort. These differences in ADC and T_2 metrics were region-specific, i.e., PCC was the most sensitive region in both measurements, followed by OCC and then PFC.

The diffusivity of several neurochemicals and tissue water was faster in the healthy older vs. younger brains. This is the first study reporting an increase in the ADC values for the major metabolites in older adults and is in disagreement with a previous study (Zheng et al., 2012) where a decrease in ADC of at least 25% was observed for NAA, tCr, and tCho. This discrepancy in metabolites' diffusivities might be related to several differences between the studies, such as the echo time, diffusion times, and VOI selection. The VOI used in the previous study contained mostly WM, while in the present study all VOIs contained a mixture of WM and GM. Another possibility is that the cross-term effects due to the gradients were not taken into account in the previous work, which might bias the ADC measurements. The increase in ADC of tissue water observed in the older adults in our study is in agreement with literature values obtained previously using DWI (Klimas et al., 2013; Watanabe et al., 2013). It is known that during aging there is a reduction in total dendritic measure (e.g., length, surface area, densities, spine numbers and branches) (Morrison and Hof, 2003). In addition, it was recently reported using diffusion simulation with data from mouse and macaque brains that a reduction in total dendritic measure would result in decreased ADC values of metabolites (Palombo et al., 2016; Palombo et al., 2017), i.e. more restricted diffusion. In contrast, ADC values increase (Palombo et al., 2018a) with a loss in dendritic spines only (Dickstein et al., 2013) where the spine space accounts for a small fraction of the total cellular volumes explored by the diffusing metabolites. Interestingly, there is a major reduction in spine densities during aging of the brain (Benavides-Piccione et al., 2012). Based on these findings, we hypothesize that the increase in ADC could in part be attributed to the large spine density decrease with age compared to the other dendritic measures (Palombo et al., 2020; Palombo et al., 2016; Palombo et al., 2017).

Another possible factor which might explain the increase in ADC findings could be related to the loss of mitochondria (López-Otín et al., 2013) and demyelination (Bennett et al., 2010) with aging. Recently, it has been shown that a small fraction (~5 to 10%) of the NAA pool is confined in a highly restricted compartments (Palombo et al., 2017), e.g., mitochondrial or myelin pool. During the normal aging process, the relative volume fraction of these restricting structures might decrease e.g., cellular aging leads to loss of mitochondria while demyelination leads to the thinning of the myelin sheath, such that both processes release more NAA, which diffuses freely in the intra-cellular space. Hence, the reduction in volume fraction will lead to an increase in the ADC of NAA consistent with the simulation findings observed in animals brain (Palombo et al., 2017). The higher ADC of mIns with age could be associated with higher relative number of glial cells (Cotrina and Nedergaard, 2002).

The observed age-related differences in ADC values are not readily attributable to differences in the VOI composition between young and older adults. The measured ADC values are averaged values of metabolites contained in GM and WM since CSF does not contain the investigated metabolites. In all VOIs, the percentage of GM was lower and the percentage of WM was higher in older adults than in young adults. In PCC, for instance,

GM is 5% lower and WM 5% higher in older adults relative to young adults. If known ADC values of neurochemicals in GM and WM (Kan et al., 2012; Najac et al., 2016) were used, the differences in ADC due only to difference in VOI composition would be relatively small. For instance, ADC of NAA in PCC would only be 2% higher in older group compared to young group. However, the measured ADC of NAA was 8% higher in the older group (Table 1). As such, difference in tissue composition could only account for a small contribution to the differences in ADC values. Whether age-related ADC differences arise more from GM or WM is beyond the scope of the current study.

T_2 values of metabolites and tissue water were faster and region-specific in the older brains. These findings are consistent with two previous studies (Kirov et al., 2008; Marja ska et al., 2013) which reported shorter apparent T_2 relaxation times for NAA, tCr, and tCho with comparable age-span to the young and older cohorts used in the present study. Similarly, shorter apparent T_2 value for mIns was observed in our older cohort. It was possible to measure this J -coupled metabolite due to the optimized T_E , high SNR of the spectra, and by taking into account the J -modulation at each T_E during the spectral fitting (Deelchand et al., 2018a). Since mIns is exclusively located in the astrocytes, the age-related difference in T_2 of mIns can be used to monitor metabolic change due to astrogliosis in the aging brain (Cotrina and Nedergaard, 2002).

The shorter T_2 values in older brains might be caused by several factors. This could be due to a change in iron content in the brain, since a strong correlation between iron deposition in the human brain and age exists (House et al., 2007; Mitumori et al., 2009). Another factor affecting T_2 relaxation could be a reduction of water content in the older adult brain mostly due to atrophy which in turn results in neuronal shrinkage (Chang et al., 1996; Dickstein et al., 2007). Myelin degeneration with age might also account for the loss in T_2 values in late life. The identification of the exact cause of the shorter T_2 in the older brain is beyond the scope of this paper. In our study, the apparent T_2 measured could be affected by the differences in the tissue composition between two groups since T_2 of metabolites differ between GM and WM (Mlynarik et al., 2001).

PCC was found to be the most sensitive region for detecting differences in cellular microenvironment with both the ADC and T_2 measurements. This was followed by the OCC and PFC. The fact that PCC shows more advanced differences with age agrees with our recent work where significant neurochemical concentration differences were observed in PCC compared to OCC (Marja ska et al., 2017). PCC is the central hub in the default mode network and is connected to PFC, and decrease in functional connectivity was shown to occur between both regions during aging (Andrews-Hanna et al., 2007; Hafkemeijer et al., 2012). These findings suggest that since different modalities examine different aspects of neurochemicals and water tissue in the brain, it is not straightforward to interpret and link the data together.

The apparent T_2 relaxation time data were found to be more sensitive than the ADC data. This is reflected by the fact that more significant differences in T_2 were observed between metabolites in the studied regions compared to the diffusivity data. One possible explanation is the low number of b -values used in the current study. Using a larger range of b -values

might minimize the bias and increase the accuracy in measuring ADC values of metabolites as previously demonstrated in T_2 measurements (Brief et al., 2005). However, this potentially means longer scan time in order to have a reasonable spectral SNR at different b -values when using conventional diffusion sequences with cross-term effects. Another possibility is to use cross-term free sequences (Ligneul et al., 2017) to reduce scan time.

Due to the imposed linewidth criteria of 10 Hz (lower data quality), a small sample size was used in the analysis of the PFC region. This has resulted in not finding significant age-dependent differences in the cellular microenvironment of neurochemicals in PFC between the young and older groups. PFC is known to be a highly vulnerable region to aging based on cellular changes compared to PCC and OCC (Fjell and Walhovd 2010). In addition, the normal brain ages from the front to the back of the head as evident from previous volumetric MRI (Fjell et al., 2009; Mrak et al., 1997), connectivity maps (Dennis and Thompson, 2014), and glucose uptake measurements (Zuendorf et al., 2003). However, substantial degradation of the spectral linewidth and therefore compromised SNR was observed in most of the acquired PFC data compared to PCC and OCC data. PFC is particularly susceptible to large B_0 variations due to the proximity of the frontal sinuses and nasal cavity. To fully compensate for these B_0 inhomogeneities, higher (2nd and 3rd) order shim corrections are required (Koch et al., 2009). However, commercially available clinical scanners as used in the current study are limited to only having 2nd order shims and therefore the B_0 inhomogeneities could not be fully compensated in PFC. We hope that high quality PFC spectra, consistent with current PCC and OCC data quality, will be possible with the availability of higher order shimming hardware in the near future.

Conclusion

Intracellular microenvironment difference of neurochemicals in young vs. older normal human brains are region-specific as reflected by the ADC and T_2 data. The diffusivity of several metabolites were higher while their apparent T_2 relaxation time constants were shorter in older adults. In conclusion, this study shows that it is feasible to detect differences in the cellular microenvironment of neurochemicals in the normal aging brain and opens up the possibility to investigate the aging process over time.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors would like to thank: Andrew Oliver and Sarah Bedell for study coordination, Akshay Patke for neuropsychological testing, and Emily Kittelson and Andrea Grant, Ph.D. for technical support.

Funding

This work was supported by funding from the National Institutes of Health grants (R21AG045606, P41 EB015894, P30 NS076408).

Abbreviations

ADC	apparent diffusion coefficients
Glu	glutamate
mIns	<i>myo</i> -inositol
NAA	<i>N</i> -acetyl-aspartate
OCC	occipital cortex
PCC	posterior cingulate cortex
PFC	prefrontal cortex
RF	radiofrequency
T₂	transverse relaxation
tCr	total creatine (i.e. creatine plus phosphocreatine)
tCho	total choline (i.e. phosphorylcholine plus glycerophosphorylcholine)
VOI	volumes-of-interest

References

- Andrews-Hanna JR, Snyder AZ, Vincent JL, Lustig C, Head D, Raichle Marcus E., Buckner RL, 2007 Disruption of Large-Scale Brain Systems in Advanced Aging. *Neuron* 56(5), 924–935. [PubMed: 18054866]
- Benavides-Piccione R, Fernaud-Espinosa I, Robles V, Yuste R, DeFelipe J, 2012 Age-Based Comparison of Human Dendritic Spine Structure Using Complete Three-Dimensional Reconstructions. *Cerebral Cortex* 23(8), 1798–1810. [PubMed: 22710613]
- Bennett IJ, Madden DJ, Vaidya CJ, Howard DV, Howard JH Jr, 2010 Age-related differences in multiple measures of white matter integrity: A diffusion tensor imaging study of healthy aging. *Human Brain Mapping* 31(3), 378–390. [PubMed: 19662658]
- Benton A, Hamsher K, Sivan AB, 1994 Multilingual Aphasia Examination. Iowa City: AJA Associates;.
- Brief EE, Whittall KP, Li DK, MacKay AL, 2005 Proton T₂ relaxation of cerebral metabolites of normal human brain over large TE range. *NMR Biomed* 18(1), 14–18. [PubMed: 15455460]
- Chang L, Ernst T, Poland RE, Jenden DJ, 1996 In vivo proton magnetic resonance spectroscopy of the normal aging human brain. *Life Sciences* 58(22), 2049–2056. [PubMed: 8637436]
- Christensen KJ, Moye J, Armson RR, Kern TM, 1992 Health screening and random recruitment for cognitive aging research. *Psychology and Aging* 7(2), 204–208. [PubMed: 1610509]
- Christiansen P, Henriksen O, Stubgaard M, Gideon P, Larsson HBW, 1993 In vivo quantification of brain metabolites by 1H-MRS using water as an internal standard. *Magn Reson Imaging* 11(1), 107–118. [PubMed: 8423713]
- Cotrina ML, Nedergaard M, 2002 Astrocytes in the aging brain. *Journal of Neuroscience Research* 67(1), 1–10. [PubMed: 11754075]
- de Graaf RA, 2007 In Vivo NMR Spectroscopy. John Wiley & Sons, Ltd.
- Deelchand DK, Auerbach EJ, Kobayashi N, Marja ska M, 2018a Transverse relaxation time constants of the five major metabolites in human brain measured in vivo using LASER and PRESS at 3 T. *Magn Reson Med* 79(3), 1260–1265. [PubMed: 28691380]

- Deelchand DK, Auerbach EJ, Marja ska M, 2018b Apparent diffusion coefficients of the five major metabolites measured in the human brain in vivo at 3T. *Magn Reson Med* 79(6), 2896–2901. [PubMed: 29044690]
- Delis DC, Kramer JH, Kaplan E, Ober BA, 2000 California Verbal Learning Test. San Antonio, TX: Psychological Corporation.
- Dennis EL, Thompson PM, 2014 Functional Brain Connectivity Using fMRI in Aging and Alzheimer's Disease. *Neuropsychology Review* 24(1), 49–62. [PubMed: 24562737]
- Dickstein DL, Kabaso D, Rocher AB, Luebke JI, Wearne SL, Hof PR, 2007 Changes in the structural complexity of the aged brain. *Aging Cell* 6(3), 275–284. [PubMed: 17465981]
- Dickstein DL, Weaver CM, Luebke JI, Hof PR, 2013 Dendritic spine changes associated with normal aging. *Neuroscience* 251, 21–32. [PubMed: 23069756]
- Fjell AM, Walhovd KB, 2010 Structural Brain Changes in Aging: Courses, Causes and Cognitive Consequences. *Reviews in the Neurosciences* 21(3), 187. [PubMed: 20879692]
- Fjell AM, Walhovd KB, Fennema-Notestine C, McEvoy LK, Hagler DJ, Holland D, Brewer JB, Dale AM, 2009 One-Year Brain Atrophy Evident in Healthy Aging. *The Journal of Neuroscience* 29(48), 15223. [PubMed: 19955375]
- Golden C, 1978 Stroop Experimental Uses. Chicago: Stoelting.
- Govind V, Young K, Maudsley AA, 2015 Corrigendum: Proton NMR chemical shifts and coupling constants for brain metabolites. Govindaraju V, Young K, Maudsley AA, *NMR Biomed*. 2000; 13: 129–153. *NMR Biomed* 28(7), 923–924. [PubMed: 26094860]
- Grieve SM, Williams LM, Paul RH, Clark CR, Gordon E, 2007 Cognitive Aging, Executive Function, and Fractional Anisotropy: A Diffusion Tensor MR Imaging Study. *American Journal of Neuroradiology* 28(2), 226. [PubMed: 17296985]
- Grober E, Sliwinski M, Korey SR, 1991 Development and Validation of a Model for Estimating Premorbid Verbal Intelligence in the Elderly. *Journal of Clinical and Experimental Neuropsychology* 13(6), 933–949. [PubMed: 1779032]
- Hafkemeijer A, van der Grond J, Rombouts SARB, 2012 Imaging the default mode network in aging and dementia. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* 1822(3), 431–441. [PubMed: 21807094]
- Henry PG, Marja ska M, Walls JD, Valette J, Gruetter R, Ugurbil K, 2006 Proton-observed carbon-edited NMR spectroscopy in strongly coupled second-order spin systems. *Magn Reson Med* 55(2), 250–257. [PubMed: 16402370]
- House MJ, St. Pierre TG, Kowdley KV, Montine T, Connor J, Beard J, Berger J, Siddaiah N, Shankland E, Jin L-W, 2007 Correlation of proton transverse relaxation rates (R2) with iron concentrations in postmortem brain tissue from alzheimer's disease patients. *Magn Reson Med* 57(1), 172–180. [PubMed: 17191232]
- Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM, 2012 FSL. *NeuroImage* 62(2), 782–790. [PubMed: 21979382]
- Kan HE, Techawiboonwong A, van Osch MJP, Versluis MJ, Deelchand DK, Henry P-G, Marja ska M, van Buchem MA, Webb AG, Ronen I, 2012 Differences in apparent diffusion coefficients of brain metabolites between grey and white matter in the human brain measured at 7 T. *Magn Reson Med* 67(5), 1203–1209. [PubMed: 22083562]
- Kaplan E, Goodglass H, Weintraub S, 1983 The Boston Naming Test. Philadelphia: Lea and Febiger;.
- Kirov II, Fleysher L, Fleysher R, Patil V, Liu S, Gonen O, 2008 Age dependence of regional proton metabolites T2 relaxation times in the human brain at 3 T. *Magn Reson Med* 60(4), 790–795. [PubMed: 18816831]
- Klimas A, Drzazga Z, Kluczevska E, Hartel M, 2013 Regional ADC measurements during normal brain aging in the clinical range of b values: a DWI study. *Clinical Imaging* 37(4), 637–644. [PubMed: 23462734]
- Koch KM, Rothman DL, de Graaf RA, 2009 Optimization of static magnetic field homogeneity in the human and animal brain in vivo. *Progress in Nuclear Magnetic Resonance Spectroscopy* 54(2), 69–96. [PubMed: 20126515]

- Kreis R, Slotboom J, Hofmann L, Boesch C, 2005 Integrated data acquisition and processing to determine metabolite contents, relaxation times, and macromolecule baseline in single examinations of individual subjects. *Magn Reson Med* 54(4), 761–768. [PubMed: 16161114]
- Ligneul C, Palombo M, Hernández-Garzón E, Carrillo-de Sauvage M-A, Flament J, Hantraye P, Brouillet E, Bonvento G, Escartin C, Valette J, 2019 Diffusion-weighted magnetic resonance spectroscopy enables cell-specific monitoring of astrocyte reactivity in vivo. *NeuroImage* 191, 457–469. [PubMed: 30818026]
- Ligneul C, Palombo M, Valette J, 2017 Metabolite diffusion up to very high b in the mouse brain in vivo: Revisiting the potential correlation between relaxation and diffusion properties. *Magn Reson Med* 77(4), 1390–1398. [PubMed: 27018415]
- López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G, 2013 The Hallmarks of Aging. *Cell* 153(6), 1194–1217. [PubMed: 23746838]
- Marja ska M, Deelchand DK, Hodges JS, McCarten JR, Hemmy LS, Grant A, Terpstra M, 2018 Altered macromolecular pattern and content in the aging human brain. *NMR in Biomedicine* 31(2), e3865.
- Marja ska M, Emir UE, Deelchand DK, Terpstra M, 2013 Faster Metabolite 1H Transverse Relaxation in the Elder Human Brain. *PLOS ONE* 8(10), e77572. [PubMed: 24098589]
- Marja ska M, McCarten JR, Hodges J, Hemmy LS, Grant A, Deelchand DK, Terpstra M, 2017 Region-specific aging of the human brain as evidenced by neurochemical profiles measured noninvasively in the posterior cingulate cortex and the occipital lobe using 1H magnetic resonance spectroscopy at 7 T. *Neuroscience* 354, 168–177. [PubMed: 28476320]
- McIntyre DJO, Charlton RA, Markus HS, Howe FA, 2007 Long and short echo time proton magnetic resonance spectroscopic imaging of the healthy aging brain. *J Magn Reson Imaging* 26(6), 1596–1606. [PubMed: 17968966]
- Meyers JE, Meyers KR, 1995 Rey Complex Figure Test and Recognition Trial: Professional Manual. Lutz, FL: Psychological Assessment Resources.
- Mitsumori F, Watanabe H, Takaya N, 2009 Estimation of brain iron concentration in vivo using a linear relationship between regional iron and apparent transverse relaxation rate of the tissue water at 4.7T. *Magn Reson Med* 62(5), 1326–1330. [PubMed: 19780172]
- Mlynarik V, Gruber S, Moser E, 2001 Proton T₁ and T₂ relaxation times of human brain metabolites at 3 Tesla. *NMR Biomed* 14(5), 325–331. [PubMed: 11477653]
- Morrison JH, Hof PR, 2003 Changes in cortical circuits during aging. *Clinical Neuroscience Research* 2(5), 294–304.
- Mrak RE, Griffin WST, Graham DI, 1997 Aging-associated Changes in Human Brain. *Journal of Neuropathology & Experimental Neurology* 56(12), 1269–1275. [PubMed: 9413275]
- Najac C, Branzoli F, Ronen I, Valette J, 2016 Brain intracellular metabolites are freely diffusing along cell fibers in grey and white matter, as measured by diffusion-weighted MR spectroscopy in the human brain at 7 T. *Brain Struct Funct* 221(3), 1245–1254. [PubMed: 25520054]
- Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, Cummings JL, Chertkow H, 2005 The Montreal Cognitive Assessment, MoCA: A Brief Screening Tool For Mild Cognitive Impairment. *Journal of the American Geriatrics Society* 53(4), 695–699. [PubMed: 15817019]
- Öngür D, Prescott AP, Jensen JE, Rouse ED, Cohen BM, Renshaw PF, Olson DP, 2010 T2 relaxation time abnormalities in bipolar disorder and schizophrenia. *Magn Reson Med* 63(1), 1–8. [PubMed: 19918902]
- Palombo M, Ianus A, Guerreri M, Nunes D, Alexander DC, Shemesh N, Zhang H, 2020 SANDI: A compartment-based model for non-invasive apparent soma and neurite imaging by diffusion MRI. *NeuroImage* 215, 116835. [PubMed: 32289460]
- Palombo M, Ligneul C, Hernandez-Garzon E, Valette J, 2018a Can we detect the effect of spines and leaflets on the diffusion of brain intracellular metabolites? *NeuroImage* 182, 283–293.
- Palombo M, Ligneul C, Najac C, Le Douce J, Flament J, Escartin C, Hantraye P, Brouillet E, Bonvento G, Valette J, 2016 New paradigm to assess brain cell morphology by diffusion-weighted MR spectroscopy in vivo. *Proceedings of the National Academy of Sciences* 113(24), 6671.

- Palombo M, Ligneul C, Valette J, 2017 Modeling diffusion of intracellular metabolites in the mouse brain up to very high diffusion-weighting: Diffusion in long fibers (almost) accounts for non-monoexponential attenuation. *Magnetic Resonance in Medicine* 77(1), 343–350. [PubMed: 27851876]
- Palombo M, Shemesh N, Ronen I, Valette J, 2018b Insights into brain microstructure from in vivo DW-MRS. *NeuroImage* 182, 97–116. [PubMed: 29155183]
- Peters R, 2006 Ageing and the brain. *Postgraduate Medical Journal* 82(964), 84. [PubMed: 16461469]
- Provencher SW, 1993 Estimation of metabolite concentrations from localized in vivo proton NMR spectra. *Magn Reson Med* 30(6), 672–679. [PubMed: 8139448]
- Reitan RM, 1979 Manual for Administration of Neuropsychological Test Batteries for Adults and Children. Tucson, AZ: Neuropsychology Press.
- Ronen I, Valette J, 2015 Diffusion-Weighted Magnetic Resonance Spectroscopy, eMagRes. John Wiley & Sons, Ltd, pp. 733–750.
- Valette J, Ligneul C, Marchadour C, Najac C, Palombo M, 2018 Brain Metabolite Diffusion from Ultra-Short to Ultra-Long Time Scales: What Do We Learn, Where Should We Go? *Frontiers in Neuroscience* 12, 2. [PubMed: 29403347]
- van der Kouwe AJW, Benner T, Fischl B, Schmitt F, Salat DH, Harder M, Sorensen AG, Dale AM, 2005 On-line automatic slice positioning for brain MR imaging. *NeuroImage* 27(1), 222–230. [PubMed: 15886023]
- Walhovd KB, Westlye LT, Amlie I, Espeseth T, Reinvang I, Raz N, Agartz I, Salat DH, Greve DN, Fischl B, Dale AM, Fjell AM, 2011 Consistent neuroanatomical age-related volume differences across multiple samples. *Neurobiology of Aging* 32(5), 916–932. [PubMed: 19570593]
- Watanabe M, Sakai O, Ozonoff A, Kussman S, Jara H, 2013 Age-related Apparent Diffusion Coefficient Changes in the Normal Brain. *Radiology* 266(2), 575–582. [PubMed: 23143020]
- Wechsler D, 1987 Wechsler Memory Scale-Revised. San Antonio, TX: Psychological Corporation/Harcourt Brace Jovanovich.
- Wechsler D, 1997 Wechsler Adult Intelligence Scale-Third Edition. San Antonio, TX: The Psychological Corporation.
- Whittall KP, MacKay AL, Graeb DA, Nugent RA, Li DK, Paty DW, 1997 In vivo measurement of T₂ distributions and water contents in normal human brain. *Magn Reson Med* 37(1), 34–43. [PubMed: 8978630]
- Zheng DD, Liu ZH, Fang J, Wang XY, Zhang J, 2012 The Effect of Age and Cerebral Ischemia on Diffusion-Weighted Proton MR Spectroscopy of the Human Brain. *American Journal of Neuroradiology* 33(3), 563. [PubMed: 22081680]
- Zuendorf G, Kerrouche N, Herholz K, Baron J-C, 2003 Efficient principal component analysis for multivariate 3D voxel-based mapping of brain functional imaging data sets as applied to FDG-PET and normal aging. *Human Brain Mapping* 18(1), 13–21. [PubMed: 12454908]

Highlights:

- Diffusivity of neurochemicals is faster in older adults
- Apparent transverse relaxation times of neurochemicals are faster in older adults
- Changes in ADC and T_2 are region-dependent
- Feasible to detect changes in the cellular microenvironment in normal aging brain
- ADC and T_2 are sensitive to changes in the cellular microenvironment during aging

Figure 1.

Diffusion-weighted spectra from one young (20 year-old female) adult and one older (70 year-old female) adult acquired in the PCC using STEAM ($T_R/T_E = 3000/21.2$ ms) at 3 T. Spectra acquired at null b -value (16 averages, top) and at high b -value in all six directions (48 averages per direction, bottom) are shown. Note the excellent spectral quality, with no lipid contamination, flat baseline, and the high SNR at the high b -value.

Figure 2.

LASER spectra ($T_R = 3$ s with different number of averages, NA) acquired at six T_E s from one young (22 F) adult and one older (73 F) adult in the PCC at 3 T. For display purposes, the spectra were normalized such that at the shortest T_E , the height of NAA peak was identical. The dashed lines show that the apparent T_2 of NAA and other singlets are shorter in older adult brain.

Figure 3.

Means and standard deviations of WM, GM and CSF content measured in three VOIs from T_1 -weighted images acquired during the T_2 measurements. * represents statistically significant differences ($P < 0.001$) between young and older adults.

Trace/3 ADC values (mean $\pm$ SD) of metabolites and tissue water in the two age groups in three brain regions. Bold values denote statistical significance, i.e. $p < 0.0033$ for metabolites and $p < 0.0167$ for water.

Compound	Trace/3 ADC ($\mu\text{m}^2/\text{ms}$)							
	PCC			OCC			PFC	
	Young	Older	<i>p</i> -value	Young	Older	<i>p</i> -value	Young	Older
tNAA	0.12 ± 0.01	0.13 ± 0.02	0.0014	0.14 ± 0.01	0.14 ± 0.02	0.4573	0.13 ± 0.01	0.13 ± 0.03
tCr	0.12 ± 0.01	0.14 ± 0.02	0.0008	0.13 ± 0.01	0.14 ± 0.01	0.0010	0.13 ± 0.01	0.13 ± 0.03
tCho	0.11 ± 0.01	0.13 ± 0.03	0.0019	0.11 ± 0.02	0.13 ± 0.02	0.0242	0.11 ± 0.04	0.13 ± 0.05
Glu	0.14 ± 0.02	0.15 ± 0.03	0.0133	0.15 ± 0.02	0.15 ± 0.02	0.7391	0.14 ± 0.02	0.13 ± 0.05
mIns	0.12 ± 0.01	0.13 ± 0.02	0.0011	0.12 ± 0.01	0.13 ± 0.02	0.0329	0.12 ± 0.04	0.13 ± 0.05
Tissue water	0.65 ± 0.02	0.69 ± 0.04	4.5 × 10⁻⁷	0.62 ± 0.03	0.64 ± 0.03	0.0039	0.69 ± 0.03	0.72 ± 0.04

T_2 values (mean $\pm$ SD) of metabolites, tissue water and water in CSF measured in the two age groups in three brain regions. Bold values denote statistical significance, i.e. $p < 0.0027$ for metabolites and $p < 0.0083$ for water.

Table 2

Compound	T_2 (ms)									
	PCC				OCC				PFC	
	Young	Older	p -value	Young	Older	p -value	Young	Older	p -value	
NAA singlet	322 $\pm$ 16	286 $\pm$ 14	9.8×10^{-16}	304 $\pm$ 13	279 $\pm$ 17	5.4×10^{-8}	355 $\pm$ 17	316 $\pm$ 25	1.3×10^{-6}	
tCr-CH ₃	183 $\pm$ 7	174 $\pm$ 7	7.3×10^{-7}	178 $\pm$ 6	171 $\pm$ 7	0.0004	194 $\pm$ 7	198 $\pm$ 11	0.1876	
tCr-CH ₂	148 $\pm$ 7	132 $\pm$ 7	6.1×10^{-14}	140 $\pm$ 6	129 $\pm$ 6	1.7×10^{-10}	158 $\pm$ 9	144 $\pm$ 11	0.0004	
tCho	360 $\pm$ 40	332 $\pm$ 20	0.0011	323 $\pm$ 25	314 $\pm$ 30	0.3156	442 $\pm$ 47	423 $\pm$ 33	0.2605	
Glu	177 $\pm$ 10	171 $\pm$ 8	0.0266	177 $\pm$ 10	168 $\pm$ 13	0.0040	194 $\pm$ 11	190 $\pm$ 24	0.5993	
mIns	206 $\pm$ 14	195 $\pm$ 10	0.0007	184 $\pm$ 12	174 $\pm$ 11	0.0018	235 $\pm$ 19	227 $\pm$ 17	0.0900	
Tissue water	62 $\pm$ 3	53 $\pm$ 6	1.4×10^{-8}	56 $\pm$ 3	50 $\pm$ 6	2.7×10^{-5}	65 $\pm$ 2	56 $\pm$ 4	2.3×10^{-13}	
CSF water	250 $\pm$ 32	308 $\pm$ 102	0.0082	225 $\pm$ 28	271 $\pm$ 34	5.0×10^{-7}	280 $\pm$ 31	325 $\pm$ 43	0.0004	