CA, USA, ¹²University of Washington, Seattle, WA, USA, ¹³Children's National, Washington, DC, USA, ¹⁴Northwestern University, Chicago, IL, USA, ¹⁵Brigham and Women's Hospital, Boston, MA, USA, ¹⁶New York Presbyterian Hospital/Columbia University Irving Medical Center, New York, NY, USA, ¹⁷Dana-Farber Cancer Institute, Boston, MA, USA

H3 K27M-mutant diffuse midline gliomas (DMG) have a dismal prognosis. We report an integrated analysis for ONC201, a DRD2 antagonist and ClpP agonist, in patients with recurrent H3 K27M DMG administered as monotherapy in 3 clinical studies (NCT03295396, n=20; NCT02525692, n=6; or expanded access, n=4) by independent, central radiology review among 30 patients. At baseline, patients had measurable contrast-enhancing disease by RANO criteria, KPS>60, and were >90 days from prior radiation. Patients with primary lesions that involved the pons or spinal cord or had evidence of leptomeningeal or cerebrospinal fluid dissemination were excluded. ONC201 was orally administered at 625 mg (scaled by body weight for patients < 18 years old) weekly in 29 patients and every 3 weeks in 1 patient. Median age was 31 years (range 8–70; two patients < 18). There were 16 women and 14 men. Tumors were predominantly thalamic (73%), with other locations including the cerebellum (5%), brainstem (non-pons) (3%), basal ganglia (3%), and midbrain (3%). Median time from prior radiation was 7.5 months. The most frequent drug-related adverse events were low grade nausea (10%) and fatigue (10%). Nine patients (30%) had >50% regression of T1 contrast enhancement and 11 (36.7%) patients had regression of T2/FLAIR. Objective response rate and progression-free survival by RANO criteria, as well overall survival will be reported. Among 8 patients with sustained radiographic regressions, 6 were tapered off dexamethasone and 4 had improvement in KPS. Five patients (16.7%) remain on treatment with a median of 9.5 months (range 7.3–12). In conclusion, single agent ONC201 is well tolerated and clinically active in recurrent H3 K27M DMG patients.

CTNI-38. PAMIPARIB IN COMBINATION WITH RADIATION THERAPY (RT) AND/OR TEMOZOLOMIDE (TMZ) IN PATIENTS WITH NEWLY DIAGNOSED (ND) OR RECURRENT/REFRACTORY (R/R) GLIOBLASTOMA (GBM); PHASE 1B/2 STUDY UPDATE

Anna Piotrowski¹, Vinay Puduvalli², Patrick Wen³, Howard Colman⁴, Jian Campian⁵, Michael Pearlman⁶, Nicholas Butowski⁷, Timothy Cloughesy⁸, James Battiste⁹, Jon Glass¹⁰, David Schiff¹¹, Martin van den Bent¹², Tobias Walbert¹³, Manmeet Ahluwalia¹⁴, Michael Badruddoja¹⁵, Amandeep Kalra¹⁶, Robert Pelham¹⁷, Kathy Zhang¹⁷, Katie Wood¹⁷, Michael Weller¹⁸, and Kent Shih¹⁹; ¹Memorial Sloan Kettering Cancer Center, New York, NY, USA, ²The Ohio State University, Columbus, OH, USA, ³Dana-Farber Cancer Institute, Boston, MA, USA, ⁴Huntsman Cancer Institute, Salt Lake City, UT, USA, ⁵Washington University School of Medicine, St. Louis, MO, USA, ⁶Sarah Cannon Research Institute at Health One, Denver, CO, USA, ⁷University of California at San Francisco, San Francisco, CA, USA, ⁸University of California at Los Angeles, Neuro-Oncology, Los Angeles, CA, USA, ⁹Stephenson Cancer Center at the University of Oklahoma, Oklahoma City, OK, USA, ¹⁰Thomas Jefferson University, Philadelphia, PA, USA, ¹¹University of Virginia Health Systems, Emory Clinic Clinical Cancer Center, Charlottesville, VA, USA, ¹²Erasmus MC Cancer Institute, Rotterdam, Netherlands, ¹³Henry Ford Hospital, Detroit, MI, USA, ¹⁴Cleveland Clinic, Cleveland, OH, USA, ¹⁵Center for Neurosciences, Tucson, AZ, USA, ¹⁶Health Midwest Ventures Group, LLC, Kansas City, MO, USA, ¹⁷BeiGene USA, Inc., San Mateo, CA, USA, ¹⁸UniversitätsSpital Zürich - Klinik für Neurologie, Zurich, Switzerland, ¹⁹Sarah Cannon Research Institute, Tennessee Oncology, Nashville, TN, USA

Pamiparib, an investigational, oral PARP 1/2 inhibitor, demonstrated pre-clinical brain penetration and synergistic cytotoxicity with TMZ. We report updated safety and antitumor data for pamiparib plus RT and/or TMZ in ND-GBM or R/R-GBM (SNO 2019, ACTR-39). This dose-escalation/expansion study includes three arms: A, pamiparib (2, 4, or 6 weeks) plus RT (6–7 weeks) in ND-GBM with unmethylated MGMT promoter (unmethylated-GBM); B, pamiparib (6 weeks) plus RT and increasing TMZ doses in Weeks 1 and 5 of RT in unmethylated ND-GBM; and C, pamiparib plus increasing TMZ doses in methylated/unmethylated R/R-GBM. Most patients in Arms A (expansion) and B received maintenance pamiparib plus TMZ after post-RT rest period at Arm C expansion. As of April 10, 2020, enrollment was complete (N=116; A, n=60; B, n=9; C, n=47). Median study follow-up was 11.3 mo (A/B) and 7.1 mo (C). Common grade ≥3 AEs were anemia (10%) in Arm A; decreased neutrophil and white blood cell count (each 22%) in B; anemia, fatigue, and decreased lymphocyte count (each 11%) in C. Brain edema (A) and pneumonia (C) (n=1 each) were fatal treatment-unrelated AEs. In ND-GBM, modified disease control rate (DCR following post-RT rest period) was 69.8% (95% CI, 55.7–81.7) overall, 68.8% (50.0–83.9) in A, and 80.0% (28.4–99.5) in B. Median duration of response was 5.1 mo (overall), 3.8 mo (A), and NE (B). In Arms A/B, median progression-free survival (PFS) and median overall survival (OS) were 4.4 mo and 12.7 mo, respectively; 12-mo OS rate, 54% (95% CI, 40–66). In R/R-GBM (Arm C),

confirmed ORR was 9.1% (95% CI, 2.5–21.7); median PFS and OS were 1.9 mo and 7.3 mo, respectively; 6-mo PFS rate, 19% (95% CI, 9–32). These results showed a manageable safety profile for pamiparib +/- RT +/-TMZ; response and survival results support further evaluation of these combinations in GBM.

CTNI-39. PHASE 1B CLINICAL TRIAL OF OKN-007 IN RECURRENT MALIGNANT GLIOMA

James Battiste¹, Alexandra Ikeguchi¹, Sukyung Woo², Satish Sharan³, Yan Zhao³, Andrew Cohoon³, Sarah Sung¹, Deborah Wright¹, April Teague¹, Randy Jensen⁴, Eun Ha Kim⁵, Won Yang⁵, and Rheel Townner⁶; ¹Stephenson Cancer Center at the University of Oklahoma, Oklahoma City, OK, USA, ²The State University of New York at Buffalo, Buffalo, NY, USA, ³University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA, ⁴University of Utah School of Medicine, Salt Lake City, UT, USA, ⁵Oblato, Inc., Princeton, NJ, USA, ⁶Oklahoma Medical Research Foundation, Oklahoma City, OK, USA

BACKGROUND: The nitrone compound OKN-007 is a novel anti-cancer agent. In glioblastoma xenografts, OKN-007 reduces cell proliferation and angiogenesis, and increases apoptosis. Here we report on the safety, efficacy, and pharmacokinetics (PK) of OKN-007 in adults with recurrent glioma. **METHODS:** NCT01672463 is a phase 1b trial of OKN-007 in adults with recurrent gliomas previously treated with standard therapy. OKN-007 was administered by IV. The study comprised a 3 + 3 dose escalation design followed by an expansion cohort at the maximum tolerated dose (MTD). The dose escalation drug levels were 20 (n = 3), 40 (n = 3), and 60 mg/kg (n = 3), treating on a schedule of thrice weekly for 4 weeks, then twice weekly for 4 weeks, then once weekly until progression. Drug PK was determined in the dose escalation cohorts. The expansion cohort was treated with 60 mg/kg thrice weekly for 12 weeks, then twice weekly for 12 weeks, then once weekly until progression (n = 6). Kaplan-Meier analysis was used to determine progression-free (PFS) and overall survival (OS). **RESULTS:** Median age was 51 years (range, 25–62). No dose-limiting toxicities were observed. The expansion dose was 60 mg/kg. Of 123 adverse events (AEs), 34 were deemed probably (1.6%) or possibly (26%) treatment-emergent (TEAE). The most commonly-occurring TEAEs were fatigue (4.1%) and headache (3.3%). Grade 3 TEAEs included headache, urinary tract infection, and increased prothrombin time (0.8% each). Two grade 1 AEs, hypokalemia and dizziness, were considered probably attributable to OKN-007. In patients receiving 60 mg OKN-007/kg, median PFS was 1.4 months and OS was 21 months (log rank p = 0.08 for dose comparison across doses). Systemic PK exposure was dose proportional. The average half-life of OKN-007 is 2.8 hours. **CONCLUSIONS:** OKN-007 appears safe and, compared to standard therapy, may prolong OS in recurrent glioma.

CTNI-41. PHASE I STUDY OF RUXOLITINIB WITH RADIATION AND TEMOZOLOMIDE IN PATIENTS WITH NEWLY DIAGNOSED GRADE III GLIOMAS AND GLIOBLASTOMA

Yasmeen Rauf¹, Rachel Hufsey², Kathy Robinson², John Suh², Samuel Chao², Erin Murphy², Jennifer Yu², David Peereboom², and Manmeet Ahluwalia²; ¹Cleveland Clinic, Dublin, OH, USA, ²Cleveland Clinic, Cleveland, OH, USA

BACKGROUND: Ruxolitinib is a novel, potent, and selective inhibitor of JAK1 (Janus kinase 1) and JAK2 with modest to marked selectivity against TYK2 (tyrosine kinase 2) and JAK3, respectively. Dysregulation of the JAK/STAT pathway has been associated with several types of cancer and increased proliferation and survival of malignant cells. **METHODS:** Newly diagnosed patients with unmethylated MGMT Glioblastoma or grade III glioma were recruited to Arm I. Every patient received ruxolitinib and 60 Gy radiation for 6 weeks over 6 weeks (2Gy x 30). The dose of Ruxolitinib was administered given the 3 + 3 design. Level 1 or starting dose was 10 mg twice daily, level 2 was 15 mg twice daily, level 3 was 20 mg twice daily and level -1 was 5 mg twice daily. Arm two was started once safe dose was established for Arm 1 for each dose level. Patients with methylated MGMT glioblastoma or grade III glioma were eligible for Arm 2. Every patient received ruxolitinib + radiation x 60 Gy + daily temozolomide at 75 mg/m² for 6 weeks over 6 weeks. **RESULTS:** 30 patients were randomized to Arm I and 16 to arm II. There was one partial response in the methylated Arm and one partial response in the unmethylated arm. The maximum tolerated dose of Ruxolitinib was 20 mg BID in both methylated and unmethylated arm. There were 14 incidences of Grade 3 toxicity including respiratory distress, seizure, gait disturbance, weakness, thrombocytopenia, cognitive disturbance, urinary retention, and bacterial meningitis. There were 4 incidences of Grade 4 toxicity including seizure, respiratory distress, somnolence and thromboembolic events. **CONCLUSION:** This is an ongoing clinical trial and final results will be presented once the trial is complete. We will perform single cell sequencing on good and poor responders, that data will be presented in the meeting.