Published in final edited form as:

Nat Prod Rep. 2020 November 01; 37(11): 1511–1531. doi:10.1039/d0np00060d.

The quest for supernatural products: The impact of total synthesis in complex natural products medicinal chemistry

Zhi-Chen Wu^a, Dale L. Boger^{a,*}

^aDepartment of Chemistry and The Skaggs Institute for Chemical Biology, The Scripps Research Institute, 10550 N. Torrey Pines Road, La Jolla, CA 92037, USA

Abstract

This review presents select recent advances in the medicinal chemistry of complex natural products that are prepared by total synthesis. The underlying studies highlight enabling divergent synthetic strategies and methods that permit the systematic medicinal chemistry studies of key analogues bearing deep-seated structural changes not readily accessible by semisynthetic or biosynthetic means. Select and recent examples are detailed where the key structural changes are designed to improve defined properties or to overcome an intrinsic limitation of the natural product itself. In the examples presented, the synthetic efforts provided supernatural products, a term first introduced by our colleague Ryan Shenvi (*Synlett*, 2016, **27**, 1145–1164), with properties superseding the parent natural product. The design principles and approaches for creating the supernatural products are highlighted with an emphasis on the properties addressed that include those that improve activity or potency, increase selectivity, enhance durability, broaden the spectrum of activity, improve chemical or metabolic stability, overcome limiting physical properties, add mechanisms of action, enhance PK properties, overcome drug resistance, and/or improve *in vivo* efficacy. Some such improvements may be regarded by some as iterative enhancements whereas others, we believe, truly live up to their characterization as supernatural products. Most such efforts are also accompanied by advances in synthetic organic chemistry, inspiring the development of new synthetic methodology and providing the supernatural products with improved synthetic accessibility.

Graphical Abstract

* dale.boger@outlook.com.

⁵.Conflicts of interest

There are no conflicts of interest to declare

This review summarizes and highlights recent advances in medicinal chemistry of natural products enabled by total synthesis that provide “supernatural products” with improved properties superseding the natural products themselves.

1. Introduction

The opportunities for natural products to address problems in biology and medicine have grown as the methods to identify their mechanisms of action and biological targets have improved and as the techniques used to directly determine their target bound structures have advanced.^{1–3} In addition to the natural compounds themselves and because of advances in organic synthesis, the examination of key synthetic partial structures and intermediates, compounds that contain deep-seated structural modifications, their unnatural diastereomers, and even their enantiomers have provided a powerful complement to such studies. Well-designed structural modifications in the natural products may be used to address the structural basis of their interaction with biological targets and to define fundamental relationships between structure and activity, functional reactivity, and biological properties.^{4–9} Because natural products integrate a constellation of properties into a highly functionalized, compact molecule, each structural feature and substituent are often thought to contribute to the biological activity. This is especially true when their biological properties are directly related to their emergence in nature where they have undergone natural selection optimization. However, we recently highlighted examples where even single atom changes to a series of natural products were found to substantially improve on their biological properties.¹⁰ In such work, a challenging problem with natural products is first to understand the subtle design elements that nature has provided and then to rationally extend them to provide more selective, more efficacious, or more potent compounds. Shenvi et al. first introduced the term “supernatural products” to describe structurally simplified asmarine analogues with improved properties that deprioritized the synthesis of the natural products themselves.¹¹ These efforts resulted in the generation of simple, potent analogues that surpassed the natural asmarines as tools for further biological examination. These “supernatural products”, delmarine in particular, provided the basis for discovery of the mechanism of action of the asmarines.¹² Although introduced with tongue and cheek humor just before Halloween by Shenvi, we have become enamoured by the power of the term “supernatural product” to succinctly describe such advances for not only the experts in the field but also for a general audience who may not appreciate the importance and magnitude of the accomplishments. Herein, we highlight select examples of supernatural products obtained through total synthesis. This restriction to selected examples of the many accomplished to date is sure to disappoint many whose successes we have not covered and for that we apologize. Similarly, although semisynthetic and biosynthetic methods for elaboration of natural products and for the generation of improved analogues are integral to drug discovery,^{13–18} we do not present examples derived from such efforts in this review.

Central to the studies and a major challenge for natural products is the development of synthetic strategies and new methodology tailored to their structures that permit their preparation and that of key analogues, including implementation of enabling divergent synthetic strategies.¹⁹ The collective efforts summarized herein have provided a stunning

series of unique, non-obvious total syntheses capable of not only affording the targeted natural products, but also key or comprehensive sets of structural analogues. Often, this was used to explore the origin of the biological properties and the mechanism of action of a natural product as well as discover compounds with improved activity. Consequently, the significant improvements highlighted herein not only represent unusual instances of productive deep-seated modifications of natural products that were accessed by total synthesis but are also remarkable in the magnitude of the advances and the underlying science responsible for the supernatural product.

In some cases, natural products may already possess suitable properties for use directly in the clinic, including high potency, selectivity and *in vivo* efficacy, reasonable pharmacokinetics (PK) and low toxicity. Such examples include the inhibition of peptidoglycan biosynthesis by β -lactams and glycopeptide antibiotics, the inhibition of bacterial ribosomal protein synthesis by tetracyclines, the binding and inhibition of tubulin polymerization by the *Vinca* alkaloids, and the immunosuppressive effects of cyclosporin, to name a few. Moreover, the quest for supernatural products is not new and has been a focus of drug discovery from its very beginning. W. Bachmann, a pioneer in steroid synthesis and who carried out the first total synthesis of a steroid hormone (equilenin) with Alfred L. Wilds in 1939, simultaneously pursued the total synthesis of analogues with the intension of improving on its properties.^{20–25} Without trying to be exhaustive in chronicling these, classic examples include the chemical evolution of β -lactam antibiotics from penicillins to cephalosporins and synthetic carbapenems to improve the activity, broaden the antimicrobial spectrum and overcome antibiotic resistance, the development of ergot alkaloids with improved receptor selectivity now used to treat migraine and Parkinson's disease, the seminal discovery and development of semisynthetic steroid analogues of progesterone (isolated in trace quantities and not orally active) as orally active contraceptives, the stunning number of prostaglandin analogues examined, and the introduction of efficacious synthetic statins as lipid-lowering medications. Advances in synthetic chemistry has accompanied the exploration of natural products and contributed to their structure determination, generation of sufficient material for biological studies for those available in low-natural abundance and has led to improved analogues. Such improvements can occur in a range of ways, and we highlight select examples (2000–2020) of total synthesis derived supernatural products that: (1) improve the activity, potency or target selectivity, (2) broaden the spectrum as antibiotics, (3) overcome a mechanism of drug resistance, (4) add additional synergistic mechanisms of action not found in the natural products that increase durability, (5) improve chemical or metabolic stability, or (6) improve *in vivo* efficacy. Many represent efforts that improve more than one such aspect of natural product properties and often incorporate modifications that additionally overcome limiting physical properties or improve *in vivo* PK properties. We originally considered and organized the review placing the examples in subsections representing the properties improved. However, we found this did a disservice to the selected examples since most conduct a multi-dimension optimization that covers multiple features. Rather, we elected to summarize the range of properties often targeted here in the introduction as well as in the conclusions and to treat each example individually, highlighting the properties the work focused on and improved in the text and providing subheading summarizing the major advance. We hope this review will not only

serve as a representative collection of past accomplishments but also highlight opportunities for new discoveries.

2. Discovery of supernatural products

2.1 Halichondrins: From total synthesis to clinic

The work of Kishi et al. on the total synthesis of halichondrin B (**1**) and its related analogues highlight the power of synthesis even for remarkably complex natural products, addressing the issue of limited material supply for drug discovery and development. Halichondrins are a family of structurally complex polyether macrolides originally isolated from a marine sponge.²⁶ The most potent member, halichondrin B, exhibits nanomolar *in vitro* activity and efficacious *in vivo* antitumor activity²⁷ by induction of non-productive tubulin aggregates, suppression of microtubule dynamics, and suppression of spindle microtubule growth events, representing a mechanism of action different from other tubulin-active natural products like paclitaxel and the *Vinca* alkaloids. Despite the outstanding potency as well as novel mechanism of action, the exploration of halichondrin B was limited by its extremely low natural availability (12.5 mg of **1** from 600 kg sponge) and structural complexity. Adopting a highly convergent strategy, Kishi et al. completed the first total synthesis²⁸ of **1** and norhalichondrin B (**2**), a similarly active analogue. In subsequent extensive efforts, they improved the synthetic route, shortened the longest linear sequence (42 to 25 steps), improved the overall yield, and discovered efficacious analogues with improved *in vivo* efficacy.^{29–31} SAR studies were initially conducted with polyether subunit analogues or macrolactone analogues, where the C1–C38 fragment **3** was observed to exhibit the *in vitro* cell growth inhibitory activity of halichondrin B within an order of magnitude but was not active *in vivo*. Further optimization of the *in vitro* activity as well measurement of their ability to maintain the mitotic block resulted in the discovery of macrocyclic ketone E7389 (**4**) as a structurally simplified but equipotent analogue of halichondrin B (**1**) (Fig. 1).³² Although **4** is structurally related to **3** but not halichondrin B (**1**), both **1** and **4** shared low- to sub-nanomolar tubulin polymerization inhibitory activity and cell growth inhibitory activity. The scalable synthesis of **4** not only provided sufficient material for preclinical and clinical studies, but also helped elucidate its halichondrin-like mechanism of action. In fact, it was the studies on **4** that shed light on the mechanism of action of halichondrins. The efforts resulted in the approval of E7389 (**4**, eribulin) by the FDA in 2010 to treat breast cancer and later in 2016 for the treatment of liposarcoma. Soon after the discovery of eribulin, the synthetic route to the intact halichondrins and their analogues was scaled up to provide decagram quantities of E7130 (**5**, Fig. 1), a C52-amine analogue of **2**, exhibiting sub-nanomolar *in vitro* antiproliferative activity (IC₅₀ 0.1–0.01 nM) against human cancer cell lines and potent *in vivo* antitumor activity.³³ Importantly, the *in vivo* efficacy of E7130 is highlighted by a more sustained tumor regression and amelioration of the tumor microenvironment by increasing intratumoral microvessel density for improved drug delivery and location. Currently a Phase I study to investigate E7130 in a solid tumor is underway in Japan by Eisai Co., Ltd.

2.2. Vancomycin: Binding pocket redesign to overcome resistance and peripheral modifications that provide newly added independent, synergistic mechanisms of action

Vancomycin (**6**) was the first and one of the most important members of glycopeptide natural products. They bind and sequester bacteria cell wall biosynthesis precursors, including lipid II, and inhibit the cross-linking transpeptidation step in cell wall synthesis. They are regarded as the antibiotics of last resort for the treatment of infections by drug-resistant Gram-positive pathogens, including methicillin-resistant *S. aureus* (MRSA).³⁴ After 30 years of clinical use, resistance against vancomycin emerged initially in enterococci, and more recently in *S. aureus*. The mechanism of resistance entails an intricate late-stage remodelling of the C-terminus of cell wall synthesis precursors from D-Ala-D-Ala to D-Ala-D-Lac, resulting in a 1000-fold loss in affinity between vancomycin and its binding target and a corresponding 1000-fold loss in antimicrobial activity.³⁵

Second-generation glycopeptide antibiotics including telavancin, dalbavancin and oritavancin are semi-synthetic drugs with added mechanisms of action and improved activity. They do not bind D-Ala-D-Lac or act by the original mechanism of action in vancomycin-resistant strains. In order to directly address the underlying antimicrobial resistance against vancomycin on a molecular basis, Boger et al. identified the contributing effects responsible for the loss in affinity, and recognized that removal of lone pair repulsion and/or restoration of a H-bond would provide vancomycin analogues that overcome the intrinsic resistance mechanism.³⁶ Based on the foundation of the total synthesis of vancomycin aglycon (**7**),^{37–40} Boger et al. synthesized binding pocket analogues (**8** and **9**) and established their binding affinity with model ligands of D-Ala-D-Ala (**10**) and D-Ala-D-Lac (**11**) (Fig. 2).^{10,41–45} Compounds **8** and **9** exhibited improved binding affinity with the model D-Ala-D-Lac ligand **11** where the former benefited from removal of the lone pair electrostatic repulsion (**8**) and the latter additionally from re-introduction of a H-bond (**9**). They displayed not only improved antimicrobial activities against vancomycin-resistant strains (**9** > **8**), but more importantly exhibited dual D-Ala-D-Ala/D-Ala-D-Lac binding properties, allowing them to remain active against vancomycin-sensitive strains. Most notably, **9** matched the activity of vancomycin against sensitive strains and reinstated this full activity against vancomycin-resistant organisms. The recent total syntheses of their peripherally modified derivatives^{46–52} that provide up to two additional mechanisms of action not found in the natural products themselves have provided glycopeptide antibiotic analogues including **12** that (1) contain synergistic binding pocket and up to two added peripheral modifications, (2) that are endowed with up to three independent mechanisms of action, only one of which is dependent upon the dual D-Ala-D-Ala/D-Lac binding, (3) that display broad spectrum activity (e.g.; MRSA, VRSA, VanA and VanB VRE) at remarkable potencies (MICs = 0.01–0.005 µg/mL), and (4) that exhibit especially durable antimicrobial activity, failing to elicit resistance (Fig. 2).⁵³ These latter features are not found in the naturally occurring glycopeptide antibiotics and those discovered above that express three synergistic independent mechanisms of action are expected to display clinical lifetimes measured not in decades, or even the half a century of vancomycin, but perhaps will remain effective for centuries. It constitutes an example of what we like to think of as a prototypical supernatural product.

2.3. Discodermolide: SAR and discovery of improved analogues

(+)-Discodermolide (**13**) is a linear polyketide natural product with high tubulin binding affinity and potent antitumor activity that is derived through microtubule stabilization and cell cycle arrest. In the face of the supply limitations in the natural product and its synthetic complexity, initial^{54–58} and subsequent scalable total syntheses of discodermolide by Smith et al.⁵⁹ and by Novartis⁶⁰ provided not only sufficient material for pre-clinical studies and subsequent clinical trials, but also access to synthetic analogues. This provided analogues with improved activity and potency and with greater synthetic accessibility.⁶¹

Smith et al. prepared a series of lactone analogues of discodermolide through a divergent total synthesis based on a key Wittig reaction to construct C8-C9 olefin linkage and established the requirements of the lactone region for antiproliferative potency.^{62,63} As a result, the C3 hydroxyl group was identified as non-essential, and 2,3-anhydrodiscodermolide (**14**) and its analogues including **15** represent compounds with improved potency against sensitive cancer cell lines and against a multi-drug resistant cancer cell line (NCI/ADR) (Fig. 3). It was also concluded that ring substituents on the lactone that contributed to a twist boat conformation with all substituents in pseudoequatorial positions is optimal for its potency. Other important SAR studies provided further valuable insights, including observations that alkylation or acylation of C7 hydroxyl group improved the potency of **13** and **14**,⁶⁴ removal of C16 methyl group is well tolerated⁶⁵ and that C19 carbamate⁶⁶ and C14 nor-methyl⁶⁷ analogues are similarly potent against sensitive cell lines but are better substrates for efflux pumps including P-glycoprotein. The collection of such efforts, reviewed by Shaw in detail,⁶⁸ helped to define the critical requirements for high activity and provided opportunities to develop clinical candidates with improved potency and PK properties.

2.4. Tetracyclines: Synthetic, broad-spectrum antibiotics with improved potency and resistance profiles

Tetracyclines are one of the most important class of broad-spectrum antibiotics in human history.⁶⁹ As a result of extensive clinical use, widespread resistance to the natural tetracyclines emerged. The development of new therapeutics ensued focused on improving their potency and overcoming resistance derived from overexpression of efflux pumps, ribosomal binding site mutations, and the expression of ribosomal protection proteins.⁷⁰ Traditional tetracycline drugs are either natural products or their semi-synthetic derivatives and the current front-line drug, tigecycline (**17**), is a broad-spectrum semisynthetic 7,9-disubstituted tetracycline analogue that extended the antimicrobial spectrum of tetracyclines, especially toward tetracycline-resistant bacteria. Nonetheless, limitations in the development of tetracycline analogues persisted due the lack of an effective approach that could introduce chemical modifications at C7, C8 and C9 on the D-ring, which has proven to be crucial for the activity of the tetracycline antibiotics.²

Myers et al. developed an “AB + D” strategy for the modular, convergent preparation of synthetic tetracycline analogues. The three-step assembly enlists the construction of C-ring through a tandem, stereocontrolled Michael-Claisen cyclization^{71–73} (Fig. 4). The scalable preparation of AB fragment, the broad scope and functional group tolerance on D-ring, and

an efficient deprotection strategy enabled a highly diversifiable and scalable synthesis of tetracycline analogues and a dramatic expansion of accessible chemical space. More than 3000 tetracycline analogues have been synthesized in the commercialized tetracycline synthesis platform at Tetraphase Pharmaceuticals.² This led to the discovery of the 7-fluorotetracyclines eravacycline (**18**),⁷⁴ TP-271 (**19**)⁷⁵ and TP-6076 (**20**)⁷⁶ that display potent broad-spectrum antimicrobial activity, including activity against multidrug resistant strains outperforming tigecycline, favorable pharmacokinetics, improved oral bioavailability and low incidence of adverse events. As the first fully synthetic tetracycline antibiotic, eravacycline was approved by the FDA for the treatment of complicated intra-abdominal infections and TP-271 as well as TP-6076 are currently in Phase I clinical trials.

Notably, Myers et al. utilized a similar modular synthetic strategy and developed a platform for the discovery of new macrolide antibiotics from key building blocks that could be prepared on large scale. Many of the resulting synthetic macrolides exhibit antimicrobial activity comparable with erythromycin and some exhibit improved activity over the semisynthetic macrolides including telithromycin and overcome drug resistance.^{77,78}

2.5. Bryostatin: Analogues with improved potency, altered functional activity, and structural simplifications

The naturally occurring bryostatins, isolated from marine invertebrates, are a family of highly oxygenated macrolides. The interest in the bryostatins as antitumor drug candidates, as agents to treat Alzheimer's disease, and as synergistic components in the treatment of AIDS is derived from their nanomolar protein kinase C (PKC) binding affinity. Because of their potent biological activities, the clinical interest in their use, and their scarce natural abundance, significant efforts focused on the total synthesis of bryostatins, allowing the scalable preparation of the parent natural products for clinical development and opportunities to synthesize bryostatin analogues.^{79–83}

With the objective of discovering structurally simplified bryostatin 1 (**21**) analogues, Wender et al. reported the synthesis of >40 analogues and defined their SAR for PKC binding. Among the analogues examined, compound **22**, with simplified A- and B-ring systems and removal of C26 methyl group, displayed a greater PKC binding affinity than bryostatin 1 (Fig. 5). More importantly and exceeding the improvement in PKC affinity itself, **2** was found to be 10 to >100-fold more potent in a panel of human cancer cell lines.⁸⁴ As a bonus to this improvement, its structural simplification provided increased synthetic accessibility, allowing the preparation of **22** in gram quantities for biological studies. In addition, an analogue with comparable potency (**23**) in the series examined exhibited selectivity in mediating cytosol-to-cellular membrane translocation (isozyme PKC δ and ϵ favored over PKC β I), an interesting feature that is not exhibited by bryostatin 1.⁸⁵ Most recently, Wender et al. reported that some simplified bryostatin 1 analogues also serve as potent anti-Chikungunya virus agents (EC_{50} = 0.35–12 μ M), where bryostatin 1 is inactive (EC_{50} >96 μ M).⁸⁶

A collection of bryostatin analogues, named Merle compounds, have been reported by Keck et al. The relationship between the structure and PKC binding affinity as well as their functional bryostatin-like or tumor-promoting phorbol 12-myristate 13acetate (PMA)-like

biological behavior were defined.^{87–90} Replacement of the C7 acetate with an exocyclic olefin, removal of the C8 methyl groups, removal of labile C9 hydroxyl group and removal of C13 ester were found to be well tolerated and the resulting Merle compounds exhibited improved PKC binding affinity compared with bryostatin 1. Such structural changes also introduced differences in the biological functional behavior of the compounds. Some analogues such as Merle 23 (Fig. 5) exhibited PMA-like behavior where they function as a PKC activator, while other analogues such as Merle 28 (Fig. 5) exhibited bryostatin-like behavior, acting as functional antagonists. The progress in the synthesis of bryostatin analogues with improved PKC binding affinity and functional activity has been summarized in detailed reviews by Wender et al.⁷⁶ and Zhang et al.⁷⁸

2.6. Novobiocin: Redesign to provide anticancer agents

Novobiocin (**26**) is a well-known clinically used antibiotic introduced in the 1960s that acts as a DNA gyrase B inhibitor. It was found to exhibit weak antitumor activity (700 μM) through Heat Shock Protein (Hsp90) inhibition by Neckers et al.⁹¹ Albeit displaying initial weak binding affinity to its target, the unique, previously unrecognized, Hsp90 C-terminus binding site of novobiocin provided a new opportunity to inhibit and modulate the protein folding machinery. Blagg et al. detailed systematic SAR studies of synthetic novobiocin analogues to improve their antiproliferative activity and to generate redesigned novobiocin-based anticancer drug candidates. Although critical for DNA gyrase inhibition, the 4-hydroxyl and 3'-carbamate were found to be detrimental to Hsp90 inhibition. This was revealed through examination of an initial library of novobiocin analogues focused on the coumarin core and the noviose sugar moiety, where their removal provided a significant improvement (compound **27**) in antiproliferative activity ($\text{IC}_{50} = 0.5 \mu\text{M}$, >1000-fold increase, SKBr3 cell line) over that of the natural product itself (Fig. 6).⁹² Introduction of a polar biaryl moiety or 2-indolyl group (compound **28**), serving as an H-bond donor, was found to be optimal as replacements for the benzamide substituent against a panel of cancer cell lines.⁹³ The attachment of a tertiary amine-containing moiety to the noviose sugar as well as the introduction of a constrained lactam ring (compound **29**) maintained the activity of the analogues (Fig. 6).^{94,95} After systematic optimization, a set of potent novobiocin analogues that exhibit mid-nanomolar antitumor activity (1000- to 3000-fold increase) was discovered with the SAR and flexibility in structural changes⁹⁶ fully defined.

2.7. Diazonamide A: Discovery of DZ-2384

Diazonamide A (**30**) is a marine natural product isolated in 1991 that exhibits nanomolar anti-mitotic activity.⁹⁷ Its unusual structure, potent activity, and lack of neurotoxicity attracted efforts on its total synthesis by Nicolaou et al.^{98,99} and Harran et al.¹⁰⁰ This led to a reassignment of its structure¹⁰¹ as well as the discovery of its mechanism of action.^{102,103} In contrast to other microtubule-targeting agents including paclitaxel and the *Vinca* alkaloids, its biological target is not tubulin itself, but ornithine δ -amino transferase (OAT), a protein essential for proper spindle assembly. Despite the synthetic accomplishments on diazonamide A, including its structure reassignment, the presence of bis-macrocyclic core limited the preparation of analogues as well as detailed SAR studies. In 2015, Harran et al. reported a highly divergent approach for the preparation of structurally simplified analogues of diazonamide A, enlisting a key oxidative macrocyclization step. This methodology

allowed the preparation of a library of diazonamide A analogues with structural changes in both the core and periphery for biological evaluation.^{104,105} As a result of these efforts, DZ-2384 (**31**) emerged as a simplified structure with fully retained diazonamide anti-mitotic characteristics (Fig. 7). Importantly, DZ-2384 displayed a 10 to 50-fold improved *in vivo* efficacy due to improved PK properties and exhibited a similar lack of neurotoxicity as found with diazonamide A. Interestingly, subsequent observations suggest that the mechanism of DZ-2384 may be different from the parent natural product, where it binds to the *Vinca* alkaloid binding site of tubulin and triggers a change in curvature of tubulin protofilaments.¹⁰⁶ The application of oxidative macrocyclization strategy in the large-scale preparation of DZ-2384 provided a highly efficient and scalable route and DZ-2384 has been under development by Diazon Pharmaceuticals.

2.8. Vinblastine: Improvements in potency, overcoming resistance and ultrapotent vinblastines

Vinblastine (**32**) and vincristine (**33**) represent the most important members of the *Vinca* alkaloid family natural products.^{107,108} They exhibit potent antitumor activity derived from inhibition and perturbation of microtubule formation and dynamics. They are among the first and most important plant-derived natural product drugs and are used extensively today as key components in combination cancer therapies. Because of their clinical importance, structural complexity, low natural abundance and unique mechanism of action, vinblastine and vincristine continue to be the subject of extensive total synthesis, semisynthetic and biosynthetic studies.^{109–117} In efforts to develop analogues with further improved tubulin binding properties and antitumor activity, Boger et al. developed a tandem intramolecular [4 + 2]/[3 + 2] oxadiazole cycloaddition cascade for the concise total synthesis of vindoline¹¹⁸ as well as a biomimetic Fe(III)-promoted coupling of vindoline with catharanthine^{119–123} and *in situ* C20' oxidation. This has allowed not only the convergent total synthesis of vinblastine itself but related analogues with deep-seated modifications.¹²⁴ Among the many studied, a C10' fluorine substitution was observed to provide analogues **34** and **35** with a uniquely 8 to 30-fold improved activity (0.9–0.06 nM) against human tumor cell lines that may be attributed to a perfect steric match of the fluorine atom (vs hydrogen atom, a single heavy atom change) in a sterically-confined hydrophobic binding pocket within the tubulin binding site (Fig. 8).¹²⁵ An additional effort that targeted the SAR of C20' ethyl group resulted in the discovery of **36** as a vinblastine analogue with an additional C15'-C20' six-membered ring system *cis*-fused to the vinblastine core structure. The added molecular complexity was designed to both constrain the side chain and further occupy the C20' ethyl hydrophobic binding pocket. Compound **36** displayed a 10-fold increase in activity (0.7–0.6 nM) compared with vinblastine itself and an increased tubulin binding affinity (Fig. 8).¹²⁶ The latter improvement represents a case where added benign complexity (ABC) to the core structure of natural product, which is only accessible through total synthesis, further improved its activity by virtue of enhanced target binding while maintaining the original interaction mode and conformation.

In systematic SAR studies on the vinblastine C20' site, Boger et al. discovered and subsequently developed a hydrogen atom transfer (HAT) free-radical functionalization reaction¹²⁷ that allowed the introduction of C20' azide, amine, amides¹²⁸ and ureas.^{129,130}

Although the vinblastine C20' amine (**37**) displayed diminished activity (ca. 100-fold), the unsubstituted urea **38** approached the potency of vinblastine. The subsequent introduction of C20' ureas substituted with lipophilic groups were found to be beneficial where the pyrrolidine urea **39** and isoindoline urea **40** exhibited improved activity toward sensitive cell lines (10-fold). More importantly, the introduction of the isoindoline group in **40** provided a further improvement against the isogenic vinblastine-resistant cell line HCT116/VM46 ($IC_{50} = 7.5$ mM, 100-fold) attributable to partially overcoming overexpressed P-glycoprotein (Pgp) transport^{129,130} while simultaneously improving tubulin binding (Fig. 9). While an additional benzamide moiety in **41** is well-tolerated, the introduction of heterocyclic diazines further improved the activity of the C20' isoindoline urea analogues (**42–44**) (Fig. 9).¹³¹ Such derivatives were found to be ultrapotent vinblastine analogues that not only exhibit remarkable activity against sensitive cell lines ($IC_{50} = 70$ pM, 100-fold improvement), but also significantly further improve activity against a vinblastine resistant cell line ($IC_{50} = 0.8–0.9$ nM, 1000-fold improvement). The newly discovered ultrapotent vinblastines bind tubulin with much higher affinity and further disrupt the tubulin head-to-tail α/β dimer–dimer interaction by virtue of the strategic placement of an added conformationally well-defined, rigid, and extended C20' urea along the adjacent continuing protein–protein interface (Fig. 9). In this case, the added molecular complexity was used to markedly enhance target binding and functional biological activity (100-fold), likely representing a general approach to improving the properties of other natural products targeting a protein–protein interaction, while simultaneously disrupting Pgp binding and efflux. Just as stunning, the compounds are readily available in three steps from commercial vindoline (\$36/g) and catharanthine (\$16/g) by using the two-step, one-pot Fe(III)-promoted coupling and HAT functionalization reactions developed in their labs.

Despite their extensive clinical use, the only significant resistance to vinblastine and vincristine that has emerged as a limitation to their continued use is not derived from alteration of the drug target tubulin, but rather is derived from overexpression of a cell-surface drug efflux transporter, P-glycoprotein (Pgp), lowering their intracellular concentration. The identification of analogues that address this mechanism of resistance, also responsible for tumor multidrug resistance (MDR), has been a major focus of the field for decades and would represent a major advance for oncology therapeutics. Boger et al. optimized vinblastine C20' amides (180 analogues) based on fundamental medicinal chemistry principles and discovered a set that maintained or improved activities compared with vinblastine itself. This could be attributed to the H-bond acceptor ability of the added amide carbonyl that in turn proportionally increase the tubulin binding affinity.¹²⁸ In these studies and with a series of benzamide derivatives, a linear correlation between the Hammett σ_p constant of a *p*-substituent and $-\log IC_{50}$ revealed that the more electron-rich C20' amides predictably improve potency (Fig. 10). In addition, the lipophilic character (cLogP) of the resulting benzamide analogues (e.g., **45–55**) was established to directly correlate with diminished Pgp binding and efflux and resistance derived from Pgp overexpression (Fig. 11). These not only provided predictable correlations as a guide for optimization of analogues, but also led to a subset of analogues that matched the activity of vinblastine in sensitive cell lines (HCT116, L1210) and are equally active against an isogenic vinblastine-

resistant cell line (HCT116/VM46), providing attractive drug candidates not subject to resistance derived from Pgp overexpression.

2.9. Arylomycins: Development into broad-spectrum antibiotics

The arylomycins are a family of lipopeptide natural product antibiotics that inhibit type I signal peptidases (SPases) of both Gram-positive and Gram-negative bacteria *in vitro*.¹³² However, they exhibited a very narrow spectrum of activity and possessed no activity against important ESKAPE pathogens.¹³³ Romesberg et al. pioneered the exploration of arylomycins as broad-spectrum antibiotics and found that their SPase inhibition activity and their antibacterial potency are masked by the nearly ubiquitous presence of a proline residue in SPase (Pro²⁹ in *S. aureus*, Pro⁸⁴ in *E. coli* and *P. aeruginosa*) over other amino acid residues.¹³⁴ The presence of the proline residue in the binding site of the arylomycin N-terminus removes a crucial H-bond interaction. The mutation of this proline to serine in all strains tested was found to be sufficient to confer a high degree of sensitivity to arylomycins. The observation demonstrated the potential of arylomycin N-terminus analogues as broad-spectrum SPase targeting antibiotics, where initial extension of the N-terminal serine by one or two methylenes improved the antimicrobial activity of arylomycin A₁₆ against resistant strains.¹³⁵ Subsequent SAR studies by Romesberg et al. established that a precisely positioned C-terminus aldehyde modification (**57**)¹³⁶ improved the SPase binding affinity of arylomycin A₁₆ (Fig. 12). Based on the initial reported total syntheses of arylomycins by Romesberg and a later improved total synthesis of Romesberg and Baran,^{137,138} Smith et al. at Genentech focused on the structural optimization of arylomycins targeting the N-terminus, C-terminus, and the phenol hydroxyl groups. As a result of these efforts, the introduction of an N-terminus aromatic lipophilic group, the simplification of the N-terminus to a diaminobutyric acid residue, the C-terminus modification to a nitrile, and appending primary amines to the phenols resulted in G0775 (**58**), exhibiting potent antimicrobial activity against a broad spectrum of Gram-negative bacteria, most of which are resistant to the parent arylomycin family (Fig. 12).¹³⁹ Finally, Wong et al. at Genentech developed and reported an efficient and scalable synthesis of structurally related GDC-5338 (**59**) as an exciting, broad-spectrum synthetic arylomycin analogue active against Gram-negative bacterial infections.¹⁴⁰

2.10. Epothilone B: Stable and improved analogues

Epothilones are naturally occurring cytotoxic macrolides exhibiting highly potent antitumor activity derived from tubulin binding, stabilization, and inhibition of microtubule disassembly.¹⁴¹ Initial *in vivo* studies of epothilone suggested potential liabilities, including a low stability under physiological conditions as well as a narrow therapeutic window.¹⁴²

Accomplished initially by total synthesis and later semi-synthesis, Borzilleri et al. at BMS developed ixabepilone (**61**), the macrolactam analogue of epothilone B (**60**).¹⁴³ Ixabepilone was found to exhibit comparable *in vitro* tumor cell growth inhibition and microtubule-polymerization and stabilization activity but a remarkably improved metabolic stability and improved PK properties. This resulted in enhanced *in vivo* potencies against both paclitaxel-sensitive and -resistant tumors (Fig. 13).^{142,144} It was approved by the FDA in 2007 for treatment of aggressive metastatic or locally advanced breast cancer.

The synthesis and optimization of epothilone B analogues with improved properties as candidate antitumor agents by Danishefsky et al. represent a beautiful series of discoveries where synthetic intermediates used to access the natural products could serve as supernatural products or key compounds to understand its SAR. In the course of the development of a scalable synthesis of epothilone B and its analogues,^{145,146} intermediate desoxyepothilone B (**62**) was found to exhibit a significant decrease in toxicity as indicated by an improvement in survival rate, although this was accompanied by a slight decrease in antitumor potency (Fig. 13).¹⁴⁷ A next generation total synthesis of **62**, utilizing a ring closure metathesis strategy, provided dehydellone **63** initially as the product of olefin metathesis. Compound **63** and its trifluoromethyl derivative fludelone (**64**) were observed to exhibit highly promising *in vitro* and *in vivo* potencies. They were also observed to maintain the diminished toxicity of desoxyepothilone, improve the metabolic stability compared with **60** and **62** and displayed enhanced water solubility advantageous to formulation (Fig. 13).^{148,149} Finally, a further optimization resulted in the introduction of the C17-isooxazole in analogues **65** and **66**, with a remarkable improvement in stability and further improved potency and water solubility. The series represent stable, safe and potent epothilone analogues as promising antitumor agents¹⁵⁰ for clinical development (iso-fludelone, **66**) at Memorial Sloan Kettering Cancer Center.

2.11. Duocarmycins, CC-1065 and yatakemycin: Improvements in activity, tolerability, and *in vivo* efficacy

Duocarmycin SA (**67**) and CC-1065 (**68**) are two of the most widely studied antitumor agents within the family of natural products that also include duocarmycin A and yatakemycin.^{151,152} They exhibit exceptionally potent picomolar biological activities derived from a sequence-selective DNA alkylation reaction. In the course of establishment of the SAR of this family of natural products, Boger et al. completed the total syntheses^{153–157} of each of the natural products and prepared >2000 analogues with deep-seated structural changes. These were used to define their DNA noncovalent binding selectivity preferentially in the narrower, deeper AT-rich minor groove and identify the source of DNA alkylation catalysis that arises from a binding-induced conformational change that disrupts the alkylation subunit stabilizing vinyllogous amide conjugation. They defined the basis of the alkylation selectivity including that of their unnatural and often equally active enantiomers, quantitated DNA alkylation rates, their reversibility and its stereoelectronically-controlled reaction selectivity. The cumulative studies established a fundamental parabolic relationship between the alkylation subunit reactivity/stability and the cytotoxic potency as a predictive tool for the development of optimized analogues with improved potency and stability.^{158–167} Notable is the example where the replacement of the alkylation subunit pyrrole NH with a sulfur atom in CC-1065 and a structurally simplified analogue (**69**) reaches the pinnacle of the parabolic relationship, providing analogues (**70** and **71**) which are both 6-fold more stable and up to 10-fold more potent compared to their pyrrole equivalent (Fig. 14).¹⁶⁸

Based on the highly potent natural products and their simplified yet optimized analogues, structurally unique prodrugs have been designed. Some have been found to be preferentially converted to the active species in the tumor environment and serve as a powerful way to

improve their therapeutic window and *in vivo* efficacy. In recent studies, Boger et al. introduced a series of cyclic or acyclic N-acyl O-amino phenol prodrugs that were designed to take advantage of the hypoxic tumor environment and the increased concentration of reducing nucleophiles, including glutathione.^{169,170} The concurrent optimization of both the intrinsic stability and its reactivity toward reductive activation provided **73** as the prodrug of free phenol **72**. Prodrug **73** exhibits a reduced *in vitro* activity but a greater stability and possesses a much larger therapeutic window (200 to >2500 µg/kg) compared with **72** (30 to 100 µg/kg) without apparent toxicity or dose-limiting myelotoxicity (Fig. 14). Remarkably, **73** is much more efficacious *in vivo* than the free drug itself, providing long-term survivors (1 year) in animal tumor models. This duocarmycin analogue is in development as a clinical candidate at Cairn Therapeutics.

2.12. Migrastatin: Core analogues as antimetastatic agents

Migrastatin (**74**) was isolated from the culture broth of *Streptomyces* in 2000. It was found to inhibit human tumor cell migration, serving as a small molecule lead as a modulator of tumor cell relocation and metastasis.^{171,172} Based on their migrastatin total synthesis,¹⁷³ Danishefsky et al. reported a series of analogues through a diverted and divergent synthetic strategy and established their biological activity as tumor cell migration inhibitors. Among the analogues examined, the migrastatin core **75** and its reduced version **76** exhibited substantial (1000-fold, mouse tumor cell line 4T1) improvement in tumor cell migration inhibition (Fig. 15).^{174,175} However, a decrease in stability accompanied the removal of the side chain and the lactone was found to be labile to hydrolysis. As a result, lactam **77**, macrocyclic ketone **78**, and later the macrocyclic ether **79** were synthesized.^{174–176} Such analogues, bearing only a single-atom change, were found to be stable in plasma while maintaining most of the improvements in tumor cell migration inhibition observed with **75** and **76**. Moreover, the structurally simplified analogue **79** was found to be highly selective for inhibition of mouse or human tumor cell migration (IC₅₀ = 0.30–0.47 µM, 1000-fold) over normal human mammary epithelial MCF10A cells (IC₅₀ = 408 µM), a welcomed property not exhibited by migrastatin (**74**) and its core analogues **75–78** (Fig. 15). Finally, **79** and its more potent analogue **80** exhibited potent tumor cell migration inhibition activity *in vivo* in a human lung cancer model as promising antimetastatic agents for cancer therapy.¹⁷⁷

2.13. Thromboxane: Discovery of stable analogues

Thromboxane A₂ (**84**) and prostaglandin endoperoxides PGG₂ and PGH₂ (**81**) are produced enzymatically from arachidonic acid with important biological functions, including response to tissue injury, promoting hemostasis, vasoconstriction, and wound healing.^{178–180} However, their instability under physiological conditions (**81**, *t*_{1/2} = ~ 5 min; **84**, *t*_{1/2} = 32 s) precludes their use as therapeutics and present an opportunity for the preparation of stable analogues with similar biological properties. Early contributions by Corey et al. include the development of stereocontrolled divergent total syntheses of the prostaglandins through the well-known Corey lactone intermediate^{181,182} and the synthesis of a norbornene-containing structure (**82**) as stable prostaglandin endoperoxide analogue with similar platelet aggregation activity (Fig. 16).^{183,184} The former work is perhaps the best known and among the earliest examples of the power of a divergent synthetic strategy. This pioneering work

was followed by the synthesis of U46619 (**83**) by Bundy as a stable ether analogue of PGH₂ with similarly potent smooth muscle-stimulation activity compared with PGH₂¹⁸⁵ and similar platelet aggregation activity compared with thromboxane A₂ (**84**),¹⁸⁶ and the synthesis of F₂-thromboxane A₂ (**85**) by Fried et al. which is a fluorinated analogue of **84** with much improved stability ($t_{1/2}$ = 46 weeks), and an increased platelet aggregation EC₅₀, but a compromised platelet aggregation efficacy (Fig. 16).¹⁸⁷ Most recently, Aggarwal et al. prepared F-thromboxane A₂ (**86**) as a closer analogue of **84** and established the stereoelectronic origin of the improvement on the stability of fluorinated thromboxane A₂ analogues.¹⁸⁸ F-thromboxane A₂ displayed biological properties resembling **83** and **85**, including a greater platelet aggregation efficacy than **85** as determined by integrin $\alpha_{IIb}\beta_3$ activation. Importantly, the stability of **86** is only slightly lower than **85** ($t_{1/2}$ = 20 days, 10-fold) but much improved (10⁵-fold) when compared with thromboxane A₂ (Fig. 16).

2.14. Salvinorin A: Improvement in configurational stability

Salvinorin A (**87**), a diterpene plant metabolite isolated in 1982, was found to act as a potent and selective κ -opioid receptor (KOR) agonist.^{189,190} Unlike conventional amine-containing natural product-derived analgesics, its neutral structure under physiological conditions allows salvinorin A to effectively pass through the blood-brain barrier. Combined with its selective receptor binding behavior, interest focused on salvinorin A as a potential lead compound for pain therapeutics with minimized side effects. However, the intrinsic chemical liability, arising from the configurational instability of the C8 stereocenter, limited semisynthetic and total synthetic approaches toward salvinorin A and its analogues.^{191,192} Under mild basic conditions, salvinorin A epimerizes to C8-*epi*-salvinorin A with the equilibrium favoring the unnatural and inactive epimeric *cis* isomer in a 3:7 ratio.

Shenvi et al. recognized that removal of C20 methyl group would release conformational strain derived from the 1,3-diaxial interaction and tune the equilibrium favoring the natural *trans* fused isomer. They reported an efficient synthesis of C20-nor-salvinorin A (**88**) (Fig. 17)¹⁹³ and found that it retained the highly selective KOR binding behavior, albeit displaying slightly less affinity and efficacy compared with salvinorin A itself (EC₅₀ for cAMP accumulation = 6.1 nM vs 0.9 nM). Importantly, a reversed C8 epimerization equilibrium was observed with **88**, now favoring the natural *trans* isomer in a 7:3 ratio. To further improve the equilibrium favoring the natural *trans* isomer, Shenvi et al. designed O6C-20-nor-salvinorin A (**89**) with a single heavy atom change from the lactone to a cyclohexanone (Fig. 17).¹⁹⁴ Compound **89** benefited from the removal of the epimerization prone semi planar lactone conformation in the *trans* isomer **88** by adopting a strain-released chair conformation. It exhibited complete stability to epimerization while maintaining selective and potent KOR agonist activity (EC₅₀ for cAMP accumulation = 3.3 nM). In addition to the improvement in configurational stability and as a bonus of removing the C20 methyl group, analogues **88** and **89** are synthetically more accessible (10–12 steps) compared with the parent natural product (20–29 steps).

2.15. Meayamycins: More stable and more potent analogues of FR901464

FR901464 (**90**) was isolated from a Japanese soil sample in 1996, and exhibited potent nanomolar cytotoxic activity against a panel of human tumor cell lines.¹⁹⁵ The mechanism

of action and biological target of FR901464 and its analogues were first revealed by Yoshida et al.^{196,197} where they bind the splicing factor 3b (SF3b) complex and inhibit spliceosome pre-mRNA splicing and maturation to mRNA. Because of the limited natural product supply, total syntheses of FR901464 were first pursued and reported by Jacobsen et al.¹⁹⁸ and Kitahara et al.¹⁹⁹ and later Koide et al.²⁰⁰ This provided synthetic FR901464, confirming its structure and establishing its absolute configuration, and provided analogues with improved chemical and metabolic stability and potency.

The instability of FR901464 in organic solvents as well as under physiological conditions limited investigation of this highly potent natural product. In order to directly address the instability of FR901464, Jacobsen et al. examined the impact of the presence of C1 hemiketal and the epoxide and found that the problematic instability was due to the hemiketal.¹⁹⁸ In their studies, the C1 deshydroxy analogue **91** was found to be chemically stable and displayed an improvement in cell growth inhibition potency, revealing that the hemiketal is not contributing to its biological activity (Fig. 18). Based on these studies, Koide et al. later reported the C1 gem-dimethyl analogue **92**, named meayamycin, which displayed the improved stability and exhibited a significant further improvement ($GI_{50} = 10\text{--}20\text{ pM}$, 100-fold) in antiproliferative potency (Fig. 18).²⁰¹ Subsequent SAR studies by Koide et al. revealed that meayamycin B (**93**) with a morpholine carbamate group further improved inhibition of SF3b, the antiproliferative activity, and stability in mouse serum.^{202,203} The discovery of meayamycins as more stable and more potent analogues of FR901464 provided highly potent antiproliferative small molecules as promising leads with a novel therapeutic target as well as widely used chemical probes in pre-mRNA splicing studies.

2.16. Ramoplanin: Improvement in stability

Ramoplanin is a lipoglycopeptide antibiotic with potent activity against Gram-positive bacteria, consisting from three closely related compounds differing only in the lipophilic side chain (ramoplanin A1–A3). It binds to the bacteria cell wall biosynthetic precursors lipid I and lipid II, inhibiting the conversion of lipid I to lipid II as well as the subsequent transglycosylation step in peptidoglycan biosynthesis.²⁰⁴ As a result of the unique mechanism of action as an enzyme substrate binder like vancomycin, ramoplanin is an especially attractive antimicrobial agent. It exhibits sub $\mu\text{g/mL}$ activity against both sensitive and drug-resistant bacterial strains and does not elicit resistance as a result of the invariant target structure. It is approved by the FDA as a treatment for drug-resistant *C. difficile* infections in the gastrointestinal tract and used off label for topical treatment of refractory skin wound infections.²⁰⁵ A limitation to its use, ramoplanin and its aglycon are not stable in plasma ($t_{1/2} < 5\text{ min}$) or in mild acidic or basic conditions as a result of rapid hydrolysis of the depsipeptide ester (HAsn² residue) to yield an inactive linear peptide.²⁰⁶ Such instability results in limitations in the clinical applications of ramoplanin for systemic infections.

By following an approach similar to the route used for a ramoplanin total synthesis^{207–209} and based on rational design, Boger et al. reported [L-Dap²]ramoplanin A2 aglycon (**95**) as a ramoplanin A2 aglycon analogue (**94**) that bears a stable amide in place of the labile depsipeptide ester.²¹⁰ This subtle single atom structural change slightly improved antimicrobial activity and resulted in substantial improvements in chemical and metabolic

stability without disrupting interaction with its biological target (Fig. 19). Importantly, the unaltered macrocyclic ring size in the 49-membered macrocyclic lactone/lactam was found to be crucial to the activity of **95**, where the attempt to expand or contract the macrocycle by one carbon atom did not provide active analogues. A subsequent alanine scan of the structure (17 analogues) prepared by using the triply convergent approach allowed an assessment of the importance of each residue side chain to the antimicrobial activity and was conducted on this stable ramoplanin analogue.²¹¹

2.17. Polymyxins: Reduction in toxicity

Polymyxins, including polymyxin B (**96**) and polymyxin E (colistin), are antibiotics that have been used in the clinic for 50 years. They have become increasingly important in the last 10–15 years as last-line therapeutics for infections caused by multidrug resistant Gram-negative bacteria.^{212–215} They exhibit their antimicrobial activity through binding to the lipid A component of lipopolysaccharide (LPS) in the outer membrane, causing membrane disruption and cell leakage. The major clinical limitation of the polymyxins is their nephrotoxicity. This led to the gradual abandonment of the intravenous formulation in 1980s and represents a need for the development of less toxic polymyxin analogues as therapeutics for the systemic treatment of multidrug resistant Gram-negative bacterial infections.^{216,217}

By targeting the nephrotoxicity of polymyxins, a set of novel polymyxin derivatives with decreased positive charge was developed by Vaara et al. at Northern Antibiotics with a focused SAR study of the exocyclic residues.^{218,219} A deletion of N-terminus diaminobutyric acid (Dab¹) and the change of Dab³ to a D-serine residue provided an analogue NAB739 (**97**) that exhibits activity similar to polymyxin B but displayed a decreased toxicity in a porcine renal proximal tubular cell line (LLC-PK₁). This suggested the possibility of directly addressing the nephrotoxicity by decreasing the net positive charge in the polymyxins. In a systematic SAR study of the polymyxins, including the macrocyclic ring system, Cooper et al.²²⁰ established the residues critical for activity, including Dab⁵ and Dab⁹ and found a similar trend where the replacement of other positively charged Dab residues (Dab¹, Dab³, Dab⁸) maintained the antimicrobial activity and lowered nephrotoxicity (6-fold). Interestingly, the N-terminus lipophilic chain analogues **98** and **99** with a biphenyl or biphenyl ether group were found to be the least nephrotoxic analogues among the series examined, with antimicrobial activity similar to polymyxin B against sensitive strains, improved activity against polymyxin-resistant strains, and a similarly low toxicity against the mammalian HepG2 cell line (Fig. 20). The results should guide the development of exciting polymyxin analogues with improved antimicrobial activity and reduced toxicity.

2.18. Sansanmycin: Analogues as anti-tuberculosis agents

Sansanmycins are a family of uridyl peptide natural products^{221,222} initially isolated from *Streptomyces* in 2007 which share a similar core structure with the pacidamycins,²²³ mureidomycins,²²⁴ and napsamycins.²²⁵ Like other uridyl peptide natural products, sansanmycins bind and inhibit bacterial phospho-MurNAc-pentapeptide translocase (MraY, MurX in mycobacteria), the integral membrane enzyme responsible for the biosynthesis of lipid I. As a result, they exhibit interesting antibiotic activity against multidrug resistant *M.*

tuberculosis and Gram-negative pathogens including *P. aeruginosa*. Because of their attractive biological properties, Payne et al. reported a detailed SAR study, targeting sansanmycin analogues with improved activity, and discovered dihydrosansanmycins as a class of anti-tuberculosis agents.²²⁶ Dihydrosansanmycins (**102** and **103**) were initially designed to address the potential metabolic instability of the embedded enamide, but were also observed to exhibit a substantial 30-fold improvement in the activity against a virulent *M. tuberculosis* strain (H37Rv) compared with sansanmycin A (**100**) and B (**101**) (Fig. 21). Subsequent optimization in the peptide subunit resulted in a cyclohexyl analogue (**104**) with further improved functional activity (10-fold, MIC = 0.037 μ M) derived from an enhanced binding and inhibition of MurX. They also displayed similar activity intracellularly in infected cells, serving as leads for the treatment of tuberculosis.

3. Additional examples of natural product analogues with improved properties developed prior to 2000

Although not included in detail in this review, we would like to highlight a few notable efforts made in the discovery of supernatural products prior to 2000 (Fig. 22). Pettit et al. developed the auristatins as dolastatin 10 analogues that act as tubulin-binding antitumor agents and display decreased toxicity and improved *in vivo* efficacy.^{227–229} Auristatin E was later incorporated in monoclonal antibody-drug conjugates by Seattle genetics²³⁰ and served as the payload of brentuximab vedotin (Adcetris) that was approved by FDA to treat Hodgkin lymphoma and systemic anaplastic large cell lymphoma (in 2011) and later to treat cutaneous T-cell lymphoma (in 2017). Schreiber et al. and Nicolaou et al. recognized the binding domain and effector domain of FK506 and rapamycin, established the relationships between structure and functional immunosuppressive activity, and developed analogues that bind FKBP (FK506 binding protein) but serve as antagonists of the immunosuppressive effects of FK506 and rapamycin.^{231–233} Mitsui et al. synthesized about 150 camptothecin analogues and developed the hexacyclic analogue exatecan as an antitumor agent with improved activity and *in vivo* efficacy.^{234,235} This compound was later incorporated as the payload in the antibody-drug conjugate trastuzumab deruxtecan (Enhertu) by Daiichi Sankyo Co. Ltd, approved for the treatment of unresectable or metastatic HER2-positive breast cancer by the FDA in 2019. Danishefsky et al. identified a benzyl analogue of spirotryprostatin B and its stereoisomer as tubulin-active antitumor agents with a 10⁴-fold improvement in potency compared with the parent natural product.²³⁶

4. Summary and Outlook

A select set of recent examples of supernatural products, analogues that supersede the parent natural product properties, derived by total synthesis are summarized. The advances in synthetic chemistry that has made this possible, the individual creativity and technical expertise expressed in the efforts, the fundamental design principles applied, and the underlying science are stunning. Enhancements in potency (by as much as 1000-fold), increases in target selectivity and reductions in toxicity, fundamental solutions to overcoming mechanisms of drug resistance, newly added synergistic mechanisms of action embedded into the compound structures, improvements in poor physical properties,

mitigation of chemical or metabolic instability, and more efficacious *in vivo* activity have been achieved in the efforts. Combined with advances in chemical biology for target identification, assessment of functional activity and target validation, and with the application of fundamental medicinal chemistry principles, the examples highlight that the opportunities for the design and total synthesis of supernatural products have never been more exciting.

Acknowledgements

We gratefully acknowledge the financial support of the National Institutes of Health (CA041101, CA042056, CA208669, and DA15648; D.L.B.) through the years and a JITRI-Scripps Fellowship (Z.-C.W.).

7. References

1. Butler MS, J. Nat. Prod, 2004, 67, 2141–2153. [PubMed: 15620274]
2. Cragg GM and Newman DJ, Biochim. Biophys. Acta, Gen. Subj, 2013, 1830, 3670–3695.
3. Newman DJ and Cragg GM, J. Nat. Prod, 2020, 83, 770–803.
4. Szpilman AM and Carreira EM, Angew. Chem., Int. Ed, 2010, 49, 9592–9628.
5. Itoh H and Inoue M, Chem. Rev, 2019, 119, 10002–10031. [PubMed: 30945851]
6. Crane EA and Gademann K, Angew. Chem., Int. Ed, 2016, 55, 3882–3902.
7. Huffman BJ and Shenvi RA, J. Am. Chem. Soc, 2019, 141, 3332–3346. [PubMed: 30682249]
8. Wender PA, Verma VA, Paxton TJ and Pillow TH, Acc. Chem. Res, 2008, 41, 40–49. [PubMed: 18159936]
9. Hong J, Chem. Eur. J 2014, 20, 10204–10212. [PubMed: 25043880]
10. Boger DL, J. Org. Chem, 2017, 82, 11961–11980. [PubMed: 28945374]
11. Wan KK and Shenvi RA, Synlett, 2016, 27, 1145–1164.
12. Lambrecht MJ, Kelly J,W, and Shenvi RA, ACS Chem. Biol, 2018, 13, 1299–1306. [PubMed: 29653046]
13. Wright PM, Seiple IB and Myers AG, Angew. Chem., Int. Ed, 2014, 53, 8840–8869.
14. Lourenco AM, Ferreira LM and Branco PS, Curr. Pharm. Des, 2012, 18, 3979–4046. [PubMed: 22632756]
15. Yñíguez-Gutiérrez AE and Bachmann BO, J. Med. Chem 2019, 62, 8412–8428. [PubMed: 31026161]
16. DeCorte BL, J. Med. Chem, 2016, 59, 9295–9304. [PubMed: 27331414]
17. Kennedy J, Nat. Prod. Rep, 2008, 25, 25–34. [PubMed: 18250896]
18. Koehn FE, Med. Chem. Commun, 2012, 3, 854–865.
19. Boger DL and Brotherton CE, J. Org. Chem, 1984, 49, 4050–4055.
20. Bachmann WE, Cole W and Wilds AL, J. Am. Chem. Soc, 1940, 62, 824–839.
21. Bachmann WE, Cole W and Wilds AL, J. Am. Chem. Soc, 1939, 61, 974–975.
22. Bachmann WE and Wilds AL, J. Am. Chem. Soc, 1940, 62, 2084–2088.
23. Bachmann WE and Holmes DW, J. Am. Chem. Soc, 1941, 63, 2592–2598.
24. Bachmann WE and Thomas DG, J. Am. Chem. Soc, 1942, 64, 94–97.
25. Bachmann WE, Kushner S and Stevenson AC, J. Am. Chem. Soc, 1942, 64, 974–981.
26. Hirata Y and Uemura D, Pure Appl. Chem, 1986, 58, 701–710.
27. Fodstad O, Breistøl K, Pettit GR, Shoemaker RH and Boyd MR, J. Exp. Ther. Oncol, 1996, 1, 119–125. [PubMed: 9414395]
28. Aicher TD, Buszek KR, Fang FG, Forsyth CJ, Jung SH, Kishi Y, Matelich MC, Scola PM, Spero DM and Yoon SK, J. Am. Chem. Soc, 1992, 114, 3162–3164.
29. Yahata K, Ye N, Ai Y, Iso K and Kishi Y, Angew. Chem., Int. Ed, 2017, 56, 10791–10795.

30. Fukuyama T, Kishi Y, Ai Y, Ye N, Wang Q, Yahata K, Iso K, Naini SR, Yamashita S and Lee J, WO2019010363A1, 2019.
31. Jackson KL, Henderson JA and Phillips AJ, *Chem. Rev*, 2009, 109, 3044–3079. [PubMed: 19534494]
32. Yu MJ, Kishi Y and Littlefield BA, in *Anticancer Agents from Natural Products*, ed. Cragg GM, Kingston DGI and Newman DJ, Taylor & Francis Group, Boca Raton, FL, 2005, pp. 241–265.
33. Kawano S, Ito K, Yahata K, Kira K, Abe T, Akagi T, Asano M, Iso K, Sato Y, Matsuura F, Ohashi I, Matsumoto Y, Isomura M, Sasaki T, Fukuyama T, Miyashita Y, Kaburagi Y, Yokoi A, Asano O, Owa T and Kishi Y, *Sci. Rep.*, 2019, 9, 8656. [PubMed: 31209263]
34. Kahne D, Leimkuhler C, Lu W, and Walsh C, *Chem. Rev*, 2005, 105, 425–448. [PubMed: 15700951]
35. Walsh CT, *Science*, 1993, 261, 308–310. [PubMed: 8392747]
36. McComas CC, Crowley BM and Boger DL, *J. Am. Chem. Soc.*, 2003, 125, 9314–9315. [PubMed: 12889959]
37. Okano A, Isley NA and Boger DL, *Chem. Rev*, 2017, 117, 11952–11993. [PubMed: 28437097]
38. Boger DL, Miyazaki S, Kim SH, Wu JH, Castle SL, Loiseleur O and Jin Q, *J. Am. Chem. Soc.*, 1999, 121, 10004–10011.
39. Boger DL, Kim SH, Mori Y, Weng J–H, Rogel O, Castle SL and McAtee JJ, *J. Am. Chem. Soc.*, 2001, 123, 1862–1871. [PubMed: 11456806]
40. Crowley BM, Mori Y, McComas CC, Tang D and Boger DL, *J. Am. Chem. Soc.*, 2004, 126, 4310–4317. [PubMed: 15053621]
41. Boger DL, *Med. Res. Rev.*, 2001, 21, 356–381. [PubMed: 11579438]
42. Crowley BM and Boger DL, *J. Am. Chem. Soc.*, 2006, 128, 2885–2892. [PubMed: 16506767]
43. Xie J, Okano A, Pierce JG, James RC, Stamm S, Crane CM and Boger DL, *J. Am. Chem. Soc.*, 2012, 134, 1284–1297. [PubMed: 22188323]
44. Xie J, Pierce JG, James RC, Okano A and Boger DL, *J. Am. Chem. Soc.*, 2011, 133, 13946–13949. [PubMed: 21823662]
45. James RC, Pierce JG, Okano A, Xie J and Boger DL, *ACS Chem. Biol.*, 2012, 7, 797–804. [PubMed: 22330049]
46. Nakayama A, Okano A, Feng Y, Collins JC, Collins KC, Walsh CT and Boger DL, *Org. Lett.*, 2014, 16, 3572–3575. [PubMed: 24954524]
47. Moore MJ, Qu S, Tan C, Cai Y, Mogi Y, Keith DJ, and Boger DL, *J. Am. Chem. Soc.*, 2020, 142, 16039–16050.
48. Okano A, Nakayama A, Wu K, Lindsey EA, Schammel AW, Feng Y, Collins KC and Boger DL, *J. Am. Chem. Soc.*, 2015, 137, 3693–3704. [PubMed: 25750995]
49. Okano A, Nakayama A, Schammel AW and Boger DL, *J. Am. Chem. Soc.*, 2014, 136, 13522–13525. [PubMed: 25211770]
50. Okano A, Isley NA and Boger DL, *Proc. Natl. Acad. Sci. USA*, 2017, 114, E5052–E5061. [PubMed: 28559345]
51. Wu Z-C, Isley NA, Okano A, Weiss WJ and Boger DL, *J. Org. Chem.*, 2020, 85, 1365–1375. [PubMed: 31670958]
52. Wu Z-C, Cameron MD and Boger DL, *ACS Infect. Dis.*, 2020, 6, 2169–2180.
53. Wu Z-C and L Boger D, *Acc. Chem. Res.*, 2020, 53, in press.
54. Nerenberg JB, Hung DT, Somers PK and Schreiber SL, *J. Am. Chem. Soc.*, 1993, 115, 12621–12622.
55. Smith III AB, Qiu Y, Jones DR and Kobayashi K, *J. Am. Chem. Soc.*, 1995, 117, 12011–12012.
56. Harried SS, Yang G, Strawn MA and Myles DC, *J. Org. Chem.*, 1997, 62, 6098–6099.
57. Marshall JA and Johns BA, *J. Org. Chem.*, 1998, 63, 7885–7892.
58. Paterson I, Florence GJ, Gerlach K and Scott J, *Angew. Chem., Int. Ed.*, 2000, 39, 377–380.
59. Smith III AB, Beauchamp TJ, LaMarche MJ, Kaufman MD, Qiu Y, Arimoto H, Jones DR and Kobayashi K, *J. Am. Chem. Soc.*, 2000, 122, 8654–8664.

60. Mickel SJ, Niederer D, Daeffler R, Osmani A, Kuesters E, Schmid E, Schaer K, Gamboni R, Chen W, Loeser E, Kinder FR Jr., Konigsberger K, Prasad K, Ramsay TM, Repic O, Wang RM, Florence G, Lyothier I and Paterson I, *Org. Process Res. Dev.*, 2004, 8, 122–130.
61. Smith III AB and Freeze BS, *Tetrahedron*, 2007, 64, 261–298. [PubMed: 21113402]
62. Smith III AB, Freeze BS, LaMarche MJ, Sager J, Kinzler KW and Vogelstein B, *Bioorg. Med. Chem. Lett.*, 2005, 15, 3623–3626. [PubMed: 15979874]
63. Shaw SJ, Sundermann KF, Burlingame MA, Myles DC, Freeze BS, Xian M, Brouard I and Smith III AB, *J. Am. Chem. Soc.*, 2005, 127, 6532–6533. [PubMed: 15869264]
64. Gunasekera SP, Longley RE and Isbrucker RA, *J. Nat. Prod.*, 2001, 64, 171–174. [PubMed: 11429994]
65. Hung DT, Nerenberg JB and Schreiber SL, *J. Am. Chem. Soc.*, 1996, 118, 11054–11080.
66. Smith III AB, Freeze BS, LaMarche MJ, Hirose T, Brouard I, Rucker PV, Xian M, Sundermann KF, Shaw SJ, Burlingame MA, Horwitz SB and Myles DC, *Org. Lett.*, 2005, 7, 311–314. [PubMed: 15646985]
67. Martello LA, LaMarche MJ, He L, Beauchamp TJ, Smith III AB and Horwitz SB, *Chem. Biol.*, 2001, 8, 843–855. [PubMed: 11564553]
68. Shaw SJ, *Mini Rev. Med. Chem.*, 2008, 8, 276–284. [PubMed: 18336347]
69. Chopra I and Roberts M, *Microbiol. Mol. Biol. Rev.*, 2001, 65, 232–260. [PubMed: 11381101]
70. Connell SR, Tracz DM, Nierhaus KH and Taylor DE, *Antimicrob. Agents Chemother.*, 2003, 47, 3675–3681. [PubMed: 14638464]
71. Sun C, Wang Q, Brubaker JD, Wright PM, Lerner CD, Noson K, Charest M, Siegel DR, Wang YM and Myers AG, *J. Am. Chem. Soc.*, 2008, 130, 17913–17927. [PubMed: 19053822]
72. Kummer DA, Li D, Dion A and Myers AG, *Chem. Sci.*, 2011, 2, 1710–1718. [PubMed: 21949588]
73. Liu F and Myers AG, *Curr. Opin. Chem. Biol.*, 2016, 32, 48–57. [PubMed: 27043373]
74. Zhanel GG, Cheung D, Adam H, Zelenitsky S, Golden A, Schweizer F, Gorityala B, Lagacé-Wiens PR, Walkty A, Gin AS, Hoban DJ and Karlowsky JA, *Drugs*, 2016, 76, 567–588. [PubMed: 26863149]
75. Sakoulas G, Kumaraswamy M, Kousha A and Nizet V, *mSphere*, 2017, 2, e0004–17.
76. Sun C, Deng Y, Hunt D, Fyfe C, Kerstein K and Xiao X, poster presented at ASM Microbe, New Orleans, 6, 2017.
77. Seiple IB, Zhang Z, Jakubec P, Langlois-Mercier A, Wright PM, Hog DT, Yabu K, Allu SR, Fukuzaki T, Carlsen PN, Kitamura Y, Zhou X, Condakes ML, Szczypiński FT, Green WD and Myers AG, *Nature*, 2016, 533, 338–345. [PubMed: 27193679]
78. Hogan PC, Chen CL, Mulvihill KM, Lawrence JF, Moorhead E, Rickmeier J and Myers AG, *J. Antibiot.*, 2018, 71, 318–325.
79. Wender PA, in *Products in Medicinal Chemistry*, ed. Hanessian S, Wiley-VCH, Weinheim, 2014, pp. 475–543.
80. Wender PA, Baryza JL, Hilinski MK, Horan JC, Kan C and Verma VA, in *Drug Discovery Research: New Frontiers in the PostGenomic Era*, ed. Huang Z, Wiley-VCH, Weinheim, 2007, pp. 127–162.
81. Wu R, Chen H, Chang N, Xu Y, Jiao J and Zhang H, *Chem. Eur. J.*, 2020, 26, 1166–1195.
82. Keck GE, Poudel YB, Cummins TJ, Rudra A and Covell JA, *J. Am. Chem. Soc.*, 2011, 133, 744–747. [PubMed: 21175177]
83. Wender PA, Hardman CT, Ho S, Jeffreys MS, Maclaren JK, Quiroz RV, Ryckbosch SM, Shimizu AJ, Sloane JL and Stevens MC, *Science*, 2017, 358, 218–223. [PubMed: 29026042]
84. Wender PA, Baryza JL, Bennett CE, Bi FC, Brenner SE, Clarke MO, Horan JC, Kan C, Lacôte E, Lippa B, Nell PG and Turner TM, *J. Am. Chem. Soc.*, 2002, 124, 13648–13649. [PubMed: 12431074]
85. Wender PA and Verma VA, *Org. Lett.*, 2006, 8, 1893–1896. [PubMed: 16623578]
86. Staveness D, Abdelnabi R, Schrier AJ, Loy BA, Verma VA, DeChristopher BA, Near KE, Neyts J, Delang L, Leyssen P and Wender PA, *J. Nat. Prod.*, 2016, 79, 675–679. [PubMed: 26900625]

87. Keck GE, Kraft MB, Truong AP, Li W, Sanchez CC, Kedei N, Lewin NE and Blumberg PM, *J. Am. Chem. Soc.*, 2008, 130, 6660–6661. [PubMed: 18452293]
88. Keck GE, Poudel YB, Rudra A, Stephens JC, Kedei N, Lewin NE, Peach ML and Blumberg PM, *Angew. Chem., Int. Ed.*, 2010, 49, 4580–4584.
89. E Keck G, Poudel YB, Welch DS, Kraft MB, Truong AP, Stephens JC, Kedei N, Lewin NE and Blumberg PM, *Org. Lett.*, 2009, 11, 593–596. [PubMed: 19113896]
90. Keck GE, Li W, Kraft MB, Kedei N, Lewin NE and Blumberg PM, *Org. Lett.*, 2009, 11, 2277–2280. [PubMed: 19419164]
91. Marcu MG, Schulte TW and Neckers L, *J. Natl. Cancer Inst.*, 2000, 92, 242–248. [PubMed: 10655441]
92. Burlison JA, Neckers L, Smith AB, Maxwell A and Blagg BS, *J. Am. Chem. Soc.*, 2006, 128, 15529–15536. [PubMed: 17132020]
93. Burlison JA, Avila C, Vielhauer G, Lubbers DJ, Holzbeierlein J and Blagg BS, *J. Org. Chem.*, 2008, 73, 2130–2137. [PubMed: 18293999]
94. Zhao H, Donnelly AC, Kusuma BR, Brandt GE, Brown D, Rajewski RA, Vielhauer G, Holzbeierlein J, Cohen MS and Blagg BS, *J. Med. Chem.*, 2011, 54, 3839–3853. [PubMed: 21553822]
95. Byrd KM, Subramanian C, Sanchez J, Motiwala HF, Liu W, Cohen MS, Holzbeierlein J and Blagg BS, *Chem. Eur. J.*, 2016, 22, 6921–6931. [PubMed: 27037933]
96. Garg G, Zhao H and Blagg BS, *ACS Med. Chem. Lett.*, 2015, 6, 204–209. [PubMed: 25699150]
97. Lindquist N, Fenical W, Van Duyne GD and Clardy J, *J. Am. Chem. Soc.*, 1991, 113, 2303–2304.
98. Nicolaou KC, Hao J, Reddy MV, Rao PB, Rassias G, Snyder SA, Huang X, Chen DY-K, Brenzovich WE, Giuseppone N, O'Brate A and Giannakakou P, *J. Am. Chem. Soc.*, 2004, 126, 12897–12906. [PubMed: 15469287]
99. Nicolaou KC, Bella M, Chen DY-K, Huang X, Ling T and Snyder SA, *Angew. Chem., Int. Ed.*, 2002, 41, 3495–3499.
100. Burgett WG, Li Q, Wei Q and Harran PG, *Angew. Chem., Int. Ed.*, 2003, 42, 4961–4966.
101. Li J, Burgett AW, Esser L, Amezcua C and Harran PG, *Angew. Chem., Int. Ed.*, 2001, 40, 4770–4773.
102. Wang G, Shang L, Burgett AW, Harran PG and Wang X, *Proc. Natl. Acad. Sci. USA*, 2007, 104, 2068–2073. [PubMed: 17287350]
103. Williams NS, Burgett AW, Atkins AS, Wang X, Harran PG and McKnight SL, *Proc. Natl. Acad. Sci. USA*, 2007, 104, 2074–2079. [PubMed: 17287337]
104. Ding H, DeRoy PL, Perreault C, Larivée A, Siddiqui A, Caldwell CG, Harran S and Harran PG, *Angew. Chem., Int. Ed.*, 2015, 54, 4818–4822.
105. Wei Q, Zhou M, Xu X, Caldwell C, Harran S and Wang L, *US8592469B2*, 2013.
106. Wieczorek M, Tcherkezian J, Bernier C, Prota AE, Chaaban S, Rolland Y, Godbout C, Hancock MA, Arezzo JC, Ocal O and Rocha C, *Sci. Transl. Med.*, 2016, 8, 365ra159.
107. Pearce HL, in *The Alkaloids: Chemistry and Pharmacology*, ed. Brossi A and Suffness M, Academic Press: San Diego, CA, 1990, pp. 145–204.
108. Neuss N and Neuss MN, in *The Alkaloids: Chemistry and Pharmacology*, ed. Brossi A and Suffness M, Academic Press: San Diego, CA, 1990, pp. 229–240.
109. Kuehne ME and Marko I, in *The Alkaloids: Chemistry and Pharmacology*, ed. Brossi A and Suffness M, Academic Press: San Diego, CA, 1990, pp. 77–132.
110. Fahy J, *Curr. Pharm. Des.*, 2001, 7, 1181–1197. [PubMed: 11472261]
111. Kutney JP, Hibino T, Jahngen E, Okutani T, Ratcliffe AH, Treasurywala AM and Wunderly S, *Helv. Chim. Acta*, 1976, 59, 2858–2882. [PubMed: 1017976]
112. Potier P, Langlois N, Langlois Y and Gueritte F, *J. Chem. Soc., Chem. Commun.*, 1975, 670–671.
113. Langlois N, Gueritte F, Langlois Y and Potier P, *J. Am. Chem. Soc.*, 1976, 98, 7017–7024. [PubMed: 965661]
114. Kutney JP, Ratcliffe AH, Treasurywala AM and Wunderly S, *Heterocycles*, 1975, 3, 639–649.
115. Kuehne ME, Matson PA and Bornmann WG, *J. Org. Chem.*, 1991, 56, 513–528.

116. Bornmann WG and Kuehne ME, *J. Org. Chem*, 1992, 57, 1752–1760.
117. Yokoshima T, Ueda, Kobayashi S, Sato A, Kuboyama T, Tokuyama H and Fukuyama T, *J. Am. Chem. Soc.*, 2002, 124, 2137–2139. [PubMed: 11878966]
118. Sears JE and Boger DL, *Acc. Chem. Res.*, 2016, 49, 241–251. [PubMed: 26813287]
119. Ishikawa H, Elliott GI, Velcicky J, Choi Y and Boger DL, *J. Am. Chem. Soc.*, 2006, 128, 10596–10612. [PubMed: 16895428]
120. Ishikawa H, Colby DA and Boger DL, *J. Am. Chem. Soc.*, 2008, 130, 420–421. [PubMed: 18081297]
121. Ishikawa H, Colby DA, Seto S, Va P, Tam A, Kakei H, Rayl TJ, Hwang I and Boger DL, *J. Am. Chem. Soc.*, 2009, 131, 4904–4916. [PubMed: 19292450]
122. Gotoh H, Sears JE, Eschenmoser A and Boger DL, *J. Am. Chem. Soc.*, 2012, 134, 13240–13243. [PubMed: 22856867]
123. Boon BA and Boger DL, *J. Am. Chem. Soc.*, 2019, 141, 14349–14355. [PubMed: 31442047]
124. Sears JE and Boger DL, *Acc. Chem. Res.*, 2015, 48, 653–662. [PubMed: 25586069]
125. Gotoh H, Duncan KK, Robertson WM and Boger DL, *ACS Med. Chem. Lett.*, 2011, 2, 948–952. [PubMed: 22247789]
126. Allemann O, Brutsch M, Lukesh III JC, Brody DM and Boger DL, *J. Am. Chem. Soc.*, 2016, 138, 8376–8379. [PubMed: 27356080]
127. Leggans EK, Barker TJ, Duncan KK and Boger DL, *Org. Lett.*, 2012, 14, 1428–1431. [PubMed: 22369097]
128. Lukesh III JC, Carney DW, Dong H, Cross RM, Shukla V, Duncan KK, Yang S, Brody DM, Brutsch MM, Radakovic A and Boger DL, *J. Med. Chem.*, 2017, 60, 7591–7604. [PubMed: 28857558]
129. Leggans EK, Duncan KK, Barker TJ, Schleicher KD and Boger DL, *J. Med. Chem.*, 2013, 56, 628–639. [PubMed: 23244701]
130. Barker TJ, Duncan KK, Otrubova K and Boger DL, *ACS Med. Chem. Lett.*, 2013, 4, 985–988.
131. Carney DW, Lukesh III JC, Brody DM, Brutsch MM and Boger DL, *Proc. Natl. Acad. Sci. USA*, 2016, 113, 9691–9698. [PubMed: 27512044]
132. Schimana J, Gebhardt K, Hoeltzel A, Schmid DG, Suessmuth R, Mueller J, Pukall R and Fiedler HP, *J. Antibiot.*, 2002, 55, 565–570.
133. Kulanthaivel P, Kreuzman AJ, Strege MA, Belvo MD, Smitka TA, Clemens M, Swartling JR, Minton KL, Zheng F, Angleton EL, Mullen D, Jungheim LN, Klimkowski VL, Nicas TI, Thompson RC and Peng SB, *J. Biol. Chem.*, 2004, 279, 36250–36258. [PubMed: 15173160]
134. Smith PA, Roberts TC and Romesberg FE, *Chem. Biol.*, 2010, 17, 1223–1231. [PubMed: 21095572]
135. Tan YX and Romesberg FE, *Med. Chem. Commun.*, 2012, 3, 916–925.
136. Liu J, Smith PA, Steed DB and Romesberg FE, *Bioorg. Med. Chem. Lett.*, 2013, 23, 5654–5659. [PubMed: 24012184]
137. Roberts TC, Smith PA, Cirz RT and Romesberg FE, *J. Am. Chem. Soc.*, 2007, 129, 15830–15838. [PubMed: 18052061]
138. Peters DS, Romesberg FE and Baran PS, *J. Am. Chem. Soc.*, 2018, 140, 2072–2075. [PubMed: 29381350]
139. Smith PA, Koehler MF, Girgis HS, Yan D, Chen Y, Chen Y, Crawford JJ, Durk MR, Higuchi RI, Kang J, Murray J, Paraselli P, Park S, Phung W, Quinn JG, Roberts TC, Rougé L, Schwarz JB, Skippington E, Wai J, Xu M, Yu Z, Zhang H, Tan MW and Heise CE, *Nature*, 2018, 561, 189–194. [PubMed: 30209367]
140. Wong N, Petronijevic F, Hong AY, Linghu X, Kelly SM, Hou H, Cravillion T, Lim NK, Robinson SJ, Han C, Molinaro C, Sowell C and Gosselin F, *Org. Lett.*, 2019, 21, 9099–9103. [PubMed: 31668077]
141. Bollag DM, McQueney PA, Zhu J, Hensens O, Koupal L, Liesch J, Goetz M, Lazarides E and Woods CM, *Cancer Res.*, 1995, 55, 2325–2333. [PubMed: 7757983]
142. Lee FY, Borzilleri R, Fairchild CR, Kim SH, Long BH, Reventos-Suarez C, Vite GD, Rose WC and Kramer RA, *Clin. Cancer Res.*, 2001, 7, 1429–1437. [PubMed: 11350914]

143. Borzilleri RM, Zheng X, Schmidt RJ, Johnson JA, Kim SH, DiMarco JD, Fairchild CR, Gougoutas JZ, Lee FY, Long BH and Vite GD, *J. Am. Chem. Soc.*, 2000, 122, 8890–8897.
144. Lee FY, Borzilleri R, Fairchild CR, Kamath A, Smykla R, Kramer R and Vite G, *Cancer Chemother. Pharmacol.*, 2008, 63, 157–166. [PubMed: 18347795]
145. Su DS, Meng D, Bertinato P, Balog A, Sorensen EJ, Danishefsky SJ, Zheng YH, Chou TC, He L and Horwitz SB, *Angew. Chem., Int. Ed. Engl.*, 1997, 36, 757–759.
146. Harris CR, Kuduk SD, Balog A, Savin K, Glunz PW and Danishefsky SJ, *J. Am. Chem. Soc.*, 1999, 121, 7050–7062.
147. Chou TC, Zhang XG, Balog A, Su DS, Meng D, Savin K, Bertino JR and Danishefsky SJ, *Proc. Natl. Acad. Sci. USA*, 1998, 95, 9642–9647. [PubMed: 9689134]
148. Rivkin A, Yoshimura F, Gabarda AE, Chou TC, Dong H, Tong WP and Danishefsky SJ, *J. Am. Chem. Soc.*, 2003, 125, 2899–2901. [PubMed: 12617656]
149. Rivkin A, Yoshimura F, Gabarda AE, Cho YS, Chou TC, Dong H and Danishefsky SJ, *J. Am. Chem. Soc.*, 2004, 126, 10913–10922. [PubMed: 15339176]
150. Chou TC, Zhang X, Zhong ZY, Li Y, Feng L, Eng S, Myles DR, Johnson R, Wu N, Yin YI, Wilson RM and Danishefsky SJ, *Proc. Natl. Acad. Sci. USA*, 2008, 105, 13157–13162. [PubMed: 18755900]
151. Ichimura M, Ogawa T, Takahashi K, Kobayashi E, Kawamoto I, Yasuzawa T, Takahashi I and Nakano H, *J. Antibiot.*, 1990, 43, 1037–1038.
152. Martin DG, Biles C, Gerpheide SA, Hanka LJ, Krueger WC, McGovren JP, Mizsak SA, Neil GL, Stewart JC and Visser J, *J. Antibiot.*, 1981, 34, 1119–1125.
153. Boger DL, Boyce CW, Garbaccio RM and Goldberg JA, *Chem. Rev.*, 1997, 97, 787–828. [PubMed: 11848889]
154. Boger DL and Coleman RS, *J. Am. Chem. Soc.*, 1988, 110, 4796–4807.
155. Boger DL, Machiya K, Hertzog DL, Kito PA and Holmes D, *J. Am. Chem. Soc.*, 1993, 115, 9025–9036.
156. Boger DL, McKie JA, Nishi T and Ogiku T, *J. Am. Chem. Soc.*, 1997, 119, 311–325.
157. Tichenor MS, Trzuppek JD, Kastrinsky DB, Shiga F, Hwang I and Boger DL, *J. Am. Chem. Soc.*, 2006, 128, 15683–15696. [PubMed: 17147378]
158. Boger DL and Garbaccio RM, *Acc. Chem. Res.*, 1999, 32, 1043–1052.
159. Parrish JP, Hughes TV, Hwang I and Boger DL, *J. Am. Chem. Soc.*, 2004, 126, 80–81. [PubMed: 14709069]
160. MacMillan KS and Boger DL, *J. Med. Chem.*, 2009, 52, 5771–5780. [PubMed: 19639994]
161. Boger DL and Johnson DS, *Angew. Chem., Int. Ed. Engl.*, 1996, 35, 1438–1474.
162. Boger DL, *Acc. Chem. Res.*, 1995, 28, 20–29.
163. Boger DL and Johnson DS, *Proc. Natl. Acad. Sci. USA*, 1995, 92, 3642–3649. [PubMed: 7731958]
164. Wolkenberg SE and Boger DL, *Chem. Rev.*, 2002, 102, 2477–2496. [PubMed: 12105933]
165. Tse WC and Boger DL, *Chem. Biol.*, 2004, 11, 1607–1617. [PubMed: 15610844]
166. Boger DL, Johnson DS and Yun W, *J. Am. Chem. Soc.*, 1994, 116, 1635–1656.
167. Tichenor MS and Boger DL, *Nat. Prod. Rep.*, 2008, 25, 220–226. [PubMed: 18389136]
168. Tichenor MS, MacMillan KS, Stover JS, Wolkenberg SE, Pavani MG, Zanella L, Zaid AN, Spalluto G, Rayl TJ, Hwang I, Baraldi PG and Boger DL, *J. Am. Chem. Soc.*, 2007, 129, 14092–14099. [PubMed: 17948994]
169. Wolfe AL, Duncan KK, Parelkar NK, Brown D, Vielhauer GA and Boger DL, *J. Med. Chem.*, 2013, 56, 4104–4115. [PubMed: 23627265]
170. Jin W, Trzuppek JD, Rayl TJ, Broward MA, Vielhauer GA, Weir SJ, Hwang I and Boger DL, *J. Am. Chem. Soc.*, 2007, 129, 15391–15397. [PubMed: 18020335]
171. Nakae K, Yoshimoto Y, Sawa T, Homma Y, Hamada M, Takeuchi T and Imoto M, *J. Antibiot.*, 2000, 53, 1130–1136.
172. Nakae K, Yoshimoto Y, Ueda M, Sawa T, Takahashi Y, Naganawa H, Takeuchi T and Imoto M, *J. Antibiot.*, 2000, 53, 1228–1230.

173. Gaul C, Njardarson JT and Danishefsky SJ, *J. Am. Chem. Soc.*, 2003, 125, 6042–6043. [PubMed: 12785819]
174. Gaul C, Njardarson JT, Shan D, Dorn DC, Wu KD, Tong WP, Huang XY, Moore MA and Danishefsky SJ, *J. Am. Chem. Soc.*, 2004, 126, 11326–11337. [PubMed: 15355116]
175. Njardarson JT, Gaul C, Shan D, Huang XY and Danishefsky SJ, *J. Am. Chem. Soc.*, 2004, 126, 1038–1040. [PubMed: 14746469]
176. Oskarsson T, Nagorny P, Krauss JJ, Perez L, Mandal M, Yang G, Ouerfelli O, Xiao D, Moore MA, Massagué J and Danishefsky SJ, *J. Am. Chem. Soc.*, 2010, 132, 3224–3228. [PubMed: 20155906]
177. Lecomte N, Njardarson JT, Nagorny P, Yang G, Downey R, Ouerfelli O, Moore MA and Danishefsky SJ, *Proc. Natl. Acad. Sci. USA*, 2011, 108, 15074–15078. [PubMed: 21808037]
178. Hamberg M and Samuelsson B, *Proc. Natl. Acad. Sci. USA*, 1973, 70, 899–903. [PubMed: 4514999]
179. Kuehl FA, Humes JL, Egan RW, Ham EA, Beveridge GC and Van Arman CG, *Nature*, 1977, 265, 170–173. [PubMed: 834259]
180. Hamberg M, Svensson J and Samuelsson B, *Proc. Natl. Acad. Sci. USA*, 1975, 72, 2994–2998. [PubMed: 1059088]
181. Corey EJ, Weinshenker NM, Schaaf TK and Huber W, *J. Am. Chem. Soc.*, 1969, 91, 5675–5677. [PubMed: 5808505]
182. Corey EJ, Schaaf TK, Huber W, Koelliker U and Weinshenker NM, *J. Am. Chem. Soc.*, 1970, 92, 397–398. [PubMed: 5411057]
183. Corey EJ, Wlodawer P, Samuelsson B and Albonico SM, *J. Am. Chem. Soc.*, 1971, 93, 2815–2816. [PubMed: 5573726]
184. Corey EJ, Shibasaki M, Nicolaou KC, Malmsten CL and Samuelsson B, *Tetrahedron Lett*, 1976, 17, 737–740.
185. Bundy GL, *Tetrahedron Lett*, 1975, 16, 1957–1960.
186. Coleman RA, Humphrey PP, Kennedy I, Levy GP and Lumley P, *Br. J. Pharmacol.*, 1980, 68, 127P–128P.
187. Fried J, John V, Szwedo MJ Jr, Chen CK, O'Yang C, Morinelli TA, Okwu AK and Halushka PV, *J. Am. Chem. Soc.*, 1989, 111, 4510–4511.
188. Jing C, Mallah S, Kriemen E, Bennett SH, Fasano V, Lennox AJ, Hers I and Aggarwal VK, *ACS Cent. Sci.*, 2020, 6, 995–1000. [PubMed: 32607446]
189. Ortega A, Blount JF and Manchand PS, *J. Chem. Soc. Perkin Trans 1*, 1982, 2505–2508.
190. Roth BL, Baner K, Westkaemper R, Siebert D, Rice KC, Steinberg S, Ernsberger P and Rothman RB, *Proc. Natl. Acad. Sci. USA*, 2002, 99, 11934–11939. [PubMed: 12192085]
191. Sherwood AM, Crowley RS, Paton KF, Biggerstaff A, Neuenswander B, Day VW, Kivell BM and Prinszano TE, *J. Med. Chem.*, 2017, 60, 3866–3878. [PubMed: 28376298]
192. Roach JJ and Shenvi RA, *Bioorg. Med. Chem. Lett*, 2018, 28, 1436–1445. [PubMed: 29615341]
193. Roach JJ, Sasano Y, Schmid CL, Zaidi S, Katritch V, Stevens RC, Bohn LM and Shenvi RA, *ACS Cent. Sci.*, 2017, 3, 1329–1336. [PubMed: 29296674]
194. Hirasawa S, Cho M, Brust TF, Roach JJ, Bohn LM and Shenvi RA, *Bioorg. Med. Chem. Lett*, 2018, 28, 2770–2772. [PubMed: 29426768]
195. Nakajima H, Sato B, Fujita T, Takase S, Terano H and Okuhara M, *J. Antibiot.*, 1996, 49, 1196–1203.
196. Kaida D, Motoyoshi H, Tashiro E, Nojima T, Hagiwara M, Ishigami K, Watanabe H, Kitahara T, Yoshida T, Nakajima H, Tani T, Horinouchi S and Yoshida M, *Nat. Chem. Biol.*, 2007, 3, 576–583. [PubMed: 17643111]
197. Albert BJ, McPherson PA, O'Brien K, Czaicki NL, DeStefino V, Osman S, Li M, Day BW, Grabowski PJ, Moore MJ, Vogt A and Koide K, *Mol. Cancer Ther.*, 2009, 8, 2308–2318. [PubMed: 19671752]
198. Thompson CF, Jamison TF and Jacobsen EN, *J. Am. Chem. Soc.*, 2001, 123, 9974–9983. [PubMed: 11592876]
199. Horigome M, Motoyoshi H, Watanabe H and Kitahara T, *Tetrahedron Lett*, 2001, 42, 8207–8210.

200. Albert BJ, Sivaramakrishnan A, Naka T and Koide K, J. Am. Chem. Soc, 2006, 128, 2792–2793. [PubMed: 16506745]
201. Albert BJ, Sivaramakrishnan A, Naka T, Czaicki NL and Koide K, J. Am. Chem. Soc, 2007, 129, 2648–2659. [PubMed: 17279752]
202. Gao Y, Vogt A, Forsyth CJ and Koide K, Chembiochem, 2013, 14, 49–52. [PubMed: 23172726]
203. Osman S, Albert BJ, Wang Y, Li M, Czaicki NL and Koide K, Chem. Eur. J, 2011, 17, 895–904. [PubMed: 21226105]
204. Walker S, Chen L, Hu Y, Rew Y, Shin D and Boger DL, Chem. Rev, 2005, 105, 449–476. [PubMed: 15700952]
205. Fulco P and Wenzel RP, Expert Rev. Anti Infect. Ther, 2006, 4, 939–945. [PubMed: 17181409]
206. Parenti F, Ciabatti R, Cavalleri B and Kettenring J, Drugs Exp. Clin. Res, 1990, 16, 451–455. [PubMed: 2100246]
207. Jiang W, Wanner J, Lee RJ, Bounaud PY and Boger DL, J. Am. Chem. Soc, 2003, 125, 1877–1887. [PubMed: 12580615]
208. Jiang W, Wanner J, Lee RJ, Bounaud PY and Boger DL, J. Am. Chem. Soc, 2002, 124, 5288–5290. [PubMed: 11996568]
209. Shin D, Rew Y and Boger DL, Proc. Natl. Acad. Sci. USA, 2004, 101, 11977–11979. [PubMed: 15175429]
210. Rew Y, Shin D, Hwang I and Boger DL, J. Am. Chem. Soc, 2004, 126, 1041–1043. [PubMed: 14746470]
211. Nam J, Shin D, Rew Y and Boger DL, J. Am. Chem. Soc, 2007, 129, 8747–8755. [PubMed: 17592838]
212. Falagas ME, Kasiakou SK and Saravolatz LD, Clin. Infect. Dis, 2005, 40, 1333–1341. [PubMed: 15825037]
213. Zavascki AP, Goldani LZ, Li J and Nation RL, J. Antimicrob. Chemother, 2007, 60, 1206–1215. [PubMed: 17878146]
214. Kadar B, Kocsis B, Nagy K and Szabo D, Curr. Med. Chem, 2013, 20, 3759–3773. [PubMed: 23895690]
215. Kassamali Z, Rotschafer JC, Jones RN, Prince RA and Danziger LH, Clin. Infect. Dis, 2013, 57, 877–883. [PubMed: 23728146]
216. Hartzell JD, Neff R, Ake J, Howard R, Olson S, Paolino K, Vishnepolsky M, Weintrob A and Wortmann G, Clin. Infect. Dis, 2009, 48, 1724–1728. [PubMed: 19438394]
217. Falagas ME and Kasiakou SK. Crit. Care, 2006, 10, R27. [PubMed: 16507149]
218. Vaara M, Fox J, Loidl G, Siikanen O, Apajalahti J, Hansen F, Frimodt-Møller N, Nagai J, Takano M and Vaara T, Antimicrob. Agents Chemother, 2008, 52, 3229–3236. [PubMed: 18591267]
219. Mingeot-Leclercq MP, Tulkens PM, Denamur S, Vaara T and Vaara M, Peptides, 2012, 35, 248–252. [PubMed: 22504013]
220. Gallardo-Godoy A, Muldoon C, Becker B, Elliott AG, Lash LH, Huang JX, Butler MS, Pelingon R, Kavanagh AM, Ramu S, Phetsang W, Blaskovich MAT and Cooper MA, J. Med. Chem, 2016, 59, 1068–1077. [PubMed: 26734854]
221. Xie Y, Chen R, Si S, Sun C and Xu H, J. Antibiot, 2007, 60, 158–161.
222. Xie Y, Cai Q, Ren H, Wang L, Xu H, Hong B, Wu L and Chen R, J. Nat. Prod, 2014, 77, 1744–1748. [PubMed: 24964393]
223. Chen RH, Buko AM, Whittern DN and McAlpine JB, J. Antibiot, 1989, 42, 512–520.
224. Isono F, Inukai M, Takahashi S, Haneishi T, Kinoshita T and Kuwano H, J. Antibiot, 1989, 42, 667–673.
225. Chatterjee S, Nadkarni SR, Vijayakumar EK, Patel MV, Ganguli BN, Fehlhaber HW and Vertesy L, J. Antibiot, 1994, 47, 595–598.
226. Tran AT, Watson EE, Pujari V, Conroy T, Dowman LJ, Giltrap AM, Pang A, Wong WR, Linington RG, Mahapatra S, Saunders J, Charman SA, West NP, Bugg TDH, Tod J, Dowson CG, Roper DI, Crick DC, Britton WJ and Payne RJ, Nat. Commun, 2017, 8, 1–9. [PubMed: 28232747]

227. Pettit GR, Srirangam JK, Barkoczy J, Williams MD, Durkin KP, Boyd MR, Bai R, Hamel E, Schmidt JM and Chapuis JC, *Anticancer Drug Des*, 1995, 10, 529–544. [PubMed: 7495477]
228. Pettit GR, Srirangam JK, Barkoczy J, Williams MD, Boyd MR, Hamel E, Pettit RK, Hogan F, Bai R, Chapuis JC, McAllister SC and Schmidt JM, *Anticancer Drug Des*, 1998, 13, 243–277. [PubMed: 9627667]
229. Mohammad RM, Limvarapuss C, Wall NR, Hamdy N, Beck FW, Pettit GR and Al-Katib A, *Int. J. Oncol*, 1999, 15, 367–439. [PubMed: 10402249]
230. Doronina SO, Toki BE, Torgov MY, Mendelsohn BA, Cervený CG, Chace DF, DeBlanc RL, Gearing RP, Bovee TD, Siegall CB, Francisco JA, Wahl AF, Meyer DL and Senter PD, *Nat. Biotechnol*, 2003, 21, 778–784. [PubMed: 12778055]
231. Bierer BE, Somers PK, Wandless TJ, Burakoff SJ and Schreiber SL, *Science*, 1990, 250, 556–559. [PubMed: 1700475]
232. Somers PK, Wandless TJ and Schreiber SL, *J. Am. Chem. Soc*, 1991, 113, 8045–8056.
233. Chakraborty TK, Weber HP and Nicolaou KC, *Chem. Biol*, 1995, 2, 157–161. [PubMed: 9383417]
234. Mitsui I, Kumazawa E, Hirota Y, Aonuma M, Sugimori M, Ohsuki S, Uoto K, Ejima A, Terasawa H and Sato K, *Jpn. J. Cancer Res*, 1995, 86, 776–782. [PubMed: 7559102]
235. Joto N, Ishii M, Minami M, Kuga H, Mitsui I and Tohgo A, *Int. J. Cancer*, 1997, 72, 680–686. [PubMed: 9259410]
236. Edmondson S, Danishefsky SJ, Sepp-Lorenzino L and Rosen N, *J. Am. Chem. Soc*, 1999, 121, 2147–2155.

Fig. 1.
Halichondrin B and its C1-C38 fragment, norhalichondrin B, eribulin and E7130.

Fig. 2.

Top: Structure, ligand binding affinity and activity of pocket modified vancomycin aglycons.
 Bottom: Representative synthetic vancomycin analogue with three independent synergistic mechanisms of action.

Fig. 3.
Structure and activity of discodermolide and select analogues.

Fig. 4. Tetracycline, tigecycline and synthetic tetracyclines prepared by the Myers modular total synthesis.

Fig. 5.
Bryostatin 1 and improved, simplified analogues.

Fig. 7.
Diazonamide A (**30**) and DZ-2384 (**31**), and the oxidative macrocyclization strategy.

32, vinblastine
 X = H, R = CH₃
33, vincristine
 X = H, R = CHO
34, C10'-fluorovinblastine
 X = F, R = CH₃
35, C10'-fluorovincristine
 X = F, R = CHO

Compd	IC ₅₀ (nM)	
	L1210	HCT116
32	6.0	6.8
34	0.7	0.8
33	6.0	7.0
35	0.8	1.5
36	0.6	0.7

Fig. 8.
 Vinblastine and vincristine analogues with improved activity.

vinblastine C20' amine and urea analogues:

(>60 urea analogues investigated)

vinblastine C20' isoindoline urea analogues:

Compound	IC ₅₀ (nM)	
	HCT116	HCT116/VM46
32	6.8	600
37	600	>10000
38	7.5	4400
39	0.7	50
40	0.5	4.0
41	0.38	5.6
42	0.074	0.88
43	0.067	0.87
44	0.075	0.83

Fig. 9.

Ultrapotent vinblastines. Left: structure and activity of vinblastine C20' ureas. Right: model of the ultrapotent vinblastine **44** in the vinblastine binding site of tubulin.

Fig. 10.
Predictable modulation of activity of vinblastine C20' benzamide analogues by a p -substituent electronic effect.

Fig. 11. Vinblastine C20' amide analogues **45**–**55** and plot of cLogP versus differential activity (IC_{50} ratio) for isogenic HCT116 sensitive and resistant (Pgp overexpression) cell lines.

IC ₅₀ (SPase, nM)			56 , arylomycin A ₁₆ , X = OH 57 , X = NHCH ₂ CHO
Compd	<i>E. coli</i>	<i>S. aureus</i>	
56	820	16000	
57	15	49

MIC (μg/mL)					
Compd	<i>E. coli</i>	<i>K. pneumoniae</i>	<i>E. cloacae</i>	<i>A. baumannii</i>	<i>P. aeruginosa</i>
56	> 64	> 64	> 64	> 64	> 64
58	0.125	0.125	0.5	1	2

Fig. 12.
Arylomycin A₁₆ and representative analogues.

Compd	CCRF-CEM IC ₅₀ (nM)	mouse plasma stability t _{1/2} (min)	solubility (H ₂ O, µg/mL)
60	0.53	57.5	-
62	3.26	46	10.7
63	0.27	84	26.5
64	0.71	212	32.1
65	0.16	> 1440	75.9
66	0.27	> 1440	61.9

Fig. 13.
 Evolution of stable and potent epothilone B analogues as antitumor agents.

Fig. 14.

Top: Duocarmycin SA and CC-1065 analogues. Bottom: Structure and *in vivo* efficacy of a structurally related prodrug.

Compound	IC ₅₀ (4T1, μ M)	stability ($t_{1/2}$, mouse plasma)
74	29	stable
75	0.022	20 min
76	0.024	< 5 min
77	0.26	stable
78	0.10	stable
79	0.47	stable
80	0.26	stable

Fig. 15.
Antimetastatic activity and stability of migrastatin and selected analogues.

Stability	
compd	t _{1/2} (pH 7.4)
81	5 min
82	stable
83	stable
84	32 s
85	320 d
86	20 d

Platelet aggregation activity		
compd	EC ₅₀ (μM)	E _{max} (%)
83	1.4	84.1
85	0.5	69.3
86	1.6	77.0

Fig. 16.

Structure, stability and activity of PGH₂, thromboxane A₂ and stable analogues.

Fig. 17.

Structure, activity and stability of salvinorin A and analogues.

Compd	antiproliferative activity GI ₅₀ (nM)	SF3b inhibition ED ₅₀ (nM)	pH 7.4 buffer t _{1/2} (h)	mouse serum t _{1/2} (h)
90	1.1	3.5	4	-
92	0.01-0.02	1.6	48	2
93	0.008	0.23	-	13

Fig. 18.
 FR901464 and stable analogues.

Compound	Authentic residue	MIC ($\mu\text{g/mL}$, <i>S. aureus</i>)
94 , ramoplanin A2 aglycon		0.11
95 , [Dap ²]ramoplanin A2 aglycon		0.7
[Dap ² , D-Ala ³]ramoplanin A2 aglycon	D-Hpg ³	5.2
[Dap ² , D-Ala ⁴]ramoplanin A2 aglycon	D-Orn ⁴	3.1
[Dap ² , D-Ala ⁵]ramoplanin A2 aglycon	D-aThr ⁵	0.16
[Dap ² , Ala ⁶]ramoplanin A2 aglycon	Hpg ⁶	0.9
[Dap ² , D-Ala ⁷]ramoplanin A2 aglycon	D-Hpg ⁷	3.7
[Dap ² , Ala ⁸]ramoplanin A2 aglycon	aThr ⁸	2.5
[Dap ² , Ala ⁹]ramoplanin A2 aglycon	Phe ⁹	0.6
[Dap ² , D-Ala ¹⁰]ramoplanin A2 aglycon	D-Orn ¹⁰	38
[Dap ² , Ala ¹¹]ramoplanin A2 aglycon	Hpg ¹¹	2.5
[Dap ² , D-Ala ¹²]ramoplanin A2 aglycon	D-aThr ¹²	0.7
[Dap ² , Ala ¹³]ramoplanin A2 aglycon	Hpg ¹³	1.4
[Dap ² , Ala ¹⁵]ramoplanin A2 aglycon	Leu ¹⁵	3.0
[Dap ² , Ala ¹⁷]ramoplanin A2 aglycon	Chp ¹⁷	0.3
[Dap ² , Hpg ¹⁷]ramoplanin A2 aglycon	Chp ¹⁷	0.09

Fig. 19.

Assessment of the importance of each residue side chain of ramoplanin based on a stable analogue (**95**) of ramoplanin A2 aglycon.

$$R = X - \text{CH}_2 - \text{CH}(\text{CH}_3) - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 -$$

(X = H or CH₃)

Compd	MIC ($\mu\text{g/mL}$) <i>K. pneumoniae</i>	LDH release ^a at 1 mM	CC ₅₀ ($\mu\text{g/mL}$) LDH release ^b	CC ₅₀ (μM) HepG2
96	0.5-1	61%	23	>300
97	1-2	5%	-	-
98	1	-	>128	298
99	0.5	-	>128	287

Fig. 20.
Less toxic polymyxins.

Fig. 21. Sansanmycin A and B, and dihydrosansanmycin analogues with improved anti-tuberculosis activity.

Fig. 22.
Select examples of supernatural products prior to 2000.