REVIEW

Unique and contrasting effects of light and temperature cues on plant transcriptional programs

Mai Jarad^a, Rea Antoniou-Kourounioti^b, Jo Hepworth^b, and Julia I. Qüesta ^a

^aCentre for Research in Agricultural Genomics (CRAG), CSIC-IRTA-UAB-UB, Campus UAB, Barcelona, Spain; ^bJohn Innes Centre, Norwich Research Park, Norwich, UK

ABSTRACT

Plants have adapted to tolerate and survive constantly changing environmental conditions by reprogramming gene expression in response to stress or to drive developmental transitions. Among the many signals that plants perceive, light and temperature are of particular interest due to their intensely fluctuating nature which is combined with a long-term seasonal trend. Whereas specific receptors are key in the light-sensing mechanism, the identity of plant thermosensors for high and low temperatures remains far from fully addressed. This review aims at discussing common as well as divergent characteristics of gene expression regulation in plants, controlled by light and temperature. Light and temperature signaling control the abundance of specific transcription factors, as well as the dynamics of co-transcriptional processes such as RNA polymerase elongation rate and alternative splicing patterns. Additionally, sensing both types of cues modulates gene expression by altering the chromatin landscape and through the induction of long non-coding RNAs (lncRNAs). However, while light sensing is channeled through dedicated receptors, temperature can broadly affect chemical reactions inside plant cells. Thus, direct thermal modifications of the transcriptional machinery add another level of complexity to plant transcriptional regulation. Besides the rapid transcriptome changes that follow perception of environmental signals, plant developmental transitions and acquisition of stress tolerance depend on long-term maintenance of transcriptional states (active or silenced genes). Thus, the rapid transcriptional response to the signal (Phase I) can be distinguished from the long-term memory of the acquired transcriptional state (Phase II – remembering the signal). In this review we discuss recent advances in light and temperature signal perception, integration and memory in *Arabidopsis thaliana*, focusing on transcriptional regulation and highlighting the contrasting and unique features of each type of cue in the process.

ARTICLE HISTORY

Received 30 April 2020

Revised 26 August 2020

Accepted 31 August 2020

KEYWORDS

Light and temperature sensing; plant transcriptional regulation; chromatin dynamics; long non-coding RNAs; epigenetic memory; stress memory

1. Introduction

1.1. Light and temperature cues induce stress and developmental responses in plants

Plants are constantly exposed to varying environmental cues that drive their growth and developmental processes. Some of the more prevalent abiotic stresses that plants are subjected to are drought (water scarcity), flooding (water overload), soil salinity, harmful radiation, toxic compounds in the soil, severe temperatures (freezing, cold, heat), and light fluctuations. Among them, light and temperature interact to control the timing of all the transitions of plant development, from germination, through the juvenile and floral transitions, to seed set and senescence.

Light may vary in intensity and wavelength and, like temperature, fluctuates widely in both predictable patterns (photoperiod) and unpredictable ones (cloud cover, shading from other plants). Plants have evolved the capacity to perceive all these complex combinations of light and temperature cues and utilize these signals to adjust their response. For instance, in the Brassicaceae and some winter cereals, spring flowering requires the combination of appropriate day length, memory of past prolonged low temperatures (vernalization), and recent warm temperatures (thermomorphogenesis) [1–3]. Thus, the intertwining of both light and temperature sensing, together with the underlying signaling pathways, allow plants access to vital information that affects their growth and

CONTACT Julia I. Qüesta julia.questa@cragenomics.es Centre for Research in Agricultural Genomics (CRAG), CSIC-IRTA-UAB-UB, Campus UAB, Barcelona, Spain; Jo Hepworth jo.hepworth@jic.ac.uk John Innes Centre, Norwich Research Park, Norwich NR4 7UH, UK; Rea Antoniou-Kourounioti rea.antoniou-kourounioti@jic.ac.uk John Innes Centre, Norwich Research Park, Norwich NR4 7UH, UK

© 2020 Informa UK Limited, trading as Taylor & Francis Group

developmental responses. In addition, plants have developed mechanisms to counteract exposure to harmful radiation and to extreme temperatures, driving stress acclimation responses for survival.

Perception and integration of light and temperature signals, both developmental triggers and stressors, result in transcriptional reprogramming. Decades of work have deciphered numerous Transcription Factors (TFs, activators, and repressors) that are responsive to light and/or temperature. However, recent efforts have started to shed light into other players such as the chromatin landscape, non-coding transcripts and components of the transcriptional machinery. In this review, we summarize the different levels of transcriptional regulation in plants and discuss how they can be affected by light and temperature cues. Our aim is to highlight common and distinct features of light and temperature modes of action on transcriptional regulation in plants.

1.2. Transcriptional regulation in plants

In eukaryotes, transcription of mRNAs, long non-coding RNAs (lncRNAs), miRNA precursors and enhancer RNAs (eRNAs) start with recruitment of the RNA POLYMERASE II (RNAPII) to proximal or distal regulatory regions of genes (promoters and enhancers, Figure 1, [4,5,6]). RNAPII recruitment and assembly at gene promoters located in chromatin accessible regions of the genome (Figure 1(a)) is generally facilitated by sequence-specific TFs, as well as the regulatory role of enhancers (Figure 1(b)). In turn, binding of these “pioneer” TFs may also increase chromatin accessibility for RNAPII recruitment, and thus a feedback mechanism between chromatin and TFs is established. Once engaged, RNAPII switches to transcriptional elongation moving along the gene body (Figure 1(c)), the nascent transcript is co-transcriptionally spliced (Figure 1(d)), and transcription is terminated upon cleavage and polyadenylation at the 3' end of the gene (Figure 1(e)). Although research on transcriptional regulation has mostly focused on TFs triggering transcriptional initiation (Figure 1(b)), detailed mechanistic studies have revealed that transcription consists of multiple regulated steps including RNAPII pausing and stalling along genes, splicing, cleavage, and polyadenylation [7]. The degree of

chromatin accessibility adds another level of regulation, influencing not only TF binding dynamics [8] but also efficiency of the co-transcriptional processes such as splicing [9]. Together, these mechanisms determine the speed of transcript production, as well as the epigenetic state of genes: ACTIVE (ON) or INACTIVE (OFF) [10]. Post-transcriptional mechanisms including transcript stability, nuclear export, and inactivation by RNA silencing pathways (for example, the miRNA pathway) will determine transcript fate.

Plant transcriptional responses to environmental threats mostly happen rapidly and transiently upon sensing of the triggering signal. Initial perception is integrated into signaling pathways that confluence in the induction of a set of pioneer TFs. The function of those TFs is to reprogramme the transcriptome by activating or repressing genes (transcriptional activators and repressors, respectively). However, plant acclimation to stress and induction of developmental transitions require that the transcriptional status acquired by specific genes (ON or OFF) is maintained for prolonged periods of time even in the absence of the triggering signal. Epigenetic mechanisms, such as DNA methylation and histone post-translational modifications, have the capacity to lock genes in either the ON or OFF state, and perpetuate these states through DNA replication and cell divisions [11,12,13,14]. In the following sections, we will discuss the impact of light and temperature on different levels of transcriptional regulation, transiently, and long-term.

2. Light and temperature control of transcription factors

Light and temperature sensing triggers a cascade of transcription factors that control growth, acclimation, and other responses. Many of these downstream factors are common to the two signals, thereby facilitating integration of environmental information. Furthermore, some of the photoreceptors have recently been found to also be sensitive to temperature. In the following sections we describe the known receptors of the two signals and the downstream pathways that control signaling, highlighting the interconnections between them.

Figure 1. Transcription in eukaryotes.

a, Chromatin accessibility affects the ability of the RNA Polymerase II (RNAPII) and other proteins to transcribe genes. b, RNAPII is recruited to the gene by transcription factors at promoters and enhancers for transcription initiation. c, RNAPII can stall at gene boundaries. After release from promoter-proximal stalling, there is transcriptional elongation and 5' capping of the nascent transcript. The chromatin is directly affected by the activity of the RNAPII through changes in supercoiling ahead of and behind the polymerase. d, Splicing determines the sequence of the transcript, and often alternative transcripts can arise from the same gene as a result of this process. e, The transcript is released from RNAPII by cleavage and the template-independent Poly (A) Polymerase polyadenylates its 3' end.

2.1. Light-triggered transcriptional responses

2.1.1. Plant photoreceptors

Relatively few reactions are light-sensitive within the spectra and intensity of light at the Earth's surface. Therefore, in order to sense light, plants have evolved to produce specific receptors for light of different wavelengths, most of which require a light-sensitive cofactor molecule for their activity. Five families of photoreceptors are known to date in *Arabidopsis*: the phytochromes (far red and red light), the cryptochromes and members of the Zeitlupe protein family (both blue and UV-A), phototropins (blue, UV-A and UV-B) and the UV-B photoreceptor UVR8 (Figure 2).

The phytochromes form a family of five proteins, phyA-phyE, whose activity depends on the light-responsive, wavelength-reversible conformation of a bilin cofactor. In the case of phyB, red

light switches the phytochrome from its nascent inactive form (Pr) to its active form (Pfr) while far red light reverts Pfr to Pr again. In the active form, phytochromes move from the cytosol into the nucleus to mediate their effects through interaction with other proteins, including other photoreceptors such as cryptochromes [15,16]. Although they are partially redundant, the different phytochromes react differently to control distinct responses depending on light intensity, wavelength and their reversion dynamics [16,17]. For example, in phyB, the active form “relaxes” back to the inactive form in the absence of red light. This “dark reversion” is temperature-sensitive, giving phyB the properties of a temperature sensor in conditions with low red light [18,19].

Like the phytochromes, the two cryptochromes are partially redundant. They require a flavin

Figure 2. Multiple light signaling pathways are sensed by receptors and converge on transcription factors in the nucleus.

From the left, UV-B is received by UVR8 and triggers its monomerization, association with COP1, and movement to the nucleus, which results in TF binding and HY5 stabilization. Phototropins (blue, UV-A and UV-B) sit at the plasma membranes and light triggers their kinase activity. The members of the Zeitlupe family (ZTL, FKF1, LKP2) respond to blue light via changes in their affinities for different protein cofactors and their E3 ligase activity to affect the circadian clock (three-arrow circle), which has effects at all levels of cell activity, including the activity of PIFs. Blue and UV-A are sensed by CRYs, resulting in oligomerization and TF binding in the nucleus as well as HY5 stabilization. Phytochromes are activated and deactivated by red and far red light and respond by moving to the nucleus, destabilizing PIF TFs and preventing PIF DNA binding. From the right: light fluence and quality activates the photosynthetic machinery, and results in biogenic and stress responses, communicated to the nucleus by a variety of routes. These include the cleavage of the chloroplast-envelope bound PHD transcription factor PTM, which moves to the nucleus itself to effect transcriptional responses. Key: Light split by a prism enters the cell (pale yellow rounded box), pale blue rounded box: nucleus, pale green rounded box: chloroplast. Lightning arrows: effect of light activation. Solid arrows: protein movement. Dotted lines: processes including multiple metabolites & proteins. Policeman black/white: COP1 active/inactive. Rectangular boxes in nucleus, grey/orange/green/white: TFs, HY5/PIFs/cleaved PTM/unspecified. Photoreceptors in thick black outlines are also defined as thermosensors.

adenosine dinucleotide as cofactor. When activated by light, the cryptochromes undergo conformational change and protein homo-oligomerization to activate binding to targets, which, like the phytochromes, includes direct interaction with target TFs [20,21].

Phototropins are serine/threonine kinases that also have flavin chromophores, binding flavin mononucleotides through two LIGHT, OXYGEN OR VOLTAGE (LOV) domains. Unlike the other photoreceptors, which are nuclear or cytosolic, the two phototropins are plasma membrane targeted, though not integral membrane proteins. Photoactivation by blue, UV-A and UV-B light results in autophosphorylation and phosphorylation of target proteins, leading to movement responses at the intracellular (chloroplast movement), cellular (stomatal opening) and whole plant level (phototropism) to maximize light reception [22,23]. In common with *Arabidopsis* phyB, the phototropins have also been implicated in temperature sensing (see thermosensors section).

A single LOV domain is present in the Zeitlupe family of three flavin-binding proteins: ZTL itself, FLAVIN-BINDING, KELCH REPEAT, F-BOX1 (FKF1) and LOV KELCH REPEAT PROTEIN 2 (LKP2). All three mediate blue light cues to the circadian clock via their F-box domains as part of E3 ubiquitin ligase complexes, marking target proteins, such as CYCLING DOF TFs, for degradation [24,25,26].

Most recently discovered is the UV-B photoreceptor UV RESISTANCE LOCUS 8 (UVR8) [27,28]. Ultraviolet photon absorption by UVR8 causes monomerization from the inactive homodimer [29] leading to translocation to the nucleus and interaction with CONSTITUTIVE PHOTOMORPHOGENIC 1 (COP1) [28,30]. Uniquely, UVR8 does not require a chromophore cofactor, with light sensitivity being provided by tryptophan residues within the protein itself [31].

In addition to these dedicated receptors, light reactions in the chloroplast lead to transcriptional responses both within the chloroplast

itself and, by a variety of retrograde signaling pathways, in the nucleus. These signaling pathways include the direct outputs of photosynthetic function such as reactive oxygen species (e.g., singlet oxygen) [32,33], metabolites required for chloroplast function such as tetrapyrroles [34], and proteins from both nucleus and plastid, including transcription factors such as PHD-TYPE TRANSCRIPTION FACTOR WITH TRANSMEMBRANE DOMAINS (PTM) and the integrator TF ABSCISIC ACID INSENSITIVE4 (ABI4) [17,35,36,37,38,39]. Moreover, as the reactions of photosynthesis are inherently heat-sensitive, this retrograde signaling is also linked to temperature responses [40,41]. Similarly, circadian dynamics are a vital part of the modulation of light and temperature signaling, and are also controlled by light (e.g., via the *Zeitlupe* family) [42,43].

2.1.2. Transcription factors in the primary response to light

Light inputs from three of five photoreceptor families converge on the activity of a small number of transcription factors (Figure 2). In particular, two families of TFs dominate light responses: the basic Helix-Loop-Helix (bHLH) PHYTOCHROME INTERACTING FACTOR (PIF) family and the basic ZIPper (bZIP) TFs ELONGATED HYPOCOTYL5 (HY5) and its orthologue HY5 HOMOLOG (HYH). Light-induced changes in PIF and HY5 activity act through relief-from-repression mechanisms at different stages of the transduction route. Both families are key to plant development, particularly photomorphogenesis. Most information on light signaling, retrograde or otherwise, has been derived from study of this process, in which seedlings undergo major transcription, cellular and morphological changes from an autotrophic etiolated “soil-grown” state to green, phototropic, open-cotyledon phase.

HY5 and HYH generally act partially redundantly to promote transcription in response to red, blue and UV light signaling (Figure 2). HY5 itself lacks a transcriptional activation domain, and

recent work has further supported the hypothesis that HY5 relies on interacting TFs to upregulate its targets, which trigger a cascade of indirect transcriptional responses [44]. In the dark, HY5 and HYH, along with several other factors that promote photomorphogenesis, are ubiquitinated and marked for degradation by the 26S proteasome, through the action of the master light integrator, the E3 ligase COP1-SUPPRESSOR OF PHYA-105 (COP1/SPA) complex [45,46,47]. COP1-SPA complex action is in turn inhibited by the binding of activated (Pfr) phytochromes and cryptochrome oligomers to SPA proteins, resulting in the stabilization of HY5 in the light and promotion of photomorphogenesis [17,20]. Active UVR8 monomers also bind to COP1 in the nucleus in a light-dependent manner, outcompeting COP1 binding of HY5 [27,28,30,48]. Although several different photoreceptors converge on HY5 via COP1, they do so in different manners and have different effects. Unlike PIF and CRY signaling, UVR8 requires COP1 as a positive factor [49], with the resulting stabilization of HY5 mediated through interplay between antagonistic E3 ligases downstream of UVR8 action [50]. UVR8 also binds to WRKY DNA-BINDING PROTEIN 36 (WRKY36) to inhibit its binding to and repression of the *HY5* promoter, resulting in increased *HY5* transcription [51]. In turn, CRYs antagonize UVR8 responses over the light spectra that both receptors share [52]. HY5 also directly promotes expression of its negative regulators *COP1*, *SPA1*, and *SPA4* in a negative feedback loop [44].

The PIF family are key pioneer TFs for light signaling with critical roles in plant development, integrating light responses with temperature, hormone signaling and circadian dynamics [16,17]. In contrast to the HYs, PIFs act as negative regulators of photomorphogenesis by binding to G-box motifs in target promoters, antagonizing HY5 action at shared targets [53]. In the dark, PIF proteins are stabilized by the binding of COP1 (PIF3 and PIF5 [54,55]) and DE-ETIOLATED 1 (DET1 [56]). DET1 is a classic negative regulator of photomorphogenesis, like COP1, and similarly provides substrate specificity to a CULLIN4 E3-ligase complex [57,58,59,60]. PIF repression of photomorphogenesis is relieved in light by phytochrome-triggered phosphorylation and degradation of PIFs, by the

action of phyB in reducing PIF1 and PIF3 binding to DNA at target promoters, and by UVR8-mediated reduction in COP1-PIF5 binding, leading to PIF5 destabilization [16,61,62,63,64,65]. Although partially redundant, different PIFs have different affinities for the various phytochromes, particularly the “light labile” phyA, and for different heterodimerization partner TFs [16,17,66]. As well as their interaction with phytochromes, CRYs also interact directly with PIF4 and PIF5 at the chromatin in limiting blue light and repress PIF transcriptional activity in high temperature responses [67,68].

As well as the PIFs and HYs, all the photoreceptors, except the phototropins, are known to interact directly with other TFs to mediate their specific responses. As a few examples: phyB triggers the light-dependent degradation of EIN3 to reduce ethylene signaling [69]. UVR8, CRYs and possibly phyB interact directly with transcription factors such as BRI1-EMS-SUPPRESSOR 1 to integrate light signaling directly with the brassinosteroid signaling cascade [70,71,72]. Separately CRY2 binds with CRYPTOCHROME-INTERACTING bHLH1 (CIB1) to directly promote *FT* expression and flowering [73], while varying effects on flowering time due to different degradation targets characterizes the separate members of the Zeitlupe family [74]. Some of the protein–protein interactions in these pathways may also be directly temperature-sensitive [75]. As well as WRKY36, UVR8 also functions in root development via association with MYB73 and 77 [76].

In summary, although the various photoreceptors have distinct roles in development, they overlap in both their spectral sensitivities and target TFs. This results in a complicated crosstalk in natural, multispectral light conditions, which provides plants with the possibility of fine tuning their responses. Moreover, both photoreceptors and their interacting TFs additionally have roles in mediating temperature responses.

2.2. Influence of temperature on transcriptional initiation

2.2.1. Plant thermosensors

A thermosensor detects changes in temperature cues by inducing changes in its structure and/or

activity to adapt to the perceived signal. At the same time, temperature affects multiple processes in the plant and appropriate coupling of these temperature sensitivities can lead to compensation [77] or amplification [78] of the signal. As a thermosensor stands at the core of any thermosensing mechanism, identifying specific thermosensors has been the target of much research effort [79].

Temperature fluctuations are known to affect cellular membrane fluidity, which in turn affects the activity of membrane-bound proteins as well as their structure [80]. These changes in membrane fluidity are speculated to be the first step in the thermosensing process. Ca^{2+} channels are membrane-bound proteins that exhibit changes due to cold temperatures, triggering Ca^{2+} influx. This influx is probably sensed via the Calmodulin Binding Transcription Activator (CAMTA) family of TFs to govern expression of the cold-responsive *C-REPEAT BINDING FACTOR* (*CBF*) genes. However, as of yet, there is no clear explanation of how Ca^{2+} influx is regulated under cold stress [81,82,83,84]. Heat stress (HS) has also been shown to affect Ca^{2+} channels and swiftly trigger Ca^{2+} influx to induce thermotolerance [85,86]. Further characterization of the role of Ca^{2+} channels in the thermosensory mechanism will be needed in order for them to be considered as thermosensors themselves.

Temperature fluctuations affect protein activity and/or structure, therefore thermosensing can occur through protein conformational changes under the influence of temperature. A good example is phyB. Red light induces the transition from active Pr to inactive Pfr by altering the structure of its chromophore, with warm temperatures inducing the relaxation of the chromophore back to the Pr state [18,19]. Other phytochromes have also been proposed as thermosensors since it has been observed that temperature-induced hypocotyl elongation in the quintuple phytochrome mutant (*phyabcde*) is lost [18,19]. In addition, the blue-light photoreceptor phototropin has been shown to sense cold signals in the liverwort *Marchantia polymorpha* through its temperature-dependent reversion dynamics, and is required for the cold-avoidance movement of chloroplasts [87,88]. This temperature-sensitive role may be mediated by

PHOT2 in Arabidopsis [88]. Further investigation is required to determine the viability of phototropin as a thermosensor in plants.

Moreover, high temperatures can lead to protein misfolding. Under the influence of high temperatures, HEAT SHOCK PROTEINS (HSPs) bind misfolded proteins, triggering in turn the release of *HEAT STRESS TRANSCRIPTION FACTORS* (HSFs) that regulate the thermotolerance response through a set of *HEAT STRESS RESPONSIVE* (HSRs) genes [89,90]. In addition to proteins, RNA molecules can also be sensitive to conformational changes upon temperature variation. Recent work has provided an example of an RNA thermostat in plants. The 5'-UTR of the *PIF7* RNA folds into a hairpin structure, which adopts two distinct conformations in a temperature-dependent manner [91]. This molecular switch adopts a more relaxed, yet distinct conformation at warmer temperatures, resulting in enhanced translation.

Pioneering work on the effect of temperature on chromatin dynamics proposed the histone variant H2A.Z as a potential thermosensor [92]. In *Arabidopsis thaliana*, nucleosomes containing H2A.Z are evicted from chromatin as ambient temperature rises [92]. The question of whether temperature could directly affect H2A.Z nucleosomes remained open for several years and led to much speculation. However, follow-up work demonstrated that H2A.Z nucleosome eviction requires the function of the HSFA1 clade of the Arabidopsis HSFs [93]. Indeed, in *Brassica rapa*, a close relative of Arabidopsis, H2A.Z controls transcriptional response to temperature but in the opposite direction, as H2A.Z levels increase with increasing temperature [94].

A new physical mechanism of temperature sensing has recently been described, whereby the relative abundance of a protein changes in response to temperature, but without an equivalent change in expression or stability. The TF NTL8 accumulates over long periods of cold through a dilution-mediated mechanism, due the temperature-dependence of growth [95]. Whether similar mechanisms underlie other temperature responses is unknown. Despite this recent progress, plant thermosensors, unlike light sensors, remain largely uncharacterized.

2.2.2. Temperature-responsive transcription factors

Several signaling pathways have been proposed to regulate cellular responses to both elevated and low environmental temperatures. HS affects several processes during plant growth and development, such as hypocotyl and petiole elongation [96], while elevated ambient temperature has a positive effect on flowering [2,97] and a negative effect on stomata formation [98]. Several genes contribute to thermotolerance. Li and colleagues systematically analyzed the transcriptional regulatory network in Arabidopsis plants after exposure to HS at 37°C using RNA sequencing (RNA-seq) [99]. They determined that HEAT SHOCK FACTOR A1 family proteins (HSFA1s), and the circadian clock proteins REVEILLE4 (RVE4) and REVEILLE8 (RVE8), are among the primary TFs that mediate the first wave of HS-induced gene expression, providing insight into how the circadian clock helps plants adapt to high temperatures during the day. During HS, HSFA1s regulate transcriptional responses leading to thermotolerance (Figure 3(a-b)). However, their activity is suppressed under normal temperature environments through the action of HEAT SHOCK PROTEIN 70 (HSP70) and HSP90 (Figure 3(a-b)) [100,101,102]. Not considered HS, high temperatures between ~29–36°C (supraoptimal) independently impact plant developmental responses. One example of this is inhibition of seed germination at supraoptimal high temperatures (thermoinhibition) [103]. High temperatures during seed imbibition activate expression of *SOMNUS* (*SOM*), which prevents seed germination by increasing ABA seed content and reducing GA biosynthesis [104]. Similarly, although warm temperatures accelerate flowering (see below), supraoptimal temperatures delay it through upregulation of *FLOWERING LOCUS C*, a MADS-box TF that represses flowering. This upregulation is due to higher protein stability of the chromatin modifier JUMONJI 30 (JM30) at 29°C, which prevents the accumulation of H3 lysine 27 di- and trimethylation (H3K27me_{2/3}) at *FLC* [105]. These joint mechanisms of delay of flowering and germination (thermoinhibition) by supraoptimal conditions may have important implications to

seed set and survival when under adverse environmental conditions.

The developmental response to warm temperatures, such as longer hypocotyls and petioles and more rapid flowering (thermomorphogenesis, ~20–28°C), is a separate process to HS (>37°C) and the supraoptimal high temperature response. *PIF4*, *PIF5*, and *PIF7*, discussed above for their role in light signaling, play a vital role in the positive regulation of plant thermomorphogenesis [106,107]. Rapid induction of *PIF4* expression has been observed to occur during early evenings due to warm temperatures and it is negatively regulated by the evening complex: EARLY FLOWERING 3 (ELF3), ELF4, and LUX ARRHYTHMO (LUX) (Figure 3(c)) [108]. Another set of negative regulators of *PIF4* are POWERDRESS (PWR) and its interactor HISTONE DEACETYLASE 9 (HDA9), which control thermomorphogenesis through the induction of H3 deacetylation (Figure 3(c)) [109]. On the other hand, SHORT HYPOCOTYL UNDER BLUE 1 (SHB1) is recruited by the core components of the circadian clock CIRCADIAN CLOCK-ASSOCIATED 1 (CCA1) and LATE ELONGATED HYPOCOTYL (LHY) to positively regulate the expression of *PIF4* (Figure 3(c)) [110]. Further, the TFs TEOSINTE BRANCHED 1/CYCLOIDEA/PCF 5 (TCP5) and BRASSINAZOLE-RESISTANT 1 (BZR1) have recently been demonstrated to play a role in the positive regulation of *PIF4* in the thermomorphogenic response (Figure 3(c)) [111,112].

In addition, the UV-B receptor UVR8 and the light signaling TF *HY5* have separately been demonstrated to impair the thermomorphogenic response through negative regulation of *PIF4* expression (Figure 3(c)) [113,114]. During hypocotyl thermomorphogenesis, COP1 is imported into the nucleus where it reduces the negative regulation of *HY5* on hypocotyl growth (Figure 3(d)). *cop1* mutants have been shown to be defective in *PIF4* expression and warmth-induced hypocotyl elongation [114,115]. Another key light-signaling component that plays a role in thermomorphogenesis is DET1. Together with COP1, DET1 is essential for the transcriptional regulation of *PIF4* through *HY5* (Figure 3(d)) [116]. *PIF4* then regulates hypocotyl elongation through auxin biosynthesis genes (e.g., *TAA1*, *CYP79B2*

Figure 3. Temperature driven transcription initiation.

a, Heat stress (HS) halts the inhibitory activity of HSP70 and HSP90 on *HSFA1s*, leading to their disassociation from HSP70 and HSP90 and subsequent activation. b, Following the activation of *HSFA1s* they in turn regulate the transcriptional initiation of several *HSR* genes such as *HSFA2* that trigger thermotolerance. c, Warm temperatures induce the expression of *PIF4* by inhibiting the expression of several of its negative regulators such as the evening complex (ELF3, ELF4, LUX), PWR/HAD9, UVR8 and HY5. Warm temperatures also lead to the positive regulation of *PIF4* expression under the influence of several positive regulators such as CCA1/LHY/SHB1, TCP5, and BZR1. d, Warmth causes both COP1 and DET1 to actively attenuate HY5 stability leading to its degradation and the subsequent induction of expression of *PIF4*. PIF controls auxin responsive proteins (e.g., IAA19) as well as auxin biosynthesis genes (e.g., YUC8), indicated as ARP and ABG respectively in the schematic. e, Cold temperatures induce the expression of *HY5* and increase its stabilization through nuclear depletion of COP1 and in turn initiate cold stress responses. f, Under cold and freezing stress, ICE1 is phosphorylated and stabilized by the protein kinase OST1 which interrupts ICE1 ubiquitination and degradation by the E3 ligase HOS1. Stabilization of ICE1 leads to the initiation of the transcription of several *COR* genes leading to the triggering of cold acclimation and freezing tolerance. g, During vernalization, perception of winter cold temperatures (Phase I) triggers the quantitative inactivation of *FLC* transcript (green solid line). In the absence of the cold signal during spring, epigenetic memory (Phase II) maintains *FLC* at low levels. *FLC* silent state correlates with the accumulation of the H3K27me3 repressive mark at *FLC* chromatin. h, Transcriptional memory of HS can also be divided into distinct phases. Perception of high temperatures induces activation of HS memory genes (Phase I). Upon removal of the heat signal (Phase II or primed state), HS memory genes can exhibit sustained (green solid line) or baseline (green-dotted line) expression levels, with their chromatin coated by the H3K4me2/3 active marks. HS memory genes show higher or faster activation following recurrent heat stress (Phase III) conferring on the plant the ability to prevent damaging effects.

[117] and YUC8 [118]) and auxin-responsive proteins (e.g., IAA19, IAA29 [119]). Auxin biosynthesis in response to PIF4 occurs primarily in the epidermis of cotyledons and auxin acts as a mobile signal, traveling to the hypocotyl where it induces elongation in combination with local signals [120,121].

HY5 expression can also be induced by cold stress where it plays a positive regulatory role in the expression of the *COLD REGULATED (COR)* genes via the function of the Z-box and other cis-acting elements in their promoters (Figure 3(e)) [122]. The expression of *HY5* under cold stress is

independent of the other major cold-stress pathway [122]. This latter involves the three *CBF* genes, *CBF1*, *CBF2*, and *CBF3*, and their transcriptional regulatory pathway (Figure 3(f)) [123,124]. Low temperatures bring on the swift induction of *CBF* gene expression and the consequential activation of *COR* gene expression. Another TF that plays a role in cold acclimation is the bHLH INDUCER OF CBF EXPRESSION 1 (*ICE1*) and its homolog *ICE2*, with both of them affecting the response to deep freezing through a *CBF*-dependent pathway in *A. thaliana* plants (Figure 3(f)) [125,126]. The role of *ICE1* in freezing tolerance is modulated through its phosphorylation by OPEN STOMATA 1 (*OST1*) [127]. *OST1* is a SNF1-related protein kinase that positively regulates freezing tolerance in plants through inhibition of the interaction between the E3 ubiquitin ligase HIGH EXPRESSION OF OSMOTICALLY RESPONSIVE GENE1 (*HOS1*) and *ICE1*, leading to a more stable form of *ICE1* and therefore a sustained freezing tolerance response [127,128]. Furthermore, *OST1* increases protein stability of *CBF* proteins through the phosphorylation of BASIC TRANSCRIPTION FACTOR 3 proteins (Figure 3(f)) [129].

Other factors also play a role in plant mechanistic responses to cold stress. For instance, cold stress has been found to activate several regulatory responses, such as the CYTOPLASMIC RECEPTOR-LIKE KINASE 1 (*CRPK1*) phosphorylation of 14-3-3 proteins, which induces their interaction with and stabilization of *CBFs* [130]. Phosphorylation of *ICE1* by the MAP kinases *MPK3* and *MPK6*, as well as by BRASSINOSTEROID INSENSITIVE 2 (*BIN2*), is also induced by cold stress, leading to negative regulation of *CBF* expression [131,132]. Nonetheless, these extensive investigations of cold responses so far fall short of providing a comprehensive understanding of the full pathway from cold-sensing to response in plant cold and freezing acclimation.

As well as cold stress and freezing, plants also sense low temperatures over long periods as a seasonal signal, in a process termed vernalization (Figure 3(g)). Vernalization is a prevalent trait of plants in temperate zones and it means that exposure to long periods of cold promotes the

induction of flowering [133]. In *Arabidopsis*, the regulation of vernalization occurs through the action of the floral transcriptional repressor *FLC* which inhibits *FT*, a component of the flowering-inducing factor florigen [134,135]. *FLC* expression is repressed by long-term cold, and this process needs the PHD protein VERNALIZATION INSENSITIVE3 (*VIN3*), which is quantitatively upregulated during cold. *VIN3* seems to be regulated at the level of transcriptional initiation by three independent temperature inputs, a fast temperature response (minutes-hours), a short-term response to warmth (hours-days) and a slow response to cold (weeks) [78,136]. The latter was recently found to be mediated by *NTL8* protein accumulation in long periods of cold [95], but the faster thermosensors remain unknown. *FLC* shut-down promotes flowering thermomorphogenesis, driven by both *FT* and other flowering regulating genes that are activated [3,137].

3. Divergent effects of light and temperature on plant transcription

Light sensing is translated into both cytoplasmic and chloroplast retrograde signaling pathways that converge in specific TFs to regulate transcriptional initiation. Temperature-induced transcriptional regulation diverges from light because it will not only influence the set of genes being turned ON or OFF by specific TFs, but may also directly affect the physical properties of nascent RNA transcript processing and RNAPII kinetics. In the following sections, we summarize the direct and indirect effects of temperature and light, respectively, on the regulation of co-transcriptional processes in plants.

3.1. Direct effect of temperature on co-transcriptional processes

It has been recently proposed that temperature sensing in plants is distributed across regulatory networks, with multiple temperature inputs combined to control each process [78]. Such a temperature sensing system could be controlling the abundance of the TFs described previously, but it could also be directly affecting mechanisms relating to transcription,

following transcriptional initiation. Co-transcriptional processes such as elongation, RNAPII stalling at gene borders, splicing, termination, and polyadenylation may also be sensitive to temperature (Figure 1(b-d)). In warm-blooded mammals, it has been proposed that cooling stabilizes secondary structures of pre-mRNAs affecting splice site selection [138]. In plants, given their sessile nature, physical perturbations on macromolecules caused by temperature variation are likely to be even more dramatic.

Recent studies have started to elucidate distinct features of nascent transcription in plants [139–143] with particular attention given to how cooling influences co-transcriptional processes [144]. A detailed time series during cold acclimation in *Arabidopsis* revealed dynamic changes in expression of several genes, including induction of previously characterized cold-inducible genes, as well as massive changes in alternative splicing (AS) patterns [145]. Over a quarter of genes whose expression changes significantly in the cold also undergo AS. Among the differentially spliced genes identified, some had previous evidence of involvement in cold responses, such as *REGULATOR OF CBF EXPRESSION 1* (*RCF1*), *PIF7*, and *PHYB*. In addition to protein-coding genes, hundreds of long non-coding RNAs (lncRNAs) are differentially expressed and alternatively spliced in response to cold [146]. Variation in gene splicing patterns occurs rapidly upon cold exposure and can be sensitive to temperature reductions as small as 2°C [145]. AS makes a major contribution to changes in the transcriptome, amplifying the ability of temperate plants to grow in a wide range of fluctuating temperatures, as influencing the spliceform generated from each locus increases the potential transcriptomic variation that a signal cascade can induce. Temperature-induced AS could be caused by either a direct effect on the splicing components and the secondary structure of the nascent RNA, or the result of the variation of RNAPII kinetics as temperature decreases. Indeed, the application of a NET-seq approach in *Arabidopsis* (plaNET-seq) has shown transient molecular adaptations of RNAPII transcription in response to cold [144,147]. RNAPII

kinetics are modulated as temperature decreases leading to changes in transcriptional elongation and termination. Interestingly, the dynamics of the splicing reaction are initially affected during early stages of cold acclimation but later recover to normal levels, reinforcing the influence of AS in rapid responses to cold.

Elevated ambient temperature and HS can also impact co-transcriptional processes in plants. Temperature-induced differential accumulation of two *FLOWERING LOCUS M* (*FLM*) splice variants, *FLM* β , and *FLM* δ , regulates *Arabidopsis* floral transition [148]. Whereas low ambient temperature favors accumulation of *FLM* β that actively represses flowering, rising temperature increases the *FLM* δ /*FLM* β ratio to release floral repression. In addition to splicing, *FLM* levels are also reduced by the Nonsense-Mediated mRNA Decay (NMD) pathway under elevated temperatures resulting in floral induction [149]. HS also has a strong effect on mRNA splicing. The U5-snRNP-interacting protein STABILIZED1 (*STA1*) is a heat inducible gene involved in the splicing of heat shock factors and *HSP* transcripts during high temperature stress [150]. Interestingly, *STA1* had originally been found to regulate the splicing of cold responsive genes such as *COR15A* under low temperature stress [151].

Widely present in mammals and metazoans, the process of RNAPII stalling over gene boundaries was recently demonstrated in plants [139,142]. Engaged RNAPII initially transcribes 20–60 nucleotides before undergoing promoter-proximal (5' end) pausing, a regulatory checkpoint for execution of transcription programs [152]. In *Arabidopsis*, promoter-proximal stalling of RNAPII transiently increases at early stages of cold acclimation to adjust transcription to low temperatures [144]. In addition to promoter-proximal stalling, RNAPII stalls near 3' ends of *Arabidopsis* genes [140,141,142]. Cold exposure significantly decreases the 3' end peak of RNAPII observed when the polymerase is stalled downstream of the Polyadenylation Site (PAS), suggesting a major change in transcription dynamics associated with termination in response to low temperatures [144].

3.2. Photoreceptor-mediated effects on co-transcriptional responses

Light also controls co-transcriptional processes such as choice of Transcription Start Site (TSS), pre-mRNA- and alternative- splicing of large numbers of genes, with mechanisms distinct from those controlling transcriptional responses [147,153–162]. In the case of light-induced alternative TSS use, this leads to changes in protein fate. Ushijima and colleagues found that red-light-activated phyB alters TSS usage for over 2000 loci, enriched in plastid-targeted proteins, and the resulting changes in N-terminal signal sequence leads to changes in subcellular protein localization for nearly a fifth of these genes [162]. In contrast, Kurihara et al. investigated blue light, finding that a small (~200) subset of target genes have TS sites both upstream and downstream of short open reading frames in the 5'UTR of the main coding sequence [154]. These upstream open reading frames can trigger transcript degradation through the NMD pathway, or interfere with translation itself, both routes leading to reduced protein output [163]. Blue light promotes use of the downstream, protein-producing TSS in most cases, a response partially mediated via HY5, which has binding sites within many of the target promoters [154]. Like the red-light response, the blue light target set is enriched in light-responsive genes, and both responses modulate the photorespiration pathway to cope with effects of fluctuating light on the photosynthetic machinery [154,162].

Like temperature, light also induces alternative splicing, in the majority of cases promoting production of protein-encoding splice variants over those with NMD features [164]. At least two sets of photoreceptors are known to mediate this response. One reacts to white light intensity via chloroplast retrograde signaling in response to Photosystem II function [157,164,165]. The other route requires red-light sensitive phytochrome action and is conserved across the land plants [158,160,161,164,166,167,168]. These routes are independent but not mutually exclusive, and converge on serine/arginine-rich (SR) splicing factors such as RS31 [159,160].

Phytochrome-dependent AS is critical for phy-mediated photomorphogenesis and regulates

several transcription factors, such as *PIF3*, *PIF6*, and *ELF3*, and negative regulators, such as *SPA3*, required for light-signaling itself [158,160,161,168,169]. The mechanism of phytochrome-dependent AS appears to be through direct interaction of phyB with the pre-mRNA splicing factors REDUCED RED-LIGHT RESPONSES IN CRY1CRY2 BACKGROUND1 (*RRC1*) and SPLICING FACTOR FOR PHYTOCHROME SIGNALING (*SFPS*) [161,166,168]. In Arabidopsis, the interaction between *RRC1* and phyB is light-induced but independent of the Pr/Pfr conformation of phyB [161], whereas in the moss *Physcomitrella patens*, interaction between PpPhy4, a phyB homolog, and the splicing regulator PphnRNP-F1 is red-light-specific, and induces interaction of PphnRNP-F1 with other spliceosome components [170,171]. In Arabidopsis, *RRC1* is itself alternatively spliced in an *SFPS*-dependent response to light, providing a positive feedback loop [161,164,166].

At least part of the light-driven AS response seems to occur due to enhanced RNAPII elongation in response to light [165]. It is unlikely that light will directly modulate the physical properties of RNAPII and RNA transcripts as has been suggested for temperature [144]. Instead, while the expression of the transcription elongation factor *TFIIS* itself is not affected, members of the plant POLYMERASE ASSOCIATED FACTOR COMPLEX (PAF1-C), including *EARLY FLOWERING7* (*ELF7*), are transcriptionally upregulated in response to light and may provide part of the mechanism [165]. More directly, genetic evidence links stress-induced singlet oxygen in the chloroplast to the activity of TOPOISOMERASE VI, which binds a subset of high-light-induced target promoters directly and may also enhance elongation [33]. Recent work has demonstrated a specific mechanism for integration of light into circadian-clock-regulated transcriptional activation via the NIGHT LIGHT-INDUCIBLE AND CLOCK-REGULATED (LNK) proteins [172]. LNKs are transcriptionally upregulated in response to light, probably via phytochrome signaling [173] and recruit RNAPII and the transcript elongation FACT complex to their targets via interaction with the MYB TF RVE8 [172]. Interestingly, RVE8 is also involved in temperature responses [99].

Questions remain, however, on which responses are direct and which due to secondary signaling [155,156]. Many of the light-responsive AS genes are splicing factors themselves, leading to a secondary cascade of AS throughout the plant [159–161]. The presence of the AS response in non-irradiated, chloroplast-lacking roots indicates the role of a secondary, non-cell-autonomous system [157], possibly mediated via sugar signaling [164]. As a result light, like temperature, can affect genome-wide co-transcriptional processes at the whole plant level, via secondary signaling.

In conclusion, both light and temperature generate genome-wide cascades of co-transcriptional effects. However, while in the case of light these are generally triggered through distinct receptor pathways and subsequent upregulation of the transcriptional machinery, temperature can modulate transcription kinetics directly.

3.3 Induction of long non-coding transcripts

A common phenomenon observed in plant responses to light and temperature is the induction of lncRNAs. During long cold periods, plants undergo massive transcriptional reprogramming, which also results in the activation of lncRNAs. The *FLC* locus has provided a highly informative system for studying the effects of these transcripts. *COOLAIR* lncRNAs, antisense to *FLC* locus, significantly increase in expression levels in the first couple of weeks of winter temperature [174,175]. Transcription from the antisense strand may inhibit RNAPII firing from the *FLC* sense strand, as suggested by the mutually exclusive expression of *COOLAIR* and *FLC* nascent transcript at single loci resolution [176]. Although the precise mechanism by which *COOLAIR* suppresses nascent *FLC* expression during cold remains unknown, modulation of chromatin dynamics is likely implicated in this process. Arabidopsis transgenic lines with reduced *COOLAIR* function fail to remove the H3K36me3 mark at the *FLC* locus, resulting in less efficient reduction of *FLC* nascent transcript during cold [174]. This observation suggests that cold-induced non-coding transcripts at *FLC* are required to set the chromatin landscape for transcriptional inactivation.

Another cold-induced lncRNA also represses *FLC* via modulation of local chromatin architecture. Expression of the sense lncRNA *COLDWRAP* from the sense *FLC* inactive promoter induces the formation of an intragenic gene loop within the *FLC* locus to perpetuate gene inactivation [177]. Finally, the intronic sense lncRNA *COLDIAIR* mediates PHD-PRC2 (PLANT HOMEODOMAIN- POLYCOMB REPRESSIVE COMPLEX 2) recruitment to *FLC* for stable epigenetic silencing [178]. Another vernalization-linked lncRNA, that does not map to the *FLC* gene, is the *MADS AFFECTING FLOWERING 4* (*MAF4*) antisense (*MAS*) transcript [179]. *MAS* is transcribed during cold exposure [179] mimicking the expression pattern of *MAF4* [180]. Induction of *MAS* seems to play a crucial role in the recruitment of COMPASS-like complexes to *MAF4* chromatin to activate its expression [179]. In all cases, it remains to be determined what pioneer TFs trigger transcription of all these cold-induced lncRNAs.

Temperature-induced non-coding transcription has been shown to affect transcription termination. Cold-induced transcription of the Arabidopsis *CBF* locus triggers expression of a cascade of antisense transcripts that fine-tunes the level of *CBF1* mRNA [181]. The lncRNA *SVALKKA*, located downstream of the *CBF1* gene, is transcribed from the antisense strand. *SVALKKA* transcription terminates downstream of the *CBF1* locus in plants growing under warm temperatures. During cold acclimation, however, RNAPII read-through transcription of *SVALKKA* results in a cryptic lncRNA overlapping *CBF1* on the antisense strand, termed *asCBF1*. The *SVALKKA-asCBF1* system limits the production of maximal *CBF1* mRNA levels by a RNAPII collision mechanism in the 3'-end of *CBF1* [181]. Thus, the interplay of two lncRNAs tightly regulating *CBF1* mRNA levels allows plants to appropriately acclimate to low temperatures.

Increasing evidence shows that non-coding transcription also leads to the formation of R-loops in the plant genome. R-loops are created by the invasion of the DNA duplex by complementary RNA molecules, generating an RNA-DNA helix and leaving a single-stranded section of complementary DNA. At the single gene scale,

R-loops appear to be important for transcriptional regulation [182,183,184,185]. R-loop stabilization at the promoter of *COOLAIR* is required for correct expression of both *COOLAIR* and *FLC* [186]. Invasion of the lncRNA *APOLO* (*AUXIN REGULATED PROMOTER LOOP*) into a number of genomic loci, including auxin-responsive genes, results in R-loop formation and subsequent displacement of Polycomb repressive activity leading to gene activation [187]. However, construction of a genome-wide R-loop atlas by the Sun lab revealed that R-loop presence does not strongly correlate with mRNA transcript abundance [188]. Indeed, genome-wide R-loop patterns seem to be largely conserved through many environmental perturbations, although developmental phase changes such as germination and flowering have more noticeable effects [188]. Exposure to long term HS results in a slight increase in antisense R-loops localized to the TSS. A subset of these show correlation of expression with the expression with known R-loop regulators during this process, hinting at a regulatory role. However, RNAPIII-transcribed genes, such as small nucleolar RNAs and tRNAs, are consistently associated with higher R-loop levels in germinated seedlings than germinating or dark-grown seedlings, raising the possibility that R-loop dynamics may be particularly important for light regulation of RNAPIII transcription [188]. Further work on individual gene dynamics and R-loop regulators will shed more light on the mechanistic basis of these responses and their biological role.

Similar genome-wide studies of lncRNAs in response to light have focused on natural antisense transcripts (NATs) in de-etiolating seedlings [189]. A majority of light-induced “concordant” NAT (those changing expression in the same direction as the coding transcript) are principally expressed in cotyledons and include members of the light-sensing pathway such as *SPA1* and *HYH*. Light also induces a number of long intergenic non-coding RNAs of unknown functional significance [164]. Further detailed work is required to functionally characterize the mechanism and roles of these light-induced lncRNAs.

3.4. Both light and temperature impact plant chromatin dynamics

Plants monitor temperature changes both diurnally and seasonally. Temperature may physically alter the interaction of DNA with nucleosomes and thus have a direct effect on chromatin compaction [190]. Not surprisingly, plant genes may display unique chromatin signatures in response to temperature cues [191]. As mentioned above, elevated temperatures trigger a universal response which involves the rapid upregulation of heat shock proteins (HSPs) [192]. HS, which for Arabidopsis usually means exposing plants to temperatures above 36°C, induces substantial changes to plant nuclear architecture, including decrease of heterochromatin compaction [193] and decondensation of chromocenters [194]. DNase-seq studies revealed that the degree of chromatin accessibility changes dynamically upon heating [195]. The most highly accessible genes following HS encode HSPs and several heat-stress-related TFs, which correlates with their rapid activation. Similarly, HS results in the reduction of nucleosome density, mediated by *HEAT INTOLERANT 4* (*HIT4*) [196].

Moving from dark to light also results in major changes in nuclear organization such as decondensation of chromocenters and decompaction of heterochromatin, especially during photomorphogenesis [191,197,198,199,200]. As well as the large-scale transcriptional reprogramming associated with the developmental aspects of photomorphogenesis, high light is a cellular stress, and UV irradiation causes DNA damage. Responses to all these aspects are united by the multifunctional roles of DET1 [60]. As well as its role in stabilization of PIF proteins [56], DET1 also has roles in the repair of UV-induced DNA damage through the global genome repair pathway [59]. In addition, DET1 binds directly to non-acetylated tails of histone H2B [201] and is required for the global increase in H2B ubiquitination (H2Bub) that is induced by light [202], via DET1-mediated degradation of a deubiquitination module of the SAGA complex [203]. Correct H2Bub is required for light-induced transcription change and seems particularly important for genes that respond especially rapidly [202].

Besides HS, more moderate variation in ambient temperature also impacts plant chromatin and transcription. In fact, temperature changes as little as 1°C may lead to gene reprogramming, and can also affect crop yield [204]. As part of the thermomorphogenic response, *A. thaliana* nucleosomes containing the histone variant H2A.Z are evicted from the chromatin of thermo-responsive genes when temperature rises from 22°C to 27°C [92] resulting in transcriptional change. H2A.Z nucleosomes may act to enhance the transcriptional dynamic range by keeping genes transcriptionally repressed under non-inductive conditions via restriction of promoter and gene accessibility. Histone deacetylation by HDA9 also facilitates H2A.Z nucleosome eviction from the promoter of the thermo-responsive gene *YUC8* under warmer temperature [205] resulting in transcriptional activation. This process, likely occurring downstream of HSFs, extends our view of histone deacetylation, which is generally associated with transcriptional repression. In addition, H3K36me3 affects temperature-induced alternative splicing, potentially by facilitating the recruitment of splicing factors to the chromatin during transcription [206]. Whether H3K36me3 dynamics could be acting as a thermosensor needs further validation. More generally, the ability of any chromatin structure to directly sense a temperature change to control thermal responses remains an open question [79].

Increases in histone acetylation appear to be a general part of light-induced chromatin changes that are associated with the transcriptional upregulation of many genes, including *HY5* and *HYH* [207]. Indeed, addition and removal of acetylation at *HY5* and *PIF* target genes may form part of the antagonistic actions of these light regulators on transcription. For example, Peng et al. find that *PIF* recruits HISTONE ACETYL TRANSFERASEs (HATs) to regulate shade response [208], while Yang et al. find *HY5* recruits HDA9 to repress target genes involved in autophagy in the dark [209]. HISTONE DEACETYLASE 15 (*HDA15*) is also recruited to targets by both *PIF1* and *PIF3* to mediate repression by deacetylation of target genes [210,211].

PIF1 additionally recruits another form of histone modifier to its targets: the SWI/SNF complex

member BRAHMA (BRM) [212]. BRM acts with *PIF1* to reduce H3K4 methylation and expression of chlorophyll biosynthesis genes in the dark, preventing photobleaching during subsequent irradiation. *PIF1* itself is under chromatin regulation, with normal *PIF1* expression requiring the H3K36 methylase SDG8 [213].

Downstream of light-regulated transcription itself, several chromatin modifiers have been characterized as necessary for light response. Among the chromatin modifiers themselves transcriptionally regulated in response to light are several Jumonji-domain proteins. *JMJ20* and *JMJ22* are upregulated in response to red light, and enhance seed germination by demethylating histone H4 arginine residues and thereby derepressing gibberellin synthesis loci [214]. The H3K27me3 demethylase *JMJ30/JMJD5*, which acts at *FLC* in response to high temperatures, is also alternatively spliced in response to light, although the functional significance of this is unknown [159]. *JMJ30* is involved in temperature compensation of the circadian clock [215], potentially providing another route for light and temperature integration at the chromatin level. In addition, the ATP-dependent chromatin remodeling factor PICKLE (PKL) acts downstream of both light and temperature signaling to promote Arabidopsis growth [216,217] thus providing further integration of light and temperature signals through chromatin regulation.

3.5. Long-term memory of environmental stimuli perception

So far, we have discussed how perception of specific cues results in transient transcriptional regulation. However, many transcriptional reprogramming events, as is the case for acquisition of stress tolerance and developmental switches, should persist for prolonged periods of time even in the absence of the triggering signal [218,219]. In those cases, two distinct phases can be distinguished in the response (Figure 3(g-h)): initial and rapid up- and down-regulation of genes induced by perception and integration of a specific cue from the surroundings (Phase I); and perpetuation of the acquired ACTIVE or INACTIVE potential of those genes for extended periods of

time by epigenetic mechanisms, in the absence of the inductive cue (Phase II).

The best characterized example of such an epigenetic process is the silencing of the gene *FLC* in vernalization, and the memory of the silencing following winter (Figure 3(g)). The *FLC* gene is highly expressed in vernalization-requiring accessions of *Arabidopsis*, until the plants experience prolonged cold [220]. Increasing weeks of cold temperature trigger the inactivation of *FLC* transcription (Phase I) and this is accompanied by the buildup of the epigenetic silencing apparatus at *FLC* chromatin. Subsequently, Phase II secures the maintenance of the *FLC* silent state once the cold signal has disappeared (Figure 3(g)).

During Phase I, exposure to cold temperatures induces epigenetic silencing of *FLC* by the action of lncRNAs (*COOLAIR*, *COLD AIR*, *COLDWRAP*, as discussed above), by the sequence-specific transcriptional repressors (VAL1 and VAL2 proteins, [221,222]) and by the PRC2 combined with Plant Homeodomain (PHD) proteins (PHD-PRC2 complex; [223,224]). PHD-PRC2 assembles at *FLC* chromatin during cold and deposits the H3K27me3 histone repressive mark at individual *FLC* loci [225,226,227,228,229] while histone demethylases remove the chromatin active marks (H3K36me3 and H3K4me3) that were present in those nucleosomes [230].

After the cold signal is gone, the individually silenced *FLC* copies remain transcriptionally inactive allowing the plants to remember the cold period and quantify its duration (Phase II, Figure 3(g)). Upon return to warm temperatures, H3K27me3 spreads from the nucleation region to cover the *FLC* locus (Figure 3(g)) [224,225,227–229].

Cold-sensing is not just a signature for floral induction. Plants are able to remember cold stress to enhance survival when they are exposed to low temperatures a second time [231,232]. Before cold stress, H3K27me3 coats the genomic region of a subset of early-cold inducible genes required to trigger cold acclimation [233,234,235]. Therefore, PRC2-mediated repression likely prevents activation of these loci under non-stress conditions [235]. However, H3K27me3 does not seem to control short-term cold-stress memory [235] as it does for the long-term maintenance of *FLC* silencing

during vernalization. Future work will reveal the chromatin components underlying cold stress transcriptional memory.

A well-documented example of stress memory is thermotolerance, where perception of moderate HS allows plants to subsequently withstand high temperatures that would otherwise be lethal [236]. A number of genes, including *HSP22.0* and *ASCORBATE PEROXIDASE 2* (*APX2*), are initially activated during the priming HS and their expression levels are either sustained or downregulated to basal levels in the absence of the inductive cue [237] (Figure 3(h)). Either way, they show an increased response after a recurring HS days after the primary HS, consistent with the definition of transcriptional memory (Figure 3(h)) [238]. HS transcriptional memory at the memory-related loci *HSP22.0* and *APX2* depends on the sustained accumulation of H3K4me3 and H3K4me2 [238]. Both HS transcriptional memory and the sustained H3K4me3 and H3K4me2 depend on the HEAT SHOCK TRANSCRIPTION FACTOR A2 (*HSFA2*) [238]. In addition, the *Arabidopsis* homolog of the metazoan Strawberry Notch protein FORGETTER 1 (*FGT1*) is required for HS memory. *FGT1* associates with the promoters of HS memory-related genes to ensure proper nucleosome remodeling during the acquisition of thermotolerance [239]. By analogy to vernalization, plant HS memory can also be divided into two phases (Figure 3(h)) corresponding to “perception” (Phase I) and “memory” or “primed state” (Phase II). Additionally, stress memory exhibits a unique Phase III that correlates with the rapid or altered transcriptional responses to the recurrence of the triggering signal (Figure 3(h)).

Another example of transcriptional stress memory occurs during recurring dehydration stresses [240]. In this case, not only does H3K4me3 act as a persistent epigenetic mark, but in addition stalling of the phosphorylated serine 5 form of RNAPII (Ser5P) is associated with transcriptional memory [240]. Interestingly, light signaling via phytochromes is required for transcriptional memory of the salt- and dehydration-stress inducible gene *Δ1-PYRROLINE-5-CARBOXYLATE SYNTHASE 1* (*P5CS1*, [241]). *HY5* and *HYH* are not required for initial upregulation of *P5CS1*, but *HY5* binding to the promoter is necessary for

maintenance of H3K4me3 levels at the locus for enhanced response to repeated stimuli [241].

Undoubtedly, chromatin dynamics play a key role in plant adaption to light fluctuations [191,199,242]. However, the existence of mechanisms of transcriptional stress memory in the field of light stress responses remains an open question. Exposure to high light results in priming, or photoacclimation, as well as in the induction of a systemic-acquired response that renders non-exposed leaf tissues less susceptible to recurrent exposure to light stress [243,244,246,245]. However, efforts to try to link epigenetic modifications, e.g. DNA methylation, with the acquisition of a “primed state” following excess light stress have not been successful [246,247]. Follow up studies will be necessary to reveal what are the chromatin components taking part in these mechanisms that prime plants to rapidly respond to harmful irradiation.

4. Concluding remarks

Light and temperature are the two most important environmental signals to determine seasonal progression and correct timing of development for most plants. At the same time, they are linked to the plant's ability to produce energy, capture CO₂ and survive other stresses in direct ways, as well as being potential stressors themselves. Therefore, it is not surprising that plants have a complex way of responding to each of these signals separately, and that they also become integrated in many responses. The pathways of environmental signaling have multiple stages, starting with signal perception by “sensors” (Phase I), followed by downstream transcriptional cascades and in some cases long-term memory of the signal (Phase II/III). The presence and duration of this memory will depend on the nature of the signal and the desired response. For example, in stress response, the experience of a stress now is likely to indicate that the same or a related stress will occur again in the near future. However, the opposite is true in more long-term signaling that determines seasonal progression such as vernalization, which tells the plant that winter has passed and is unlikely to occur again for a long time.

In the perception of environmental signals, it is very hard to distinguish direct, sensor-mediated responses and indirect, downstream transcriptional cascades, and so the search to identify the primary sensors has been a difficult one. For light sensors, which have more distinctive properties, many have been known for a long time. Temperature instead has more direct effects in biochemistry and so most processes can contribute to its perception, making temperature sensing a question of integration of (often subtle) inputs from multiple “thermosensors”. Thus, plant thermosensors, unlike light sensors, remain largely unexplored. Furthermore, though the mechanism of its significance is still unknown for most cases, the fluctuating nature of the temperature signal has a strong effect on its perception [136,248,249]. This highlights the pressing need for further knowledge on the subject, considering the important role of thermosensing in plant growth, development and temperature stress acclimation. Under a scenario of global temperature increase, in combination with an increase in extreme events, knowledge on how to modulate plant thermosensing mechanisms would provide great advantages for crop breeding and growth.

Regarding transcriptional regulation, light and temperature have similar yet divergent effects. Following perception, the underlying signaling pathways converge in the function of numerous TFs, some of which are common to both cues. In addition, temperature affects the photosensitive reactions of several of the known light receptors, thus signaling is integrated from the start. The application of novel transcriptomic technologies has now allowed the focus to shift from TFs to the analysis of effects on splicing, as well as on RNAPII elongation rate and stalling in plants. In the case of light, thus far these effects seem to be mediated through TF control of the components of the transcriptional and splicing complexes. However, this has not yet been found in analyses of temperature effects. These initial key findings will pave the way for future studies to determine whether the components of the transcriptional machinery (RNAPII, spliceosome, nascent RNA) are themselves thermosensors. Given the highly structured nature of RNA molecules, the possibility remains that nascent RNAs may adopt

dynamic conformational changes directly modulating co-transcriptional processes. Despite advances in recent years, research on RNA *in vivo* structures in plants is still in its early days. Efforts in this direction should focus on investigating perturbations to RNA molecules in plants growing in natural habitats, to ensure their biological relevance.

Both light and temperature signaling converge in modulation of chromatin states. These dynamic changes in chromatin conformation become themselves part of the signaling process by controlling expression of numerous genes both short and long term. Chromatin changes that follow light perception are very likely a consequence of the action of photoreceptor-induced TFs. Conversely, temperature may directly perturb physical conformations of chromatin, but conclusive *in vivo* examples of such processes are still missing. Work is required to define to what extent nuclear chromatin is responsive to temperature variation. In parallel, possible thermal effects on the kinetics of the catalytic activities of chromatin “readers” and “writers” should also be evaluated.

In this context, precise, quantitative analysis of the responses to temperature and light levels are required to dissect the contribution of each mechanism to the overall response. Moreover, the sheer complexity of these interacting, simultaneous processes will likely require mathematical and computational approaches to elucidate non-intuitive results. Such approaches, very informative in dissecting the mechanism of photomorphogenesis [250], have already begun to bear fruit in the identification of the elusive plant thermosensors.

Acknowledgments

We are grateful to Paloma Mas and Jordi Moreno-Romero for critical comments on the manuscript. We apologise for not citing all the relevant primary research articles and reviews due to space limitation.

Disclosure statement

The authors declare no conflicts of interest.

Funding

This work was supported by the Junior Leader Fellowship [LCF/BQ/PI19/11690003], from “la Caixa,” Foundation [ID 100010434] awarded to J.I.Q; M.J. and J.I.Q. were supported by the Severo Ochoa Excellence Programme for Centres in R&D 2016-2019 (SEV-2015-0533). J. H. and RLAK were supported by Biotechnology and Biological Sciences Research Council grants: BB/P020380/1 (R.L.A.K.), Brassica rapeseed and vegetable optimisation strategic Lola (BB/P003095/1) (J.H.); and the “Genes in the environment” Institute Strategic Programme (BB/P013511/1) (J.H. and R. L.A.K.).

ORCID

Julia I. Qüesta <http://orcid.org/0000-0002-0462-8423>

References

- [1] Blázquez MA, Ahn JH, Weigel D. A thermosensory pathway controlling flowering time in *Arabidopsis thaliana*. *Nat Genet.* 2003;33:168–171.
- [2] Balasubramanian S, Sureshkumar S, Lempe J, et al. Potent induction of *Arabidopsis thaliana* flowering by elevated growth temperature. *PLoS Genet.* 2006;2:e106.
- [3] Andrés F, Coupland G. The genetic basis of flowering responses to seasonal cues. *Nat Rev Genet.* 2012;13:627–639.
- [4] Lee Y, Kim M, Han J, et al. MicroRNA genes are transcribed by RNA polymerase II. *Embo J.* 2004;23:4051–4060.
- [5] de Santa F, Barozzi I, Mietton F, et al. A large fraction of extragenic RNA Pol II transcription sites overlap enhancers. *PLoS Biol.* 2010;8:e1000384.
- [6] Guttman M, Amit I, Garber M, et al. Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature.* 2009;458:223–227.
- [7] Wissink EM, Vihervaara A, Tippens ND, et al. Nascent RNA analyses: tracking transcription and its regulation. *Nat Rev Genet.* 2019;20:705–723.
- [8] Brahma S, Henikoff S. Epigenome regulation by dynamic nucleosome unwrapping. *Trends Biochem Sci.* 2020;45:13–26.
- [9] Fiszbein A, Godoy Herz MA, Gomez Acuña LI, et al. Interplay between chromatin and splicing. Chapter 8 - Chromatin regulation and dynamics. Academic Press, Elsevier Inc.; 2017. p. 191–209. DOI:10.1016/B978-0-12-803395-1.00008-3
- [10] Berry S, Dean C, Howard M. Slow chromatin dynamics allow polycomb target genes to filter fluctuations in transcription factor activity. *Cell Syst.* 2017;4:445–457. e8.

- [11] Steffen PA, Ringrose L. What are memories made of? how polycomb and trithorax proteins mediate epigenetic memory. *Nat Rev Mol Cell Biol.* 2014;15:340–356.
- [12] Yu J-R, Lee C-H, Oksuz O, et al. PRC2 is high maintenance. *Genes Dev.* 2019;33:903–935.
- [13] Reinberg D, Vales LD. Chromatin domains rich in inheritance. *Science.* 2018;361:33–34.
- [14] Stewart-Morgan KR, Petryk N, Groth A. Chromatin replication and epigenetic cell memory. *Nat Cell Biol.* 2020;22:361–371.
- [15] Galvão VC, Fankhauser C. Sensing the light environment in plants: photoreceptors and early signaling steps. *Curr Opin Neurobiol.* 2015;34:46–53.
- [16] Franklin KA, Quail PH. Phytochrome functions in Arabidopsis development. *J Exp Bot.* 2010;61:11–24.
- [17] Legris M, Ince YÇ, Fankhauser C. Molecular mechanisms underlying phytochrome-controlled morphogenesis in plants. *Nat Commun.* 2019;10:1–15.
- [18] Jung J-H, Domijan M, Klose C, et al. Phytochromes function as thermosensors in *Arabidopsis*. *Science.* 2016;354:886–889.
- [19] Legris M, Klose C, Burgie ES, et al. Phytochrome B integrates light and temperature signals in *Arabidopsis*. *Science.* 2016;354:897–900.
- [20] Wang Q, Lin C. Mechanisms of cryptochrome-mediated photoresponses in plants. *Annu Rev Plant Biol.* 2020;71:103–129.
- [21] Yang Z, Liu B, Su J, et al. Cryptochromes orchestrate transcription regulation of diverse blue light responses in plants. *Photochem Photobiol.* 2017;93:112–127.
- [22] Vanhaelewyn L, Schumacher P, Poelman D, et al. REPRESSOR OF ULTRAVIOLET-B PHOTOMORPHOGENESIS function allows efficient phototropin mediated ultraviolet-B phototropism in etiolated seedlings. *Plant Sci.* 2016;252:215–221.
- [23] Christie JM, Blackwood L, Petersen J, et al. Plant flavoprotein photoreceptors. *Plant Cell Physiol.* 2015;56:401–413.
- [24] Somers DE, Schultz TF, Milnamow M, et al. ZEITLUPE encodes a novel clock-associated PAS protein from Arabidopsis. *Cell.* 2000;101:319–329.
- [25] Imaizumi T, Tran HG, Swartz TE, et al. FKF1 is essential for photoperiodic-specific light signalling in Arabidopsis. *Nature.* 2003;426:302–306.
- [26] Kim WY, Fujiwara S, Suh -S-S, et al. ZEITLUPE is a circadian photoreceptor stabilized by GIGANTEA in blue light. *Nature.* 2007;449:356–360.
- [27] Heijde M, Ulm R. UV-B photoreceptor-mediated signalling in plants. *Trends Plant Sci.* 2012;17:230–237.
- [28] Favory -J-J, Stec A, Gruber H, et al. Interaction of COP1 and UVR8 regulates UV-B-induced photomorphogenesis and stress acclimation in Arabidopsis. *Embo J.* 2009;28:591–601.
- [29] Rizzini L, Favory -J-J, Cloix C, et al. Perception of UV-B by the Arabidopsis UVR8 protein. *Science.* 2011;332:103–106.
- [30] Kaiserli E, Jenkins GI. UV-B promotes rapid nuclear translocation of the Arabidopsis UV-B-specific signaling component UVR8 and activates its function in the nucleus. *Plant Cell.* 2007;19:2662–2673.
- [31] Di W, Hu Q, Yan Z, et al. Structural basis of ultraviolet-B perception by UVR8. *Nature.* 2012;484:214–219.
- [32] Keun PL, Kim C, Landgraf F, et al. EXECUTER1- and EXECUTER2-dependent transfer of stress-related signals from the plastid to the nucleus of Arabidopsis thaliana. *Proc Natl Acad Sci U S A.* 2007;104:10270–10275.
- [33] Šimková K, Moreau F, Pawlak P, et al. Integration of stress-related and reactive oxygen species-mediated signals by Topoisomerase VI in Arabidopsis thaliana. *Proc Natl Acad Sci U S A.* 2012;109:16360–16365.
- [34] Strand Å, Asami T, Alonso J, et al. Chloroplast to nucleus communication triggered by accumulation of Mg-protoporphyrinix. *Nature.* 2003;421:79–83.
- [35] Koussevitzky S, Nott A, Mockler TC, et al. Multiple signals from damaged chloroplasts converge on a common pathway to regulate nuclear gene expression. *Science.* 2007;316:715–719.
- [36] Sun X, Feng P, Xu X, et al. A chloroplast envelope-bound PHD transcription factor mediates chloroplast signals to the nucleus. *Nat Commun.* 2011;2:1–10.
- [37] de Souza A, Wang J-Z, Dehesh K. Retrograde signals: integrators of interorganellar communication and orchestrators of plant development. *Annu Rev Plant Biol.* 2017;68:85–108.
- [38] Zhao X, Huang J, Chory J. Unraveling the linkage between retrograde signaling and RNA metabolism in plants. *Trends Plant Sci.* 2020;25:141–147.
- [39] Chan KX, Phua SY, Crisp P, et al. Learning the languages of the chloroplast: retrograde signaling and beyond. *Annu Rev Plant Biol.* 2016;67:25–53.
- [40] Fang X, Zhao G, Zhang S, et al. Chloroplast-to-nucleus signaling regulates microRNA biogenesis in Arabidopsis. *Dev Cell.* 2019;48:371–382.e4.
- [41] Sun AZ, Guo FQ. Chloroplast retrograde regulation of heat stress responses in plants. *Front Plant Sci.* 2016;7:398.
- [42] Greenham K, McClung CR. Integrating circadian dynamics with physiological processes in plants. *Nat Rev Genet.* 2015;16:598–610.
- [43] Ronald J, Davis SJ. Focusing on the nuclear and sub-nuclear dynamics of light and circadian signalling. *Plant Cell Environ.* 2019;42:2871–2884.
- [44] Burko Y, Seluzicki A, Zander M, et al. Chimeric activators and repressors define HY5 activity and reveal a light-regulated feedback mechanism. *Plant Cell.* 2020;32:tpc.00772.2019.
- [45] Holm M, Ma LG, Qu LJ, et al. Two interacting bZIP proteins are direct targets of COP1-mediated control of light-dependent gene expression in Arabidopsis. *Genes Dev.* 2002;16:1247–1259.

- [46] Osterlund MT, Hardtke CS, Wei N, et al. Targeted destabilization of HY5 during light-regulated development of Arabidopsis. *Nature*. 2000;405:462–466.
- [47] Huang X, Ouyang X, Deng XW. Beyond repression of photomorphogenesis: role switching of COP/DET/FUS in light signaling. *Curr Opin Plant Biol*. 2014;21:96–103.
- [48] Lau K, Podolec R, Chappuis R, et al. Plant photoreceptors and their signaling components compete for COP1 binding via VP peptide motifs. *Embo J*. 2019;38. DOI:10.15252/embj.2019102140
- [49] Oravecz A, Baumann A, Máté Z, et al. CONSTITUTIVELY PHOTOMORPHOGENIC1 is required for the UV-B response in Arabidopsis. *Plant Cell*. 2006;18:1975–1990.
- [50] Ren H, Han J, Yang P, et al. Two E3 ligases antagonistically regulate the UV-B response in Arabidopsis. *Proc Natl Acad Sci U S A*. 2019;116:4722–4731.
- [51] Yang Y, Liang T, Zhang L, et al. UVR8 interacts with WRKY36 to regulate HY5 transcription and hypocotyl elongation in Arabidopsis. *Nat Plants*. 2018;4:98–107.
- [52] Rai N, O'Hara A, Farkas D, et al. The photoreceptor UVR8 mediates the perception of both UV-B and UV-A wavelengths up to 350 nm of sunlight with responsiveness moderated by cryptochromes. *Plant Cell Environ*. 2020;pce.13752. DOI:10.1111/pce.13752
- [53] Toledo-Ortiz G, Johansson H, Lee KP, et al. The HY5-PIF regulatory module coordinates light and temperature control of photosynthetic gene transcription. *PLoS Genet*. 2014;10:e1004416–e1004416.
- [54] Ling JJ, Li J, Zhu D, et al. Noncanonical role of Arabidopsis COP1/SPA complex in repressing BIN2-mediated PIF3 phosphorylation and degradation in darkness. *Proc Natl Acad Sci U S A*. 2017;114:3539–3544.
- [55] Pham VN, Kathare PK, Huq E. Dynamic regulation of PIF5 by COP1-SPA complex to optimize photomorphogenesis in Arabidopsis. *Plant J*. 2018;96:260–273.
- [56] Dong J, Tang D, Gao Z, et al. Arabidopsis DE-ETIOLATED1 represses photomorphogenesis by positively regulating phytochrome-interacting factors in the dark. *Plant Cell*. 2014;26:3630–3645.
- [57] Pepper A, Delaney T, Washburnt T, et al. DET1, a negative regulator of light-mediated development and gene expression in arabidopsis, encodes a novel nuclear-localized protein. *Cell*. 1994;78:109–116.
- [58] Schroeder DF, Gahrtz M, Maxwell BB, et al. De-etiolated 1 and damaged DNA binding protein 1 interact to regulate Arabidopsis photomorphogenesis. *Curr Biol*. 2002;12:1462–1472.
- [59] Castells E, Molinier J, Benvenuto G, et al. The conserved factor DE-ETIOLATED 1 cooperates with CUL4-DDB1^{DDB2} to maintain genome integrity upon UV stress. *Embo J*. 2011;30:1162–1172.
- [60] Fonseca S, Rubio V. Arabidopsis CRL4 complexes: surveying chromatin states and gene expression. *Front Plant Sci*. 2019;10:1095.
- [61] Park E, Kim J, Lee Y, et al. Degradation of phytochrome interacting factor 3 in phytochrome-mediated light signaling. *Plant Cell Physiol*. 2004;45:968–975.
- [62] Bauer D, Viczián A, Kircher S, et al. Constitutive photomorphogenesis 1 and multiple phytoresectors control degradation of phytochrome interacting factor 3, a transcription factor required for light signaling in arabidopsis. *Plant Cell*. 2004;16:1433–1445.
- [63] Sharma A, Sharma B, Hayes S, et al. UVR8 disrupts stabilisation of PIF5 by COP1 to inhibit plant stem elongation in sunlight. *Nat Commun*. 2019;10:1–10.
- [64] Al-Sady B, Ni W, Kircher S, et al. Photoactivated phytochrome induces rapid PIF3 phosphorylation prior to proteasome-mediated degradation. *Mol Cell*. 2006;23:439–446.
- [65] Park E, Park J, Kim J, et al. Phytochrome B inhibits binding of phytochrome-interacting factors to their target promoters. *Plant J*. 2012;72:537–546.
- [66] Shen H, Zhu L, Castillon A, et al. Light-induced phosphorylation and degradation of the negative regulator PHYTOCHROME-INTERACTING FACTOR1 from arabidopsis depend upon its direct physical interactions with photoactivated phytochromes. *Plant Cell*. 2008;20:1586–1602.
- [67] Pedmale UV, Huang -S-S, Zander M, et al. Cryptochromes interact directly with PIFs to control plant growth in limiting blue light. *Cell*. 2016;164:233–245.
- [68] Ma D, Li X, Guo Y, et al. Cryptochrome 1 interacts with PIF4 to regulate high temperature-mediated hypocotyl elongation in response to blue light. *Proc Natl Acad Sci U S A*. 2016;113:224–229.
- [69] Shi H, Shen X, Liu R, et al. The red light receptor phytochrome B directly enhances substrate-E3 ligase interactions to attenuate ethylene responses. *Dev Cell*. 2016;39:597–610.
- [70] Liang T, Mei S, Shi C, et al. UVR8 Interacts with BES1 and BIM1 to Regulate Transcription and Photomorphogenesis in Arabidopsis. *Dev Cell*. 2018;44:512–523.e5.
- [71] Wu J, Wang W, Xu P, et al. phyB Interacts with BES1 to regulate brassinosteroid signaling in Arabidopsis. *Plant Cell Physiol*. 2019;60:353–366.
- [72] Wang W, Lu X, Li L, et al. Photoexcited CRYPTOCHROME1 interacts with dephosphorylated bes1 to regulate brassinosteroid signaling and photomorphogenesis in arabidopsis. *Plant Cell*. 2018;30:1989–2005.
- [73] Liu H, Yu X, Li K, et al. Photoexcited CRY2 interacts with CIB1 to regulate transcription and floral initiation in Arabidopsis. *Science*. 2008;322:1535–1539.
- [74] Lee CM, Fekete A, Li MW, et al. Decoys untangle complicated redundancy and reveal targets of circadian clock F-BOX proteins. *Plant Physiol*. 2018;177:1170–1186.
- [75] Kim TS, Wang L, Kim YJ, et al. Compensatory mutations in GI and ZTL may modulate temperature

- compensation in the circadian clock. *Plant Physiol.* 2020;182:1130–1141.
- [76] Yang Y, Zhang L, Chen P, et al. UV-B photoreceptor UVR8 interacts with MYB73/MYB77 to regulate auxin responses and lateral root development. *Embo J.* 2020;39. DOI:10.15252/embj.2019101928
- [77] Gould PD, Locke JCW, Larue C, et al. The molecular basis of temperature compensation in the arabidopsis circadian clock. *Plant Cell.* 2006;18:1177–1187.
- [78] Antoniou-Kourounioti RL, Hepworth J, Heckmann A, et al. Temperature sensing is distributed throughout the regulatory network that controls FLC epigenetic silencing in vernalization. *Cell Syst.* 2018;7:643–655.e9.
- [79] Vu LD, Gevaert K, De Smet I. Feeling the heat: searching for plant thermosensors. *Trends Plant Sci.* 2019;24:210–219.
- [80] Martinière A, Shvedunova M, Thomson AJW, et al. Homeostasis of plasma membrane viscosity in fluctuating temperatures. *New Phytol.* 2011;192:328–337.
- [81] Knight H, Trewavas AJ, Knight MR. Cold calcium signaling in Arabidopsis involves two cellular pools and a change in calcium signature after acclimation. *Plant Cell.* 1996;8:489–503.
- [82] Gong M, van der Luit AH, Knight MR, et al. Heat-shock-induced changes in intracellular Ca^{2+} level in tobacco seedlings in relation to thermotolerance. *Plant Physiol.* 1998;116:429–437.
- [83] Guo X, Liu D, Chong K. Cold signaling in plants: insights into mechanisms and regulation. *J Integr Plant Biol.* 2018;60:745–756.
- [84] Ding Y, Shi Y, Yang S. Molecular regulation of plant responses to environmental temperatures. *Mol Plant.* 2020;13:544–564.
- [85] Saidi Y, Finka A, Muriset M, et al. The heat shock response in moss plants is regulated by specific calcium-permeable channels in the plasma membrane. *Plant Cell.* 2009;21:2829–2843.
- [86] Finka A, Cuendet AFH, Maathuis FJM, et al. Plasma membrane cyclic nucleotide gated calcium channels control land plant thermal sensing and acquired thermotolerance. *Plant Cell.* 2012;24:3333–3348.
- [87] Casal JJ, Qüesta JI. Light and temperature cues: multitasking receptors and transcriptional integrators. *New Phytol.* 2018;217:1029–1034.
- [88] Fujii Y, Tanaka H, Konno N, et al. Phototropin perceives temperature based on the lifetime of its photo-activated state. *Proc Natl Acad Sci U S A.* 2017;114:9206–9211.
- [89] Li B, Gao K, Ren H, et al. Molecular mechanisms governing plant responses to high temperatures. *J Integr Plant Biol.* 2018;60:757–779.
- [90] Åkerfelt M, Morimoto RI, Sistonen L. Heat shock factors: integrators of cell stress, development and lifespan. *Nat Rev Mol Cell Biol.* 2010;11:545–555.
- [91] Chung BYW, Balcerowicz M, Di Antonio M, et al. An RNA thermoswitch regulates daytime growth in Arabidopsis. *Nat Plants.* 2020;6:522–532.
- [92] Kumar SV, Wigge PA. H2A.Z-containing nucleosomes mediate the thermosensory response in Arabidopsis. *Cell.* 2010;140:136–147.
- [93] Cortijo S, Charoensawan V, Brestovitsky A, et al. Transcriptional regulation of the ambient temperature response by H2A.Z nucleosomes and HSF1 transcription factors in Arabidopsis. *Mol Plant.* 2017;10:1258–1273.
- [94] Del Olmo I, Poza-Viejo L, Piñeiro M, et al. High ambient temperature leads to reduced *FT* expression and delayed flowering in *Brassica rapa* via a mechanism associated with H2A.Z dynamics. *Plant J.* 2019;100:343–356.
- [95] Zhao Y, Antoniou-Kourounioti RL, Calder G, et al. Temperature-dependent growth contributes to long-term cold sensing. *Nature.* 2020;583:825–829.
- [96] Quint M, Delker C, Franklin KA, et al. Molecular and genetic control of plant thermomorphogenesis. *Nat Plants.* 2016;2:1–9.
- [97] Kumar SV, Lucyshyn D, Jaeger KE, et al. Transcription factor PIF4 controls the thermosensory activation of flowering. *Nature.* 2012;484:242–245.
- [98] Crawford AJ, McLachlan DH, Hetherington AM, et al. High temperature exposure increases plant cooling capacity. *Curr Biol.* 2012;22:R396–R397.
- [99] Li B, Gao Z, Liu X, et al. Transcriptional profiling reveals a time-of-day-specific role of REVEILLE 4/8 in regulating the first wave of heat shock-induced gene expression in Arabidopsis. *Plant Cell.* 2019;31:2353–2369.
- [100] Yamada K, Fukao Y, Hayashi M, et al. Cytosolic HSP90 regulates the heat shock response that is responsible for heat acclimation in *Arabidopsis thaliana*. *J Biol Chem.* 2007;282:37794–37804.
- [101] Hahn A, Buback D, Schleiff E, et al. Crosstalk between Hsp90 and Hsp70 chaperones and heat stress transcription factors in tomato. *Plant Cell.* 2011;23:741–755.
- [102] Ohama N, Sato H, Shinozaki K, et al. Transcriptional regulatory network of plant heat stress response. *Trends Plant Sci.* 2017;22:53–65.
- [103] Toorop PE, Barroco RM, Engler G, et al. Differentially expressed genes associated with dormancy or germination of Arabidopsis thaliana seeds. *Planta.* 2005;221:637–647.
- [104] Lim S, Park J, Lee N, et al. ABA-INSENSITIVE3, ABA-INSENSITIVE5, and DELLAs interact to activate the expression of SOMNUS and other high-temperature-inducible genes in imbibed seeds in arabidopsis. *Plant Cell.* 2013;25:4863–4878.
- [105] Gan ES, Xu Y, Wong J-Y, et al. Jumonji demethylases moderate precocious flowering at elevated temperature via regulation of FLC in Arabidopsis. *Nat Commun.* 2014;5:1–13.
- [106] Koini MA, Alvey L, Allen T, et al. High temperature-mediated adaptations in plant architecture require the bhlh transcription factor PIF4. *Curr Biol.* 2009;19:408–413.

- [107] Fiorucci A-S, Galvão VC, Ince YÇ, et al. PHYTOCHROME INTERACTING FACTOR 7 is important for early responses to elevated temperature in *Arabidopsis* seedlings. *New Phytol.* 2020;226:50–58.
- [108] Ezer D, Jung J-H, Lan H, et al. The evening complex coordinates environmental and endogenous signals in *Arabidopsis*. *Nat Plants.* 2017;3:17087.
- [109] Tasset C, Singh Yadav A, Sureshkumar S, et al. POWERDRESS-mediated histone deacetylation is essential for thermomorphogenesis in *Arabidopsis thaliana*. *PLOS Genet.* 2018;14:e1007280–e1007280.
- [110] Sun Q, Wang S, Xu G, et al. SHB1 and CCA1 interaction desensitizes light responses and enhances thermomorphogenesis. *Nat Commun.* 2019;10:3110.
- [111] Ibañez C, Delker C, Martinez C, et al. Brassinosteroids dominate hormonal regulation of plant thermomorphogenesis via BZR1. *Curr Biol.* 2018;28:303–310.e3.
- [112] Han X, Yu H, Yuan R, et al. *Arabidopsis* transcription factor TCP5 controls plant thermomorphogenesis by positively regulating PIF4 activity. *iScience.* 2019;15:611–622.
- [113] Hayes S, Sharma A, Fraser DP, et al. UV-B perceived by the UVR8 photoreceptor inhibits plant thermomorphogenesis. *Curr Biol.* 2017;27:120–127.
- [114] Delker C, Sonntag L, James G, et al. The DET1-COP1-HY5 pathway constitutes a multipurpose signaling module regulating plant photomorphogenesis and thermomorphogenesis. *Cell Rep.* 2014;9:1983–1989.
- [115] Park Y-J, Lee H-J, Ha J-H, et al. COP1 conveys warm temperature information to hypocotyl thermomorphogenesis. *New Phytol.* 2017;215:269–280.
- [116] Gangappa SN, Berriri S, Kumar SV. PIF4 coordinates thermosensory growth and immunity in *Arabidopsis*. *Curr Biol.* 2017;27:243–249.
- [117] Franklin KA, Lee SH, Patel D, et al. PHYTOCHROME-INTERACTING FACTOR 4 (PIF4) regulates auxin biosynthesis at high temperature. *Proc Natl Acad Sci U S A.* 2011;108:20231–20235.
- [118] Sun J, Qi L, Li Y, et al. PIF4-mediated activation of YUCCA8 expression integrates temperature into the auxin pathway in regulating *arabidopsis* hypocotyl growth. *PLoS Genet.* 2012;8:e1002594.
- [119] Sun J, Qi L, Li Y, et al. PIF4 and PIF5 transcription factors link blue light and auxin to regulate the phototropic response in *Arabidopsis*. *Plant Cell.* 2013;25:2102–2114.
- [120] Bellstaedt J, Trenner J, Lippmann R, et al. A mobile auxin signal connects temperature sensing in cotyledons with growth responses in hypocotyls. *Plant Physiol.* 2019;180:757–766.
- [121] Kim S, Hwang G, Kim S, et al. The epidermis coordinates thermoresponsive growth through the phyB-PIF4-auxin pathway. *Nat Commun.* 2020;11:1–13.
- [122] Catala R, Medina J, Salinas J. Integration of low temperature and light signaling during cold acclimation response in *Arabidopsis*. *Proc Natl Acad Sci U S A.* 2011;108:16475–16480.
- [123] Jia Y, Ding Y, Shi Y, et al. The *cbfs* triple mutants reveal the essential functions of CBFs in cold acclimation and allow the definition of CBF regulons in *Arabidopsis*. *New Phytol.* 2016;212:345–353.
- [124] Zhao C, Zhang Z, Xie S, et al. Mutational evidence for the critical role of CBF genes in cold acclimation in *Arabidopsis*. *Plant Physiol.* 2016;00533.2016. DOI:10.1104/pp.16.00533
- [125] Chinnusamy V. ICE1: a regulator of cold-induced transcriptome and freezing tolerance in *Arabidopsis*. *Genes Dev.* 2003;17:1043–1054.
- [126] Fursova OV, Pogorelko GV, Tarasov VA. Identification of ICE2, a gene involved in cold acclimation which determines freezing tolerance in *Arabidopsis thaliana*. *Gene.* 2009;429:98–103.
- [127] Ding Y, Li H, Zhang X, et al. OST1 kinase modulates freezing tolerance by enhancing ICE1 stability in *arabidopsis*. *Dev Cell.* 2015;32:278–289.
- [128] Ding Y, Lv J, Shi Y, et al. EGR2 phosphatase regulates OST 1 kinase activity and freezing tolerance in *Arabidopsis*. *Embo J.* 2019;38(1):e99819.
- [129] Ding Y, Jia Y, Shi Y, et al. OST 1-mediated BTF 3L phosphorylation positively regulates CBFs during plant cold responses. *Embo J.* 2018;37. DOI:10.15252/embj.201798228
- [130] Liu Z, Jia Y, Ding Y, et al. Plasma membrane CRPK1-mediated phosphorylation of 14-3-3 proteins induces their nuclear import to fine-tune CBF signaling during cold response. *Mol Cell.* 2017;66:117–128 e5.
- [131] Li H, Ding Y, Shi Y, et al. MPK3- and MPK6-mediated ICE1 phosphorylation negatively regulates ICE1 stability and freezing tolerance in *Arabidopsis*. *Dev Cell.* 2017;43:630–642.e4.
- [132] Ye K, Li H, Ding Y, et al. BRASSINOSTEROID-INSENSITIVE2 negatively regulates the stability of transcription factor ICE1 in response to cold stress in *Arabidopsis*. *Plant Cell.* 2019;tpc.00058.2019. DOI:10.1105/tpc.19.00058
- [133] Bouché F, Woods DP, Amasino RM. Winter memory throughout the plant kingdom: different paths to flowering. *Plant Physiol.* 2017;173:27–35.
- [134] Michaels SD, Amasino RM. FLOWERING LOCUS C encodes a novel MADS domain protein that acts as a repressor of flowering. *Plant Cell.* 1999;11:949–956.
- [135] Sheldon CC, Conn AB, Dennis ES, et al. Different regulatory regions are required for the vernalization-induced repression of FLOWERING LOCUS C and for the epigenetic maintenance of repression. *Plant Cell.* 2002;14:2527–2537.
- [136] Hepworth J, Antoniou-Kourounioli RL, Bloomer RH, et al. Absence of warmth permits epigenetic memory of winter in *Arabidopsis*. *Nat Commun.* 2018;9. DOI:10.1038/s41467-018-03065-7
- [137] Hepworth J, Dean C. Flowering Locus C's lessons: conserved chromatin switches underpinning

- developmental timing and adaptation. *Plant Physiol.* 2015;168:1237–1245.
- [138] Hiller M, Zhang Z, Backofen R, et al. Pre-mRNA secondary structures influence exon recognition. *PLoS Genet.* 2007;3:e204.
- [139] Erhard KF, Talbot JERB, Deans NC, et al. Nascent transcription affected by RNA polymerase IV in *Zea mays*. *Genetics.* 2015;199:1107–1125.
- [140] Hetzel J, Duttke SH, Benner C, et al. Nascent RNA sequencing reveals distinct features in plant transcription. *Proc Natl Acad Sci U S A.* 2016;113:12316–12321.
- [141] Liu W, Duttke SH, Hetzel J, et al. RNA-directed DNA methylation involves co-transcriptional small-RNA-guided slicing of polymerase ν transcripts in *Arabidopsis*. *Nat Plants.* 2018;4:181–188.
- [142] Zhu J, Liu M, Liu X, et al. RNA polymerase II activity revealed by GRO-seq and pNET-seq in *Arabidopsis*. *Nat Plants.* 2018;4:1112–1123.
- [143] Leng X, Ivanov M, Kindgren P, et al. Organismal benefits of transcription speed control at gene boundaries. *EMBO Rep.* 2020;21. DOI:10.1525/embr.201949315
- [144] Kindgren P, Ivanov M, Marquardt S. Native elongation transcript sequencing reveals temperature dependent dynamics of nascent RNAPII transcription in *Arabidopsis*. *Nucleic Acids Res.* 2019. DOI:10.1093/nar/gkz1189
- [145] Calixto CPG, Guo W, James AB, et al. Rapid and dynamic alternative splicing impacts the *Arabidopsis* cold response transcriptome[CC-BY]. *Plant Cell.* 2018;30:1424–1444.
- [146] Calixto CPG, Tzioutziou NA, James AB, et al. Cold-dependent expression and alternative splicing of *Arabidopsis* long non-coding RNAs. *Front Plant Sci.* 2019;10:235.
- [147] Mano S, Hayashi M, Nishimura M. Light regulates alternative splicing of hydroxypyruvate reductase in pumpkin. *Plant J.* 1999;17:309–320.
- [148] Pose D, Verhage L, Ott F, et al. Temperature-dependent regulation of flowering by antagonistic FLM variants. *Nature.* 2013;503:414–417.
- [149] Sureshkumar S, Dent C, Seleznev A, et al. Nonsense-mediated mRNA decay modulates FLM-dependent thermosensory flowering response in *Arabidopsis*. *Nat Plants.* 2016;2:1–7.
- [150] Kim G-D, Yoo S-D, Cho Y-H. STABILIZED1 as a heat stress-specific splicing factor in *Arabidopsis thaliana*. *Plant Signal Behav.* 2018;13:e1432955.
- [151] Lee BH, Kapoor A, Zhu J, et al. Stabilized1, a stress-upregulated nuclear protein, is required for pre-mRNA splicing, mRNA turnover, and stress tolerance in *Arabidopsis*. *Plant Cell.* 2006;18:1736–1749.
- [152] Vihervaara A, Duarte FM, Lis JT. Molecular mechanisms driving transcriptional stress responses. *Nat Rev Genet.* 2018;19:385–397.
- [153] Simpson CG, Fuller J, Maronova M, et al. Monitoring changes in alternative precursor messenger RNA splicing in multiple gene transcripts. *Plant J.* 2007;53:1035–1048.
- [154] Kurihara Y, Makita Y, Kawashima M, et al. Transcripts from downstream alternative transcription start sites evade uORF-mediated inhibition of gene expression in *Arabidopsis*. *Proc Natl Acad Sci U S A.* 2018;115:7831–7836.
- [155] Cheng Y-L, Tu S-L. Alternative splicing and cross-talk with light signaling. *Plant Cell Physiol.* 2018;59:1104–1110.
- [156] Tanabe N, Yoshimura K, Kimura A, et al. Differential expression of alternatively spliced mRNAs of *Arabidopsis* SR protein homologs, atSR30 and atSR45a, in response to environmental stress. *Plant Cell Physiol.* 2007;48:1036–1049.
- [157] Petrillo E, Godoy Herz MA, Fuchs A, et al. A chloroplast retrograde signal regulates nuclear alternative splicing. *Science.* 2014;344:427–430.
- [158] Tognacca RS, Servi L, Hernando CE, et al. Alternative splicing regulation during light-induced germination of *Arabidopsis thaliana* seeds. *Front Plant Sci.* 2019;10:1076.
- [159] Mancini E, Sanchez SE, Romanowski A, et al. Acute effects of light on alternative splicing in light-grown plants. *Photochem Photobiol.* 2016;92:126–133.
- [160] Shikata H, Hanada K, Ushijima T, et al. Phytochrome controls alternative splicing to mediate light responses in *Arabidopsis*. *Proc Natl Acad Sci U S A.* 2014;111:18781–18786.
- [161] Xin R, Kathare PK, Huq E. Coordinated regulation of pre-mRNA splicing by the SFPS-RRC1 complex to promote photomorphogenesis. *Plant Cell.* 2019;31:2052–2069.
- [162] Ushijima T, Hanada K, Gotoh E, et al. Light controls protein localization through phytochrome-mediated alternative promoter selection. *Cell.* 2017;171:1316–1325.e12.
- [163] Zhang H, Wang Y, Function LJ. Evolution of upstream ORFs in eukaryotes. *Trends Biochem Sci.* 2019;44:782–794.
- [164] Hartmann L, Drewe-Boß P, Wießner T, et al. Alternative splicing substantially diversifies the transcriptome during early photomorphogenesis and correlates with the energy availability in *Arabidopsis*. *Plant Cell.* 2016;28:2715–2734.
- [165] Godoy Herz MA, Kubaczka MG, Brzyżek G, et al. Light regulates plant alternative splicing through the control of transcriptional elongation. *Mol Cell.* 2019;73:1066–1074.e3.
- [166] Shikata H, Shibata M, Ushijima T, et al. The RS domain of *Arabidopsis* splicing factor RRC1 is required for phytochrome B signal transduction. *Plant J.* 2012;70:727–738.
- [167] Wu HP, Su Y-S, Chen H-C, et al. Genome-wide analysis of light-regulated alternative splicing mediated by

- photoreceptors in *Physcomitrella patens*. *Genome Biol.* 2013;15:R10.
- [168] Xin R, Zhu L, Salomé PA, et al. SPF45-related splicing factor for phytochrome signaling promotes photomorphogenesis by regulating pre-mRNA splicing in *Arabidopsis*. *Proc Natl Acad Sci U S A.* 2017;114:E7018–E7027.
- [169] Dong J, Chen H, Deng XW, et al. Phytochrome B induces intron retention and translational inhibition of PHYTOCHROME-INTERACTING FACTOR3. *Plant Physiol.* 2020;182:159–166.
- [170] Shih CJ, Chen H-W, Hsieh H-Y, et al. Heterogeneous nuclear ribonucleoprotein H1 coordinates with phytochrome and the U1 snRNP complex to regulate alternative splicing in *Physcomitrella patens*. *Plant Cell.* 2019;31:2510–2524.
- [171] Lin BY, Shih CJ, Hsieh HY, et al. Phytochrome coordinates with a hnRNP to regulate alternative splicing via an exonic splicing silencer. *Plant Physiol.* 2020;182:243–254.
- [172] Ma Y, Gil S, Grasser KD, et al. Targeted recruitment of the basal transcriptional machinery by LNK clock components controls the circadian rhythms of nascent RNAs in *Arabidopsis*. *Plant Cell.* 2018;30:907–924.
- [173] Rugnone ML, Faigon Soverna A, Sanchez SE, et al. LNK genes integrate light and clock signaling networks at the core of the *Arabidopsis* oscillator. *Proc Natl Acad Sci U S A.* 2013;110:12120–12125.
- [174] Csorba T, Questa JI, Sun Q, et al. Antisense COOLAIR mediates the coordinated switching of chromatin states at FLC during vernalization. *Proc Natl Acad Sci U S A.* 2014;111:16160–16165.
- [175] Swiezewski S, Liu F, Magusin A, et al. Cold-induced silencing by long antisense transcripts of an *Arabidopsis* polycomb target. *Nature.* 2009;462:799–802.
- [176] Rosa S, Duncan S, Dean C. Mutually exclusive sense-antisense transcription at FLC facilitates environmentally induced gene repression. *Nat Commun.* 2016;7:13031.
- [177] Kim DH, Sung S. Vernalization-triggered intragenic chromatin loop formation by long noncoding RNAs. *Dev Cell.* 2017;40:302–312 e4.
- [178] Heo JB, Sung S. Vernalization-mediated epigenetic silencing by a long intronic noncoding RNA. *Science.* 2011;331:76–79.
- [179] Zhao X, Li J, Lian B, et al. Global identification of *Arabidopsis* lncRNAs reveals the regulation of MAF4 by a natural antisense RNA. *Nat Commun.* 2018;9:1–12.
- [180] Kim DH, Sung S. Coordination of the vernalization response through a VIN3 and FLC gene family regulatory network in *Arabidopsis*. *Plant Cell.* 2013;25:454–469.
- [181] Kindgren P, Ard R, Ivanov M, et al. Transcriptional read-through of the long non-coding RNA SVALKA governs plant cold acclimation. *Nat Commun.* 2018;9:1–11.
- [182] Niehrs C, Luke B. Regulatory R-loops as facilitators of gene expression and genome stability. *Nat Rev Mol Cell Biol.* 2020;21:167–178.
- [183] Hegazy YA, Fernando CM, Tran EJ. The balancing act of R-loop biology: the good, the bad, and the ugly. *J Biol Chem.* 2020;295:905–913.
- [184] Crossley MP, Bocek M, Cimprich KA. R-loops as cellular regulators and genomic threats. *Mol Cell.* 2019;73:398–411.
- [185] Belotserkovskii BP, Tornaletti S, D'Souza AD, et al. R-loop generation during transcription: formation, processing and cellular outcomes. *DNA Repair (Amst).* 2018;71:69–81.
- [186] Sun Q, Csorba T, Skourti-Stathaki K, et al. R-loop stabilization represses antisense transcription at the *Arabidopsis* FLC locus. *Science.* 2013;340:619–621.
- [187] Ariel F, Lucero L, Christ A, et al. R-loop mediated trans action of the APOLO long noncoding RNA. *Mol Cell.* 2020;77:1055–1065.e4.
- [188] Xu W, Li K, Li S, et al. The R-loop atlas of *Arabidopsis* development and responses to environmental stimuli. *Plant Cell.* 2020;32:tpc.00802.2019.
- [189] Wang Y, Fan X, Lin F, et al. *Arabidopsis* noncoding RNA mediates control of photomorphogenesis by red light. *Proc Natl Acad Sci U S A.* 2014;111:10359–10364.
- [190] McClung CR, Davis SJ. Ambient thermometers in plants: from physiological outputs towards mechanisms of thermal sensing. *Curr Biol.* 2010;20:R1086–R1092.
- [191] Kaiserli E, Perrella G, Davidson ML. Light and temperature shape nuclear architecture and gene expression. *Curr Opin Plant Biol.* 2018;45:103–111.
- [192] Lindquist S. The heat-shock response. *Annu Rev Biochem.* 1986;55:1151–1191.
- [193] Pecinka A, Dinh HQ, Baubec T, et al. Epigenetic regulation of repetitive elements is attenuated by prolonged heat stress in *Arabidopsis*. *Plant Cell.* 2010;22:3118–3129.
- [194] Wang L-C, Wu J-R, Chang W-L, et al. *Arabidopsis* HIT4 encodes a novel chromocentre-localized protein involved in the heat reactivation of transcriptionally silent loci and is essential for heat tolerance in plants. *J Exp Bot.* 2013;64:1689–1701.
- [195] Sullivan AM, Arsovski A, Lempe J, et al. Mapping and dynamics of regulatory DNA and transcription factor networks in *A. thaliana*. *Cell Rep.* 2014;8:2015–2030.
- [196] Wang LC, Wu JR, Hsu YJ, et al. *Arabidopsis* HIT4, a regulator involved in heat-triggered reorganization of chromatin and release of transcriptional gene silencing, relocates from chromocenters to the nucleolus in response to heat stress. *New Phytol.* 2015;205:544–554.
- [197] Tessadori F, Schulkes RK, van Driel R, et al. Light-regulated large-scale reorganization of chromatin during the floral transition in *Arabidopsis*. *Plant J.* 2007;50:848–857.
- [198] Bourbousse C, Mestiri I, Zabulon G, et al. Light signaling controls nuclear architecture reorganization during

- seedling establishment. *Proc Natl Acad Sci U S A*. 2015;112:E2836–E2844.
- [199] Bourbousse C, Barneche F, Laloi C. Plant chromatin catches the sun. *Front Plant Sci*. 2020;10:1728.
- [200] Perrella G, Zioutopoulou A, Headland LR, et al. The impact of light and temperature on chromatin organisation and plant adaptation. *J Exp Bot*. 2020;71:5247–5255.
- [201] Benvenuto G, Formiggini F, Laflamme P, et al. The photomorphogenesis regulator DET1 binds the amino-terminal tail of histone H2B in a nucleosome context. *Curr Biol*. 2002;12:1529–1534.
- [202] Bourbousse C, Ahmed I, Roudier F, et al. Histone H2B monoubiquitination facilitates the rapid modulation of gene expression during *Arabidopsis* photomorphogenesis. *PLoS Genet*. 2012;8:e1002825.
- [203] Nassrallah A, Rougée M, Bourbousse C, et al. DET1-mediated degradation of a SAGA-like deubiquitination module controls H2Bub homeostasis. *Elife*. 2018;7. DOI:10.7554/eLife.37892
- [204] Zhao C, Liu B, Piao S, et al. Temperature increase reduces global yields of major crops in four independent estimates. *Proc Natl Acad Sci U S A*. 2017;114:9326–9331.
- [205] Van Der Woude LC, Perrella G, Snoek BL, et al. HISTONE DEACETYLASE 9 stimulates auxin-dependent thermomorphogenesis in *Arabidopsis thaliana* by mediating H2A.Z depletion. *Proc Natl Acad Sci U S A*. 2019;116:25343–25354.
- [206] Pajoro A, Severing E, Angenent GC, et al. Histone H3 lysine 36 methylation affects temperature-induced alternative splicing and flowering in plants. *Genome Biol*. 2017;18:102.
- [207] Charron JBF, He H, Elling AA, et al. Dynamic landscapes of four histone modifications during deetiolation in *Arabidopsis*. *Plant Cell*. 2009;21:3732–3748.
- [208] Peng M, Li Z, Zhou N, et al. Linking phytochrome-interacting factor to histone modification in plant shade avoidance. *Plant Physiol*. 2018;176:1341–1351.
- [209] Yang C, Shen W, Yang L, et al. HY5-HDA9 module transcriptionally regulates plant autophagy in response to light-to-dark conversion and nitrogen starvation. *Mol Plant*. 2020;13:515–531.
- [210] Liu X, Chen C-Y, Wang K-C, et al. PHYTOCHROME INTERACTING FACTOR3 associates with the histone deacetylase HDA15 in repression of chlorophyll biosynthesis and photosynthesis in etiolated *Arabidopsis* seedlings. *Plant Cell*. 2013;25:1258–1273.
- [211] Gu D, Chen C-Y, Zhao M, et al. Identification of HDA15-PIF1 as a key repression module directing the transcriptional network of seed germination in the dark. *Nucleic Acids Res*. 2017;45:7137–7150.
- [212] Zhang D, Li Y, Zhang X, et al. The SWI2/SNF2 chromatin-remodeling ATPase BRAHMA regulates chlorophyll biosynthesis in *Arabidopsis*. *Mol Plant*. 2017;10:155–167.
- [213] Lee N, Kang H, Lee D, et al. A histone methyltransferase inhibits seed germination by increasing PIF1 mRNA expression in imbibed seeds. *Plant J*. 2014;78:282–293.
- [214] Cho JN, Ryu J-Y, Jeong Y-M, et al. Control of seed germination by light-induced histone arginine demethylation activity. *Dev Cell*. 2012;22:736–748.
- [215] Jones MA, Morohashi K, Grotewold E, et al. *Arabidopsis* JMJD5/JMJ30 acts independently of LUX ARRHYTHMO within the plant circadian clock to enable temperature compensation. *Front Plant Sci*. 2019;10:57.
- [216] Zha P, Jing Y, Xu G, et al. PICKLE chromatin-remodeling factor controls thermosensory hypocotyl growth of *Arabidopsis*. *Plant Cell Environ*. 2017;40:2426–2436.
- [217] Jing Y, Zhang D, Wang X, et al. *Arabidopsis* chromatin remodeling factor PICKLE interacts with transcription factor HY5 to regulate hypocotyl cell elongation. *Plant Cell*. 2013;25:242–256.
- [218] Friedrich T, Faivre L, Bäurle I, et al. Chromatin-based mechanisms of temperature memory in plants. *Plant Cell Environ*. 2019;42:762–770.
- [219] Costa S, Dean C. Storing memories: the distinct phases of Polycomb-mediated silencing of *Arabidopsis* FLC. *Biochem Soc Trans*. 2019;47:1187–1196.
- [220] Shindo C, Lister C, Crevillen P, et al. Variation in the epigenetic silencing of FLC contributes to natural variation in *Arabidopsis* vernalization response. *Genes Dev*. 2006;20:3079–3083.
- [221] Questa JI, Song J, Geraldo N, et al. *Arabidopsis* transcriptional repressor VAL1 triggers Polycomb silencing at FLC during vernalization. *Science*. 2016;353:485–488.
- [222] Yuan W, Luo X, Li Z, et al. A cis cold memory element and a trans epigenome reader mediate Polycomb silencing of FLC by vernalization in *Arabidopsis*. *Nat Genet*. 2016;48:1527–1534.
- [223] De Lucia F, Crevillen P, Jones AM, et al. PHD-polycomb repressive complex 2 triggers the epigenetic silencing of FLC during vernalization. *Proc Natl Acad Sci U S A*. 2008;105:16831–16836.
- [224] Finnegan EJ, Dennis ES. Vernalization-induced trimethylation of histone H3 lysine 27 at FLC is not maintained in mitotically quiescent cells. *Curr Biol*. 2007;17:1978–1983.
- [225] Angel A, Song J, Dean C, et al. Polycomb-based switch underlying quantitative epigenetic memory. *Nature*. 2011;476:105–108.
- [226] Berry S, Hartley M, Olsson TS, et al. Local chromatin environment of a Polycomb target gene instructs its own epigenetic inheritance. *Elife*. 2015;4. DOI:10.7554/eLife.07205
- [227] Yang H, Berry S, Olsson TSG, et al. Distinct phases of Polycomb silencing to hold epigenetic memory of cold in *Arabidopsis*. *Science*. 2017;357:1142–1145.

- [228] Angel A, Song J, Yang H, et al. Vernalizing cold is registered digitally at FLC. *Proc Natl Acad Sci U S A*. 2015;112:4146–4151.
- [229] Qüesta JI, Antoniou-Kourounioti RL, Rosa S, et al. Noncoding SNPs influence a distinct phase of Polycomb silencing to destabilize long-term epigenetic memory at Arabidopsis FLC. *Genes Dev*. 2020;34:446–461.
- [230] Yang H, Howard M, Dean C. Antagonistic roles for H3K36me3 and H3K27me3 in the cold-induced epigenetic switch at Arabidopsis FLC. *Curr Biol*. 2014;24:1793–1797.
- [231] Byun YJ, Koo MY, Joo HJ, et al. Comparative analysis of gene expression under cold acclimation, deacclimation and reacclimation in Arabidopsis. *Physiol Plant*. 2014;152:256–274.
- [232] Zuther E, Schaarschmidt S, Fischer A, et al. Molecular signatures of increased freezing tolerance due to low temperature memory in Arabidopsis. *Plant Cell Environ*. 2018;42:pce.13502.
- [233] Kwon CS, Lee D, Choi G, et al. Histone occupancy-dependent and -independent removal of H3K27 trimethylation at cold-responsive genes in Arabidopsis. *Plant J*. 2009;60:112–121.
- [234] Schubert D, Primavesi L, Bishopp A, et al. Silencing by plant Polycomb-group genes requires dispersed trimethylation of histone H3 at lysine 27. *Embo J*. 2006;25:4638–4649.
- [235] Vyse K, Faivre L, Romich M, et al. Transcriptional and post-transcriptional regulation and transcriptional memory of chromatin regulators in response to low temperature. *Front Plant Sci*. 2020;11:39.
- [236] Mittler R, Finka A, Goloubinoff P. How do plants feel the heat? *Trends Biochem Sci*. 2012;37:118–125.
- [237] Stief A, Altmann S, Hoffmann K, et al. Arabidopsis miR156 regulates tolerance to recurring environmental stress through SPL transcription factors. *Plant Cell*. 2014;26:1792–1807.
- [238] Lämke J, Brzezinka K, Altmann S, et al. A hit-and-run heat shock factor governs sustained histone methylation and transcriptional stress memory. *Embo J*. 2016;35:162–175.
- [239] Brzezinka K, Altmann S, Czesnick H, et al. Arabidopsis FORGETTER1 mediates stress-induced chromatin memory through nucleosome remodeling. *Elife*. 2016;5. DOI:10.7554/eLife.17061
- [240] Ding Y, Fromm M, Avramova Z. Multiple exposures to drought ‘train’ transcriptional responses in Arabidopsis. *Nat Commun*. 2012;3:1–9.
- [241] Feng XJ, Li JR, Qi SL, et al. Light affects salt stress-induced transcriptional memory of P5CS1 in Arabidopsis. *Proc Natl Acad Sci U S A*. 2016;113:E8335–E8343.
- [242] Martínez-García JF, Moreno-Romero J. Shedding light on the chromatin changes that modulate shade responses. *Physiol Plant*. 2020;169:407–417.
- [243] Gordon MJ, Carmody M, Albrecht V, et al. Systemic and local responses to repeated HL stress-induced retrograde signaling in Arabidopsis. *Front Plant Sci*. 2013;3:303.
- [244] Rossel JB, Wilson PB, Hussain D, et al. Systemic and intracellular responses to photooxidative stress in Arabidopsis. *Plant Cell*. 2007;19:4091–4110.
- [245] Szechyńska-Hebda M, Kruk J, Górecka M, et al. Evidence for light wavelength-specific photoelectrophysiological signaling and memory of excess light episodes in Arabidopsis. *Plant Cell*. 2010;22:2201–2218.
- [246] Ganguly DR, Crisp PA, Eichten SR, et al. Maintenance of pre-existing DNA methylation states through recurring excess-light stress. *Plant Cell Environ*. 2018;41:1657–1672.
- [247] Crisp PA, Ganguly DR, Smith AB, et al. Rapid recovery gene downregulation during excess-light stress and recovery in arabidopsis. *Plant Cell*. 2017;29:1836–1863.
- [248] Topham AT, Taylor RE, Yan D, et al. Temperature variability is integrated by a spatially embedded decision-making center to break dormancy in Arabidopsis seeds. *Proc Natl Acad Sci U S A*. 2017;114:6629–6634.
- [249] Burghardt LT, Edwards BR, Donohue K. Multiple paths to similar germination behavior in *Arabidopsis*. *Thaliana New Phytol*. 2016;209:1301–1312.
- [250] Chew YH, Smith RW, Jones HJ, et al. Mathematical models light up plant signaling. *Plant Cell*. 2014;26:5–20.