Received:
01 May 2020

Revised:
14 June 2020

Accepted:
25 August 2020

<https://doi.org/10.1259/bjr.20200489>

Cite this article as:
Chandramohan A, Bhat TA, John R, Simon B. Multimodality imaging review of complex pelvic lesions in female pelvis. *Br J Radiol* 2020;
93: 20200489.

REVIEW ARTICLE

Multimodality imaging review of complex pelvic lesions in female pelvis

ANURADHA CHANDRAMOHAN, MD, FRCR, TAMEEM AHMED BHAT, MD, DNB, REETU JOHN, DMRD, MD and BETTY SIMON, DMRD, MD

Department of Radiology, Christian Medical College, Vellore, India

Address correspondence to: Dr Anuradha Chandramohan
E-mail: anuradhachandramohan@gmail.com

ABSTRACT:

Complex pelvic lesions can originate from various anatomical structures in the pelvis and pose a diagnostic dilemma due to a wide range of possible diagnoses. Accurate characterisation of these lesions would often require an algorithmic approach, which incorporates clinical findings, sequential use of multiple imaging modalities and a multiparametric approach. This approach usually aims at identifying key imaging features, which aid in anatomical localisation, morphology and tissue characterisation. There have been various attempts to standardise the lexicon used for describing adnexal masses in female patients; stratify their risk of cancer and suggest appropriate next steps in the management pathway. Through this review, we extend this approach to complex pelvic masses in female pelvis in general and will focus on optimal use of different imaging modalities to arrive at definitive diagnosis or meaningful differential diagnosis. We will also discuss potential pitfalls of imaging diagnosis and common mimics.

INTRODUCTION

Complex pelvic masses in females are thought to arise most commonly from the genital tract. But the differential diagnosis for pelvic masses in a female patient is vast. We defined complex pelvic masses in females as masses centered in the pelvis that requires more than one imaging modality, either for characterisation or for obtaining further information pertaining to determining its origin or extent, in order to optimally manage the patient. Often, it is the odds of malignancy in a complex pelvic mass that drives the management strategy and the referral pathway for the patient. Imaging plays a central role in the diagnosis, staging and surgical planning of complex pelvic masses. One or more modalities such as ultrasound, CT and MRI are used in a sequential fashion to obtain information relevant to management. Ultrasound is the first modality used for characterisation and cancer risk stratification. MRI is used for evaluation of indeterminate masses and to determine local extent. CT and positron emission tomography (PET)-CT are used largely for staging of malignant pelvic masses. While accurate diagnosis is vital for optimal management, the proximity of anatomical structures in the pelvis and the overlapping appearances in different imaging modalities pose diagnostic challenges in the evaluation of pelvic masses. Thus, a systematic approach with one or more imaging modalities is often needed for these lesions.

Through this review, we provide an overview of the multimodality approach to complex pelvic masses, roles of each imaging modality and discuss potential mimics and pitfalls in the diagnosis.

IMAGING MODALITIES AND THEIR ROLES

Ultrasound

Both transabdominal and transvaginal ultrasound with colour Doppler examination is used for evaluation of the female pelvis. Ultrasound helps in quick triage of patients by determining the presence of a pelvic mass. Around 90% of detected adnexal masses can be successfully characterised by ultrasound and Doppler examination alone.¹ Different ultrasound-based methods of stratifying the risk of malignancy such as O-RADS, GIRADS, IOTA simple rules are based on recognising ultrasound appearances associated with benign and malignant lesions. Using ultrasound, definitive diagnosis of benign lesions like a haemorrhagic cyst, dermoid cyst, endometrioma, simple ovarian or paraovarian cyst, peritoneal inclusion cyst and hydrosalpinx can be made with 94 to 100% specificity and 77 to 86% sensitivity.^{2,3} Ultrasound features which indicate high probability of malignancy are solid papillary excrescences (≥ 3 mm); solid mass or thick septa with strong colour flow; irregular margins; ascites and peritoneal nodules.^{4,5} But following an initial ultrasound, 18 to 31% of pelvic

Figure 1. Leiomyoma. (a) T2 oblique coronal MRI along the point of maximum contact between the mass and the uterus shows bridging vessels (arrow heads) coursing from the uterus to the mass. (b) T2 sagittal MRI shows 'clawing' of normal myometrium (arrow heads) around an intramural fibroid.

lesions remain indeterminate.^{6–11} Majority of these sonologically indeterminate lesions are benign and a misdiagnosis of malignancy in these patients will lead to unnecessary complications and morbidity of an extensive cancer surgery.¹² On the other hand, misdiagnosis of malignant mass as benign can lead to incomplete surgery and poor oncological outcome. Also, large pelvic masses, especially those arising from intra- and extraperitoneal non-adnexal structures, obesity, virginity and operator dependency and bowel gas pose practical difficulties for optimal ultrasound evaluation of pelvic masses.

Magnetic resonance imaging

MRI is used when ultrasound features are equivocal for malignancy; for evaluating indeterminate solid adnexal masses and to determine the origin of large complex pelvic masses.¹³ Characterisation of the pelvic masses using MRI involves two steps: first to anatomically localise the mass and second to determine its morphology and tissue composition.

Anatomical localisation of complex pelvic masses

The pelvic cavity can be divided into intra- and extraperitoneal compartments by the peritoneal reflection, which dips into the pelvic cavity. Displacement of the normal anatomic structures provides a clue to the compartment of origin. Intra- and extraperitoneal pelvic masses from anterior and middle compartment causes lateral and posterolateral displacement of the pelvic ureter. Extraperitoneal pelvic masses from the lateral and posterior compartment cause medial and anterior displacement of the ureters and the iliac vessels, may cause effacement and encasement of the external iliac vessels and loss of fat plane with the muscles of the pelvic side-wall. Presacral and retro-rectal masses displaces the rectum anteriorly and anterolaterally.¹⁴

Intraperitoneal masses can mimic an ovarian mass due to the intraperitoneal location of ovaries. Understanding the blood supply of the masses helps in identifying their origin. The ovarian blood vessels reach the mesovarium through the suspensory ligament, which suspends the ovary from the pelvic side-wall.¹⁵ The ovarian vein can be traced along its length leading to a mass arising from the ovary. This is described as an 'ovarian vascular pedicle sign' and can be seen in 93% of masses of ovarian origin and few large uterine and tubo-ovarian masses.¹⁶ Ovarian mass can cause non-visualisation of the ipsilateral

ovary (phantom organ sign) or cause the ovary to be embedded in the mass (embedded organ sign) or compressed by the mass (ovarian crescent sign) with few peripheral follicles.¹⁷ To assess if an adnexal mass is of uterine or ovarian origin, T2 oblique MRI is obtained either in a plane along the long axis of the uterus, which best shows the ovaries, or along the point of maximum contact between the mass and the uterus, in order to best the bridging vessels coursing from the uterus to the mass confirming the uterine origin (Figure 1a).¹⁸ Also, mass can deform the edges of the organ of origin like a 'beak' or a 'claw' (Figure 1b).¹⁷

Characterising indeterminate adnexal masses

The updated version of European Society of Uroradiology (ESUR) guidelines for MRI characterisation of indeterminate adnexal masses classifies MRI sequences as basic and problem-solving sequences.¹³ The basic sequences such as T1 and T2 large field of view axial MRI and T2 sagittal MRI help in getting an overview of the pelvis, and in assessing the basic signal characteristics of the pelvic mass. The problem-solving sequences include T1 fat suppressed MRI, T2 oblique MRI, diffusion-weighted imaging (DWI) and dynamic contrast-enhanced MRI (DCE-MRI).¹³ Table 1 summarises the sequences used for evaluation of adnexal masses and their specific roles.

The signal intensity of the mass on T₁ and T₂ weighted MRI is compared with the pelvic muscles. T1 'bright' adnexal masses are further evaluated with fat suppressed images to distinguish hemorrhage and fat, and these two sequences must be along the same plane to allow accurate comparison. The T2 'dark' solid components of an adnexal mass are further evaluated with DWI. The optimal b-value for interpretation of DWI findings is determined by observing the b-value that best suppresses the bladder contents, and this is usually between 800–1000 s/mm².^{13,19} T2 'dark' masses which are low signal on DWI (T2 black out effect) are definitely benign and DWI adds to the diagnostic confidence in these lesions.^{19,20} On the other hand, masses that are intermediate or bright on DWI may be benign or malignant and would need further evaluation with DCE-MRI. DWI should not be used in T1 'bright' masses since contents of T1 'bright' masses like teratoma, endometrioma and tubo-ovarian abscesses can show restricted diffusion.²¹ Also, DWI appearance of T2 heterogeneous masses and T2 hyperintense masses can be confusing due to both restricted diffusion and T2 shine-through effect. Thus, DWI should not be used on its own merit for characterising adnexal masses.

Solid components of a T1 'bright' mass, T2 'dark' masses with intermediate or high DWI signal, non-T2 'dark' solid and solid cystic masses and heterogeneous masses are evaluated with DCE-MRI. Images are obtained as a three-dimensional isotropic volume acquisition in the axial plane matching the plane of T2 and DWI to allow comparison and cross-reference. Following injection of 0.1 mmol/Kg of extracellular gadolinium-based contrast agent, the images are acquired every 15 s starting 30 s prior to contrast injection until 3 min after injection.²⁰ Time-intensity curve (TIC) of the maximally enhancing portion of the solid component of the adnexal mass is compared with the outer myometrium. While none of benign lesions show rapid and higher intensity enhancement

Table 1. MRI protocol for imaging patients with indeterminate adnexal masses adapted from ESUR recommendation.¹³

Sequences	Specific role of the sequence
T2 FSE large FOV axial MRI T2 sagittal MRI	To provide an overview of the pelvis. To assess basic signal characteristics of the mass. To look for associated features like lymphadenopathy and hydronephrosis.
T2 oblique high-resolution MRI without fat suppression along long axis of uterus or site of maximum contact with the uterus	To assess if the mass is T2 'bright' or 'dark'. To assess the origin and extent of the mass.
T1 axial	To detect T1 hyperintense foci, which could be from fat, blood, protein or mucin
T1 fat suppressed MRI	To distinguish fat from hemorrhage in T1 hyperintense lesions
Diffusion-weighted imaging (b-value 800–1000 s/mm ²)	To assess the need for dynamic contrast enhanced MRI (DCE-MRI). No contrast for masses with no restricted diffusion. Intermediate DWI and lesions with restricted diffusion is further evaluated with DCE-MRI.
T1 fat suppressed DCE-MRI with subtraction	To characterise solid component-based degrees of contrast enhancement with time shown on the time intensity curve. Type one curve with delayed low-level peak enhancement is benign Type two curve with intermediate level enhancement are indeterminate may be benign or malignant Type three curve with rapid high-level enhancement are malignant

ESUR, European Society of Uroradiology; FOV, field of view; FSE, fast spin echo.

Table 2. Differential diagnosis of complex pelvic masses in females

Gynecological masses	Non-gynecological masses
Uterus Benign: Leiomyoma/ Adenomyoma Malignant uterine masses Congenital cervico-uterine anomalies Pyometra/ hydrometra/hematometra	Gastrointestinal/ mesenteric Gastrointestinal stromal tumor Appendicular mucocoele Bowel malignancy/ lymphoma Pelvic abscess related to appendicitis or diverticulitis Peritoneal masses Peritoneal inclusion cyst Cystic mesothelioma Pseudomyxoma peritonei Post-pancreatitis pseudocyst Mesenteric or omental cyst Peritoneal hydatidosis Peritoneal carcinomatosis
Cervix Benign: Leiomyoma, adenomyoma Malignant: squamous cell carcinoma, adenocarcinoma, lymphoma, metastases Tunnel cluster and cervicitis	Presacral/ ischio-rectal masses <i>Developmental</i> : Benign: Tail gut cyst, rectal duplication cyst, epidermoid/dermoid cyst Malignant: Teratoma, teratocarcinoma, yolk sac tumor. <i>Mesenchymal</i> Benign: Myelolipoma, Solitary fibrous tumor, Castleman disease, IgG4 related pseudotumors, hemangioma, Lymphangioma Malignant: Soft tissue sarcoma, lymphoma, gastrointestinal stromal tumour <i>Neurogenic</i> Benign: Schwannoma/ Neurofibroma, neuroblastoma, anterior sacral meningocele Malignant: Neurofibrosarcoma, chordoma, malignant schwannoma <i>Osteochondral lesions</i> Metastases
Tubo-ovarian masses : Tubo-ovarian infections Endometriosis Ectopic pregnancy Malignancy	Bladder Over distended bladder Urachal cyst Bladder tumours Large diverticulum
Broad ligament : Pedunculated/ parasitic Fibroid Leiomyomatosis Paraovarian cyst	Others : Pelvic kidney/ Pancake kidney Aneurysms/ AVM of pelvic vessels Lymphadenopathy Ectopic spleen Foreign bodies
Vulvo-vaginal : Hydrocolpos/ hematocolpos Vulvo-vaginal cysts	Iatrogenic Lymphocele Hematoma Gossypiboma Collections

compared to myometrium (Type 3 TIC), 58% of malignant masses show Type 3 TIC due to increased angiogenesis, higher VEGFR-2 expression and pericyte coverage index.^{20,22–25} Benign adnexal masses show Type 1 TIC with slow progressive low-level contrast enhancement compared to the myometrium. In Type 2 TIC, there is moderate enhancement of the lesion compared to myometrium followed by plateau, and these lesions may be either benign or malignant.^{23–25} Some of the common challenges in the MRI interpretation of pelvic masses are caused by bowel gas and hip prosthesis-related susceptibility artefacts, which affects the quality of DWI and the bowel peristalsis-related movement artefacts, which affects

the quality of DCE-MRI. These can be addressed by using anti-spasmodics and by routinely emptying bladder and bowel 1 h before the study.

For extraperitoneal non-adnexal masses, the primary purpose of imaging is to determine the origin and the anatomical extent. The plane of imaging is tailored to gather maximum information on the anatomical extent of these masses, and this is achieved with T2 non-fat suppressed MRI obtained in oblique axial, sagittal and coronal planes. DWI and DCE-MRI do not have much role in the evaluation of extraperitoneal non-gynaecologic pelvic masses.

Table 3. Classification of pelvic masses based on anatomical compartment and morphology

	Anterior compartment	Middle compartment	Posterior compartment
Solid	TCC Urachal malignancy Leiomyoma	Solid ovarian masses: Fibroma/ fibrothecoma Brenner tumour Dysgerminomas Sertoli-stromal tumors/ Leydig cell tumors Granulosa cell cancer Steroid cell tumors Embryonal cancer High grade serous ovarian cancer Krukenburg's Fibroids Cervico-uterine malignancies Ileal GIST Ectopic kidney/ spleen	Lymphadenopathy Retroperitoneal fibrosis IgG4 disease Nerve sheath tumors Chordoma Desmoplastic small round cell tumor Castleman disease Recto-sigmoid malignancy/ GIST Metastases Lymphoma Leiomyomatosis
Cystic	Over distended bladder Urachal cyst Bladder diverticulum	Tubo-ovarian abscess Ectopic pregnancy Endometriosis Benign and malignant ovarian masses Post- operative collections Lymphocele Peritoneal inclusion cyst Hydrocele of canal of Nuck Other vulvo-vaginal cysts	Anterior sacral meningocele Dermoid cyst Epidermoid cyst Tail gut cyst Rectal duplication cyst Lymphangioma Abscess
Fat	Pelvic lipomatosis	Mature ovarian teratoma Immature teratoma Lipoleiomyoma	Lipoma Liposarcoma Myelolipoma Extramedullary hematopoiesis
Calcification/ ossification	Hemangioma Lymphangioma	Calcified fibroid Low grade serous ovarian cancer Ovarian teratoma Aneurysm Hydatidosis Calcified nodes	Osteosarcoma Ewing's sarcoma Chondrosarcoma Teratoma Hemangioma Lymphangioma Aneurysms/ AVM of pelvic vessels
Hemorrhagic	Organised hematoma Scar endometriosis Bladder paraganglioma	Fibroid with red degeneration Ovarian torsion Endometrioma Hemorrhagic cyst Hematometra Hematocolpos Cystic adenomyoma	Aneurysmal bone cyst Giant cell tumor Schwannomas Aneurysms/ AVM of pelvic vessels Large necrotic malignant masses
Myxoid		Myxoid degeneration of fibroid	Nerve sheath tumors Myxoid liposarcoma Aggressive angiomyxoma Ganglioneuroma Malignant fibrous histiocytoma Myxoma

MRI has an overall accuracy of 93% for characterising complex adnexal masses as benign or malignant.²⁶ There are robust scoring systems such as ADNEX-MR and O-RADS-MRI, which are over 90% sensitive and specific for characterising sonologically indeterminate pelvic masses.^{22,27} The rate of malignancy among all the females evaluated with MRI for sonologically indeterminate pelvic mass was 18%.²⁷ At the end of a good quality MRI evaluation, 11% of adnexal masses may remain indeterminate and may need surgical assessment or close follow-up.²⁷

Figure 2. Line diagram showing the anatomical compartments of the pelvis and the list of differentials for pelvic masses in these compartments.

Computed tomography

Contrast-enhanced computed tomography (CECT) is mainly indicated for staging of an obviously malignant mass detected on ultrasound. Usually, CT of the thorax, abdomen and the pelvis is used for this purpose in order to detect peritoneal, nodal and distant metastases. CECT is also the modality of choice in acute pelvic conditions presenting as a pelvic mass. Though fat containing lesions like ovarian dermoid can be seen easily, CT has little role in characterising pelvic masses. But 4.1% of CT scans performed for unrelated causes such as CT colonography can harbour indeterminate adnexal masses and would need further evaluation with ultrasound or MRI.²⁸ PET-CT has no proven role in characterising complex pelvic masses.

DIFFERENTIAL DIAGNOSIS OF COMPLEX PELVIC MASSES IN FEMALE

Pelvic masses can be broadly categorised based on the organ of origin as gynaecological and non-gynaecological (Table 2), or based on the involved anatomical compartment in the pelvis as intra- and extraperitoneal masses (Figure 2).²⁹ We can also classify pelvic masses based on predominant components of the mass such as cystic, solid, fat, haemorrhage, myxoid matrix and calcification (Table 3).³⁰⁻³⁴ The classifications that are based on anatomical compartments and predominant components are more useful for reporting radiologists, in order to develop an

Figure 3. Endometrioma. MRI shows a (a) T1 'bright' multilocular mass with (b) lower T2 signal intensity compared to T1 (positive 'T2 shading' sign). Also noted are small eccentric markedly hypointense 'T2 dark spots' (arrow heads) related to haemosiderin within chronically retracted adherent clots. (c) Ultrasound of the same lesion shows a multilocular cystic lesion with fine ground glass internal echoes and corresponding to T2 dark spots seen in MRI, there are echogenic foci along the wall and septae (arrow heads).

approach for lesion characterisation and provide information relevant to management.

COMPLEX PELVIC MASSES THAT COMMONLY NEED A MULTI-MODALITY APPROACH:

i) complex cystic adnexal lesions

Majority of complex cystic masses with equivocal solid areas or colour flow patterns on ultrasound Doppler studies are benign and most commonly endometriomas, ovarian dermoids, haemorrhagic cysts and tubo-ovarian infections. Borderline ovarian neoplasms and benign ovarian stromal tumours can masquerade

Figure 4. Endometrioma with malignancy. (a) Ultrasound shows cystic mass with hyperechoic eccentric solid component (arrowhead). (b) T1 and (c) T1 fat suppressed MRI shows a T1 'bright' mass with isointense solid papillary projections (arrowhead). (d) In the T_2 weighted MRI, the solid component appears hyperintense (arrowhead) and subtle in a background of low signal intensity cyst contents (T2 shading). DCE-MRI (not shown) must be used to confirm the nature of solid components in these lesions. DCE, dynamic contrast enhanced.

Figure 5. Endometrioid carcinoma in a 40-year-old with known endometriosis. MRI (a) T1 fat suppressed and (b) T_2 weighted images show T1 'bright' cystic mass with no T2 shading and an eccentric papillary solid area (arrowhead).

as benign and malignant masses respectively. Further evaluation with MRI helps in accurate characterisation of these lesions.

Haemorrhagic cyst vs endometrioma

Ultrasound appearances of classic haemorrhagic cyst and endometrioma are straightforward. But diagnosis of large haemorrhagic cysts and endometriomas can be confusing and MRI can help characterisation. While haemorrhagic adnexal lesions can be easily identified with basic MRI sequences (T1, T2 and T1 fat suppressed MRI), the accuracy for identifying the aetiology is only modest.^{20,35,36} The T2 shading sign (Figure 3a and b) or lower T2 signal compared to T1 signal though very sensitive (93%), lacks specificity (45%) for diagnosis of endometriomas.^{35,36} Both the haemorrhagic cysts and the endometriomas are T1 'bright' and show T2 shading. But this phenomenon is more in endometrioma due to more T1 and T2 shortening caused by concentrated protein and iron contents from repeated haemorrhage. Endometriomas may also have T2 'dark spots' (Figure 3b) along the inner wall related to marked T2 shortening from haemosiderin within a chronically retracted clot. This finding was 93% specific but only 36% sensitive for distinguishing endometriomas from haemorrhagic cysts.³⁷ The corresponding finding on ultrasound is small eccentric echogenic foci (Figure 3c), which may show twinkle artifacts on colour Doppler. Also, endometriomas have lower ADC value compared to haemorrhagic cysts.³⁸ Contrast-enhanced MRI may show intense wall enhancement in a luteal cyst when compared to endometrioma.³⁹ While haemorrhagic cysts resolve at 6–12 weeks follow-up ultrasound, endometriomas tend to persist for a long time.

Endometrioma with malignant change

Malignant transformation of endometrioma is seen in 0.3–1% of patients with endometriosis. Endometrioid carcinoma and clear cell carcinoma are the most common ovarian malignancies seen in these patients.⁴⁰ Ultrasound detected endometrioma with solid areas must be evaluated further with MRI to exclude malignancy. Solid components from malignant transformation of endometrioma are T2 intermediate in signal, and this finding can be masked by the T2 shading seen in endometrioma (Figure 4). Also, rapid increase in size and disappearance of previously seen T2 shading within an endometrioma may indicate malignancy

Figure 6. Immature teratoma. (a) Ultrasound and (b) contrast enhanced CT of a 21-year-old female shows a multilocular cystic mass with thick septations, solid components, fat and scattered calcification.

(Figure 5). In these lesions, DCE-MRI with subtraction images must be used to exclude malignancy.^{13,39}

Mature vs immature teratoma

Ultrasound has very high specificity (99%), but only moderate sensitivity (58%) for diagnosis of mature ovarian teratoma.⁴¹ This is due to its inability to distinguish lipid rich teratoma from bowel and haemorrhagic cysts. MRI with T1 and T1 fat suppressed sequences aid in the accurate diagnosis by demonstrating fat within the lesion. However, DWI and DCE-MRI is not recommended in a typical mature teratoma since their epidermoid contents can show restricted diffusion, and a small proportion may show Type 3 enhancement, which can lead to wrong diagnosis of malignancy.¹³

Though it is hard to suspect immature teratomas and malignant transformation of mature teratoma on ultrasound, there is usually little doubt regarding the malignant nature due to the solid cystic nature of these lesions, heterogeneous echogenicity and scattered foci of calcifications (Figure 6). Malignant transformation of mature teratoma typically occurs in the elderly, most commonly to squamous cell carcinoma, melanoma, sarcoma or rarely thyroid cancer.^{42–44} CT is the most common next imaging done in these patients and the diagnosis is usually made with surgical histopathology. MRI characterisation of these lesions should include T1 fat suppressed, DCE-MRI with subtraction images.

Figure 7. Teratoma with ovarian torsion. (a, b) Contrast-enhanced CT images show a large mixed density mass containing fat, calcification and cystic components in the left ovary, twisted vascular pedicle (arrowhead) and free fluid.

Figure 8. Mature teratoma presenting with bleeding per rectum and rectal growth. (a, b) Contrast-enhanced CT shows a mixed density mass with fat, calcification and soft tissue in the left adnexa with a large polypoidal component in the rectal lumen.

In patients presenting with ovarian torsion from an underlying ovarian teratoma, acute presentation with pelvic pain and enlarged avascular ovary on colour Doppler ultrasound are most useful in making accurate diagnosis. However, in patients with complete ovarian necrosis and torsion detorsion sequelae, pelvic pain may no longer be present, and the vascularity of the ovary may be restored. Either CT or MRI may be used in these scenarios to aid diagnosis (Figure 7). When MRI is used in this situation, the basic T1, T2 and T1 fat suppressed MRI sequences are enough for the diagnosis. Rarely, mature ovarian teratoma can penetrate the rectum (Figure 8) or the sigmoid colon and be mistaken for bowel malignancy.

Tubo-ovarian masses

Ultrasound diagnosis of typical hydrosalpinx is straightforward when a convoluted tubular cystic mass with incomplete septae is seen. However, pelvic inflammatory disease (PID), endometriosis, ectopic pregnancy etc. present as complex cystic lesions. In patients presenting acutely, CT usually follows an inconclusive ultrasound mainly to plan treatment and exclude other non-pelvic conditions. However, MRI evaluation is very useful in patients with chronic PID and infection complicating endometriosis, especially in females who are being treated for infertility. In these patients, adherent tubal folds can masquerade as solid components on ultrasound and increased colour flow is common due to inflammation. T1 and T2 shortening, diffusion restriction can be seen in PID due to protein rich contents of the abscess and may mimic endometrioma. CECT or MRI is useful in the diagnosis of infected endometriomas, which shows thick, intensely enhancing walls (Figure 9).^{13,39}

Borderline ovarian tumour, cystadenofibroma, Brenner tumour

Mucinous and serous cystic neoplasms of the ovary present as complex cystic masses. Ultrasound diagnosis of a borderline ovarian tumour (BOT) is challenging. The only ultrasound feature significantly associated with BOT is solid papillary components within a cystic mass (Figure 10), which is seen in 40% of mucinous BOT and 78% of serous BOT.^{45–47} Mucinous BOT tends to be multilocular with fine internal echoes, and each locule typically has different echogenicity on ultrasound

Figure 9. Infected endometrioma in a 29-year-old female on treatment for infertility. (a) Ultrasound shows multilocular cyst with fine internal echoes. (b, c) Contrast-enhanced CT shows large multilocular left adnexal cyst with uniformly enhancing smooth thick walls, adjacent fluid and fat stranding.

and density on CT (Figure 10). MRI shows variable degrees of T1 and T2 shortening. They are larger in size and have a greater number of locules compared to their serous counterparts. DCE-MRI is useful in characterising indeterminate multilocular cystic masses based on the enhancement pattern of the wall and its solid components.

Ovarian cystadenofibroma are composed of variable amounts of solid fibrous component and cystic epithelial component. Based on the nature of the epithelial component, these lesions could be benign, borderline or malignant. MRI shows multi- or unilocular cystic mass with smooth external surface and T2 'dark' solid components or thick septae from the fibrous stroma (Figure 11). DWI and DCE-MRI are often needed to ascertain the nature of the lesion. While the T2 'dark' solid component of benign cystadenofibroma shows low signal on DWI (T2 'black out' effect) and very little contrast enhancement, borderline and malignant cystadenofibroma are likely to have enhancing components.

Brenner tumour is a rare benign ovarian stromal tumour seen in older females (60–80 years) and mimics malignant ovarian mass on ultrasound. Majority (83%) have calcification (Figure 12a and b). One-third of them are associated with mucinous cystadenofibroma or teratoma and very rarely associated with serous cystadenofibroma.^{48,49} MRI shows a complex cystic mass with T2 'dark' solid component, variable DWI findings and contrast enhancement (Figure 12c–e).

Struma ovarii is a rare benign stromal tumour containing thyroid tissue, which appears as multilocular cystic lesion with variable T1 'bright' signal from haemorrhage and T2 'dark' signal from thick gelatinous colloid material. Post-contrast images show intense early enhancement of the cyst wall and septae due to

Figure 10. Mucinous borderline ovarian tumour. (a) Ultrasound and (b) MRI T₂ weighted coronal image shows a multilocular cystic mass with multiple solid papillary components.

Figure 11. Serous cystadenofibroma. (a) Ultrasound and (b) T₂ weighted MRI show a large unilocular cyst with eccentric T2 'dark' solid area (arrowhead), which appeared low signal on high b-value (b800) DWI (c). DWI, diffusion-weighted imaging.

thyroid tissue mimicking a malignancy. However, progressive centripetal pattern of enhancement in a pelvic tumour seen in young females with rare clinical presentation of hyperthyroidism in 5% of them may help in the diagnosis.^{50,51}

Though accurate histological diagnosis may not be possible in the above conditions, a combination of DWI and DCE-MRI can help in correctly suggesting possibility of benign ovarian neoplasm in 16–25% of these lesions.²⁰ Cystic masses of cervico-uterine origin are easily recognised on ultrasound, however accurate characterisation of these lesions and treatment planning would need MRI evaluation. We list the cystic uterine masses in this section for completion sake without intending to discuss them in detail. These include large hydrometra, haematometra and pyometra; leiomyoma with cystic or red degeneration; cystic adenomyoma; female genital tract anomalies, tunnel cluster and arteriovenous malformation. The key to accurate MRI diagnosis of these conditions is to demonstrate the relationship between

Figure 12. Brenner tumour in two different patients. (a) Ultrasound and (b) Contrast-enhanced CT of an 87-year-old with Brenner tumour associated with mucinous cystadenoma shows a multilocular cystic mass with poorly enhancing solid area and calcification. (c) T₂ weighted MRI and (d, e) high b-value (b800) DWI of a 29-year-old with Brenner tumour associated with serous cystadenoma shows a unilocular cyst with eccentric T2 'dark' solid component, which was low signal on DWI (arrowheads). DWI, diffusion-weighted imaging.

Figure 13. Krukenberg's tumour in two different patients. (a) Coronally reconstructed contrast-enhanced CT of a colon cancer (arrow) patient shows bilateral complex cystic ovarian masses with smooth surface. (b) Coronally reconstructed contrast enhanced CT of a gastric cancer (arrow) patient shows bilateral solid smoothly lobulated ovarian masses.

the mass and the junctional zone, endometrial cavity and the myometrium. Usually routine T2 oblique axial, coronal and sagittal high-resolution MRI, T1 and T1 fat suppressed MRI sequences are enough for adequate evaluation of these lesions.

ii) Solid adnexal lesion

Clearly malignant adnexal masses are usually solid, irregular and show increased colour flow on ultrasound Doppler examinations and are mostly accompanied by ascites and peritoneal disease. Such masses require further evaluation with CECT in order to identify if the mass is a primary ovarian malignancy or an ovarian metastasis and then to stage the disease. While primary ovarian carcinoma has irregular and papillary margins, ovarian metastases have smooth, gently lobulated or bosselated margins. Krukenberg's tumours arising from breast, stomach, lymphoma and uterus are almost always solid, while those from rectum, colon, appendix and the biliary tree tend to be complex solid cystic in nature (Figure 13).⁵² Majority (80%) of Krukenberg's tumours are bilateral and solid, but 20% may be unilateral or cystic. Enlarged ovaries due to ovarian

Figure 14. Massive ovarian oedema due to partial or intermittent ovarian torsion in a 19-year-old female who presented with pelvic pain and a solid left adnexal mass on ultrasound (not shown). (a) Coronal and (b) sagittal T₂ weighted MRI shows diffusely enlarged ovary with hyperintense stroma and multiple follicles.

hyperstimulation syndrome (OHSS) and massive ovarian oedema may also present pelvic masses. The presence of bilateral enlarged ovaries with multiple follicles in a typical clinical setting of oncofertility or infertility treatment and abdominal pain helps in the diagnosis of OHSS. Asymmetry in the ovarian sizes and the presence of ascites in advanced stages of OHSS may mimic ovarian neoplasm. Massive ovarian oedema usually follows a partial torsion or repeated intermittent torsion. The unilateral nature, presence of colour flow and free fluid in the pelvis may mimic an ovarian neoplasm. However, history of pelvic pain and MRI appearance of diffusely enlarged ovary, T2 hyperintense ovarian stroma and presence of multiple peripheral follicles will help with the diagnosis (Figure 14). Solid masses that often need a multimodality approach for characterisation are those with no or low colour flow and masses of uncertain organ of origin. Ovarian fibroma, fibrothecoma and atypical leiomyomas are masses that commonly require a multimodality evaluation.

Ovarian fibroma, fibrothecoma

Ovarian fibroma and fibrothecoma are benign ovarian stromal tumours, which usually present as solid ovarian masses with low colour flow on ultrasound and Doppler. There may be associated ascites and pleural effusion from Meigs syndrome in 10–15% of them.⁵³ Around 70% of fibrothecomas are seen in post-menopausal females and they may present with post-menopausal bleeding, endometrial hyperplasia, uterine enlargement and elevated CA-125.⁵⁴ Around 25% of fibromas and fibrothecomas are homogeneously T2 'dark' with no restricted diffusion and show lesser contrast enhancement compared to myometrium or fibroid (Figure 15). The rest of them are T2 intermediate (52%) or T2 hyperintense (22%). The larger ones more than 6 cm may have peripheral cysts; heterogeneous T2 signal; central T2 hyperintensity from degeneration; diffusion restriction and less enhancement than myometrium (Figure 16).^{54,55} Thus, DCE-MRI is essential for accurate diagnosis of heterogeneous and large lesions.

Leiomyomas

Ultrasound is mostly accurate in diagnosing leiomyomas. However, large leiomyomas, pedunculated leiomyomas, degenerated leiomyomas, atypical and cellular leiomyomas and leiomyomatosis pose diagnostic challenges. Fibroids with red degeneration present with pain and are hyperdense on CT with T1 'bright' and T2 'dark' areas on MRI. Leiomyomas with cystic and myxoid degeneration are T2 hyperintense and fibroids with myxoid degeneration may show unusual laminated patterns and intense contrast enhancement (Figure 17). Atypical fibroids including cellular fibroid, stromal tumour of unknown malignant potential (STUMP) and leiomyosarcoma present as large and solitary uterine mass with heterogeneous ultrasound appearance, with or without an irregular margin and increased colour flow. MRI of these lesions show intermediate to high signal on T₂ weighted images, T1 'bright' foci from hemorrhage, high signal intensity on DWI and heterogeneous contrast enhancement (Figure 18).⁵⁶ It is important to recognise atypical fibroids due to risk of leiomyosarcoma following inadvertent morcellation of these lesions.

Figure 15. Fibroma of ovary. (a) Ultrasound shows a solid smooth mass with posterior acoustic shadowing. (b) T_2 weighted MRI shows a homogeneous, well defined, smoothly lobulated T_2 'dark' mass (asterisk). (c, d) High b-value (b800) DWI and ADC map shows low signal (asterisk) or T_2 black out effect suggestive of a fibrous lesion. ADC, apparent diffusion coefficient; DWI, diffusion-weighted imaging.

Leiomyomatosis and extrauterine leiomyomas

Diffuse peritoneal leiomyomatosis (Figure 19a) is a very rare benign disorder where numerous vascular leiomyomas grow along the submesothelial plane into the peritoneal cavity. Broad

Figure 16. Fibrothecoma of the ovary. (a) T_2 weighted MRI shows a well-defined, heterogeneous mass with T_2 intermediate peripheral signal and hyperintense center. (b) High b-value DWI (b800) shows predominantly high signal. (c, d) Early phase of dynamic contrast enhanced MRI shows intense heterogeneous enhancement and the late phase shows that the lesion (L) enhances equal to the myometrium (U). DWI, diffusion-weighted imaging.

Figure 17. Degenerated fibroid. (a) MRI of myxoid degeneration of a submucosal fibroid shows T_2 hyperintense mass with laminated appearance. (b) MRI of cystic degeneration of fibroid shows heterogeneous T_2 hyperintense well-defined mass. (c) CT of a patient with red degeneration of fibroid shows a thick walled hyperdense lesion in the uterus. (d) CT of a large calcified fibroid.

Figure 18. Atypical fibroid. (a) T_2 sagittal MRI shows a large solitary well defined, intermediate signal intensity mass in the posterior wall of the uterus. (b) T_1 weighted MRI shows an eccentric T_1 'bright' haemorrhagic focus. (c) High b-value (b800) DWI shows high signal and (d) ADC map shows low signal. Histopathology revealed a cellular fibroid. ADC, apparent diffusion coefficient; DWI, diffusion-weighted imaging.

Figure 19. Leiomyomatosis, broad ligament fibroid and parasitic leiomyoma (a) CT of a patient with leiomyomatosis shows a grossly enlarged uterus with numerous well-defined soft tissue density lesions. Similar discrete lesions were also seen in the peritoneal cavity and the retroperitoneum (arrow). (b) CT of a broad ligament fibroid in the left retroperitoneum masquerading as retroperitoneal sarcoma. (c) CT of a parasitic leiomyoma shows a well-defined homogeneous mass in the presacral region.

ligament leiomyoma can grow into the retroperitoneum and mimic retroperitoneal sarcoma (Figure 19b). A round ligament leiomyomas may be far from the uterus and mimics lymphadenopathy. Tracing the round ligament to aid accurate anatomical localisation is the key to diagnosis. Parasitic leiomyomas can be far from the uterus and have separate blood supply and may show significant interval growth following hysterectomy (Figure 19c). The typical T2 'dark' MRI appearance and enhancement like myometrium will help in clinching the diagnosis in these conditions.

iii) Non-gynaecological pelvic masses in females

Non-adnexal masses constitute 8.6% of indeterminate pelvic masses evaluated with MRI.²⁷ Intraperitoneal complex pelvic masses that mimic gynaecological masses include peritoneal diseases in the pouch of Douglas; pelvic abscesses and inflammatory masses of bowel origin such as appendicitis and sigmoid diverticulitis; malignant masses that arise from small bowel, sigmoid colon or upper rectum such as gastrointestinal stromal tumour (GIST), lymphoma and carcinoma and rare conditions like pelvic splenosis following remote splenic injury.⁵⁷

Peritoneal inclusion cyst/ cystic mesothelioma

Peritoneal inclusion cysts and cystic mesothelioma are both on the spectrum of the same disease with later being a borderline malignant condition, more commonly seen in females. They are associated with previous pelvic surgery or pelvic inflammatory disease. MRI is the modality of choice and shows multilocular

Figure 20. Peritoneal inclusion cyst. (a, b) T_2 weighted MRI shows a multilocular cystic lesion with thin septations that takes the shape of the pelvis and surrounds the uterus and the ovaries.

Figure 21. Peritoneal hydatidosis and abdominal tuberculosis (a, b) Contrast-enhanced CT of a patient with peritoneal hydatidosis shows multilocular cystic lesions with enhancing walls in the pelvis and subhepatic space. (c) Ultrasound and (d) CT of abdominal tuberculosis shows loculated ascites and diffuse, enhancing, smooth peritoneal thickening.

cystic lesions with thin non-enhancing walls and septations that take the shape of the pelvis, surrounds the ovaries and may extend into the rest of the peritoneal cavity (Figure 20). Cystic ovarian neoplasms, endometriomas, peritoneal hydatidosis (Figure 21a and b), loculated ascites from abdominal tuberculosis (Figure 21c and d) mimics peritoneal inclusion cyst, but these show wall enhancement.⁵⁸ Pseudomyxoma peritonei may closely mimic cystic mesothelioma on CT and MRI, but appears solid on ultrasound (Figure 22).

Masses of bowel origin

Large masses from GIST, lymphoma and malignancy of small bowel and rectosigmoid can masquerade as gynaecological malignancy. Nearly 4% of female pelvic masses, which were initially suspected to be a gynaecological malignancy turns out to be GIST.⁵⁹ Based on ultrasound findings, these lesions are usually diagnosed as a malignant pelvic mass and the patient is referred for staging contrast enhanced CT. On CT, GIST appears as a large heterogeneously enhancing, necrotic, exophytic mass arising from the submucosal layer of the small or large bowel (Figure 23). There may be air pockets within the mass due to ulceration and communication with bowel lumen. Bowel adenocarcinoma, especially mucinous adenocarcinoma of the rectosigmoid tend to present as large bulky low-density mass and may contiguously infiltrate or metastasis to ovaries mimicking an ovarian malignancy (Figure 24). The vascular supply of the mass usually aids in accurate identification of the origin of the mass (Figure 23). While a bowel mass is supplied by splanchnic vessels, an ovarian mass is supplied by ovarian vessels and a dilated ipsilateral ovarian vein can be traced to the mass.⁶⁰ When the diagnosis is in doubt, ultrasound-guided biopsy must be performed to confirm the diagnosis. Appendicular mucocele can mimic a

Figure 22. Pseudomyxoma peritonei. (a) Contrast-enhanced CT shows low density diffuse peritoneal process that appears solid on ultrasound (b).

right adnexal mass of ovarian or tubal origin. While appendicular mucocele presents as featureless blind ending tubular structure with or without mural calcification, a tubo-ovarian mass has incomplete septations due to thickened longitudinal folds (Figure 25a and b). Also, identification of the right ovary separate from the mass and tracing the tubular mass to the bowel will help in the diagnosis. Having said this, synchronous presence of both appendicular mucocele and ovarian mass is well known and this is most commonly due to ovarian metastases from appendicular mucinous neoplasm and very rarely due to synchronous occurrence of both appendicular and ovarian mucinous neoplasms (Figure 25c).⁶¹

Extraperitoneal non-gynaecological masses

Extra peritoneal non-gynaecological masses may arise from the bladder, pelvic mesenchymal tissue, rectum, presacral space and ischio-rectal space (Table 2). MRI plays a key role in identifying the anatomical origin and the extent of non-adnexal extraperitoneal masses. Though discussion of all these conditions is beyond the scope of the current review, some presacral masses are discussed here.

Unilocular cystic masses in the presacral space include anterior sacral meningocele, epidermoid cyst, dermoid cyst and rectal duplication cyst. Anterior sacral meningocele is a cerebrospinal fluid signal intensity, thin walled, unilocular cyst that is continuous with the spinal canal. Epidermoid cysts are thin walled, unilocular presacral or ischio-rectal cysts with T1 'bright' contents that show restricted diffusion due to protein rich epidermal contents (Figure 26). Dermoid cysts most commonly occur in

Figure 23. Rectal gastrointestinal stromal tumour mimicking ovarian cancer. (a, b) Contrast CT shows a large solid cystic pelvic mass. Though it is difficult to determine the organ of origin, dilated retroperitoneal vessels (arrowheads) give a clue to extraperitoneal origin of the mass.

Figure 24. Mucinous rectal adenocarcinoma with ovarian metastases. (a) CT shows low density annular rectal wall thickening and bilateral low-density ovarian masses. (b) MRI shows T2 hyperintense rectal wall thickening and bilateral hyperintense smooth ovarian masses.

the young as a unilocular cyst and may have fat and calcification. Rectal duplication cyst is seen as unilocular retro-rectal cyst with its wall showing gut signature and its contents may be T1 'bright' due to bleeding from ectopic gastric mucosa or may be filled with debris and air from bowel communication.

Multilocular cystic presacral or ischio-rectal masses include tail gut cyst, cystic schwannoma, cystic lymphangioma and teratoma. Tailgut cysts are congenital cysts arising from the vestigial remnant of the hindgut in the retro-rectal space, which is between the posterior wall of the rectum and the presacral fascia. Tailgut cysts are most seen in middle aged females and MRI shows multilocular thin walled cystic mass with T1 'bright' and T2 'dark' contents due to mucinous material within the cyst (Figure 27). Malignant transformation can be seen in 7–25% of patients with tailgut cyst, most commonly to adenocarcinoma or neuroendocrine carcinoma.^{62,63} This is seen as a nodular solid component within the cyst. Tail gut cysts are also prone for recurrent infection, bleeding and recurrence following surgery. Iatrogenic lymphoceles (Figure 28) and haematomas in the pelvis also appear multilocular and takes the shape of the tissue plane. History of recent surgery and short interval follow-up to demonstrate temporal resolution are key to diagnosis.

Diagnosis of solid extraperitoneal non-adnexal masses is mainly made with biopsy. Imaging is used for assessing the extent of disease and for planning treatment. However, in some rare conditions with unique imaging appearance like Castleman's disease

Figure 25. Appendicular mucocele from low grade mucinous appendicular neoplasm in two different patients. (a, b) Ruptured tubular mass in the right lower abdomen and pelvis with a rim of intramural calcification (arrows) and low-density peritoneal contents from pseudomyxoma peritonei (c) Synchronous appendicular mucocele (arrow) and mucinous ovarian neoplasm (asterisk).

Figure 26. Presacral epidermoid cyst in a 29-year-old female. (a-c) T2 and T₁ weighted MRI shows a large, thin walled, unilocular T1 'bright', T2 heterogeneous presacral and right ischio-rectal mass. (d) High b-value (b800) DWI and (e) ADC map shows restricted diffusion. ADC, apparent diffusion coefficient; DWI, diffusion-weighted imaging.

and aggressive angiomyxoma accurate histopathological diagnosis is possible. Castleman's disease is a rare benign neoplasm of lymph nodes and its occurrence in the pelvis is even rarer.⁶⁴ However, post-contrast CT or MRI appearance of Castleman's disease is classical with homogeneous and intense enhancement in a smoothly marginated mass (Figure 29). Aggressive angiomyxoma is a rare mesenchymal tumour most seen in the pelvis of middle-aged females. Due to the soft pliable nature of aggressive angiomyxoma, it is often clinically mistaken as giant Bartholin's cyst, canal of Nuck hernia or pelvic collection. Characteristic MRI appearance of T2 hyperintense multicompartamental mass in the pelvis and perineum with laminated appearance due to myxoid matrix and intense contrast enhancement helps in accurate diagnosis (Figure 30).⁶⁵

Figure 27. Tail gut cyst. (a) Sagittal and (b) axial T₂ weighted MRI of a 21-year-old female shows a large, thin walled, multilocular cystic lesion in the retro-rectal space. The contents of the cyst have varying signal intensity. Also note the partial sacral agenesis.

Figure 28. Iatrogenic lymphocele in a 45-year-old female after surgery for carcinoma endometrium which involved pelvic nodal dissection. (a) Axial and (b) coronal-enhanced CT shows thin walled, lobulated, fluid density extraperitoneal lesion.

SUMMARY AND PEARLS OF MULTIMODALITY IMAGING WORK-UP OF COMPLEX PELVIC MASSES

- (1) Majority of female pelvic masses arise from gynaecological structures and most of them are benign. Ultrasound is the first imaging modality used in their evaluation.
- (2) Clinical findings (age, examination findings, tumour markers) must be considered while providing a specific diagnosis or differential diagnosis.
- (3) Ultrasound helps in classifying pelvic masses as benign, malignant or indeterminate.
- (4) Sonologically indeterminate adnexal masses are characterised using MRI.
 - Pure unilocular cysts, teratoma and endometrioma are diagnosed on T1, T2 and T1 fat suppressed images
 - T2 'dark' masses with low DWI signal are definitely benign.
 - All sonologically indeterminate adnexal masses with solid areas that are not T2 'dark' with low signal on DWI must be evaluated with dynamic contrast enhanced MRI.
- (5) Non-adnexal intraperitoneal solid masses often end up being evaluated with CECT due to presumed gynaecological

Figure 29. Presacral Castleman's disease. (a, b) Contrast-enhanced CT shows a well-defined, homogeneous, intensely enhancing mass in the presacral region (asterisk) causing anterior and leftwards displacement of the rectum.

Figure 30. Aggressive angiomyxoma. (a) Coronal and (b) Sagittal T2 high resolution MRI shows a T2 hyperintense mass with hypointense linear lines giving a laminated appearance in the left ischioirectal fossa.

malignancy. High index of suspicion and vascular supply of

these masses aid in diagnosis.

- (6) Extraperitoneal non-gynaecological masses are evaluated with MRI to determine the location and the extent.

CONCLUSION

While it is important not to miss a diagnosis of malignant pelvic mass, it is equally vital not to overdiagnose malignancy, since this will lead to morbidity of extensive oncological surgery, unnecessary cost and misuse of limited resources available. Thus, adopting a patient centered, multimodality tailored approach using ultrasound, MRI and CT will aid in accurate diagnosis of complex pelvic masses in females.

ACKNOWLEDGEMENT

Dr Abraham Peedicayil, MD, MPH, FRCOG, Professor, Department of Gynecological Oncology, Christian Medical College, Vellore.

REFERENCES

- Valentin L, Ameye L, Jurkovic D, Metzger U, Lécure F, Van Huffel S, et al. Which extrauterine pelvic masses are difficult to correctly classify as benign or malignant on the basis of ultrasound findings and is there a way of making a correct diagnosis? *Ultrasound Obstet Gynecol* 2006; **27**: 438–44. doi: <https://doi.org/10.1002/uog.2707>
- Sokalska A, Timmerman D, Testa AC, Van Holsbeke C, Lissoni AA, Leone FPG, et al. Diagnostic accuracy of transvaginal ultrasound examination for assigning a specific diagnosis to adnexal masses. *Ultrasound Obstet Gynecol* 2009; **34**: 462–70. doi: <https://doi.org/10.1002/uog.6444>
- Levine D, Brown DL, Andreotti RF, Benacerraf B, Benson CB, Brewster WR, et al. Management of asymptomatic ovarian and other adnexal cysts imaged at US: Society of radiologists in ultrasound consensus conference statement. *Radiology* 2010; **256**: 943–54. doi: <https://doi.org/10.1148/radiol.10100213>
- Nunes N, Ambler G, Foo X, Naftalin J, Widschwendter M, Jurkovic D. Use of iota simple rules for diagnosis of ovarian cancer: meta-analysis. *Ultrasound Obstet Gynecol* 2014; **44**: 503–14. doi: <https://doi.org/10.1002/uog.13437>
- Andreotti RF, Timmerman D, Strachowski LM, Froyman W, Benacerraf BR, Bennett GL, et al. O-RADS us risk stratification and management system: a consensus guideline from the ACR Ovarian-Adnexal reporting and data system Committee. *Radiology* 2020; **294**: 168–85. doi: <https://doi.org/10.1148/radiol.2019191150>
- Froyman W, Landolfo C, De Cock B, Wynants L, Sladkevicius P, Testa AC, et al. Risk of complications in patients with conservatively managed ovarian tumours (IOTA5): a 2-year interim analysis of a multicentre, prospective, cohort study. *Lancet Oncol* 2019; **20**: 448–58. doi: [https://doi.org/10.1016/S1470-2045\(18\)30837-4](https://doi.org/10.1016/S1470-2045(18)30837-4)
- Meys EMJ, Kaijser J, Kruitwagen RFP, Slangen BFM, Van Calster B, Aertgeerts B, et al. Subjective assessment versus ultrasound models to diagnose ovarian cancer: a systematic review and meta-analysis. *Eur J Cancer* . 2016; **58**: 17–291990. doi: <https://doi.org/10.1016/j.ejca.2016.01.007>
- Van Calster B, Timmerman D, Valentin L, McIndoe A, Ghaem-Maghani S, Testa AC, et al. Triaging women with ovarian masses for surgery: observational diagnostic study to compare RCOG guidelines with an international ovarian tumour analysis (iota) group protocol. *BJOG* 2012; **119**: 662–71. doi: <https://doi.org/10.1111/j.1471-0528.2012.03297.x>
- Basha MAA, Refaat R, Ibrahim SA, Madkour NM, Awad AM, Mohamed EM, et al. Gynecology imaging reporting and data system (GI-RADS): diagnostic performance and inter-reviewer agreement. *Eur Radiol* 2019; **29**: 5981–90. doi: <https://doi.org/10.1007/s00330-019-06181-0>
- Meys EMJ, Jeelof LS, Achten NMJ, Slangen BFM, Lambrechts S, Kruitwagen RFP, et al. Estimating risk of malignancy in adnexal masses: external validation of the ADNEX model and comparison with other frequently used ultrasound methods. *Ultrasound Obstet Gynecol* 2017; **49**: 784–92. doi: <https://doi.org/10.1002/uog.17225>
- Sadowski EA, Paroder V, Patel-Lippmann K, Robbins JB, Barroilhet L, Maddox E, et al. Indeterminate adnexal cysts at US: prevalence and characteristics of ovarian cancer. *Radiology* 2018; **287**: 1041–9. doi: <https://doi.org/10.1148/radiol.2018172271>
- Jacobs IJ, Menon U, Ryan A, Gentry-Maharaj A, Burnell M, Kalsi JK, et al. Ovarian cancer screening and mortality in the UK Collaborative trial of ovarian cancer screening (UKCTOCS): a randomised controlled trial. *Lancet* 2016; **387**: 945–56. doi: [https://doi.org/10.1016/S0140-6736\(15\)01224-6](https://doi.org/10.1016/S0140-6736(15)01224-6)
- Forstner R, Thomassin-Naggara I, Cunha TM, Kinkel K, Masselli G, Kubik-Huch R, et al. ESUR recommendations for MR imaging of the sonographically indeterminate adnexal mass: an update. *Eur Radiol* 2017; **27**: 2248–57. doi: <https://doi.org/10.1007/s00330-016-4600-3>
- Foshager MC, Hood LL, Walsh JW. Masses simulating gynecologic diseases at CT and MR imaging. *Radiographics* 1996; **16**: 1085–99. doi: <https://doi.org/10.1148/radiographics.16.5.8888392>
- Saksouk FA, Johnson SC. Recognition of the ovaries and ovarian origin of pelvic masses with CT. *Radiographics* 2004; **24** Suppl 1(suppl1): S133–46. doi: <https://doi.org/10.1148/rg.24si045507>

16. Lee JH, Jeong YK, Park JK, Hwang JC. "Ovarian Vascular Pedicle" Sign Revealing Organ of Origin of a Pelvic Mass Lesion on Helical CT. *American Journal of Roentgenology* 2003; **181**: 131–7. doi: <https://doi.org/10.2214/ajr.181.1.1810131>
17. Nishino M, Hayakawa K, Minami M, Yamamoto A, Ueda H, Takasu K. Primary retroperitoneal neoplasms: CT and MR imaging findings with anatomic and pathologic diagnostic clues. *RadioGraphics* 2003; **23**: 45–57. doi: <https://doi.org/10.1148/rg.231025037>
18. Kim JC, Kim SS, Park JY. "Bridging vascular sign" in the MR diagnosis of exophytic uterine leiomyoma. *J Comput Assist Tomogr* 2000; **24**: 57–60. doi: <https://doi.org/10.1097/00004728-200001000-00012>
19. Thomassin-Naggara I, Daraï E, Cuenod CA, Fournier L, Toussaint I, Marsault C, et al. Contribution of diffusion-weighted MR imaging for predicting benignity of complex adnexal masses. *Eur Radiol* 2009; **19**: 1544–52. doi: <https://doi.org/10.1007/s00330-009-1299-4>
20. Thomassin-Naggara I, Toussaint I, Perrot N, Rouzier R, Cuenod CA, Bazot M, et al. Characterization of complex adnexal masses: value of adding perfusion- and diffusion-weighted MR imaging to conventional MR imaging. *Radiology* 2011; **258**: 793–803. doi: <https://doi.org/10.1148/radiol.10100751>
21. Fujii S, Kakite S, Nishihara K, Kanasaki Y, Harada T, Kigawa J, et al. Diagnostic accuracy of diffusion-weighted imaging in differentiating benign from malignant ovarian lesions. *J Magn Reson Imaging* 2008; **28**: 1149–56. doi: <https://doi.org/10.1002/jmri.21575>
22. Thomassin-Naggara I, Aubert E, Rockall A, Jalaguer-Coudray A, Rouzier R, Daraï E, et al. Adnexal masses: development and preliminary validation of an Mr imaging scoring system. *Radiology* 2013; **267**: 432–43. doi: <https://doi.org/10.1148/radiol.13121161>
23. Thomassin-Naggara I, Bazot M, Daraï E, Callard P, Thomassin J, Cuenod CA. Epithelial ovarian tumors: value of dynamic contrast-enhanced MR imaging and correlation with tumor angiogenesis. *Radiology* 2008; **248**: 148–59. doi: <https://doi.org/10.1148/radiol.2481071120>
24. Thomassin-Naggara I, Balvay D, Aubert E, Daraï E, Rouzier R, Cuenod CA, et al. Quantitative dynamic contrast-enhanced MR imaging analysis of complex adnexal masses: a preliminary study. *Eur Radiol* 2012; **22**: 738–45. doi: <https://doi.org/10.1007/s00330-011-2329-6>
25. Bernardin L, Dilks P, Liyanage S, Miquel ME, Sahdev A, Rockall A. Effectiveness of semi-quantitative multiphase dynamic contrast-enhanced MRI as a predictor of malignancy in complex adnexal masses: radiological and pathological correlation. *Eur Radiol* 2012; **22**: 880–90. doi: <https://doi.org/10.1007/s00330-011-2331-z>
26. Hricak H, Chen M, Coakley FV, Kinkel K, Yu KK, Sica G, et al. Complex adnexal masses: detection and characterization with MR imaging--multivariate analysis. *Radiology* 2000; **214**: 39–46. doi: <https://doi.org/10.1148/radiology.214.1.r00ja3939>
27. Thomassin-Naggara I, Poncelet E, Jalaguer-Coudray A, Guerra A, Fournier LS, Stojanovic S, et al. Ovarian-Adnexal reporting data system magnetic resonance imaging (O-RADS MRI) score for risk stratification of Sonographically indeterminate adnexal masses. *JAMA Netw Open* 2020; **3**: e1919896. doi: <https://doi.org/10.1001/jamanetworkopen.2019.19896>
28. Pickhardt PJ, Hanson ME. Incidental adnexal masses detected at low-dose unenhanced CT in asymptomatic women age 50 and older: implications for clinical management and ovarian cancer screening. *Radiology* 2010; **257**: 144–50. doi: <https://doi.org/10.1148/radiol.10100511>
29. Reiter MJ, Schwoppe RB, Lisanti CJ. Algorithmic approach to solid adnexal masses and their mimics: utilization of anatomic relationships and imaging features to facilitate diagnosis. *Abdom Imaging* 2014; **39**: 1284–96. doi: <https://doi.org/10.1007/s00261-014-0163-4>
30. Alessandrino F, Dellafiore C, Eshja E, Alfano F, Ricci G, Cassani C. . . Differential Diagnosis for Female Pelvic Masses. In: Erondur OF, editor. *Medical Imaging in Clinical Practice* [Internet]. ; 2013. Available from: <http://www.intechopen.com/books/medical-imaging-in-clinical-practice/differential-diagnosis-for-female-pelvic-masses> [cited 2020 Apr 6].
31. Fritsch H. Clinical Anatomy of the Female Pelvis. In: Hamm B, Forstner R, eds. *MRI and CT of the Female Pelvis* [Internet. Berlin, Heidelberg: Springer; 2007. pp.]. 1–24. . . https://doi.org/10.1007/978-3-540-68212-7_1 [cited 2020 Apr 11].
32. Gangadhar K, Mahajan A, Sable N, Bhargava P. Magnetic resonance imaging of pelvic masses: a compartmental approach. *Semin Ultrasound CT MR* 2017; **38**: 213–30. doi: <https://doi.org/10.1053/j.sult.2016.11.004>
33. Hain KS, Pickhardt PJ, Lubner MG, Menias CO, Bhalla S. Presacral masses: multimodality imaging of a multidisciplinary space. *Radiographics* 2013; **33**: 1145–67. doi: <https://doi.org/10.1148/rg.334115171>
34. Szklaruk J, Tamm EP, Choi H, Varavithya V. Mr imaging of common and uncommon large pelvic masses. *Radiographics* 2003; **23**: 403–24. doi: <https://doi.org/10.1148/rg.232025089>
35. Dias JL, Veloso Gomes F, Lucas R, Cunha TM. The shading sign: is it exclusive of endometriomas? *Abdom Imaging* 2015; **40**: 2566–72. doi: <https://doi.org/10.1007/s00261-015-0465-1>
36. Outwater E, Schiebler ML, Owen RS, Schnall MD. Characterization of hemorrhagic adnexal lesions with MR imaging: blinded reader study. *Radiology* 1993; **186**: 489–94. doi: <https://doi.org/10.1148/radiology.186.2.8421756>
37. Corwin MT, Gerscovich EO, Lamba R, Wilson M, McGahan JP. Differentiation of ovarian endometriomas from hemorrhagic cysts at MR imaging: utility of the T2 dark spot sign. *Radiology* 2014; **271**: 126–32. doi: <https://doi.org/10.1148/radiol.13131394>
38. Balaban M, Idilman IS, Toprak H, Unal O, Ipek A, Kocakoc E. The utility of diffusion-weighted magnetic resonance imaging in differentiation of endometriomas from hemorrhagic ovarian cysts. *Clin Imaging* 2015; **39**: 830–3. doi: <https://doi.org/10.1016/j.clinimag.2015.05.003>
39. Bazot M, Bharwani N, Huchon C, Kinkel K, Cunha TM, Guerra A, et al. European Society of urogenital radiology (ESUR) guidelines: MR imaging of pelvic endometriosis. *Eur Radiol* 2017; **27**: 2765–75. doi: <https://doi.org/10.1007/s00330-016-4673-z>
40. Matias-Guiu X, Stewart CJR. Endometriosis-Associated ovarian neoplasia. *Pathology* 2018; **50**: 190–204. doi: <https://doi.org/10.1016/j.pathol.2017.10.006>
41. Mais V, Guerriero S, Ajossa S, Angiolucci M, Paoletti AM, Melis GB. Transvaginal ultrasonography in the diagnosis of cystic teratoma. *Obstet Gynecol* 1995; **85**: 48–52. doi: [https://doi.org/10.1016/0029-7844\(94\)00323-6](https://doi.org/10.1016/0029-7844(94)00323-6)
42. Outwater EK, Siegelman ES, Hunt JL. Ovarian teratomas: tumor types and imaging characteristics. *Radiographics* 2001; **21**: 475–90. doi: <https://doi.org/10.1148/radiographics.21.2.g01mr09475>
43. Comerci JT, Licciardi F, Bergh PA, Gregori C, Breen JL. Mature cystic teratoma: a clinicopathologic evaluation of 517 cases and review of the literature. *Obstet Gynecol* 1994; **84**: 22–8.

44. Wolff EF, Hughes M, Merino MJ, Reynolds JC, Davis JL, Cochran CS, et al. Expression of benign and malignant thyroid tissue in ovarian teratomas and the importance of multimodal management as illustrated by a BRAF-positive follicular variant of papillary thyroid cancer. *Thyroid* 2010; **20**: 981–7. doi: <https://doi.org/10.1089/thy.2009.0458>
45. Vasconcelos I, Darb-Esfahani S, Sehoul J. Serous and mucinous borderline ovarian tumours: differences in clinical presentation, high-risk histopathological features, and lethal recurrence rates. *BJOG* 2016; **123**: 498–508. doi: <https://doi.org/10.1111/1471-0528.13840>
46. Gotlieb WH, Soriano D, Achiron R, Zalel Y, Davidson B, Kopolovic J, et al. Ca 125 measurement and ultrasonography in borderline tumors of the ovary. *Am J Obstet Gynecol* 2000; **183**: 541–6. doi: <https://doi.org/10.1067/mob.2000.105940>
47. Pascual MA, Tresserra F, Grases PJ, Labastida R, Dexeus S. Borderline cystic tumors of the ovary: gray-scale and color Doppler sonographic findings. *J Clin Ultrasound* 2002; **30**: 76–82. doi: <https://doi.org/10.1002/jcu.10028>
48. Kato H, Kanematsu M, Furui T, Morishige K-ichiro, Hirose Y. Ovarian mucinous cystadenoma coexisting with benign Brenner tumor: MR imaging findings. *Abdom Imaging* 2013; **38**: 412–6. doi: <https://doi.org/10.1007/s00261-012-9887-1>
49. Pschera H, Wikström B. Extraovarian Brenner tumor coexisting with serous cystadenoma. Case report. *Gynecol Obstet Invest* 1991; **31**: 185–7. doi: <https://doi.org/10.1159/000293153>
50. Matsuki M, Kaji Y, Matsuo M, Kobashi Y. Struma ovarii: MRI findings. *Br J Radiol* 2000; **73**: 87–90. doi: <https://doi.org/10.1259/bjr.73.865.10721328>
51. Matsubayashi R, Matsuo Y, Doi J, Kudo S, Matsuguchi K, Sugimori H. Sclerosing stromal tumor of the ovary: radiologic findings. *Eur Radiol* 1999; **9**: 1335–8. doi: <https://doi.org/10.1007/s003300050844>
52. Testa AC, Ferrandina G, Timmerman D, Savelli L, Ludovisi M, Van Holsbeke C, et al. Imaging in gynecological disease (1): ultrasound features of metastases in the ovaries differ depending on the origin of the primary tumor. *Ultrasound Obstet Gynecol* 2007; **29**: 505–11. doi: <https://doi.org/10.1002/uog.4020>
53. Chechia A, Attia L, Temime RB, Makhoulouf T, Koubaa A. Incidence, clinical analysis, and management of ovarian fibromas and fibrothecomas. *Am J Obstet Gynecol* 2008; **199**: 473.e1–473.e4. doi: <https://doi.org/10.1016/j.ajog.2008.03.053>
54. Wu B, Peng WJ, Gu YJ, Cheng YF, Mao J. MRI diagnosis of ovarian fibrothecomas: tumour appearances and oestrogenic effect features. *Br J Radiol* 2014; **87**: 20130634 Available from. cited 2020 Apr 26. doi: <https://doi.org/10.1259/bjr.20130634>
55. Shinagare AB, Meylaerts LJ, Laury AR, Morteale KJ. MRI features of ovarian fibroma and fibrothecoma with histopathologic correlation. *AJR Am J Roentgenol* 2012; **198**: W296–303. doi: <https://doi.org/10.2214/AJR.11.7221>
56. Arleo EK, Schwartz PE, Hui P, McCarthy S. Review of leiomyoma variants. *AJR Am J Roentgenol* 2015; **205**: 912–21. doi: <https://doi.org/10.2214/AJR.14.13946>
57. Lameiras R, Matos AP, Luz C, Palas J, Ramalho M. Pelvic splenosis—a very unusual location. *BJR Case Rep* 2017; **3**: 20160026. doi: <https://doi.org/10.1259/bjrcr.20160026>
58. Chandramohan A, Thrower A, Shah N, Mohamed F. Radiological predictors of complete cytoreduction in 59 patients with peritoneal mesothelioma treated with cytoreductive surgery and hyperthermic intraperitoneal chemotherapy at a UK referral centre. *Br J Radiol* 2017; **90**: 20170361. doi: <https://doi.org/10.1259/bjr.20170361>
59. Tao K, Zeng X, Liu W, Wang S, Gao J, Shuai X, et al. Primary gastrointestinal stromal tumor mimicking as gynecologic mass: characteristics, management, and prognosis. *J Surg Res* 2020; **246**: 584–90. doi: <https://doi.org/10.1016/j.jss.2019.09.043>
60. Lee JH, Jeong YK, Park JK, Hwang JC. “Ovarian Vascular Pedicle” Sign Revealing Organ of Origin of a Pelvic Mass Lesion on Helical CT. *American Journal of Roentgenology* 2003; **181**: 131–7. doi: <https://doi.org/10.2214/ajr.181.1.1810131>
61. Cecil T, Bunni J, Mehta A. Appendicular tumours and psuedomyxoma peritonei: A ‘paradigm’ for peritoneal disease. Chapter 1, A Practical Guide to Peritoneal Malignancy. Boca Raton: CRC Press 2020;.
62. Nicoll K, Bartrop C, Walsh S, Foster R, Duncan G, Payne C, et al. Malignant transformation of tailgut cysts is significantly higher than previously reported: systematic review of cases in the literature. *Colorectal Dis* 2019; **21**: 869–78. doi: <https://doi.org/10.1111/codi.14628>
63. Abel ME, Nelson R, Prasad ML, Pearl RK, Orsay CP, Abcarian H. Parasacrocccygeal approach for the resection of retrorectal developmental cysts. *Dis Colon Rectum* 1985; **28**: 855–8. doi: <https://doi.org/10.1007/BF02555492>
64. Lee J, Paek J, Lee YH, Kong TW, Chang S-J, Ryu H-S. Pelvic Castleman’s disease presenting as an adnexal tumor in a young woman. *ObstetGynecol Sci* 2015; **58**: 323–6.
65. Surabhi VR, Garg N, Frumovitz M, Bhosale P, Prasad SR, Meis JM. Aggressive angiomyxomas: a comprehensive imaging review with clinical and histopathologic correlation. *AJR Am J Roentgenol* 2014; **202**: 1171–8. doi: <https://doi.org/10.2214/AJR.13.11668>