Published in final edited form as:

J Cardiovasc Electrophysiol. 2020 November ; 31(11): 2833–2835. doi:10.1111/jce.14742.

Is Otsu Thresholding the Answer to Reproducible Quantification of Left Atrial Scar from Late Gadolinium-Enhancement MRI?

Suvai Gunasekaran, PhD^a, Daniel Kim, PhD^{a,b}

^aDepartment of Radiology, Northwestern University Feinberg School of Medicine, Chicago, IL, United States

^bDepartment of Biomedical Engineering, Northwestern University, Evanston, IL, United States

Keywords

LA scar; LA fibrosis; late gadolinium-enhancement; MRI

Atrial fibrillation (AF) is the most common sustained arrhythmia in adults¹. Catheter ablation targeting the pulmonary veins and other atrial sites has emerged as the best intervention for restoring and maintaining sinus rhythm; however, 1-year success rates are only 60–70% or even lower for more persistent types of AF^{2,3}. These statistics highlight an unmet clinical need to avoid an unnecessary procedure for expected non-responders (30–40%) or consider a more extensive ablative or surgical approach. Potential predictors of AF recurrence post-ablation derived from standard clinical and imaging metrics have proven to be of limited use. Left atrial (LA) fibrosis is more promising, because fibrosis plays a central role in the development of an arrhythmogenic substrate for AF and may be a marker for more extensive disease that is less amenable to standard pulmonary vein isolation (PVI)^{4–8}. In fact, LA fibrosis assessed with 3D LA late gadolinium enhancement (LGE) cardiovascular magnetic resonance (CMR)⁹, pioneered by the Utah group, has shown promise for predicting AF recurrence post-ablation^{10–16}. However, LA fibrosis quantification has garnered skepticism from the field because it has not been independently reproduced. This lack of reproducibility stems from two fundamental methodologic deficiencies of (1) data acquisition limitations: inadequate spatial resolution (1.5 mm × 1.5 mm × 2.5 to 5 mm)^{9,17–21}, contrast-to-noise ratio, and lengthy scan time (11–15 min), and (2) unreliable image analysis techniques: no robust method for quantifying of fibrosis in the thin (~ 2 mm) LA wall. These deficiencies preclude widespread adoption of LA fibrosis quantification in clinical practice.

Currently, there is no standardized method for quantifying LA fibrosis or scar from LGE images. Existing analysis tools include standard deviation (SD) above the mean LA wall signal^{11,22}, SD above the blood pool (BP)^{23,24}, image intensity ratio (IIR)^{19,20}, and visual assessment^{9,25}. Regrettably, the performance of these threshold-based methods varies among

Please send correspondence to: Daniel Kim, PhD, Department of Radiology, Northwestern University, 737 N. Michigan Avenue Suite 1600, Chicago, IL 60611, daniel.kim3@northwestern.edu, O:312.926.1733, F:312.926.5991.

Disclosures: None

different groups and lacks independent evaluation. In this issue, Kamali et. al. conducted a study to identify which LA scar quantification methods is most reproducible. They do this by quantifying LA scar using different quantification methods from two post-ablation LGE CMR scans performed three months apart, while assuming that LA scarring remains the same between those two time points. The authors performed a retrospective study of patients who underwent AF radiofrequency ablation between May 2007 to August 2017, and found 45 patients who underwent consecutive post-ablation LGE CMR scans 3 months apart, first scan 120 ± 65 days post-ablation and the second scan 162 ± 98 days post first scan. They used a previously published pulse sequence²⁶ to acquire their LGE images that had a voxel size of $1.25 \text{ mm} \times 1.25 \times 2.5 \text{ mm}$ and was interpolated to $0.625 \text{ mm} \times 0.625 \text{ mm} \times 1.25 \text{ mm}$. Full LA segmentation from the LGE imaging was performed by trained raters using previously published methods^{26,27}. From the LA wall intensities, outliers were removed, and the histograms were normalized with intensity values ranging from 0 to 1.

To determine areas of scar, the researchers performed four different methods: (1) simple thresholding (from 60th to 90th percentile), (2) Otsu thresholding^{28,29}, (3) 3.3 SD above mean BP³⁰, and (4) IIR > 1.61 ²⁰. The authors studied two methods that used LA wall intensity (simple thresholding and Otsu thresholding) and two methods that used BP normalization (SD above BP and IIR). LA wall segmentation was registered between the first and second scan. After registration, a cut-off distance of 1.25 mm was used for comparing scar and healthy tissue voxels and Dice Similarity Coefficient (DSC) was computed to determine the level of agreement.

The authors determined that the normalized histograms for LA wall intensity was very closely correlated between the first and second scans, regardless of scanner's magnetic field strength. They calculated DSC for the different scan quantification methods and found the following results: Otsu thresholding = 71.3 ± 8.3 , 3.3 SD above BP mean = 57.8 ± 21.2 , and IIR $> 1.61 = 45.8 \pm 29.5$, indicating that Otsu thresholding was significantly better than BP based methods. In addition, for all scanner type based groups, DSC for the intensity distribution was mostly flat until the 70th percentile and then increased progressively for higher deciles. Based on their results, the authors concluded three main points. First, they determined that for RF ablation, atrial scar quantification is highly reproducible from one scan to the next over a three-month span. Second, they found the reproducibility of LGE CMR for thresholds less than the 70th percentile of the wall intensity distribution is no better than the reproducibility of a random distribution. And finally, LA wall intensity-based quantification methods are more reproducible than BP based methods.

While this study systematically compared four different methods for quantification of LA scar, there are several limitations that warrant further discussion. First, the small sample size of 45 patients limits statistical significance. Second, several patients were excluded because of poor image quality, which brings into question a bias in patient selection. There are numerous factors that may influence LA intensity based thresholding, including spatial resolution, appropriate inversion time to null the normal myocardium, and image artifacts induced by cardiac and respiratory motion. Variations in such factors are likely to influence LA scar quantification. Third, the two post-ablation CMR exams were approximately 3 months apart, and there is a possibility of remodeling in both scarred and normal wall

between those time points. Thus, readers ought to interpret reproducibility results with these caveats.

Is Otsu thresholding capable of producing a moderate DSC of 71.3 the answer to reproducible quantification of left atrial scar from LGE MRI? In the absence of standardization across the field, it is our opinion that the search must continue to establish more reproducible analysis methods, perhaps involving artificial intelligence. In conclusion, the study by Kamali et. al. is a strong reminder that LA scar or fibrosis quantification is an active area of research that warrants further investigation. If a robust LA fibrosis or scar quantification method is established, then it will be extremely beneficial for future studies to understand AF and its treatment to improve patient outcomes.

Funding:

National Institutes of Health: R01HL116895, R01HL138578, R21EB024315, R21AG055954, R01HL151079, R01HL151079; American Heart Association: 19IPLOI34760317

References

1. Anter E, Jessup M, Callans DJ. Atrial fibrillation and heart failure: treatment considerations for a dual epidemic. *Circulation*. 2009;119(18):2516–2525. [PubMed: 19433768]
2. Khan AR, Khan S, Sheikh MA, Khuder S, Grubb B, Moukarbel GV. Catheter ablation and antiarrhythmic drug therapy as first- or second-line therapy in the management of atrial fibrillation: systematic review and meta-analysis. *Circulation Arrhythmia and electrophysiology*. 2014;7(5):853–860. [PubMed: 25110162]
3. Wynn GJ, Das M, Bonnett LJ, Panikker S, Wong T, Gupta D. Efficacy of catheter ablation for persistent atrial fibrillation: a systematic review and meta-analysis of evidence from randomized and nonrandomized controlled trials. *Circulation Arrhythmia and electrophysiology*. 2014;7(5):841–852. [PubMed: 25132078]
4. Dzeshka MS, Lip GY, Snezhitskiy V, Shantsila E. Cardiac Fibrosis in Patients With Atrial Fibrillation: Mechanisms and Clinical Implications. *J Am Coll Cardiol*. 2015;66(8):943–959. [PubMed: 26293766]
5. Spragg D Left Atrial Fibrosis: Role in Atrial Fibrillation Pathophysiology and Treatment Outcomes. *Journal of atrial fibrillation*. 2013;5(6):810. [PubMed: 28496835]
6. Tan AY, Zimetbaum P. Atrial fibrillation and atrial fibrosis. *J Cardiovasc Pharmacol*. 2011;57(6):625–629. [PubMed: 21633248]
7. Burstein B, Nattel S. Atrial fibrosis: mechanisms and clinical relevance in atrial fibrillation. *J Am Coll Cardiol*. 2008;51(8):802–809. [PubMed: 18294563]
8. Goldberger JJ, Arora R, Green D, et al. Evaluating the Atrial Myopathy Underlying Atrial Fibrillation: Identifying the Arrhythmogenic and Thrombogenic Substrate. *Circulation*. 2015;132(4):278–291. [PubMed: 26216085]
9. Peters DC, Wylie JV, Hauser TH, et al. Detection of pulmonary vein and left atrial scar after catheter ablation with three-dimensional navigator-gated delayed enhancement MR imaging: initial experience. *Radiology*. 2007;243(3):690–695. [PubMed: 17517928]
10. Mahnkopf C, Badger TJ, Burgon NS, et al. Evaluation of the left atrial substrate in patients with lone atrial fibrillation using delayed-enhanced MRI: implications for disease progression and response to catheter ablation. *Heart rhythm : the official journal of the Heart Rhythm Society*. 2010;7(10):1475–1481.
11. Oakes RS, Badger TJ, Kholmovski EG, et al. Detection and quantification of left atrial structural remodeling with delayed-enhancement magnetic resonance imaging in patients with atrial fibrillation. *Circulation*. 2009;119(13):1758–1767. [PubMed: 19307477]

12. Akoum N, Daccarett M, McGann C, et al. Atrial fibrosis helps select the appropriate patient and strategy in catheter ablation of atrial fibrillation: a DE-MRI guided approach. *Journal of cardiovascular electrophysiology*. 2011;22(1):16–22. [PubMed: 20807271]
13. McGann C, Akoum N, Patel A, et al. Atrial fibrillation ablation outcome is predicted by left atrial remodeling on MRI. *Circ Arrhythm Electrophysiol*. 2014;7(1):23–30. [PubMed: 24363354]
14. Sohns C, Marrouche NF. Atrial fibrillation and cardiac fibrosis. *European heart journal*. 2019.
15. Marrouche NF, Wilber D, Hindricks G, et al. Association of atrial tissue fibrosis identified by delayed enhancement MRI and atrial fibrillation catheter ablation: the DECAAF study. *JAMA*. 2014;311(5):498–506. [PubMed: 24496537]
16. Khurram IM, Habibi M, Gucuk Ipek E, et al. Left Atrial LGE and Arrhythmia Recurrence Following Pulmonary Vein Isolation for Paroxysmal and Persistent AF. *JACC Cardiovascular imaging*. 2016;9(2):142–148. [PubMed: 26777218]
17. Chubb H, Karim R, Roujol S, et al. The reproducibility of late gadolinium enhancement cardiovascular magnetic resonance imaging of post-ablation atrial scar: a cross-over study. *Journal of cardiovascular magnetic resonance : official journal of the Society for Cardiovascular Magnetic Resonance*. 2018;20(1):21. [PubMed: 29554919]
18. Chelu MG, King JB, Kholmovski EG, et al. Atrial Fibrosis by Late Gadolinium Enhancement Magnetic Resonance Imaging and Catheter Ablation of Atrial Fibrillation: 5-Year Follow-Up Data. *J Am Heart Assoc*. 2018;7(23):e006313. [PubMed: 30511895]
19. Benito EM, Carlosena-Remirez A, Guasch E, et al. Left atrial fibrosis quantification by late gadolinium-enhanced magnetic resonance: a new method to standardize the thresholds for reproducibility. *Europace*. 2017;19(8):1272–1279. [PubMed: 27940935]
20. Khurram IM, Beinart R, Zipunnikov V, et al. Magnetic resonance image intensity ratio, a normalized measure to enable interpatient comparability of left atrial fibrosis. *Heart rhythm : the official journal of the Heart Rhythm Society*. 2014;11(1):85–92.
21. Gunasekaran S Accelerated 3D Left Atrial Late Gadolinium-Enhancement in Patients with Atrial Fibrillation at 1.5 Tesla: Technical Development. *Radiol Cardiothorac Imaging*. (In Press).
22. Cochet H, Mouries A, Nivet H, et al. Age, atrial fibrillation, and structural heart disease are the main determinants of left atrial fibrosis detected by delayed-enhanced magnetic resonance imaging in a general cardiology population. *J Cardiovasc Electrophysiol*. 2015;26(5):484–492. [PubMed: 25727248]
23. Harrison JL, Sohns C, Linton NW, et al. Repeat left atrial catheter ablation: cardiac magnetic resonance prediction of endocardial voltage and gaps in ablation lesion sets. *Circ Arrhythm Electrophysiol*. 2015;8(2):270–278. [PubMed: 25593109]
24. Malcolm-Lawes LC, Juli C, Karim R, et al. Automated analysis of atrial late gadolinium enhancement imaging that correlates with endocardial voltage and clinical outcomes: a 2-center study. *Heart Rhythm*. 2013;10(8):1184–1191. [PubMed: 23685170]
25. Peters DC, Wylie JV, Hauser TH, et al. Recurrence of atrial fibrillation correlates with the extent of post-procedural late gadolinium enhancement: a pilot study. *JACC Cardiovasc Imaging*. 2009;2(3):308–316. [PubMed: 19356576]
26. Parmar BR, Jarrett TR, Burgon NS, et al. Comparison of left atrial area marked ablated in electroanatomical maps with scar in MRI. *J Cardiovasc Electrophysiol*. 2014;25(5):457–463. [PubMed: 24383404]
27. Parmar BR, Jarrett TR, Kholmovski EG, et al. Poor scar formation after ablation is associated with atrial fibrillation recurrence. *J Interv Card Electrophysiol*. 2015;44(3):247–256. [PubMed: 26455362]
28. Tao Q, Milles J, Zeppenfeld K, et al. Automated segmentation of myocardial scar in late enhancement MRI using combined intensity and spatial information. *Magn Reson Med*. 2010;64(2):586–594. [PubMed: 20665801]
29. Otsu N Threshold Selection Method from Gray-Level Histograms. *Ieee T Syst Man Cyb*. 1979;9(1):62–66.
30. Harrison JL, Jensen HK, Peel SA, et al. Cardiac magnetic resonance and electroanatomical mapping of acute and chronic atrial ablation injury: a histological validation study. *Eur Heart J*. 2014;35(22):1486–1495. [PubMed: 24419806]