Research

Cite this article: Herraiz JL, Ribé J, Botella H, Martínez-Pérez C, Ferrón HG. 2020 Use of nursery areas by the extinct megatooth shark *Otodus megalodon* (Chondrichthyes: Lamniformes). *Biol. Lett.* **16**: 20200746. <http://dx.doi.org/10.1098/rsbl.2020.0746>

Received: 19 October 2020

Accepted: 29 October 2020

Subject Areas:

palaeontology, evolution

Keywords:

Otodus megalodon, palaeoecology, sharks, nurseries, Miocene

Authors for correspondence:

Carlos Martínez-Pérez

e-mail: carlos.martinez-perez@uv.es

Humberto G. Ferrón

e-mail: humberto.ferron@bristol.ac.uk

Electronic supplementary material is available online at <https://dx.doi.org/10.6084/m9.figshare.c.5199296>.

Palaeontology

Use of nursery areas by the extinct megatooth shark *Otodus megalodon* (Chondrichthyes: Lamniformes)

Jose L. Herraiz¹, Joan Ribé², Héctor Botella¹, Carlos Martínez-Pérez^{1,3} and Humberto G. Ferrón^{1,3}

¹Institut Cavanilles de Biodiversitat i Biologia Evolutiva, Universitat de València, C/ Catedràtic José Beltrán Martínez, 2, 46980 Paterna, Spain

²Museu del Cau del Tauró de l'Arboç, Carrer de la Muralla, 23, 43720 L'Arboç, Spain

³School of Earth Sciences, University of Bristol, 24 Tyndall Avenue, Bristol BS8 1RJ, UK

CM-P, 0000-0001-7795-5997; HGF, 0000-0003-2254-8424

Nursery areas are fundamental for the success of many marine species, particularly for large, slow-growing taxa with low fecundity and high age of maturity. Here, we examine the population size-class structure of the extinct gigantic shark *Otodus megalodon* in a newly described middle Miocene locality from Northeastern Spain, as well as in eight previously known formations (Temblor, Calvert, Pisco, Gatún, Chucunague, Bahía Inglesa, Yorktown and Bone Valley). In all cases, body lengths of all individuals were inferred from dental parameters and the size-class structure was estimated from kernel probability density functions and Gaussian mixture models. Our analyses support the presence of five potential nurseries ranging from the Langhian (middle Miocene) to the Zanclean (Pliocene), with higher densities of individuals with estimated body lengths within the typical range of neonates and young juveniles. These results reveal, for the first time, that nursery areas were commonly used by *O. megalodon* over large temporal and spatial scales, reducing early mortality and playing a key role in maintaining viable adult populations. Ultimately, the presumed reliance of *O. megalodon* on the presence of suitable nursery grounds might have also been determinant in the demise of this iconic top predatory shark.

1. Introduction

Otodus megalodon is the largest macropredatory shark to have ever lived, with body length estimates of up to 15–18 m for the largest adult individuals [1–4]. This species inhabited the warm and temperate waters of all major ocean basins, spanning a range of almost 20 Myr, from the early Miocene to the Pliocene [5–7]. Most of the studies assessing the causes of the cosmopolitan distribution, the evolution of gigantism, and the extinction of otodontid sharks have focused on the impact of climatic factors and/or the abundance and migration patterns of their potential prey [1,8–11], competition with other macropredatory taxa [7,12], the availability of suitable habitats [13] and the presence of regional endothermy [6,14–16]. However, much less attention has been paid to important aspects of their reproductive biology [17].

Nursery areas are crucial for several marine groups, as they improve the probability of survival of juveniles and allow a greater persistence and recruitment to the adult populations [18,19]. Shark nurseries are usually located in geographically discrete zones with high primary productivity, being defined by a comparatively high abundance of juveniles that tend to remain or return for extended periods of time [20,21]. The use of these areas has been

documented in a number of living species [22], and may vary depending on the habitat (coastal, pelagic or deep-water nurseries) [22], the temporal patterns of occupation (permanent or seasonal) [23], or the number of taxa involved [24]. Some authors have also distinguished between primary nurseries, where gravid females give birth and neonates spend the first stages of their life, and secondary nurseries, with slightly older but still immature individuals [25].

Despite the utility of more recently proposed criteria in defining shark nursery areas, these are usually difficult to apply to extinct species and the recognition of nurseries in the fossil record is therefore challenging (see Heupel *et al.* [20,22] for a detailed discussion on the evolution of the concept for modern taxa). The existence of palaeonurseries has been recently supported by the presence of either abundant fossilized egg cases [26], biased representations of juveniles [8,27–35], or both [36,37]. Accordingly, the late Miocene Gatún Formation (Panama) has been proposed as a possible nursery area for *O. megalodon* based on the high proportion of juvenile individuals [27]. A second possible nursery area for *O. megalodon* could have been present in Borneo Island [38], but this observation, based only on three isolated juvenile teeth, is questionable. Although relevant, these sporadic findings are insufficient to evaluate the prevalence and the impact of such a reproductive strategy in the evolutionary history of *O. megalodon* and, as consequence, a more comprehensive body of evidence on nursery use patterns is required to reliably assess these questions. This is especially evident when considering the wide temporal and palaeobiogeographic distribution of this species.

In the present study, a new potential nursery of *O. megalodon* is described from the Langhian (middle Miocene) of Spain. Moreover, we analyse a comprehensive database with records of this species revealing the plausible existence of multiple additional nursery areas, widely distributed in time and space. The broader implications of these findings are then contextualized in the light of some previous hypotheses about the success and later demise of this iconic extinct top predator.

2. Material and methods

The *O. megalodon* teeth studied first-hand in the present work were collected from Langhian outcrops (late middle Miocene) in the Reverté and Vidal quarries, which are situated in the southwestern part of the Vallès-Penedès Basin, close to the lowest stratigraphic layers of the El Camp de Tarragona Basin (see electronic supplementary material, figure S1a and supplementary methods for a detailed description of the geological context of the area). These quarries are located in the municipality of Castellet i La Gornal, in the province of Tarragona (Northeastern Spain) (electronic supplementary material, figure S1b). A total of 32 teeth, deposited in the Museu del Cau del Tauró de l'Arboç (Tarragona, Spain), have been collected over the past 20 years. Of these, 25 were considered in the present study owing to their excellent preservation (figure 1; electronic supplementary material, table S1). Their assignment to *O. megalodon* is well supported based on several morphological characteristics [27,39] (see electronic supplementary material, supplementary methods).

Teeth were assigned into jaw positions based on comparison with images in Pimiento & Balk [2], as well as an artificial tooth set figured in Applegate & Espinosa-Arrubarrena [40]. The population size-class structure of *O. megalodon* from the Reverté and

Vidal quarries was inferred from dental parameters. According to the methodology described in Pimiento *et al.* [27], crown height was used to calculate total body lengths from the position-specific regressions established in the great white shark (*Carcharodon carcharias*) by Shimada [41] (electronic supplementary material, table S2 and supplementary methods).

The population size-class structure of *O. megalodon* was analysed by estimating kernel probability densities of the inferred body lengths, with violin plots generated using the package 'ggplot2' [42] implemented in R [43]. Unimodality was checked by computing Hartigans' dip statistic and its statistical significance in the R package 'dipTest' [44]. Additionally, different modal parameters (mode(s), bimodality ratio and bimodality coefficient) were calculated using the R package 'modes' [45]. The bimodality ratio captures the difference in magnitude between the two identified peaks as the ratio between the amplitudes of the right and left peaks, respectively. The bimodality coefficient can vary between 0 and 1, where a value greater than 0.56 suggests the existence of bimodality and the maximum value of 1 indicates a fully bimodal distribution. The population size-class structure of *O. megalodon* was also evaluated with Gaussian mixture models using the R package 'Mclust' [46]. The optimal model and number of clusters were determined by the maximum Bayesian information criteria (BIC).

The population size-class structure of *O. megalodon* was assessed in eight additional formations covering a wide geographical area, including sites from the Atlantic, Caribbean and Pacific basins, which range from the middle Miocene to the Pliocene (i.e. the Temblor, Calvert, Yorktown and Bone Valley formations of the USA, the Pisco Formation of Peru, the Gatún and Chucunaque formations of Panama, and the Bahía Inglesa Formation of Chile; figure 2). We analysed the dataset compiled in Pimiento & Balk [2], selecting only the formations therein containing more than 20 tooth records (see electronic supplementary material). Analyses were first performed at formation level because this provided higher sample sizes and more statistically robust results. To obtain higher temporal and spatial resolutions, all the analyses were then repeated at locality level when more than 15 teeth were available (i.e. Shark Tooth Hill, Parkers Creek, Las Lomas, Lago Bayano, Mina Fosforita and Lee Creek Mine from Temblor, Calvert, Pisco, Chucunaque, Bahía Inglesa and Yorktown formations, respectively). All the R scripts are available in the electronic supplementary material.

3. Results

Body length estimates derived from our teeth sample ranged from 2.61 m (MCTA04205, upper anterior tooth) to 13.40 m (MCTA04212, lower lateral tooth) (electronic supplementary material, table S1). Kernel probability density analysis and Hartigans' dip test (HDT) indicate that the population size-class structure of *O. megalodon* from Reverté and Vidal quarries shows a bimodal distribution, as suggested by estimated density graphs (figure 2), the high bimodality coefficient (0.73) and the comparatively low unimodality test *p*-value (0.22; although no statistical significance at 0.05 level is found). Higher densities of individuals are estimated within small and large body sizes (modes equal to 5.45 and 10.52 m, respectively) (table 1). Similarly, the Gaussian mixture model analysis (GMM) supports bimodality within the optimal model, with modes equal to 5.28 and 11.10 m, respectively (table 1).

The analysis of additional formations reveals a variety of size-class structures that can be grouped into two major types (figure 2 and table 1). The Temblor, Pisco, Bahía Inglesa and Yorktown formations show maximum density peaks in

Figure 1. *Otodus megalodon* teeth collection from the Museu del Cau del Tauró de l'Arboç (MCTA, Tarragona, Spain), figured in lingual view.

occurrences of large body sizes (modes from the HDT equal to 11.07, 11.05, 11.95 and 9.82 m, respectively; modes from the GMM equal to 11.68, 11.40, 12.47 and 11.04 m, respectively). Unimodal distributions are well supported by comparatively low bimodality coefficients (below 0.45), high unimodality test p -values (above 0.87) and BIC model selection. By contrast, the Calvert, Gatún, Chucunaque and Bone Valley formations show maximum density peaks within smaller body sizes (modes from the HDT equal to 4.62, 6.56, 6.25, 5.62 m, respectively; modes from the GMM equal to 4.29, 6.66, 7.69 and 6.71 m, respectively). Bimodality

coefficients are comparatively higher (above 0.53) and BIC values provide similar support for both unimodal and bimodal models in some instances. A second smaller peak can be recognized in the Gatún Formation (in both the HDT and the GMM), corresponding to large body sizes (modes equal to 17.15 and 14.79 m, respectively), but is negligible when compared to the main left peak (bimodality ratio equals to 0.16). A second smaller peak is also detected in the Calvert Formation (only in the GMM), corresponding to comparatively small body sizes (mode equal to 7.54 m). Despite this, unimodality is well supported in all these formations based on

Figure 2. Size-class structure of *Otodus megalodon* populations from all the formations and localities analysed in this study ($n > 20$ tooth records), including information about their temporal and spatial distribution. Size-class categorization as neonates, juveniles and adults follows criteria from Gottfried *et al.* [1]. Palaeogeographic maps from Scotese [47]. Stages: Aq, Aquitanian; Bu, Burdigalian; La, Langhian; Me, Messinian; Pi, Piacenzian; Se, Serravallian; To, Tortonian; Za, Zanclean. Localities: 1, Reverté and Vidal quarries; 2, Shark tooth Hill; 3, Parkers Creek and Scientists Cliff; 4, Cerro La Bruja and Cerro Colorado; 5, Yesera de Amara; 6, Aguada de Lomas; 7, Cerro Los Quesos; 8, Montemar; 9, Texaco; 10, Alborada; 11, Las Lomas and San Judas; 12, Isla Payardi; 13, Lago Bayano; 14, Mina Fosforita; 15, Lee Creek Mine; 16, Achan Mine, Chicora Mine, Fort Green Mine, Four Corners Mine, Kingsford Mine, North Palmetto Mine, Palmetto Mine, and Payne Creek Mine (Tro Quarry).

the unimodality test p -values (above 0.6). Results from the analyses performed at locality level are largely equivalent to those of the formations to which they belong (electronic supplementary material, figure S2 and table S3).

4. Discussion

Our results reveal, for the first time, that nursery areas were commonly used by *O. megalodon* over large temporal and spatial scales. Five of the nine formations examined in this study (i.e. Reverté and Vidal quarries, Calvert, Gatún, Chucunaque and Bone Valley formations) show size class structures that are consistent with the presence of nurseries, with the main size modes placed among body lengths typical of neonates and young juveniles (neonates approx. 4 m and juveniles less than 10.5 m [1]; figure 2 and table 1). Among these, the Calvert, Gatún, Chucunaque and Bone Valley formations show unimodal distributions that are comparable to the age distributions present in most modern shark nursery areas ([48–55] among others). The population size-class structures estimated from the Reverté and Vidal quarries have bimodal distributions, with a second-density peak among typical adult body sizes (greater than 10.5 m [1]) that might represent gravid females. This distributional pattern is present in some living shark populations occupying areas that serve as primary nurseries [56–58]; however, a change in the use of this locality over time could also explain this size distribution, with juvenile and adult populations occupying this region at different times. In any case, the high proportion of neonates and

juveniles supports the hypothesis that this area acted as a nursery ground for *O. megalodon* at some point during the Langhian. This interpretation is further supported by the fact that, during this interval, the Vallès-Penedès Basin was part of a shallow protected bay [59,60] with several taxa that could act as potential prey (i.e. marine mammals and a number fish species) and high primary productivity (as highlighted by the numerous fossil remains found in association with the teeth; electronic supplementary material, table S4). The remaining four formations (i.e. Temblor, Pisco, Bahía Inglesa and Yorktown) demonstrate size-class structures typical of populations dominated by adults (greater than 10.5 m [1]; figure 2 and table 1), suggesting that these regions might correspond to feeding or mating areas [23].

Overall, our analysis based on kernel probability density and Gaussian mixture model also supports the interpretation of the Gatún Formation as a nursery area [27] and reveals the presence of at least four additional nurseries for *O. megalodon* (i.e. Reverté and Vidal quarries (considered together as a single area), Calvert, Chucunaque and Bone Valley formations). We further validate the results from the size-class structure analysis with a more integrative approach including stratigraphic, sedimentological and palaeontological evidence (electronic supplementary material, supplementary information), which is a fundamental procedure for the reliable recognition of new nursery areas [34]. Moreover, these findings are reinforced by analyses performed at locality level, with higher temporal and spatial resolutions (electronic supplementary material, figure S2 and table S3).

Table 1. Modal parameters, Hartigans' dip test and Gaussian mixture models analysis results for estimated size-class density curves in all analysed formations. In Gaussian mixture models: E, equal variance; V, variable variance; 1, unimodal; 2, bimodal.

modal parameters and Hartigans' dip test					
	Mode 1 (m)	Mode 2 (m)	bimodality ratio	bimodality coefficient	unimodality test (<i>p</i> -value)
this study	5.45	10.52	0.92	0.73	0.22
Temblor	11.07	4.86	7.66	0.29	0.91
Calvert	4.62	—	—	0.54	0.60
Pisco	11.05	—	—	0.45	0.87
Gatún	6.56	17.15	0.16	0.60	0.98
Chucunaque	6.25	—	—	0.56	0.87
Bahía Inglesa	11.95	—	—	0.35	0.92
Yorktown	9.82	—	—	0.43	0.98
Bone Valley	5.62	—	—	0.53	0.70
Gaussian mixture models					
	Mode 1 (m)	Mode 2 (m)	Top 3 models based on the BIC values		
this study	5.28	11.10	E2 (−133.98)	E1 (−136.34)	V1 (−136.34)
Temblor	11.68	—	E1 (−187.01)	V1 (−187.01)	E2 (−194.20)
Calvert	4.29	7.54	V2 (−224.78)	E1 (−225.83)	V1 (−225.83)
Pisco	11.40	—	E1 (−133.44)	V1 (−133.44)	V2 (−139.79)
Gatún	6.66	14.79	E2 (−222.39)	V2 (−225.64)	E1 (−227.11)
Chucunaque	7.69	—	E1 (−125.57)	V1 (−125.57)	E2 (−129.23)
Bahía Inglesa	12.47	—	E1 (−283.76)	V1 (−283.76)	V2 (−292.02)
Yorktown	11.04	—	E1 (−1028.80)	V1 (−1028.80)	V2 (−1029.31)
Bone Valley	6.71	—	E1 (−186.33)	V1 (−186.33)	V2 (−186.99)

This high proportion of potential nursery areas among the analysed localities may not be surprising as nurseries are zones with a high abundance of sharks and, as consequence, huge numbers of teeth can be shed, increasing the chances of subsequent fossil discoveries.

The use of nursery areas is likely to play a key role in the evolutionary history of some shark species [20,61]. This seems to be particularly true for large, slow-growing taxa with low fecundity and high age of maturity, which can benefit from this strategy of maximizing juvenile survival [22]. Evidence from analysis of vertebral growth rings suggests that the slowing or cessation of somatic growth in *O. megalodon* occurred around 25 years of age, denoting an extremely delayed sexual maturity [14]. It therefore seems plausible that the use of nursery areas could have been essential for *O. megalodon*, in order to reduce neonate and juvenile mortality and to provide maximum recruitment, thus maintaining viable populations on a long-term temporal scale. The widespread presence of *O. megalodon* nurseries supports this idea, demonstrating that this species benefited from this strategy at least from the Langhian (middle Miocene) to the Zanclean (Pliocene) in geographically separate regions (figure 2; electronic supplementary material, figure S2). The presumed reliance of *O. megalodon* on the presence of suitable nursery grounds should be considered in future work as a potential disadvantage in light of the increasing competition

of juveniles with the great white shark [6,7] and the drastic reduction in coastal habitats in the Pliocene [13]. More broadly, this work advances our understanding of the impact of reproductive strategies on the evolution of gigantic marine top predators and sheds light on the underlying factors of their evolutionary dynamics.

Ethics. This study does not concern live animals and therefore was not subject to an ethics assessment.

Data accessibility. All raw data and code are included as electronic supplementary material.

Authors' contributions. H.G.F. and C.M.-P. conceived and coordinated the study. C.M.-P., H.G.F., H.B. and J.L.H. participated in the design of the study. J.R. collected and provided the specimens. J.L.H. collected the data. H.G.F. performed statistical analyses. J.L.H. and H.G.F. wrote the paper with input from all co-authors. All authors approved the final version of the manuscript and agree to be held accountable for the content therein.

Competing interests. We declare we have no competing interests.

Funding. H.G.F. is a recipient of a Marie Skłodowska-Curie Individual Fellowship (H2020-MSCA-IF-2018-839636).

Acknowledgements. The authors thank Manuel Gimeno Peinado (València), Lluís Prieto and María Sánchez (Barcelona) for their help collecting some of the specimens, Dr John Cunningham (University of Bristol, UK) and Dr Imran Rahman (Oxford University Museum of Natural History, UK) for proofreading the article, and Dr Alberto Collareta, Dr Jürgen Kriwet and one anonymous reviewer for their helpful comments on the manuscript.

- Gottfried MD, Compagno LJV, Bowman SC. 1996 Size and skeletal anatomy of the giant 'megatooth' shark *Carcharodon megalodon*. In *Great white sharks: the biology of Carcharodon carcharias* (eds AP Klimley, DG Ainley), pp. 55–66. San Diego, CA: Academy Press.
- Pimiento C, Balk MA. 2015 Body-size trends of the extinct giant shark *Carcharocles megalodon*: a deep-time perspective on marine apex predators. *Paleobiology* **41**, 479–490. (doi:10.1017/pab.2015.16)
- Shimada K. 2019 The size of the megatooth shark, *Otodus megalodon* (Lamniformes: Otodontidae), revisited. *Hist. Biol.* **20**, 1–8. (doi:10.1080/08912963.2019.1666840)
- Cooper JA, Pimiento C, Ferrón HG, Benton MJ. 2020 Body dimensions of the extinct giant shark *Otodus megalodon*: a 2D reconstruction. *Sci. Rep.* **10**, 1–9. (doi:10.1038/s41598-020-71387-y)
- Pimiento C, Clements CF. 2014 When did *Carcharocles megalodon* become extinct? A new analysis of the fossil record. *PLoS ONE* **9**, e111086. (doi:10.1371/journal.pone.0111086)
- Pimiento C, MacFadden BJ, Clements CF, Varela S, Jaramillo C, Velez-Juarbe J, Silliman BR. 2016 Geographical distribution patterns of *Carcharocles megalodon* over time reveal clues about extinction mechanisms. *J. Biogeogr.* **43**, 1645–1655. (doi:10.1111/jbi.12754)
- Boessenecker RW, Ehret DJ, Long DJ, Churchill M, Martin E, Boessenecker SJ. 2019 The Early Pliocene extinction of the mega-toothed shark *Otodus megalodon*: a view from the eastern North Pacific. *PeerJ* **7**, e6088. (doi:10.7717/peerj.6088)
- Purdy RW. 1996 Paleocology of fossil white sharks. In *Great white sharks: the biology of Carcharodon carcharias* (eds AP Klimley, DG Ainley), pp. 67–80. San Diego, CA: Academy Press.
- Dickson KA, Graham JB. 2004 Evolution and consequences of endothermy in fishes. *Physiol. Biochem. Zool.* **77**, 998–1018. (doi:10.1086/423743)
- Diedrich CG. 2013 Evolution of white and megatooth sharks, and evidence for early predation on seals, sirenians, and whales. *Nat. Sci.* **5**, 1203–1218. (doi:10.4236/ns.2013.511148)
- Collareta A, Lambert O, Landini W, Di Celma C, Malinverno E, Varas-Malca R, Urbina M, Bianucci G. 2017 Did the giant extinct shark *Carcharocles megalodon* target small prey? Bite marks on marine mammal remains from the late Miocene of Peru. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **469**, 84–91. (doi:10.1016/j.palaeo.2017.01.001)
- McLennan L. 2018 Tooth wear, microwear and diet in Elasmobranchs. Doctoral dissertation, Department of Geology, University of Leicester, Leicester, UK.
- Pimiento C, Griffin JN, Clements CF, Silvestro D, Varela S, Uhen MD, Jaramillo C. 2017 The Pliocene marine megafauna extinction and its impact on functional diversity. *Nat. Ecol. Evol.* **1**, 1100–1106. (doi:10.1038/s41559-017-0223-6)
- Ehret DJ. 2010 Paleobiology and taxonomy of extinct lamnid and otodontid sharks (Chondrichthyes, Elasmobranchii, Lamniformes). Doctoral Dissertation, University of Florida, Gainesville, FL.
- Ferrón HG. 2017 Regional endothermy as a trigger for gigantism in some extinct macropredatory sharks. *PLoS ONE* **12**, e0185185. (doi:10.1371/journal.pone.0185185)
- Ferrón HG, Martínez-Pérez C, Botella H. 2018 The evolution of gigantism in active marine predators. *Hist. Biol.* **30**, 712–716. (doi:10.1080/08912963.2017.1319829)
- Shimada K, Becker MA, Griffiths ML. 2020 Body, jaw, and dentition lengths of macrophagous lamniform sharks, and body size evolution in Lamniformes with special reference to 'off-the-scale' gigantism of the megatooth shark, *Otodus megalodon*. *Hist. Biol.* (doi:10.1080/08912963.2020.1812598)
- Williams H, Schaap AH. 1992 Preliminary results of a study into the incidental mortality of sharks in gill-nets in two Tasmanian shark nursery areas. *Aust. J. Mar. Freshwater Res.* **43**, 237–250.
- Beck MW et al. 2001 The identification, conservation, and management of estuarine and marine nurseries for fish and invertebrates: a better understanding of the habitats that serve as nurseries for marine species and the factors that create site-specific variability in nursery quality will improve conservation and management of these areas. *Bioscience* **51**, 633–641. (doi:10.1641/0006-3568(2001)051[0633:TICAMO]2.0.CO;2)
- Heupel MR, Carlson JK, Simpfendorfer CA. 2007 Shark nursery areas: concepts, definition, characterization and assumptions. *Mar. Ecol. Prog. Ser.* **337**, 287–297. (doi:10.3354/meps337287)
- Heithaus MR. 2007 Nursery areas as essential shark habitats: a theoretical perspective. In *Shark nursery grounds of the Gulf of Mexico and east coast waters of the United States*. *Am. Fish. Soc. Symp.* **50** (eds CT McCandless, HL Jr Pratt, NE Kohler), pp. 3–13. Bethesda, MD: American Fisheries Society.
- Heupel MR, Kanno S, Martins AP, Simpfendorfer CA. 2019 Advances in understanding the roles and benefits of nursery areas for elasmobranch populations. *Mar. Freshwater Res.* **70**, 897–907. (doi:10.1071/MF18081)
- Castro JI. 1993 The shark nursery of Bulls Bay, South Carolina, with a review of the shark nurseries of the southeastern coast of the United States. *Environ. Biol. Fishes* **38**, 37–48.
- Simpfendorfer CA, Milward NE. 1993 Utilisation of a tropical bay as a nursery area by sharks of the families Carcharhinidae and Sphyrnidae. *Environ. Biol. Fishes* **37**, 337–345.
- Bass AJ. 1978 Problems in studies of sharks in the southwest Indian Ocean. In *Sensory biology of sharks, skates and rays* (eds ES Hodgson, RF Matheson), pp. 545–594. Arlington, VA: Office of Naval Research, Department of the Navy.
- Fischer J, Reich M. 2013 On the Early Cretaceous chondrichthyan egg capsule *Palaeoxyris jugleri*. In *6th Int. meeting on mesozoic fishes – diversification and diversity patterns* (eds C Schwartz, J Kriwet), p. 24. Vienna, Austria: Verlag Dr. Friedrich Pfeil.
- Pimiento C, Ehret DJ, MacFadden BJ, Hubbell G. 2010 Ancient nursery area for the extinct giant shark *Megalodon* from the Miocene of Panama. *PLoS ONE* **5**, e10552. (doi:10.1371/journal.pone.0010552)
- Purdy RW. 1998 Chondrichthyan fishes from the Paleocene of South Carolina. *T. Am. Phil. Soc.* **88**, 122–146.
- Grandstaff B, Parris D, Gallagher W. 2006 Implications of *Scapanorhynchus* tooth size in a Campanian (Cretaceous) estuary at Ellisdale. *NJ. In. J. Vertebr. Paleontol.* **26**, 69A. (doi:10.1080/02724634.2006.10010069)
- Everhart MJ. 2007 New stratigraphic records (Albian-Campanian) of *Rhinobatos* sp. (Chondrichthyes; Rajiformes) from the Cretaceous of Kansas. *Trans. Kans. Acad. Sci.* **110**, 225–236. (doi:10.1660/0022-8443(2007)110[225:NSRAOR]2.0.CO;2)
- Padilla A. 2008 Faunal description and paleobiology of Eocene sharks from Banks Island in the Canadian Arctic. Master's Thesis, University of Colorado, Boulder, CO.
- Kriwet J, Nunn EV, Klug S. 2009 Neoselachians (Chondrichthyes, Elasmobranchii) from the Lower and lower Upper Cretaceous of north-eastern Spain. *Zoo. J. Linn. Soc.* **155**, 316–347. (doi:10.1111/j.1096-3642.2008.00439.x)
- Cuny G, Martin JE, Sarr R. 2012 A neoselachian shark fauna from the Late Cretaceous of Senegal. *Cretac. Res.* **34**, 107–115. (doi:10.1016/j.cretres.2011.10.008)
- Landini W, Collareta A, Pesci F, Di Celma C, Urbina M, Bianucci G. 2017 A secondary nursery area for the copper shark *Carcharhinus brachyurus* from the late Miocene of Peru. *J. S. Am. Earth Sci.* **78**, 164–174. (doi:10.1016/j.jsames.2017.07.003)
- Villafañá JA, Hernandez S, Alvarado A, Shimada K, Pimiento C, Rivadeneira MM, Kriwet J. 2020 First evidence of a palaeo-nursery area of the great white shark. *Sci. Rep.* **10**, 1–8. (doi:10.1038/s41598-020-65101-1)
- Fischer J, Voigt S, Schneider JW, Buchwitz M, Voigt S. 2011 A selachian freshwater fauna from the Triassic of Kyrgyzstan and its implication for Mesozoic shark nurseries. *J. Vertebr. Paleontol.* **31**, 937–953. (doi:10.1080/02724634.2011.601729)
- Sallan LC, Coates MJ. 2014 The long-rostrum elasmobranch *Bandringa* Zangerl, 1969, and taphonomy within a Carboniferous shark nursery. *J. Vertebr. Paleontol.* **34**, 22–33. (doi:10.1080/02724634.2013.782875)

38. Razak H, Kocsis L. 2018 Late Miocene *Otodus* (*Megaelachius*) *megalodon* from Brunei Darussalam: body length estimation and habitat reconstruction. *Neues Jahrb. Geol. Paläontol. Abh.* **288**, 299–306. (doi:10.1127/njgpa/2018/0743)
39. Cappetta H. 2012 *Handbook of paleoichthyology—chondrichthyes—mesozoic and cenozoic elasmobranchii: teeth*. Stuttgart, NY: Verlag Dr. Friedrich Pfeil.
40. Applegate SP, Espinosa-Arrubarrena L. 1996 The fossil history of *Carcharodon* and its possible ancestor, *Cretolamna*: a study in tooth identification. In *Great white sharks: the biology of carcharodon carcharias* (eds AP Klimley, DG Ainley), pp. 19–36. San Diego, CA: Academy Press.
41. Shimada K. 2002 The relationship between the tooth size and total body length in the white shark. *J. Fossil Res.* **35**, 28–33.
42. Wickham H. 2016 *Ggplot2: elegant graphics for data analysis*. New York, NY: Springer-Verlag.
43. Team RC. 2019 *R: A language and environment for statistical computing*. Vienna, Austria: R foundation for statistical computing.
44. Maechler M. 2013 *Package 'dipTest' R package version 0.75–5*. Vienna, Austria: R Foundation for Statistical Computing.
45. Deevi S. 2016 *modes: Find the Modes and Assess the Modality of Complex and Mixture Distributions, Especially with Big Datasets*. R package version 0.7.0.
46. Scrucca L, Fop M, Murphy TB, Raftery AE. 2016 mclust 5: clustering, classification and density estimation using Gaussian finite mixture models. *R J.* **8**, 289–317.
47. Scotese CR. 2014 *Atlas of Neogene Paleogeographic Maps (Mollweide Projection)*. Maps 1–7. Volume 1, The Cenozoic, PALEOMAP Atlas for ArcGIS. Evanston, IL: PALEOMAP Project.
48. Carlson JK. 1999 Occurrence of neonate and juvenile sandbar sharks, *Carcharhinus plumbeus*, in the northeastern Gulf of Mexico. *Fish. Bull.* **97**, 387–391.
49. Merson RR, Pratt HL. 2001 Distribution, movements and growth of young sandbar sharks, *Carcharhinus plumbeus*, in the nursery grounds of Delaware Bay. *Environ. Biol. Fishes* **61**, 13–24. (doi:10.1023/A:1011017109776)
50. Smale MJ. 2002 Occurrence of *Carcharias taurus* in nursery areas of the Eastern and Western Cape, South Africa. *Mar. Freshwater Res.* **53**, 551–556.
51. DeAngelis BM, McCandless CT, Kohler NE, Recksiek CW, Skomal GB. 2008 First characterization of shark nursery habitat in the United States Virgin Islands: evidence of habitat partitioning by two shark species. *Mar. Ecol. Prog. Ser.* **358**, 257–271. (doi:10.3354/meps07308)
52. Reyier EA, Adams DH, Lowers RH. 2008 First evidence of a high density nursery ground for the lemon shark, *Negaprion brevirostris*, near Cape Canaveral, Florida. *Fla. Sci.* **71**, 134–148.
53. Cartamil D, Wegner NC, Kacev D, Ben-Aderet N, Kohin S, Graham JB. 2010 Movement patterns and nursery habitat of juvenile thresher sharks *Alopias vulpinus* in the Southern California Bight. *Mar. Ecol. Prog. Ser.* **404**, 249–258. (doi:10.3354/meps08495)
54. Froeschke JT, Stunz GW, Sterba-Boatwright B, Wildhaber ML. 2010 An empirical test of the 'shark nursery area concept' in Texas bays using a long-term fisheries-independent data set. *Aquat. Biol.* **11**, 65–76. (doi:10.3354/ab00290)
55. Kinney MJ. 2011 The communal nursery area paradigm revisited: niche overlap versus niche separation among juvenile shark species of Cleveland Bay. Doctoral Dissertation, James Cook University, Australia.
56. Yokota L, Lessa RP. 2006 A nursery area for sharks and rays in Northeastern Brazil. *Environ. Biol. Fishes* **75**, 349–360. (doi:10.1007/s10641-006-0038-9)
57. Skomal GB. 2007 Shark nursery areas in the coastal waters of Massachusetts. In *Shark nursery grounds of the Gulf of Mexico and east coast waters of the United States*. Am. Fish. Soc. Symp. 50 (eds CT McCandless, HL Jr Pratt, NE Kohler), pp. 17–33. Bethesda, MD: American Fisheries Society.
58. Boldrocchi G, Kiszka J, Purkis S, Storai T, Zinzula L, Burkholder D. 2017 Distribution, ecology, and status of the white shark, *Carcharodon carcharias*, in the Mediterranean Sea. *Rev. Fish Biol. Fish.* **27**, 515–534. (doi:10.1007/s11160-017-9470-5)
59. Krijgsman W, Hilgen FJ, Raffi I, Sierro FJ, Wilson DS. 1999 Chronology, causes and progression of the Messinian salinity crisis. *Nature* **400**, 652–655. (doi:10.1038/23231)
60. Cabrera L, Calvet F. 1996 E8 Onshore Neogene record in NE Spain: Vallès-Penedès and El Camp half-grabens (NW Mediterranean). In *Tertiary basins of Spain: The stratigraphic record of crustal kinematics* (eds PF Friend, CJ Dabrio), pp. 97–105. Cambridge, UK: Cambridge University Press.
61. Branstetter S. 1990 Early life-history implications of selected carcharhinoid and lamnoid sharks of the northwest Atlantic. In *Elasmobranchs as living resources: advances in the biology, ecology, systematics, and the status of the fisheries*, NOAA tech. Rep. 90 (eds HL Pratt, SH Gruber, T Taniuchi), pp. 17–28. Silver Spring, MD: National Marine Fisheries Service.