Published in final edited form as:

J Neurogenet. 2020 ; 34(3-4): 489–499. doi:10.1080/01677063.2020.1771333.

The discovery and consequences of the central role of the nervous system in the control of protein homeostasis

Veena Prahlada,b

^aDepartment of Biology, Aging Mind and Brain Initiative, University of Iowa, Iowa City, IA, USA;

^bIowa Neuroscience Institute, University of Iowa, Iowa City, IA, USA

Abstract

Organisms function despite wide fluctuations in their environment through the maintenance of homeostasis. At the cellular level, the maintenance of proteins as functional entities at target expression levels is called protein homeostasis (or proteostasis). Cells implement proteostasis through universal and conserved quality control mechanisms that surveil and monitor protein conformation. Recent studies that exploit the powerful ability to genetically manipulate specific neurons in *C. elegans* have shown that cells within this metazoan lose their autonomy over this fundamental survival mechanism. These studies have uncovered novel roles for the nervous system in controlling how and when cells activate their protein quality control mechanisms. Here we discuss the conceptual underpinnings, experimental evidence and the possible consequences of such a control mechanism.

PRELUDE: Whether the detailed examination of parts of the nervous system and their selective perturbation is sufficient to reconstruct how the brain generates behavior, mental disease, music and religion remains an open question. Yet, Sydney Brenner's development of *C. elegans* as an experimental organism and his faith in the bold reductionist approach that 'the understanding of wild-type behavior comes best after the discovery and analysis of mutations that alter it', has led to discoveries of unexpected roles for neurons in the biology of organisms.

Keywords

Homeostasis; proteostasis; non-autonomous; neuronal; *C. elegans*

Introduction

Most protein-based biological mechanisms proceed optimally only within a narrow range of environmental conditions (Fields, Dong, Meng, & Somero, 2015; Hofmann & Somero, 1995; Somero, 1995). Despite this, organisms thrive in a variety of ecological niches. A mechanism by which organisms function despite wide fluctuations in their environment is through the maintenance of homeostasis (from Greek: ὁμοίος homoeos, 'similar' and

---

CONTACT Veena Prahlada, veena-prahlad@uiowa.edu, Department of Biology, Aging Mind and Brain Initiative, University of Iowa, Iowa City, IA, USA; Iowa Neuroscience Institute, University of Iowa, Iowa City, IA, USA.

Disclosure statement

No potential conflict of interest was reported by the author(s).

στάσις stasis, ‘standing still’), whereby the internal milieu is maintained close to constant in the face of external perturbations. This concept, first described by Claude Bernard in 1865, has provided the foundation for our understanding of macroscopic physiological processes such as the maintenance of core body temperature, pH of blood (Bernard, 1965), regulation of food intake, body mass, and energy metabolism, all of which are under homeostatic control. More recently, the concept of homeostasis has been extended to mechanisms that occur at the sub-cellular level, such as the maintenance of protein function in cells, and has become popularly known as proteostasis (Balch, Morimoto, Dillin, & Kelly, 2008; Gidalevitz, Ben-Zvi, Ho, Brignull, & Morimoto, 2006; Powers, Morimoto, Dillin, Kelly, & Balch, 2009). In general, there appear to be two distinct kinds of strategies to achieve homeostasis. The first are servomechanisms (Leow, 2007), whereby error-sensing negative feedback loops correct the performance of a system to maintain some specific feature, called a *regulated variable*, constant or close to some set-point (Figure 1(A)). Such a mechanism requires an error detector that senses deviations from the set-point, and a mechanism to perform the necessary error corrections (often called the servomotor, as this terminology grew from mechanical devices such as steam engines). While servomechanisms can precisely maintain the regulated variable at the set-point, they are triggered only *after* the set-point, and homeostasis, have been perturbed. An alternate mechanism by which biological systems achieve homeostasis is through the activation of anticipatory mechanisms. These mechanisms, termed Cephalic mechanisms, are predictive, being implemented prior to the actual perturbation of the system. For instance, temperature homeostasis in an organism is maintained through mechanisms that *prevent* the core body temperature from deviating above or below a certain range through launching preemptive/anticipatory adjustments such as sweating, *before* core body temperatures are perturbed. The essence of such mechanisms is that, by definition, the command for the correction of the deviation comes from centers that can predict the oncoming perturbation using associated information, prior to themselves becoming perturbed. Thus, cephalic mechanisms can prevent that the system be perturbed in the first place (Lechan & Fekete, 2006; Mattes, 1997); however, they are not fail-safe as they can be ‘fooled’ and triggered even in the absence of any perturbations to homeostasis.

Until about a decade ago, the mechanisms by which cells and organisms maintained protein homeostasis were best studied in isolated cells in culture or unicellular organisms such as yeast and bacteria (Akerfelt, Morimoto, & Sistonen, 2010; Balch *et al.*, 2008; Goff & Goldberg, 1987; Goldberg, 1971, 1972; Hightower, 1980; Lindquist, 1986; Morano, Liu, & Thiele, 1998; Neef, Turski, & Thiele, 2010; Ritossa, 1996). In these systems, protein homeostasis is implemented mainly through feed-back loops that act as servomechanisms (Morimoto, 1998; Voellmy, 1994, 1996). In the last decade, however, experiments that may not have been so readily feasible were it not for the ability to manipulate neurons in *C. elegans* (Avery, Bargmann, & Horvitz, 1993; Chalfie *et al.*, 1985; Walrond, Kass, Stretton, & Donmoyer, 1985; White, Southgate, Thomson, & Brenner, 1983, 1986), demonstrated that in this metazoan the nervous system exerts neuroendocrine control over the conserved cellular machinery that maintains protein quality control. Such control enables the animal to trigger these deeply conserved survival and defense mechanisms in anticipation of a threat. Here we briefly highlight how the genetic manipulation of *C. elegans* neurons enabled these insights

and discuss some of the implications of the existence of cephalic control over cellular protein homeostasis. We focus mainly on the conceptual framework, and would like to direct interested readers to more thorough reviews for details (Miles, Scherz-Shouval, & van Oosten-Hawle, 2019; Wolff, Weissman, & Dillin, 2014).

Proteostasis

A mammalian cell synthesizes on average ~10,000 proteins at any given time, most of which can assume multiple conformations, many of which are unstructured, occur in multiple copies, have widely varying half-lives, and are in the continual process of being modified, transported and degraded (Jayaraj, Hipp, & Hartl, 2020; Mogk, Bukau, & Kampinga, 2018; Tyedmers, Mogk, & Bukau, 2010; Wolff *et al.*, 2014). Despite this complexity, cells largely maintain a target expression level of any particular protein, and protein concentrations do not vary erratically. Moreover, cells can detect the presence of protein aggregates and non-native protein conformations (Geiler-Samerotte *et al.*, 2011; Vendruscolo, Knowles, & Dobson, 2011), and respond in specific and conserved ways to restore conformation or trigger degradation of the abnormal species (Bednarska, Schymkowitz, Rousseau, & Van Eldere, 2013; Hipp, Park, & Hartl, 2014; Labbadia & Morimoto, 2015; Mattoo & Goloubinoff, 2014; Miller *et al.*, 2015; Silverman *et al.*, 2015; Vendruscolo *et al.*, 2011; Walther *et al.*, 2015). Since the conformation of proteins drives their stability and function, this ability of cells to maintain proteins as functional entities at specific concentrations occurs through mechanisms that surveil protein conformation (Ananthan, Goldberg, & Voellmy, 1986; Goff & Goldberg, 1985, 1987; Goldberg, 1972; Kandrór, Busconi, Sherman, & Goldberg, 1994; Knowles, Gunn, Hanson, & Ballard, 1975; Knowles & Ballard, 1976, 1978; Prouty, Karnovsky, & Goldberg, 1975). These mechanisms are collectively called proteostasis (Balch *et al.*, 2008; Gidalevitz *et al.*, 2006). Exciting studies have demonstrated the existence of interconnected networks of surveillance and quality control mechanisms that monitor protein quality during normal protein synthesis. For instance, ribosome stalling that can occur due to stochastic perturbations of translation, mutations, or a number of biotic or abiotic fluctuations in the environment will lead to the aborted synthesis of a nascent polypeptide chains. These aborted polypeptides are recognized by the cellular surveillance machinery and modified by the addition of carboxy-terminal alanine and threonine residues (CAT tails) (Brandman *et al.*, 2012; Kostova *et al.*, 2017; Sitron, Park, & Brandman, 2017; Sitron, Park, Giafaglione, & Brandman, 2020; Sitron & Brandman, 2019). This, in turn, leads to their aggregation and targeted degradation through the sequestration of a group of specialized proteins called molecular chaperones that recognize the aberrant conformation of the CAT-tailed polypeptides (Sitron *et al.*, 2017, 2020; Sitron & Brandman, 2019). The sequestration of molecular chaperones is also thought to activate a conserved transcription factor to increase the amounts of molecular chaperone proteins available to surveil and prevent the build-up of aberrant polypeptides. Translating these concepts into homeostatic control theory (Figure 1(A,B)) protein conformation is the *regulated variable*, being maintained at some *set-point* (optimal flux, optimal concentration etc.) by the activity of *controlled variables*: protein biogenesis and protein degradation (green boxes). The *regulated variable* is maintained by the activity of PLANTS: ribosomes for protein biogenesis and proteases, the proteasome, and autophagy pathways for degradation. The *set-point* for

misfolded proteins is monitored and maintained through the activity of CONTROLLERS, the best known of which are molecular chaperones that surveil the folding state of proteins. If the CONTROLLER detects deviations from the *set-point*, signals to the PLANTS change the flux of proteins through the system restoring the *regulated variable* to its acceptable set-point (Goldstein & Kopin, 2017; Iberall & Cardon, 1964).

As with most biological processes, the levels of proteins in a cell, and therefore also misfolded proteins, varies with the types of protein synthesized, physiological needs and environmental changes. Accordingly, *controlled variables* i.e. the flux of proteins through the PLANTS, and even the concentration of PLANTS and CONTROLLERS required for maintaining protein homeostasis vary. The best understood regulatory mechanisms by which a sudden increase in the *regulated variable*, i.e. misfolded protein, impacts the PLANTS and CONTROLLERS, are the so-called heat shock response (HSR) in the cytoplasm and the unfolded protein response in the endoplasmic reticulum (Gomez-Pastor, Burchfiel, & Thiele, 2018; Vihervaara, Duarte, & Lis, 2018). First discovered as a heat-inducible transcriptional response when *Drosophila busckii* cells were accidentally exposed to high temperatures (De Maio, Santoro, Tanguay, & Hightower, 2012; Ritossa, 1996), the HSR is a highly conserved, universal transcription program executed in response to protein damage (Akerfelt *et al.*, 2010; Ananthan *et al.*, 1986; Morimoto, 1998; Voellmy, 1994). The HSR is mediated by the heat shock transcription factor 1 (HSF1) whose activity results in the rapid expression of more molecular chaperones (CONTROLLERS) to help identify, and refold or degrade the damaged proteins. Molecular chaperones accomplish this by triggering changes in the *controlled variables*, typically promoting a decrease in protein biogenesis and an increase in degradation of the damaged proteins (Goldberg, 1971; Hightower, 1980; Lindquist, 1986; Morimoto, 1998). Similarly, the Unfolded Protein Response of the Endoplasmic Reticulum (UPR^{ER}) is a conserved cellular mechanism to detect and regulate proteostasis within the lumen of the endoplasmic reticulum (ER) if protein processing stalls due to the increase in aberrant ER proteins (Gething & Sambrook, 1992; Kozutsumi, Segal, Normington, Gething, & Sambrook, 1988; Munro & Pelham, 1986, 1987; Pelham & Munro, 1993). UPR^{ER} acts through the activation of all or a subset of three stress signal transducers found within the ER membrane, Inositol-requiring enzyme 1 (IRE1), Activating transcription factor 6 (ATF6), and protein kinase R-like endoplasmic reticulum kinase (PERK), which also increase the molecular chaperones (CONTROLLERS) that act in the ER and modulate the *controlled variables* (Mori, 2009; Mori, Kawahara, Yoshida, Yanagi, & Yura, 1996; Walter, 2010; Yoshida, Matsui, Yamamoto, Okada, & Mori, 2001). Parallel mechanisms exist for ensuring protein homeostasis in mitochondria (Haynes & Ron, 2010). Until recently, activation of the transcription factors that increase the amount of CONTROLLERS (HSF1 or IRE1, ATF6 and XBP1) was thought to be *solely* triggered by the individual cells in which protein homeostasis was perturbed. Thus, both HSF1 and the transcriptional activators of the UPR^{ER} are typically inhibited unless the flux of protein misfolding exceeds some set-point or threshold because they themselves are under the negative inhibition of the CONTROLLERS. Upon sensing an increase in protein misfolding, the CONTROLLERS are kinetically titrated away, dis-inhibiting (or permitting the activation of) the transcription factors, increasing the amounts of CONTROLLERS, and restoring homeostasis (Bernales, Papa, & Walter, 2006). Indeed, this is what is thought to occur during normal protein synthesis upon ribosome

stalling. Under these conditions, the aborted synthesis of the nascent aggregation prone polypeptide titrates away the negative inhibition on HSF1 activity to promote an increase in chaperone gene transcription. This elegant model thus lay the foundation for the proteostasis network in isolated cells.

Proteostasis in *C. elegans*

C. elegans possesses dedicated neurons to sense threats in its environment and can detect environmental changes well before they are severe enough to disrupt protein homeostasis (Bargmann, 1998; Gally & Bessereau, 2003; Goodman & Sengupta, 2019; Hobert, 2005). One such sensory system that senses temperature change is the thermosensory circuitry, which involves at least three pairs of sensory neurons (AFD, AWC and ASI), that are extraordinarily sensitive to temperature changes, and a layer of interneurons (AIZ and AIY) that process and integrate the temperature information (Beverly, Anbil, & Sengupta, 2011; Clark, Biron, Sengupta, & Samuel, 2006; Goodman & Sengupta, 2018, 2019). It had been shown, once all the neurons of *C. elegans* had been catalogued and mapped, that AFD neurons can detect changes as small as 0.05 °C above ambient temperature; hence, they could arguably be excited by temperature increments well below those that cause cellular damage (Beverly *et al.*, 2011; Clark *et al.*, 2006; Colosimo *et al.*, 2004). With such specialized machinery to sense the environment, had metazoans with a nervous system evolved mechanisms to activate their protein homeostasis machinery prior to experiencing temperatures where their proteins would misfold? This was the question that motivated the first experiments that demonstrated that, in addition to the elegant servomechanism of proteostasis that existed in individual cells, in *C. elegans* there also exists cephalic control over protein homeostasis (Prahlad, Cornelius, & Morimoto, 2008). In the first experiments we used mutations that impaired the thermosensory capacity of the AFD neurons, and asked whether chaperones would be induced upon a transient increase in temperatures, if the animal did not sense the temperature upshift. Indeed, mutations affecting the ability of thermosensory AFD neurons to sense heat also delayed HSF1 dependent molecular chaperone induction throughout *C. elegans* upon heat shock (Prahlad *et al.*, 2008) (Figure 2(A)). However, mutations that impair neurons throughout the lifetime of animals could arguably have reset physiology and metabolism such that animals had altered the flux of proteins through the PLANTS or re-set their thresholds for tolerating misfolded proteins. Therefore, the more convincing demonstration that proteostasis was under cephalic control came from experiments using optogenetics to excite the AFD neurons in the absence of heat. Such a perturbation mimicked the sensation of heat and was sufficient to simulate a HSR, activate HSF1 and upregulate HSPs in distal tissues (Figure 2(A)) (Tatum *et al.*, 2015). We found that the signal for AFD-dependent upregulation of HSPs was the ancient bioamine, serotonin that was released upon the excitation of the AFD neurons. Serotonin is best studied for its roles in mood, anxiety and cognition, and in almost all organisms real or perceived threats cause the release of neural serotonin (Cruz-Corchado, Ooi, Das, & Prahlad, 2020; Tatum *et al.*, 2015). In *C. elegans*, the use of powerful genetic tools including neuronal mutants that lack all serotonin, had shown that serotonin played a pivotal role in learning and memory (Chase & Koelle, 2007; Curran & Chalasani, 2012). These studies have thus immediately linked anxiety, experience, learning—aspects that are fundamental to cephalic

processes—to cellular changes in transcription and protein quality control. Indeed, in following experiments we have shown that *C. elegans* can ‘learn’, through prior experience of a threat, to enhance chaperone expression were it to encounter the specific threat in its environment (Ooi & Pahlad, 2017) (Figure 2(B)). Moreover, the release of serotonin that occurs upon the sensing of temperature change has a rapid effect on chromatin accessibility in the germline of animals conferring stress protection on future offspring (Das *et al.*, 2020) (Figure 2(C)).

Analogous neurohormonal signaling mechanisms have since been discovered for the CONTROLLERS of proteostasis in other cellular compartments in *C. elegans* (Berendzen *et al.*, 2016; Durieux, Wolff, & Dillin, 2011; Frakes *et al.*, 2020; Taylor, Berendzen, & Dillin, 2014; Taylor & Dillin, 2013; Zhang *et al.*, 2018) (Figure 2(A)). Thus, the constitutive ectopic expression of XBP1 in the nervous system—in neurons or supporting glial cells—or the activation of the mitochondrial UPR in neurons induces a systemic UPR^{ER} or UPR^{MT} respectively, in other tissues (Durieux *et al.*, 2011; Frakes *et al.*, 2020; Taylor *et al.*, 2014). In addition, it appears that neurons may not be the only sentinels of harm: activation of the heat shock response in one tissue results in the systemic induction of this stress response (Miles *et al.*, 2019; O’Brien *et al.*, 2018; van Oosten-Hawle, Porter, & Morimoto, 2013). Thus, it appears that in a multicellular organism, changes in proteostasis in one tissue can feedback onto protein metabolism in another tissue to systemically increase the quantity of the molecular chaperones or the CONTROLLERS that monitor set points for misfolded proteins.

Many studies have also shown, directly or indirectly, that the autonomy of cells over the *controlled variables*, protein biogenesis and protein degradation, is also outsourced to neurons (Ben-Gedalya & Cohen, 2012). The nervous system controls growth, metabolism and aging in response to nutrient and energy status of the animal (Broughton & Partridge, 2009; Burkewitz *et al.*, 2015; Kawli, Wu, & Tan, 2010; Minnerly, Zhang, Parker, Kaul, & Jia, 2017; Weir *et al.*, 2017; Wolkow, Kimura, Lee, & Ruvkun, 2000; Zhang *et al.*, 2019). As a consequence, dietary restriction activates the conserved cap’n’collar transcription factor, SKN-1 in two neurons (called the ASI neurons) (Bishop & Guarente, 2007; Schmeisser *et al.*, 2013) to increase metabolic activity (Bishop & Guarente, 2007; Schmeisser *et al.*, 2013). SKN-1 is also a key regulator of the abundance of the ubiquitin-proteasome machinery within the animal (Lehrbach & Ruvkun, 2016; Li *et al.*, 2011; Steinbaugh *et al.*, 2015). Dietary restriction is thus known to increase lifespan through (amongst other mechanisms) the increase in proteasome activity and the E3 ubiquitin ligase WWP-1 (Carrano, Liu, Dillin, & Hunter, 2009). Remarkably, the smell of food is sufficient to reverse the benefits of dietary restriction and a recent study demonstrated rapid changes in protein degradation rates in response to food odors sensed by olfactory neurons (Finger *et al.*, 2019). Similarly changes in the levels of the neuromodulator dopamine control the flux of proteins by modulating protein degradation through the ubiquitin-proteasome system (Joshi, Matlack, & Rongo, 2016). Moreover, since growth and differentiation of cells within metazoa such as *C. elegans* occur in a coordinated manner, it is perhaps not surprising that neurons are not the only cell-type that exerts hierarchical control over the proteostasis machinery of another cell: cells also manage the proteostasis control systems of each other. For instance, germline stem cells limit the capacity of muscle cells, and perhaps other cells, to respond to heat stress in

multiple tissues by inhibiting the amounts of PLANTS, in this case the regulatory particle non-ATPase 6 (RPN6) proteasome subunit that these distal cells are allowed to express (Sinha & Rae, 2014; Vilchez *et al.*, 2012). Similarly, specialized cells called coelomocytes play a key role in the clearance of aggregation-prone transthyretin expressed in *C. elegans* muscle (Madhivanan *et al.*, 2018), and protein degradation in the muscle, albeit driven by neuronal activity, is the integrated outcome of signals from neurons and other tissues such as hypodermis and intestine. In turn, distal cells modulate *controlled variables* to feedback on neurons and the rest of the animal (Chikka, Anbalagan, Dvorak, Dombeck, & Prahlad, 2016; Dalton & Curran, 2018; Minnerly *et al.*, 2017).

While neuronal networks confer the ability to respond rapidly to environmental threats, we and others have found that this ability comes at the expense of the ‘normal’ proteostasis machinery (Berendzen *et al.*, 2016; Cao & Aballay, 2016; David, 2013; El-Ami *et al.*, 2014; Imanikia, Ozbey, Krueger, Casanueva, & Taylor, 2019; Joshi *et al.*, 2016; Maman *et al.*, 2013; Moll, Ben-Gedalya, Reuveni, & Cohen, 2016; O’Brien *et al.*, 2018; Prahlad & Morimoto, 2011; Ray, Zhang, Rentas, Caldwell, & Caldwell, 2014; Roitenberg *et al.*, 2018; Taylor & Dillin, 2013) (Figure 2(D)). For example, the AFD-dependent circuitry accelerates the activation of HSF1 upon sensing environmental stress but prevents HSF1 from being activated when cells chronically express aggregation prone proteins (Prahlad & Morimoto, 2011). Consequently, the activity of the AFD-thermosensory neurons needs to be inhibited in order to allow cells in the metazoan to activate HSF1 and increase the expression of CONTROLLERS needed to respond adequately to protein aggregation. Similarly, in wild-type animals, the presence of a G protein coupled receptor expressed on chemosensory neurons (*gtr-1*) is required for animals to increase chaperone expression upon exposure to heat, but its presence inhibits the ‘normal’ proteostasis machinery from rescuing the proteotoxicity of Alzheimer’s-disease-linked aggregative peptide A β_{3-42} (Maman *et al.*, 2013). A similar phenomenon is seen in studies on aging, a process that results in the systemic loss of protein quality control and the accumulation of misfolded proteins in various tissues: sensory neuronal function while allowing animals to respond to their environment, invariably inhibits the proteostasis machinery from acting on misfolded proteins (Bocchitto, Lamitina, & Kalb, 2012; Broughton & Partridge, 2009; Burkewitz *et al.*, 2015; Byrne, Wilhelm, & Richly, 2017; Jiang *et al.*, 2015; Kumsta *et al.*, 2014; Stein & Murphy, 2012; Taylor & Dillin, 2013; Yin, Liu, Yuan, Jiang, & Cai, 2014; Zhang *et al.*, 2019). In these cases, the loss of normal neuronal control is required to allow protein misfolding to activate the normal proteostasis machinery, increase CONTROLLERS, and recalibrate the flow of proteins through the PLANTS to restore health and function (Figures 1 and 2). Why might such a seemingly counterintuitive mechanism have evolved? We believe this is because neuronal signals have higher physiological priority because they orchestrate the protective response to life threatening environmental changes. Therefore, in a metazoan, the ability of neurons to antagonize the mechanisms that could normally be used by the proteostasis machinery of individual cells allows neurons to disengage this very same machinery for use upon the sensing of a threat. The control over the activation of transcription factors that are responsible for the expression of CONTROLLERS in all compartments of cells allows neurons to play this dominant role. In addition, such a control mechanism also forces the normal proteostasis machinery—the CONTROLLERS and the

PLANTS involved in protein biogenesis—to function within a range of stringent set points under normal physiological conditions even through the regulated variable, protein misfolding, itself may not be well defined. Indeed, experiments in *C. elegans* have shown that the proteostasis machinery operates within a narrow range which is not easily perturbed (Gidalevitz *et al.*, 2006).

It is interesting to speculate on what could happen were cell intrinsic and cell extrinsic control mechanisms in disagreement. In human diseases such as Alzheimer's disease and Parkinson's disease, and in animal models of these diseases, cells accumulate misfolded and aggregated proteins but fail to *naturally* activate their cell-autonomous proteostasis response. This suggests that in the absence of external stimuli, the cell-extrinsic control dominates, preventing the increase in CONTROLLERS and PLANTS even at the cost of increasing the cellular burden of misfolded proteins. Indeed, experimentally increasing CONTROLLERS such as molecular chaperones in the affected cells, by transfection or other means, ameliorates misfolding and disease pathology. However, it is also telling that the cell intrinsic increase in the abundance of CONTROLLERS and PLANTS occurring in the apparent absence of cell extrinsic control is a *prerequisite* for cancer maintenance and malignancy in mammals (Dai, Whitesell, Rogers, & Lindquist, 2007; Gaglia *et al.*, 2020; Mendillo *et al.*, 2012; Scherz-Shouval *et al.*, 2014). Since the most obvious function of cephalic control over a community of cells, where each cell has autonomous control over its own proteostasis machinery, is the maintenance of harmony and co-ordination required for multicellular function, one might image that an anarchy of cell intrinsic control over that exerted by neurons may result in defects in development and differentiation, and other such organismal functions.

Is what we have learnt from *C. elegans* more generally true?

The ability to manipulate *C. elegans* neurons has uncovered a novel role for the nervous system in the control of protein quality control mechanisms. Dysfunction of proteostasis is responsible for the devastating age-associated neurodegenerative diseases and other disorders prevalent in our societies today. Therefore, that need to understand the extent to which neurohormonal signaling pathways influence cellular proteostasis in mammals is urgent. Interventions to activate protective protein quality control mechanisms in the treatment of protein conformational diseases have, to date, been based largely on the assumption that protein misfolding, and its subsequent correction, are cell autonomous processes: namely the cells that need to be targeted are the cells within which protein misfolding occurs (Balch *et al.*, 2008; Powers *et al.*, 2009). However, if protein folding homeostasis in all multicellular organisms is under the cell non-autonomous regulation of sensory neuronal modalities, another method for treatment of neurodegenerative diseases could involve modulation of neurosensory systems: a method that can be used alone, or in combination with small molecules to effectively target protein misfolding. Evidence for such control in mammals does exist, and is growing. Prior to the studies in *C. elegans*, restraint stress had been shown to induce expression of HSP70 mRNA in the adrenal cortex of the rat through the activity of the by the hypothalamic-pituitary-adrenal (HPA) axis (Blake, Buckley, & Buckley, 1993; Blake, Udelsman, Feulner, Norton, & Holbrook, 1991; Fawcett, Sylvester, Sarge, Morimoto, & Holbrook, 1994). Notably, the control of the proteostasis

machinery of cells by the nervous system shares features with the systemic inhibition of acute inflammation seen in mammals (Tracey, 2002) and invertebrates including *C. elegans* (Anderson, Laurenson-Schafer, Partridge, Hodgkin, & McMullan, 2013; Singh & Aballay, 2012; Styer *et al.*, 2008; Sun, Liu, & Aballay, 2012; Sun, Singh, Kajino-Sakamoto, & Aballay, 2011; Yu, Zhi, Wu, Jing, & Wang, 2018), and treatments to modulate inflammatory processes are leveraging the existence of neuronal control. In mammals cholinergic activation within the efferent vagus nerve suppresses chronic inflammation, restricting the innate immune response to remain an acute response to local insults, and not flare up into chronic inflammation (Tracey, 2002). Moreover, as with the inflammatory response, recent experiments show that increased activity of the CONTROLLERS and PLANTS is protective only under a controlled regimen, and can be detrimental if unchecked (Styer *et al.*, 2008; Sun *et al.*, 2011). In mouse models of Alzheimer's Disease and Prion Disease it was the sustained activation of proteostasis machinery, rather than its dysfunction that accelerated neurodegeneration (Ma *et al.*, 2013; Moreno *et al.*, 2012). Protein misfolding within the cell, if detected and responded to by the CONTROLLERS elicits not repair, but also a change in the activity of the PLANTS such as a transient attenuation of protein translation through the phosphorylation of the α -subunit of eukaryotic initiation factor-2 (eIF2 α) (Harding *et al.*, 2003), presumably to lower the burden of nascent polypeptides to an already stressed proteome (Ma *et al.*, 2013; Moreno *et al.*, 2012). This attenuation of translation has been shown to be toxic to neurons.

Thus, in retrospect, the finding that proteostasis is systemically controlled in a metazoan could perhaps have been anticipated. The maintenance of protein homeostasis ensures the presence of adequate amounts of functional proteins in a cell and is therefore directly responsible for cell growth, differentiation and function. It is possible that the nervous system, because of its role in regulating multicellular outcomes such as aging, behavior, metabolism, longevity and reproduction, also determines the set-points of misfolded proteins in the different cells of an organism maintaining them within some 'optimal' range. A better understanding of the mechanisms responsible for determining these set-points could go a long-way in treating protein conformation diseases.

Acknowledgements

We thank the members of V.P. laboratory and Dr. Tali Gidalevitz (Drexel University) for useful comments. We apologize to all our colleagues whose work we may have omitted to cite.

Funding

This work was supported by NIH R01 AG 050653 (V.P.).

References

- Akerfelt M, Morimoto RI, & Sistonen L (2010). Heat shock factors: Integrators of cell stress, development and lifespan. *Nature Reviews. Molecular Cell Biology*, 11(8), 545–555. doi:10.1038/nrm2938 [PubMed: 20628411]
- Ananthan J, Goldberg AL, & Voellmy R (1986). Abnormal proteins serve as eukaryotic stress signals and trigger the activation of heat shock genes. *Science (New York, N.Y.)*, 232(4749), 522–524. doi:10.1126/science.3083508

- Anderson A, Laurenson-Schafer H, Partridge FA, Hodgkin J, & McMullan R (2013). Serotonergic chemosensory neurons modify the *C. elegans* immune response by regulating G-protein signaling in epithelial cells. *PLoS Pathogens*, 9(12), e1003787. doi:10.1371/journal.ppat.1003787 [PubMed: 24348250]
- Avery L, Bargmann CI, & Horvitz HR (1993). The *Caenorhabditis elegans* unc-31 gene affects multiple nervous system-controlled functions. *Genetics*, 134(2), 455–464. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8325482> [PubMed: 8325482]
- Balch WE, Morimoto RI, Dillin A, & Kelly JW (2008). Adapting proteostasis for disease intervention. *Science (New York, N.Y.)*, 319(5865), 916–919. doi:10.1126/science.1141448
- Bargmann CI (1998). Neurobiology of the *Caenorhabditis elegans* genome. *Science (New York, N.Y.)*, 282(5396), 2028–2033. doi:10.1126/science.282.5396.2028
- Bednarska NG, Schymkowitz J, Rousseau F, & Van Eldere J (2013). Protein aggregation in bacteria: the thin boundary between functionality and toxicity. *Microbiology (Reading, England)*, 159(Pt 9), 1795–1806. doi:10.1099/mic.0.069575-0
- Ben-Gedalya T, & Cohen E (2012). Quality control compartments coming of age. *Traffic (Copenhagen, Denmark)*, 13(5), 635–642. doi: 10.1111/j.1600-0854.2012.01330.x
- Berendzen KM, Durieux J, Shao L-W, Tian Y, Kim H-E, Wolff S, ... Dillin A (2016). Neuroendocrine coordination of mitochondrial stress signaling and proteostasis. *Cell*, 166(6), 1553–1563 e1510. doi:10.1016/j.cell.2016.08.042 [PubMed: 27610575]
- Bernales S, Papa FR, & Walter P (2006). Intracellular signaling by the unfolded protein response. *Annual Review of Cell and Developmental Biology*, 22, 487–508. doi:10.1146/annurev.cellbio.21.122303.120200
- Bernard C (1965). An introduction to the study of experimental medicine. *Medical Journal of Australia*, 1(4), 119–120. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/14248653>
- Beverly M, Anbil S, & Sengupta P (2011). Degeneracy and neuromodulation among thermosensory neurons contribute to robust thermosensory behaviors in *Caenorhabditis elegans*. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, 31(32), 11718–11727. doi:10.1523/JNEUROSCI.1098-11.2011 [PubMed: 21832201]
- Bishop NA, & Guarente L (2007). Two neurons mediate diet-restriction-induced longevity in *C. elegans*. *Nature*, 447(7144), 545–549. doi:10.1038/nature05904 [PubMed: 17538612]
- Blake MJ, Buckley DJ, & Buckley AR (1993). Dopaminergic regulation of heat shock protein-70 expression in adrenal gland and aorta. *Endocrinology*, 132(3), 1063–1070. doi:10.1210/endo.132.3.8095012 [PubMed: 8095012]
- Blake MJ, Udelsman R, Feulner GJ, Norton DD, & Holbrook NJ (1991). Stress-induced heat shock protein 70 expression in adrenal cortex: An adrenocorticotrophic hormone-sensitive, age-dependent response. *Proceedings of the National Academy of Sciences of the United States of America*, 88(21), 9873–9877. doi:10.1073/pnas.88.21.9873 [PubMed: 1658790]
- Bocitto M, Lamitina T, & Kalb RG (2012). Daf-2 signaling modifies mutant SOD1 toxicity in *C. elegans*. *PLoS One*, 7(3), e33494. doi:10.1371/journal.pone.0033494 [PubMed: 22457769]
- Brandman O, Stewart-Ornstein J, Wong D, Larson A, Williams CC, Li G-W, ... Weissman JS (2012). A ribosome-bound quality control complex triggers degradation of nascent peptides and signals translation stress. *Cell*, 151(5), 1042–1054. doi:10.1016/j.cell.2012.10.044 [PubMed: 23178123]
- Broughton S, & Partridge L (2009). Insulin/IGF-like signalling, the central nervous system and aging. *The Biochemical Journal*, 418(1), 1–12. doi:10.1042/BJ20082102 [PubMed: 19159343]
- Burkewitz K, Morante I, Weir HJM, Yeo R, Zhang Y, Huynh FK, ... Mair WB (2015). Neuronal CRTCL-1 governs systemic mitochondrial metabolism and lifespan via a catecholamine signal. *Cell*, 160(5), 842–855. doi:10.1016/j.cell.2015.02.004 [PubMed: 25723162]
- Byrne J, Wilhelm T, & Richly H (2017). Inhibition of neuronal autophagy mediates longevity. *Aging*, 9(9), 1953–1954. doi:10.18632/aging.101297 [PubMed: 29017148]
- Cao X, & Aballay A (2016). Neural inhibition of dopaminergic signaling enhances immunity in a cell-non-autonomous manner. *Current Biology*, 26(17), 2329–2334. doi:10.1016/j.cub.2016.06.036 [PubMed: 27524480]

- Carrano AC, Liu Z, Dillin A, & Hunter T (2009). A conserved ubiquitination pathway determines longevity in response to diet restriction. *Nature*, 460(7253), 396–399. doi:10.1038/nature08130 [PubMed: 19553937]
- Chalfie M, Sulston JE, White JG, Southgate E, Thomson JN, & Brenner S (1985). The neural circuit for touch sensitivity in *Caenorhabditis elegans*. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, 5(4), 956–964. doi:10.1523/JNEUROSCI.05-04-00956.1985 [PubMed: 3981252]
- Chase DL, & Koelle MR (2007). Biogenic amine neurotransmitters in *C. elegans*. *WormBook*, pp. 1–15. doi:10.1895/wormbook.1.132.1
- Chikka MR, Anbalagan C, Dvorak K, Dombeck K, & Prahlad V (2016). The mitochondria-regulated immune pathway activated in the *C. elegans* intestine is neuroprotective. *Cell Reports*, 16(9), 2399–2414. doi:10.1016/j.celrep.2016.07.077 [PubMed: 27545884]
- Clark DA, Biron D, Sengupta P, & Samuel AD (2006). The AFD sensory neurons encode multiple functions underlying thermotactic behavior in *Caenorhabditis elegans*. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, 26(28), 7444–7451. doi:10.1523/JNEUROSCI.1137-06.2006 [PubMed: 16837592]
- Colosimo ME, Brown A, Mukhopadhyay S, Gabel C, Lanjuin AE, Samuel AD, & Sengupta P (2004). Identification of thermosensory and olfactory neuron-specific genes via expression profiling of single neuron types. *Current Biology : CB*, 14(24), 2245–2251. doi:10.1016/j.cub.2004.12.030 [PubMed: 15620651]
- Cruz-Corchado J, Ooi FK, Das S, & Prahlad V (2020). Global transcriptome changes that accompany alterations in serotonin levels in *Caenorhabditis elegans*. *G3 (Bethesda)*, 10, 1225–1246. doi:10.1534/g3.120.401088 [PubMed: 31996358]
- Curran KP, & Chalasani SH (2012). Serotonin circuits and anxiety: What can invertebrates teach us? *Invertebrate Neuroscience : IN*, 12(2), 81–92. doi:10.1007/s10158-012-0140-y [PubMed: 22918570]
- Dai C, Whitesell L, Rogers AB, & Lindquist S (2007). Heat shock factor 1 is a powerful multifaceted modifier of carcinogenesis. *Cell*, 130(6), 1005–1018. doi:10.1016/j.cell.2007.07.020 [PubMed: 17889646]
- Dalton HM, & Curran SP (2018). Hypodermal responses to protein synthesis inhibition induce systemic developmental arrest and AMPK-dependent survival in *Caenorhabditis elegans*. *PLoS Genetics*, 14(7), e1007520. doi:10.1371/journal.pgen.1007520 [PubMed: 30020921]
- Das S, Ooi FK, Cruz Corchado J, Fuller LC, Weiner JA, & Prahlad V (2020). Serotonin signaling by maternal neurons upon stress ensures progeny survival. *eLife* 2020;9:e55246. doi:10.7554/eLife.55246
- David R (2013). Protein metabolism: Proteostasis goes global. *Nature Reviews Molecular Cell Biology*, 14(8), 461. doi:10.1038/nrm3626
- De Maio A, Santoro MG, Tanguay RM, & Hightower LE (2012). Ferruccio Ritossa's scientific legacy 50 years after his discovery of the heat shock response: A new view of biology, a new society, and a new journal. *Cell Stress Chaperones*, 17(2), 139–143. doi:10.1007/s12192-012-0320-z [PubMed: 22252402]
- Durieux J, Wolff S, & Dillin A (2011). The cell-non-autonomous nature of electron transport chain-mediated longevity. *Cell*, 144(1), 79–91. doi:10.1016/j.cell.2010.12.016 [PubMed: 21215371]
- El-Ami T, Moll L, Carvalhal Marques F, Volovik Y, Reuveni H, & Cohen E (2014). A novel inhibitor of the insulin/IGF signaling pathway protects from age-onset, neurodegeneration-linked proteotoxicity. *Aging Cell*, 13(1), 165–174. doi:10.1111/accel.12171 [PubMed: 24261972]
- Fawcett TW, Sylvester SL, Sarge KD, Morimoto RI, & Holbrook NJ (1994). Effects of neurohormonal stress and aging on the activation of mammalian heat shock factor 1. *The Journal of Biological Chemistry*, 269(51), 32272–32278. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=7798227 [PubMed: 7798227]
- Fields PA, Dong Y, Meng X, & Somero GN (2015). Adaptations of protein structure and function to temperature: There is more than one way to 'skin a cat'. *The Journal of Experimental Biology*, 218(Pt 12), 1801–1811. doi:10.1242/jeb.114298 [PubMed: 26085658]

- Finger F, Ottens F, Springhorn A, Drexel T, Proksch L, Metz S, ... Hoppe T (2019). Olfaction regulates organismal proteostasis and longevity via microRNA-dependent signaling. *Nature Metabolism*, 1(3), 350–359. doi:10.1038/s42255-019-0033-z
- Frakes AE, Metcalf MG, Tronnes SU, Bar-Ziv R, Durieux J, Gildea HK, ... Dillin A (2020). Four glial cells regulate ER stress resistance and longevity via neuropeptide signaling in *C. elegans*. *Science* (New York, N.Y.), 367(6476), 436–440. doi:10.1126/science.aaz6896
- Gaglia G, Rashid R, Yapp C, Joshi GN, Li CG, Lindquist SL, ... Santagata S (2020). HSF1 phase transition mediates stress adaptation and cell fate decisions. *Nature Cell Biology*, 22(2), 151–158. doi:10.1038/s41556-019-0458-3 [PubMed: 32015439]
- Gally C, & Bessereau JL (2003). [*C. elegans*: of neurons and genes]. *Medecine Sciences : M/S*, 19(6–7), 725–734. doi:10.1051/medsci/20031967725 [PubMed: 12942444]
- Geiler-Samerotte KA, Dion MF, Budnik BA, Wang SM, Hartl DL, & Drummond DA (2011). Misfolded proteins impose a dosage-dependent fitness cost and trigger a cytosolic unfolded protein response in yeast. *Proceedings of the National Academy of Sciences of the United States of America*, 108(2), 680–685. doi:10.1073/pnas.1017570108 [PubMed: 21187411]
- Gething MJ, & Sambrook J (1992). Protein folding in the cell. *Nature*, 355(6355), 33–45. doi:10.1038/355033a0 [PubMed: 1731198]
- Gidalevitz T, Ben-Zvi A, Ho KH, Brignull HR, & Morimoto RI (2006). Progressive disruption of cellular protein folding in models of polyglutamine diseases. *Science* (New York, N.Y.), 311(5766), 1471–1474. doi:10.1126/science.1124514
- Goff SA, & Goldberg AL (1985). Production of abnormal proteins in *E. coli* stimulates transcription of lon and other heat shock genes. *Cell*, 41(2), 587–595. doi:10.1016/S0092-8674(85)80031-3 [PubMed: 3886165]
- Goff SA, & Goldberg AL (1987). An increased content of protease La, the lon gene product, increases protein degradation and blocks growth in *Escherichia coli*. *The Journal of Biological Chemistry*, 262(10), 4508–4515. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/3549709> [PubMed: 3549709]
- Goldberg AL (1971). A role of aminoacyl-tRNA in the regulation of protein breakdown in *Escherichia coli*. *Proceedings of the National Academy of Sciences of the United States of America*, 68(2), 362–366. doi:10.1073/pnas.68.2.362 [PubMed: 5277086]
- Goldberg AL (1972). Correlation between rates of degradation of bacterial proteins in vivo and their sensitivity to proteases. *Proceedings of the National Academy of Sciences of the United States of America*, 69(9), 2640–2644. doi:10.1073/pnas.69.9.2640 [PubMed: 4560693]
- Goldstein DS, & Kopin IJ (2017). Homeostatic systems, biocybernetics, and autonomic neuroscience. *Autonomic Neuroscience : Basic & Clinical*, 208, 15–28. doi:10.1016/j.autneu.2017.09.001 [PubMed: 28918243]
- Gomez-Pastor R, Burchfiel ET, & Thiele DJ (2018). Regulation of heat shock transcription factors and their roles in physiology and disease. *Nature Reviews. Molecular Cell Biology*, 19(1), 4–19. doi:10.1038/nrm.2017.73 [PubMed: 28852220]
- Goodman MB, & Sengupta P (2018). The extraordinary AFD thermosensor of *C. elegans*. *Pflugers Archiv : European Journal of Physiology*, 470(5), 839–849. doi:10.1007/s00424-017-2089-5 [PubMed: 29218454]
- Goodman MB, & Sengupta P (2019). How *Caenorhabditis elegans* senses mechanical stress, temperature, and other physical stimuli. *Genetics*, 212(1), 25–51. doi:10.1534/genetics.118.300241 [PubMed: 31053616]
- Harding HP, Zhang Y, Zeng H, Novoa I, Lu PD, Calfon M, ... Ron D (2003). An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Molecular Cell*, 11(3), 619–633. doi:10.1016/S1097-2765(03)00105-9 [PubMed: 12667446]
- Haynes CM, & Ron D (2010). The mitochondrial UPR - protecting organelle protein homeostasis. *Journal of Cell Science*, 123(Pt 22), 3849–3855. doi:10.1242/jcs.075119 [PubMed: 21048161]
- Hightower LE (1980). Cultured animal cells exposed to amino acid analogues or puromycin rapidly synthesize several polypeptides. *Journal of Cellular Physiology*, 102(3), 407–427. doi:10.1002/jcp.1041020315 [PubMed: 6901532]

- Hipp MS, Park SH, & Hartl FU (2014). Proteostasis impairment in protein-misfolding and -aggregation diseases. *Trends in Cell Biology*, 24(9), 506–514. doi:10.1016/j.tcb.2014.05.003 [PubMed: 24946960]
- Hobert O (2005). Specification of the nervous system In *WormBook* (pp. 1–19). *WormBook*, ed. The *C. elegans* Research Community, doi:10.1895/wormbook.
- Hofmann G, & Somero G (1995). Evidence for protein damage at environmental temperatures: seasonal changes in levels of ubiquitin conjugates and hsp70 in the intertidal mussel *Mytilus trossulus*. *The Journal of Experimental Biology*, 198(Pt 7), 1509–1518. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9319406 [PubMed: 9319406]
- Iberall AS, & Cardon SZ (1964). Control in Biological systems-a physical review. *Annals of the New York Academy of Sciences*, 117, 445–518. doi:10.1111/j.1749-6632.1964.tb48202.x [PubMed: 14196660]
- Imanikia S, Ozbey NP, Krueger C, Casanueva MO, & Taylor RC (2019). Neuronal XBP-1 activates intestinal lysosomes to improve proteostasis in *C. elegans*. *Current Biology : CB*, 29(14), 2322–2338 e2327. doi:10.1016/j.cub.2019.06.031 [PubMed: 31303493]
- Jayaraj GG, Hipp MS, & Hartl FU (2020). Functional modules of the proteostasis network. *Cold Spring Harbor Perspectives in Biology*, 12(1), a033951. doi:10.1101/cshperspect [PubMed: 30833457]
- Jiang HC, Hsu JM, Yen CP, Chao CC, Chen RH, & Pan CL (2015). Neural activity and CaMKII protect mitochondria from fragmentation in aging *Caenorhabditis elegans* neurons. *Proceedings of the National Academy of Sciences of the United States of America*, 112(28), 8768–8773. doi:10.1073/pnas.1501831112 [PubMed: 26124107]
- Joshi KK, Matlack TL, & Rongo C (2016). Dopamine signaling promotes the xenobiotic stress response and protein homeostasis. *The EMBO Journal*, 35(17), 1885–1901. doi:10.15252/embj.201592524 [PubMed: 27261197]
- Kandror O, Busconi L, Sherman M, & Goldberg AL (1994). Rapid degradation of an abnormal protein in *Escherichia coli* involves the chaperones GroEL and GroES. *The Journal of Biological Chemistry*, 269(38), 23575–23582. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/7916344> [PubMed: 7916344]
- Kawli T, Wu C, & Tan MW (2010). Systemic and cell intrinsic roles of Gqalpha signaling in the regulation of innate immunity, oxidative stress, and longevity in *Caenorhabditis elegans*. *Proceedings of the National Academy of Sciences of the United States of America*, 107(31), 13788–13793. doi:10.1073/pnas.0914715107 [PubMed: 20647387]
- Knowles SE, & Ballard FJ (1976). Selective control of the degradation of normal and aberrant proteins in Reuber H35 hepatoma cells. *The Biochemical Journal*, 156(3), 609–617. doi:10.1042/bj1560609 [PubMed: 182157]
- Knowles SE, & Ballard FJ (1978). Effects of amino acid analogues on protein synthesis and degradation in isolated cells. *The British Journal of Nutrition*, 40(2), 275–287. doi:10.1079/bjn19780123 [PubMed: 212095]
- Knowles SE, Gunn JM, Hanson RW, & Ballard FJ (1975). Increased degradation rates of protein synthesized in hepatoma cells in the presence of amino acid analogues. *The Biochemical Journal*, 146(3), 595–600. doi:10.1042/bj1460595 [PubMed: 167723]
- Kostova KK, Hickey KL, Osuna BA, Hussmann JA, Frost A, Weinberg DE, & Weissman JS (2017). CAT-tailing as a fail-safe mechanism for efficient degradation of stalled nascent polypeptides. *Science (New York, N.Y.)*, 357(6349), 414–417. doi:10.1126/science.aam7787
- Kozutsumi Y, Segal M, Normington K, Gething MJ, & Sambrook J (1988). The presence of malformed proteins in the endoplasmic reticulum signals the induction of glucose-regulated proteins. *Nature*, 332(6163), 462–464. doi:10.1038/332462a0 [PubMed: 3352747]
- Kumsta C, Ching T-T, Nishimura M, Davis AE, Gelino S, Catan HH, ... Hansen M (2014). Integrin-linked kinase modulates longevity and thermotolerance in *C. elegans* through neuronal control of HSF-1. *Aging Cell*, 13(3), 419–430. doi:10.1111/ace.12189 [PubMed: 24314125]
- Labbadia J, & Morimoto RI (2015). The biology of proteostasis in aging and disease. *Annual Review of Biochemistry*, 84, 435–464. doi: 10.1146/annurev-biochem-060614-033955

- Lechan RM, & Fekete C (2006). The TRH neuron: A hypothalamic integrator of energy metabolism. *Progress in Brain Research*, 153, 209–235. doi:10.1016/S0079-6123(06)53012-2 [PubMed: 16876577]
- Lehrbach NJ, & Ruvkun G (2016). Proteasome dysfunction triggers activation of SKN-1A/Nrf1 by the aspartic protease DDI-1. *eLife*, 5, 7554. doi:10.7554/eLife.17721
- Leow MK (2007). A mathematical model of pituitary-thyroid interaction to provide an insight into the nature of the thyrotropin-thyroid hormone relationship. *Journal of Theoretical Biology*, 248(2), 275–287. doi:10.1016/j.jtbi.2007.05.016 [PubMed: 17602707]
- Li X, Matilainen O, Jin C, Glover-Cutter KM, Holmberg CI, & Blackwell TK (2011). Specific SKN-1/Nrf stress responses to perturbations in translation elongation and proteasome activity. *PLoS Genetics*, 7(6), e1002119. doi:10.1371/journal.pgen.1002119 [PubMed: 21695230]
- Lindquist S (1986). The heat-shock response. *Annual Review of Biochemistry*, 55, 1151–1191. doi:10.1146/annurev.bi.55.070186.005443
- Ma T, Trinh MA, Wexler AJ, Bourbon C, Gatti E, Pierre P, ... Klann E (2013). Suppression of eIF2 α kinases alleviates Alzheimer's disease-related plasticity and memory deficits. *Nature Neuroscience*, 16(9), 1299–1305. doi:10.1038/nn.3486 [PubMed: 23933749]
- Madhivanan K, Greiner ER, Alves-Ferreira M, Soriano-Castell D, Rouzbeh N, Aguirre CA, ... Encalada SE (2018). Cellular clearance of circulating transthyretin decreases cell-nonautonomous proteotoxicity in *Caenorhabditis elegans*. *Proceedings of the National Academy of Sciences of the United States of America*, 115(33), E7710–E7719. doi:10.1073/pnas.1801117115 [PubMed: 30061394]
- Maman M, Carvalhal Marques F, Volovik Y, Dubnikov T, Bejerano-Sagie M, & Cohen E (2013). A neuronal GPCR is critical for the induction of the heat shock response in the nematode *C. elegans*. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, 33(14), 6102–6111. doi:10.1523/JNEUROSCI.4023-12.2013 [PubMed: 23554491]
- Mattes RD (1997). Physiologic responses to sensory stimulation by food: Nutritional implications. *Journal of the American Dietetic Association*, 97(4), 406–413. doi:10.1016/S0002-8223(97)00101-6 [PubMed: 9120195]
- Mattoo RU, & Goloubinoff P (2014). Molecular chaperones are nanomachines that catalytically unfold misfolded and alternatively folded proteins. *Cellular and Molecular Life Sciences : Cmls*, 71(17), 3311–3325. doi:10.1007/s00018-014-1627-y [PubMed: 24760129]
- Mendillo ML, Santagata S, Koeva M, Bell GW, Hu R, Tamimi RM, ... Lindquist S (2012). HSF1 drives a transcriptional program distinct from heat shock to support highly malignant human cancers. *Cell*, 150(3), 549–562. doi:10.1016/j.cell.2012.06.031 [PubMed: 22863008]
- Miles J, Scherz-Shouval R, & van Oosten-Hawle P (2019). Expanding the organismal proteostasis network: Linking systemic stress signaling with the innate immune response. *Trends in Biochemical Sciences*, 44(11), 927–942. doi:10.1016/j.tibs.2019.06.009 [PubMed: 31303384]
- Miller SBM, Ho C-T, Winkler J, Khokhrina M, Neuner A, Mohamed MYH, ... Bukau B (2015). Compartment-specific aggregates direct distinct nuclear and cytoplasmic aggregate deposition. *The EMBO Journal*, 34(6), 778–797. doi:10.15252/embj.201489524 [PubMed: 25672362]
- Minnerly J, Zhang J, Parker T, Kaul T, & Jia K (2017). The cell non-autonomous function of ATG-18 is essential for neuroendocrine regulation of *Caenorhabditis elegans* lifespan. *PLoS Genetics*, 13(5), e1006764. doi:10.1371/journal.pgen.1006764 [PubMed: 28557996]
- Mogk A, Bukau B, & Kampina HH (2018). Cellular handling of protein aggregates by disaggregation machines. *Molecular Cell*, 69(2), 214–226. doi:10.1016/j.molcel.2018.01.004 [PubMed: 29351843]
- Moll L, Ben-Gedalya T, Reuveni H, & Cohen E (2016). The inhibition of IGF-1 signaling promotes proteostasis by enhancing protein aggregation and deposition. *FASEB Journal : Official Publication of the Federation of American Societies for Experimental Biology*, 30(4), 1656–1669. doi:10.1096/fj.15-281675 [PubMed: 26722006]
- Morano KA, Liu PC, & Thiele DJ (1998). Protein chaperones and the heat shock response in *Saccharomyces cerevisiae*. *Current Opinion in Microbiology*, 1(2), 197–203. doi:10.1016/S1369-5274(98)80011-8 [PubMed: 10066474]

- Moreno JA, Radford H, Peretti D, Steinert JR, Verity N, Martin MG, ... Mallucci GR (2012). Sustained translational repression by eIF2 α -P mediates prion neurodegeneration. *Nature*, 485(7399), 507–511. doi:10.1038/nature11058 [PubMed: 22622579]
- Mori K (2009). Signalling pathways in the unfolded protein response: Development from yeast to mammals. *Journal of Biochemistry*, 146(6), 743–750. doi:10.1093/jb/mvp166 [PubMed: 19861400]
- Mori K, Kawahara T, Yoshida H, Yanagi H, & Yura T (1996). Signalling from endoplasmic reticulum to nucleus: Transcription factor with a basic-leucine zipper motif is required for the unfolded protein-response pathway. *Genes to Cells : Devoted to Molecular & Cellular Mechanisms*, 1(9), 803–817. doi:10.1046/j.1365-2443.1996.d01-274.x [PubMed: 9077435]
- Morimoto RI (1998). Regulation of the heat shock transcriptional response: Cross talk between a family of heat shock factors, molecular chaperones, and negative regulators. *Genes & Development*, 12(24), 3788–3796. doi:10.1101/gad.12.24.3788 [PubMed: 9869631]
- Munro S, & Pelham HR (1986). An Hsp70-like protein in the ER: Identity with the 78 kd glucose-regulated protein and immunoglobulin heavy chain binding protein. *Cell*, 46(2), 291–300. doi:10.1016/0092-8674(86)90746-4 [PubMed: 3087629]
- Munro S, & Pelham HR (1987). A C-terminal signal prevents secretion of luminal ER proteins. *Cell*, 48(5), 899–907. doi:10.1016/0092-8674(87)90086-9 [PubMed: 3545499]
- Neef DW, Turski ML, & Thiele DJ (2010). Modulation of heat shock transcription factor 1 as a therapeutic target for small molecule intervention in neurodegenerative disease. *PLoS Biology*, 8(1), e1000291. doi:10.1371/journal.pbio.1000291 [PubMed: 20098725]
- O'Brien D, Jones LM, Good S, Miles J, Vijayabaskar MS, Aston R, ... van Oosten-Hawle P (2018). A PQM-1-mediated response triggers transcellular chaperone signaling and regulates organismal proteostasis. *Cell Reports*, 23(13), 3905–3919. doi:10.1016/j.celrep.2018.05.093 [PubMed: 29949773]
- Ooi FK, & Prahlad V (2017). Olfactory experience primes the heat shock transcription factor HSF-1 to enhance the expression of molecular chaperones in *C. elegans*. *Science Signaling*, 10(501), ea4893. doi:10.1126/scisignal.aan4893 [PubMed: 29042483]
- Pelham HR, & Munro S (1993). Sorting of membrane proteins in the secretory pathway. *Cell*, 75(4), 603–605. doi:10.1016/0092-8674(93)90479-A [PubMed: 8242736]
- Powers ET, Morimoto RI, Dillin A, Kelly JW, & Balch WE (2009). Biological and chemical approaches to diseases of proteostasis deficiency. *Annual Review of Biochemistry*, 78, 959–991. doi:10.1146/annurev.biochem.052308.114844
- Prahlad V, Cornelius T, & Morimoto RI (2008). Regulation of the cellular heat shock response in *Caenorhabditis elegans* by thermosensory neurons. *Science (New York, N.Y.)*, 320(5877), 811–814. doi:10.1126/science.1156093
- Prahlad V, & Morimoto RI (2011). Neuronal circuitry regulates the response of *Caenorhabditis elegans* to misfolded proteins. *Proceedings of the National Academy of Sciences of the United States of America*, 108(34), 14204–14209. doi:10.1073/pnas.1106557108 [PubMed: 21844355]
- Prouty WF, Karnovsky MJ, & Goldberg AL (1975). Degradation of abnormal proteins in *Escherichia coli*. Formation of protein inclusions in cells exposed to amino acid analogs. *The Journal of Biological Chemistry*, 250(3), 1112–1122. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/1089651> [PubMed: 1089651]
- Ray A, Zhang S, Rentas C, Caldwell KA, & Caldwell GA (2014). RTCB-1 mediates neuroprotection via XBP-1 mRNA splicing in the unfolded protein response pathway. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, 34(48), 16076–16085. doi:10.1523/JNEUROSCI.1945-14.2014 [PubMed: 25429148]
- Ritossa F (1996). Discovery of the heat shock response. *Cell Stress & Chaperones*, 1(2), 97–98. doi:10.1379/1466-1268(1996)001<0097:DOTHSR>2.3.CO;2 [PubMed: 9222594]
- Roitenberg N, Bejerano-Sagie M, Bocholez H, Moll L, Marques FC, Golodetzki L, ... Cohen E (2018). Modulation of caveolae by insulin/IGF-1 signaling regulates aging of *Caenorhabditis elegans*. *EMBO Reports*, 19(8), e45673. doi:10.15252/embr.201745673 [PubMed: 29945933]

- Scherz-Shouval R, Santagata S, Mendillo ML, Sholl LM, Ben-Aharon I, Beck AH, ... Lindquist S (2014). The reprogramming of tumor stroma by HSF1 is a potent enabler of malignancy. *Cell*, 158(3), 564–578. doi:10.1016/j.cell.2014.05.045 [PubMed: 25083868]
- Schmeisser S, Priebe S, Groth M, Monajembashi S, Hemmerich P, Guthke R, ... Ristow M (2013). Neuronal ROS signaling rather than AMPK/sirtuin-mediated energy sensing links dietary restriction to lifespan extension. *Molecular Metabolism*, 2(2), 92–102. doi:10.1016/j.molmet.2013.02.002 [PubMed: 24199155]
- Silverman RM, Cummings EE, O'Reilly LP, Miedel MT, Silverman GA, Luke CJ, ... Pak SC (2015). The aggregation-prone intracellular serpin SRP-2 fails to transit the ER in *Caenorhabditis elegans*. *Genetics*, 200(1), 207–219. doi:10.1534/genetics.115.176180 [PubMed: 25786854]
- Singh V, & Aballay A (2012). Endoplasmic reticulum stress pathway required for immune homeostasis is neurally controlled by arrestin-1. *The Journal of Biological Chemistry*, 287(40), 33191–33197. doi:10.1074/jbc.M112.398362 [PubMed: 22875856]
- Sinha A, & Rae R (2014). A functional genomic screen for evolutionarily conserved genes required for lifespan and immunity in germ-line-deficient *C. elegans*. *PLoS One*, 9(8), e101970. doi:10.1371/journal.pone.0101970 [PubMed: 25093668]
- Sitron CS, & Brandman O (2019). CAT tails drive degradation of stalled polypeptides on and off the ribosome. *Nature Structural & Molecular Biology*, 26(6), 450–459. doi:10.1038/s41594-019-0230-1
- Sitron CS, Park JH, & Brandman O (2017). Asc1, Hel2, and Slh1 couple translation arrest to nascent chain degradation. *RNA (New York, N.Y.)*, 23(5), 798–810. doi:10.1261/rna.060897.117
- Sitron CS, Park JH, Giafaglione JM, & Brandman O (2020). Aggregation of CAT tails blocks their degradation and causes proteotoxicity in *S. cerevisiae*. *PLoS One*, 15(1), e0227841. doi:10.1371/journal.pone.0227841 [PubMed: 31945107]
- Somero GN (1995). Proteins and temperature. *Annual Review of Physiology*, 57, 43–68. doi:10.1146/annurev.ph.57.030195.000355
- Stein GM, & Murphy CT (2012). The intersection of aging, longevity pathways, and learning and memory in *C. elegans*. *Frontiers in Genetics*, 3, 259. doi:10.3389/fgene.2012.00259 [PubMed: 23226155]
- Steinbaugh MJ, Narasimhan SD, Robida-Stubbs S, Moronetti Mazzeo LE, Dreyfuss JM, Hourihan JM, ... Blackwell TK (2015). Lipid-mediated regulation of SKN-1/Nrf in response to germ cell absence. *eLife* 2015;4:e07836. doi:10.7554/eLife.07836
- Styer KL, Singh V, Macosko E, Steele SE, Bargmann CI, & Aballay A (2008). Innate immunity in *Caenorhabditis elegans* is regulated by neurons expressing NPR-1/GPCR. *Science (New York, N.Y.)*, 322(5900), 460–464. doi:10.1126/science.1163673
- Sun J, Liu Y, & Aballay A (2012). Organismal regulation of XBP-1-mediated unfolded protein response during development and immune activation. *EMBO Reports*, 13(9), 855–860. doi:10.1038/embor.2012.100 [PubMed: 22791024]
- Sun J, Singh V, Kajino-Sakamoto R, & Aballay A, (2011). Neuronal GPCR controls innate immunity by regulating noncanonical unfolded protein response genes. *Science (New York, N.Y.)*, 332(6030), 729–732. doi:10.1126/science.1203411
- Tatum MC, Ooi FK, Chikka MR, Chauve L, Martinez-Velazquez LA, Steinbusch HWM, ... Prahlad V (2015). Neuronal serotonin release triggers the heat shock response in *C. elegans* in the absence of temperature increase. *Current Biology : CB*, 25(2), 163–174. doi:10.1016/j.cub.2014.11.040 [PubMed: 25557666]
- Taylor RC, Berendzen KM, & Dillin A (2014). Systemic stress signalling: Understanding the cell non-autonomous control of proteostasis. *Nature Reviews. Molecular Cell Biology*, 15(3), 211–217. doi: 10.1038/nrm3752
- Taylor RC, & Dillin A (2013). XBP-1 is a cell-nonautonomous regulator of stress resistance and longevity. *Cell*, 153(7), 1435–1447. doi: 10.1016/j.cell.2013.05.042 [PubMed: 23791175]
- Tracey KJ (2002). The inflammatory reflex. *Nature*, 420(6917), 853–859. doi:10.1038/nature01321 [PubMed: 12490958]

- Tyedmers J, Mogk A, & Bukau B (2010). Cellular strategies for controlling protein aggregation. *Nature Reviews. Molecular Cell Biology*, 11(11), 777–788. doi:10.1038/nrm2993 [PubMed: 20944667]
- van Oosten-Hawle P, Porter RS, & Morimoto RI (2013). Regulation of organismal proteostasis by transcellular chaperone signaling. *Cell*, 153(6), 1366–1378. doi:10.1016/j.cell.2013.05.015 [PubMed: 23746847]
- Vendruscolo M, Knowles TP, & Dobson CM (2011). Protein solubility and protein homeostasis: a generic view of protein misfolding disorders. *Cold Spring Harbor Perspectives in Biology*, 3(12), a010454. doi:10.1101/cshperspect.a010454 [PubMed: 21825020]
- Vihervaara A, Duarte FM, & Lis JT (2018). Molecular mechanisms driving transcriptional stress responses. *Nature Reviews. Genetics*, 19(6), 385–397. doi:10.1038/s41576-018-0001-6
- Vilchez D, Morante I, Liu Z, Douglas PM, Merkwirth C, Rodrigues APC, ... Dillin A (2012). RPN-6 determines *C. elegans* longevity under proteotoxic stress conditions. *Nature*, 489(7415), 263–268. doi:10.1038/nature11315 [PubMed: 22922647]
- Voellmy R (1994). Transduction of the stress signal and mechanisms of transcriptional regulation of heat shock/stress protein gene expression in higher eukaryotes. *Critical Reviews in Eukaryotic Gene Expression*, 4(4), 357–401. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/7734836> [PubMed: 7734836]
- Voellmy R (1996). Sensing stress and responding to stress. *EXS*, 77, 121–137. doi:10.1007/978-3-0348-9088-5_9 [PubMed: 8856972]
- Walrond JP, Kass IS, Stretton AO, & Donmoyer JE (1985). Identification of excitatory and inhibitory motoneurons in the nematode *Ascaris* by electrophysiological techniques. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, 5(1), 1–8. doi:10.1523/JNEUROSCI.05-01-00001.1985 [PubMed: 3965635]
- Walter P (2010). Walking along the serendipitous path of discovery. *Molecular Biology of the Cell*, 21(1), 15–17. doi:10.1091/mbc.E09-08-0662 [PubMed: 20048259]
- Walther DM, Kasturi P, Zheng M, Pinkert S, Vecchi G, Ciryam P, ... Hartl FU (2015). Widespread Proteome Remodeling and Aggregation in Aging *C. elegans*. *Cell*, 161(4), 919–932. doi:10.1016/j.cell.2015.03.032 [PubMed: 25957690]
- Weir HJ, Yao P, Huynh FK, Escoubas CC, Goncalves RL, Burkewitz K, ... Mair WB (2017). Dietary restriction and AMPK increase lifespan via mitochondrial network and peroxisome remodeling. *Cell Metabolism*, 26(6), 884–896 e885. doi:10.1016/j.cmet.2017.09.024 [PubMed: 29107506]
- White JG, Southgate E, Thomson JN, & Brenner S (1983). Factors that determine connectivity in the nervous system of *Caenorhabditis elegans*. *Cold Spring Harbor Symposia on Quantitative Biology*, 48 Pt 2, 633–640. doi:10.1101/sqb.1983.048.01.067 [PubMed: 6586380]
- White JG, Southgate E, Thomson JN, & Brenner S (1986). The structure of the nervous system of the nematode *Caenorhabditis elegans*. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 314(1165), 1–340. doi:10.1098/rstb.1986.0056 [PubMed: 22462104]
- Wolff S, Weissman JS, & Dillin A (2014). Differential scales of protein quality control. *Cell*, 157(1), 52–64. doi:10.1016/j.cell.2014.03.007 [PubMed: 24679526]
- Wolkow CA, Kimura KD, Lee MS, & Ruvkun G (2000). Regulation of *C. elegans* life-span by insulinlike signaling in the nervous system. *Science (New York, N.Y.)*, 290(5489), 147–150. doi:10.1126/science.290.5489.147
- Yin JA, Liu XJ, Yuan J, Jiang J, & Cai SQ (2014). Longevity manipulations differentially affect serotonin/dopamine level and behavioral deterioration in aging *Caenorhabditis elegans*. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, 34(11), 3947–3958. doi:10.1523/JNEUROSCI.4013-13.2014 [PubMed: 24623772]
- Yoshida H, Matsui T, Yamamoto A, Okada T, & Mori K (2001). XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor. *Cell*, 107(7), 881–891. doi:10.1016/S0092-8674(01)00611-0 [PubMed: 11779464]
- Yu Y, Zhi L, Wu Q, Jing L, & Wang D (2018). NPR-9 regulates the innate immune response in *Caenorhabditis elegans* by antagonizing the activity of AIB interneurons. *Cellular & Molecular Immunology*, 15(1), 27–37. doi:10.1038/cmi.2016.8 [PubMed: 27133473]

- Zhang Y, Lanjuin A, Chowdhury SR, Mistry M, Silva-García CG, Weir HJ, ... Mair WB (2019). Neuronal TORC1 modulates longevity via AMPK and cell nonautonomous regulation of mitochondrial dynamics in *C. elegans*. *eLife* 2019;8:e49158doi:10.7554/eLife.49158
- Zhang Q, Wu X, Chen P, Liu L, Xin N, Tian Y, & Dillin A (2018). The Mitochondrial unfolded protein response is mediated cell-non-autonomously by retromer-dependent Wnt signaling. *Cell*, 174(4), 870–883 e817. doi:10.1016/j.cell.2018.06.029 [PubMed: 30057120]

Author Manuscript
Author Manuscript
Author Manuscript
Author Manuscript

Figure 1.
(A) A brief definition of control theory terminology in relation to its use to describe proteostasis. (B) A simplified model of proteostasis. Protein conformation is the *regulated variable* that is maintained at a set-point through the regulated activities of protein biogenesis and protein degradation (controlled variables). CONTROLLERS (here mainly, molecular chaperones) that surveil and monitor protein conformation are present at specific amounts during the normal activities of the cells. They direct the activity of the *controlled variables* through PLANTS (e.g. ribosomes, proteasome, autophagy etc.) to generate homeostatic levels of the *regulated variable* (namely low levels of misfolded proteins). Extra CONTROLLERS can be generated in response to signals that change the levels of the *regulated variable*. However, as explained in the text, increasing the levels of the CONTROLLERS is under neuronal control in *C. elegans*.

Figure 2.

Consequences of neuronal and cephalic control over homeostasis in *C. elegans*. (A) Neuronal circuits through serotonergic, dopaminergic, WNT, FLP-2 and octopaminergic signaling can activate transcription factors that are the master regulators of chaperones, and degradation machinery such as the ubiquitin-proteasome system in distal tissue of *C. elegans*. (B) Prior experience of an environmental threat can enhance the expression of chaperones upon actual encounter with the threat. (C) Maternal release of serotonin upon stress encounter increases the stress tolerance of progeny. (D) Neuronal control enables rapid response to stimuli at the cost of the normal proteostasis machinery.