Published in final edited form as:

Fetal Diagn Ther. 2020 ; 47(12): 912–917. doi:10.1159/000510813.

Surviving Lambs With Myelomeningocele Repaired In Utero With Placental Mesenchymal Stromal Cells for Six Months: A Pilot Study

Kaeli Yamashiro^{1,*}, Laura Galganski¹, Jamie Peyton², Kalie Haynes², Victoria Vicuna², Priyadarsini Kumar^{1,3}, Benjamin Keller¹, James Becker¹, Christopher Pivetti^{1,3}, Sarah Stokes¹, Christina Theodorou¹, Jordan Jackson¹, Aijun Wang^{1,3,4}, Diana Farmer^{1,3}

¹Department of Surgery, University of California-Davis, Sacramento, CA, USA

²Department of Veterinary Medicine, University of California-Davis, Sacramento, CA, USA

³Shriner's Hospitals for Children Northern California, Sacramento, CA, USA

⁴Department of Biomedical Engineering, University of California-Davis, Davis, CA, USA

Abstract

Background: Fetal repair of myelomeningocele (MMC) with placental mesenchymal stromal cells (PMSCs) rescues ambulation in the ovine model up to 48hrs postnatally. Outcomes past 48hrs are unknown as MMC lambs have not been survived past this timepoint.

Objective: We aimed to survive lambs for 6mo following the fetal repair of MMC with PMSCs.

Methods: Fetal MMC lambs were repaired with PMSCs. Lambs received either no additional treatment or postnatal bracing and physical therapy (B/PT). Motor function was assessed with the sheep locomotor rating (SLR). Lambs with an SLR of 15 at birth were survived for 6mo or until a decline in SLR less than 15, whichever came first. All lambs underwent a perimortem MRI.

Results: The lambs with no postnatal treatment (n=2) had SLR declines to 7 and 13 at 29d and 65d respectively, and were euthanized. These lambs had a spinal angulation of 57° and 47°, respectively. The B/PT lamb (n=1) survived to 6mo with a sustained SLR of 15 and a lumbar angulation of 42°.

***Corresponding author:** Kaeli Yamashiro, Department of Surgery, University of California-Davis, 2335 Stockton Blvd, Room 5107, Sacramento, CA 95817, USA, Telephone: 916-734-2724, Fax: 916-734-5633, kjiyamashiro@ucdavis.edu.

Author Contributions

Study conception and design: KY, LG, JP, PK, CP, AW, DF

Data acquisition: KY, LG, KH, VV, BK, JB, CP

Analysis and data interpretation: KY, LG

Drafting of the manuscript: KY

Critical revision: LG, JP, PK, CP, SS, CT, JJ, AW, DF

Statement of Ethics

The views expressed in this material are those of the authors and do not reflect the official policy or position of the U.S. Government, the Department of Defense, or the Department of the Air Force. The animals involved in this study were procured, maintained, and used in accordance with the *Laboratory Animal Welfare Act of 1966*, as amended, and the *Guide for the Care and Use of Laboratory Animals, National Research Council*. The work reported herein was performed under the University of California Davis *Institutional Animal Care and Use Committee* protocol number 19676.

Conflict of Interest Statement

No disclosures.

Conclusion: Postnatal physical therapy and bracing counteracted the inherent morbidity of the absent paraspinal muscles in the ovine MMC model allowing for survival and maintenance of rescued motor function of the prenatally treated lamb up to 6 months.

Keywords

fetal myelomeningocele repair; fetal surgery; placental mesenchymal stromal cells

INTRODUCTION

The Management of Myelomeningocele Study established fetal repair of myelomeningocele (MMC) as the standard of care [1]. Despite significant improvement in motor function, only 55% of patients repaired prenatally were able to walk independently at the 30 month follow up [2]. This inspired a search for a novel therapy to augment the fetal MMC repair to improve distal motor function and other clinical outcomes.

The ovine model is the most commonly used large animal model to study fetal MMC repair as sheep have a robust ability to tolerate multiple fetal interventions [3–7]. It is a surgical model that is morbid by design, with removal of all lumbar paraspinal muscles and lamina to create the defect and prevent innate healing *in utero*. The morbidity of absent paraspinal muscles and protective lamina make survival of the lambs past 48 hours of life challenging. We have previously shown that augmenting the standard fetal MMC repair in the ovine model with early gestation placental mesenchymal stromal cells (PMSCs) engineered in neurogenic media rescues ambulation in lambs up to 48 hours of life [8]. This time point was previously selected as it was easily feasible across the many experiments and provided a consistent point of comparison. However, the durability of the improved motor outcomes past 48 hours of life, as the lambs grow and gain weight similar to humans with MMC, remains unknown.

To determine the durability of the augmented repair, we aimed to survive lambs who underwent fetal MMC repair with PMSCs for 6 months. We hypothesized that the improved motor outcome of the augmented repair was durable for 6 months with improved motor outcome indicated by a motor function score of 15 on the sheep locomotor rating (SLR).

METHODS

Placental mesenchymal stromal cells isolation and seeding

PMSCs were explanted from donated placentas as described by Lankford *et al* [9]. Cells were cultured in media consisting of Dulbecco's Modified Eagle Medium/High Glucose, 5% fetal bovine serum (Hyclone, Thermo Fisher Scientific), 100 units/mL penicillin/100µg/mL streptomycin (Thermo Fisher Scientific) and 20ng/mL recombinant human epidermal growth factor (R&D Systems).

The PMSCs were seeded at passage 5 onto a 6cm x 2cm piece of extracellular matrix (ECM) (Biodesign® Dural graft, Cook Biotech, West Lafayette, IN) the day prior to MMC repair. Each fetus received a different PMSC cell line. A cell density of 500,000 cells/cm² was used for the first lamb and 300,000 cells/cm² for subsequent lambs. The PMSC-ECM product

was incubated in culture media overnight the night prior to surgery. Two different imaging techniques were used. To prepare the ECM for the first two lambs, the PMSCs were transduced with green fluorescent protein (GFP)-containing lentiviral vector (University of California, Davis Stem Cell Center, California Institute of Regenerative Medicine, Sacramento, CA) at passage 4, then imaged the morning of the MMC repair to verify PMSC adherence to the ECM. To prepare the ECM for the third lamb, an additional ECM punch-out was seeded at the same time as the ECM and incubated with the PMSC-ECM product overnight, then stained with Calcein AM (Thermo Fisher Scientific) the morning of the MMC repair to verify PMSC adherence to the ECM.

Ovine myelomeningocele defect creation, repair and Cesarean delivery

Animal work was approved by the Institutional Animal Care and Use Committee (IACUC) and care was in compliance with the Guide for the Care and Use of Laboratory Animals. The facilities used to conduct this study were accredited by the Association for the Assessment and Accreditation of Laboratory Animal Care, International.

As previous described, the fetal MMC defect was created at 75 days of gestation $\pm$ 7 days [10]. The skin, paraspinal muscles, vertebral lamina and dorsal portion of the dura were resected from L1 to L6. A myelotomy was not performed because this study targeted evaluating motor function rather than hindbrain herniation. Lost amniotic fluid was replaced with normal saline and antibiotics (100mg of gentamicin and 1 million units of penicillin) were added to the amniotic fluid at the time of uterine closure.

As previously described, the MMC defect was repaired at 100 days of gestation $\pm$ 7 days [10]. The fibrinous scar over the spinal cord was removed. The PMSC-ECM product was placed on the spinal cord with the cell seeded side down. The ECM was sutured into place and the fetal skin was closed over the ECM. Lost amniotic fluid was replaced with normal saline and antibiotics (100mg of gentamicin and 1 million units of penicillin) were added to the amniotic fluid at the time of uterine closure.

The fetuses were delivered via Cesarean delivery between 142 to 146 days of gestation and, unlike previous studies, the ewes were survived to care for the lambs.

Postnatal assessment

The SLR was used to assess the motor function of the lambs. The SLR is a standardized and validated system to assess sheep locomotor function [7]. A score of 15 indicates normal locomotor function, 10–14 indicates a mild motor deficit, 5–9 indicates a moderate deficit and 4 or less indicates a severe deficit [7]. The SLR was assessed on the first and second days of life, then weekly for the first 2 months and then twice per month until 6 months. Only lambs with an SLR of 15 on the first day of life were selected for the study. If the lamb had a sustained SLR of less than 15 on assessment, it was euthanized prior to 6 months.

A perimortem lumbar MRI was performed on all of the lambs. The MRI images were processed in Horos (Nimble, Annapolis, MD). Spinal angulation was measured in Horos (Nimble, Annapolis, MD) on sagittal imaging. The area of maximum spinal cord

compression was determined on axial imaging by identifying where the spinal cord had the most decreased height and the least amount of surrounding cerebral spinal fluid (CSF).

Addition of physical therapy

The first two lambs were unable to be survived for 6 months so physical therapy and bracing were added for the subsequent lamb. Spinal x-rays of the third lamb were obtained after birth and monthly.

The physical therapy consisted of 1 hour sessions, once to twice daily for 5 days a week with the goal of improving spinal stability through core strengthening and maintaining hindlimb muscle strength. Core and pelvic limb muscle strengthening was achieved through a series of therapeutic exercises including tools such as caveletti poles, cones, wobble boards, and balance discs. To maximize effective therapy, the lamb was also trained to walk and do exercises using a harness. Bracing consisted of wearing a thoracic-lumbar-sacral brace for 8 hours per day for 5 days a week with the goal of providing external bracing support. Initially, a L'il Back Bracer (Caerus Corp, Arden Hills, MN) was used. The straps were secured to the lamb taking care to not be too tight and cause abrasions. A custom brace with the same design was used once the lamb outgrew the L'il Back Bracer (Caerus Corp, Arden Hills, MN).

Data Analysis

The motor function outcome and the MRI results were quantitatively compared between the lambs who had no postnatal therapy and the lamb who underwent postnatal physical therapy and bracing.

RESULTS

Three lambs were survived past 48 hours. Lambs 1 and 2 did not receive postnatal physical therapy and bracing, while lamb 3 did. The characteristics of the gestational ages at defect creation and repair, fetal operative times and defect lengths were similar between the three lambs (Table 1).

All of the lambs had an SLR of 15 at birth. Lamb 1's SLR declined to 7 at 29 days and lamb 2's SLR declined to 13 at 65 days. This is in contrast to the physical therapy and bracing lamb, lamb 3, who survived 6 months with a persistent SLR of 15 (Figure 1).

The amount of spinal cord compression varied between the lambs at the time of euthanasia. At the location of maximum spinal cord compression on MRI, lambs 1 and 2 had minimal CSF around the spinal cord, while lamb 3 had much more CSF around the spinal cord (Figure 2).

The spinal angulation was worse in the lambs who did not receive physical therapy and bracing (lambs 1 and 2) an earlier timepoint than the lamb who received physical therapy and bracing (lamb 3, Table 2). The spinal angulation of lamb 3 gradually increased over time (Figure 3). However, the spinal angulation of lamb 3 at 6 months was still less severe than lamb 1 at 1 month and lamb 2 at 2 months (Figure 4).

DISCUSSION/CONCLUSION

This pilot study is the first description of surviving lambs from the ovine MMC model treated *in utero* with PMSCs. When the inherent morbidity of absent paraspinal muscles in a quadruped is removed through physical therapy and bracing, lambs treated *in utero* with PMSCs can survive for at least 6 months with normal motor function. The augmented repair with PMSCs in the ovine model is durable for at least 6 months.

In addition to the known limitations of this surgical model, we found that the morbidity of absent lumbar paraspinal muscles and lamina allowed for progressive deformation and angulation of the spine limiting survival [6]. However, with the addition of postnatal physical therapy and bracing, this limitation of the model was minimized. This was evidenced by both the decreased spinal angulation of lamb 3 when compared to age-matched timepoints of lambs 1 and 2, and by the final angulation of lamb 3 at 6 months which was much less than lamb 1 at 1 month and lamb 2 at 2 months. Both physical therapy and bracing are used as postnatal treatments for human MMC. Physical therapy can improve a child with MMC's ability to ambulate up to what is expected based on the anatomic level of the defect [11]. Spinal bracing is also used to treat spinal deformities associated with MMC [12]. Physical therapy and bracing are important components of postnatal treatment in the ovine MMC model and in humans with MMC.

The improved motor function outcomes of the ovine model cannot be attributed to physical therapy and bracing alone. In humans, physical therapy maintains the patient's expected functional level based on the anatomic level of the MMC defect, but does not typically improve the patient's functional level beyond the anatomic level, as this was shown for the first time in the MOMS trial with prenatal surgery [1, 13]. All of the lambs had normal hindlimb motor function after birth, as if there were no MMC defect, which was sustained in the lamb who received postnatal physical therapy and bracing. Thus, the role of physical therapy and bracing in the ovine MMC model is to counteract the inherent musculoskeletal morbidity of the model.

There are several limitations of this study. First, only one lamb was treated postnatally with physical therapy and bracing, limiting the power of the study. Second, all of the lambs did not receive the same cell density which could confound the outcomes. Third, spinal imaging was not obtained at baseline or monthly in lambs 1 and 2, so the progressive angulation of these lambs is inferred. Additionally, the spinal imaging was not obtained with the lambs bearing weight, which would more closely approximate functional angulation while ambulating.

Now that MMC lambs can be survived up to 6 months, there are many new areas of research. Current areas of research include the outcomes of the augmented repair on motor, bowel and bladder function up to 6 months. These data are ultimately useful to support an Investigational New Drug Application with the Food and Drug Administration. Finally, human clinical trials are being designed to provide fetal MMC patients with the augmented repair.

In conclusion, this is the first description of the survival of lambs from the ovine MMC model past 48 hours of life. Fetal lambs with MMC repaired *in utero* with PMSCs can be survived for at least 6 months with physical therapy and bracing. The rescued motor function from the *in utero* MMC repair augmented with PMSCs persists for at least 6 months in the ovine model.

Acknowledgement

We would like to thank the veterinarians, technicians and volunteers who provided excellent care for our animals, and Cook Biotech for the donated ECM.

Funding Sources

This work was in part supported by California Institute for Regenerative Medicine (PC1-08103 and CLIN1-11404), Shriner's Hospital for Children (grants #85120-NCA-16, 85108-NCA-19), the NIH (grants #5R01NS100761-02, #1R03HD091601-01). Author LG and CT were also supported by the National Center for Advancing Translational Sciences, National Institutes of Health, through grant number UL1TR001860. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

REFERENCES

1. Adzick NS, et al., A randomized trial of prenatal versus postnatal repair of myelomeningocele. *N Engl J Med*, 2011 364(11): p. 993–1004. [PubMed: 21306277]
2. Farmer DL, et al., The Management of Myelomeningocele Study: full cohort 30-month pediatric outcomes. *Am J Obstet Gynecol*, 2018 218(2): p. 256.e1–256.e13.
3. Moldenhauer JS, In utero repair of spina bifida. *Am J Perinatol*, 2014 31(7): p. 595–604. [PubMed: 24819146]
4. Meuli M, et al., Creation of myelomeningocele in utero: a model of functional damage from spinal cord exposure in fetal sheep. *J Pediatr Surg*, 1995 30(7): p. 1028–32; discussion 1032–3. [PubMed: 7472926]
5. Bouchard S, et al., Correction of hindbrain herniation and anatomy of the vermis after in utero repair of myelomeningocele in sheep. *J Pediatr Surg*, 2003 38(3): p. 451–8; discussion 451–8. [PubMed: 12632366]
6. Vanover M, et al., Spinal Angulation: A Limitation of the Fetal Lamb Model of Myelomeningocele. *Fetal Diagn Ther*, 2019 46(6): p. 376–384. [PubMed: 30970373]
7. Brown EG, et al., Development of a locomotor rating scale for testing motor function in sheep. *J Pediatr Surg*, 2015 50(4): p. 617–21. [PubMed: 25840074]
8. Wang A, et al., Placental mesenchymal stromal cells rescue ambulation in ovine myelomeningocele. *Stem Cells Transl Med*, 2015 4(6): p. 659–69. [PubMed: 25911465]
9. Lankford L, et al., Manufacture and preparation of human placenta-derived mesenchymal stromal cells for local tissue delivery. *Cytotherapy*, 2017 19(6): p. 680–688. [PubMed: 28438482]
10. Galganski LA, et al., In Utero Treatment of Myelomeningocele with Placental Mesenchymal Stromal Cells - Selection of an Optimal Cell Line in Preparation for Clinical Trials. *J Pediatr Surg*, 2019.
11. Charney EB, Melchionni JB, and Smith DR, Community ambulation by children with myelomeningocele and high-level paralysis. *J Pediatr Orthop*, 1991 11(5): p. 579–82. [PubMed: 1918342]
12. Karol LA KE, The Orthopaedic Management of Myelomeningocele. *Operative Techniques in Plastic and Reconstructive Surgery*, 2000 7(2): p. 53–59.
13. Hoffer MM, et al., Functional ambulation in patients with myelomeningocele. *J Bone Joint Surg Am*, 1973 55(1): p. 137–48. [PubMed: 4570891]

Figure 1.

Survival of the lambs. Lamb 1 and 2 survived less than 6 months due to a sheep locomotor rating (SLR) of less than 15. In contrast, lamb 3, who had postnatal physical therapy and bracing, survived the entire 6 months with a consistent SLR of 15.

Figure 2.

Spinal cord compression of lambs 1–3 on MRI at the maximum point of compression. Lamb 1 (Figure 2A) and lamb 2 (Figure 2B) had much less cerebral spinal fluid surrounding the spinal cord at the maximum point of compression compared to lamb 3 (Figure 2C).

Figure 3. Monthly spinal x-rays of lamb 3. The spinal angulation progressively increased each month (mo).

Figure 4. Spinal angulation of the lambs. Lamb 1 (Figure 3A) had an angulation of 57° at 29 days (d), lamb 2 (Figure 3B) had an angulation of 47° at 65d and lamb 3 (Figure 3C) had an angulation of 42° at 6 months (mo).

Table 1.

Characteristics of the ovine myelomeningocele model. The lambs had similar gestational ages (GA), operative times and defect lengths.

	Lamb 1	Lamb 2	Lamb 3
Defect Creation			
GA at defect creation	75 days	78 days	80 days
Number of fetuses	1	1	2
Horn	Left	Left	Right
Fetal operative time at defect creation	30 min	34 min	33 min
Defect creation length	3.2 cm	3.4 cm	3.1 cm
Defect Repair			
GA at defect repair	98 days	101 days	104 days
Fetal operative time at defect repair	54 min	39 min	57 in
Defect repair length	4.5 cm	5 cm	4.4 cm

Table 2.

Spinal angulation of the lambs at age-matched timepoints. The bracing and physical therapy (B/PT) lamb had less spinal angulation than the lambs who did not have B/PT. There were no surviving lambs at 6 months who did not have B/PT.

	No B/PT	B/PT
1 month	Lamb 1 – 57°	Lamb 3 – 19°
2 months	Lamb 2 – 47°	Lamb 3 – 33°
6 months	-	Lamb 3 – 42°