Published in final edited form as:

Wound Repair Regen. 2020 November ; 28(6): 848–855. doi:10.1111/wrr.12849.

Optical imaging of collagen fiber damage to assess thermally injured human skin

Alexandra B. Schroeder, PhD^{1,2,3}, Aos Karim, MD⁴, Edgar Ocotl, BS⁴, Jesús M. Dones, MS⁵, Jenu V. Chacko, PhD³, Aiping Liu, PhD⁴, Ronald T. Raines, PhD⁵, Angela L.F. Gibson, MD, PhD⁴, Kevin W. Eliceiri, PhD^{1,2,3}

¹Department of Medical Physics, University of Wisconsin–Madison, Madison, WI, USA

²Morgridge Institute for Research, Madison, WI, USA

³Laboratory for Optical and Computational Instrumentation, University of Wisconsin–Madison, Madison, WI, USA

⁴Department of Surgery, University of Wisconsin–Madison Hospitals and Clinics, Madison, WI, USA

⁵Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA, USA

Abstract

Surgery is the definitive treatment for burn patients who sustain full thickness burn injuries. Visual assessment of burn depth is made by the clinician early after injury but is accurate only up to 70% of the time among experienced surgeons. Collagen undergoes denaturation as a result of thermal injury; however, the association of collagen denaturation and cellular death in response to thermal injury is unknown. While gene expression assays and histologic staining allow for *ex vivo* identification of collagen changes, these methods do not provide spatial or integrity information *in vivo*. Thermal effects on collagen and the role of collagen in wound repair have been understudied in human burn models due to a lack of methods to visualize both intact and denatured collagen. Hence, there is a critical need for a clinically applicable method to discriminate between damaged and intact collagen fibers in tissues. We present two complementary candidate methods for visualization of collagen structure in three dimensions. Second harmonic generation imaging offers a label-free, high-resolution method to identify intact collagen. Simultaneously, a fluorophore-tagged collagen-mimetic peptide can detect damaged collagen. Together, these methods enable the characterization of collagen damage in human skin biopsies from burn patients, as well as *ex vivo* thermally injured human skin samples. These combined methods could enhance the understanding of the role of collagen in human wound healing after thermal injury and potentially assist in clinical decision-making.

Reprint requests: Kevin W. Eliceiri, Laboratory for Optical and Computational Instrumentation, University of Wisconsin–Madison, 1675 Observatory Drive, Madison, WI 53706-1205, USA, Tel: 608-263-6288, Fax: 608-262-4570, eliceiri@wisc.edu, Angela L.F. Gibson, Department of Surgery, University of Wisconsin–Madison Hospitals and Clinics, 600 Highland Avenue, Madison, WI 53792-3236, USA, Tel: 608-265-9574, Fax: 608-252-0936, gibson@surgery.wisc.edu, Ronald T. Raines, Department of Chemistry, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139-4307, USA, Tel: 617-253-1470, Fax: 617-258-7500, rtraines@mit.edu.

INTRODUCTION

Non-fatal thermal injuries are a leading cause of morbidity worldwide, with roughly 40,000 hospitalizations a year required in the United States.¹ While considered largely preventable, burns can leave patients with debilitating injuries and scarring.² A major challenge lies in the variable healing time for burns that are less than full thickness, along with the disparity between clinical identification of burn depth, histopathologic determination of cellular and extracellular matrix (ECM) damage and the regenerative potential of the tissue.³ More severe, full thickness burns that damage the subcutaneous layer require excision and autologous skin grafts while dermal burns may heal spontaneously if thermal damage is confined to the more superficial dermis.^{2,4} Surgical removal of necrotic tissue is necessary to remove the inflammatory burden, allowing viable cells to regenerate. Conversely, over-excision of burns can result in scarring and morbidity in the form of donor site wounds, as well as increase the cost of care incurred by the patient.² In an attempt to understand the potential for wound healing, tissue excised from the initial surgical intervention may be analyzed through gene expression assays and histological staining. Gene expression can identify the types of collagen expressed but this method does not provide spatial information regarding collagen structure. Histologic stains like Picrosirius red⁵ and Masson's trichrome⁶ can identify collagen structures and organization but only on a thin, single slice of tissue *ex vivo*. These methods are not applicable for identifying damaged collagen on a patient in a clinical setting. Additionally, the exact depth of burn injury that will heal without surgery and the association between cellular and collagen damage is unknown. There is a critical need for developing robust minimally-invasive techniques that are clinically translatable to quantify the extent of thermal damage to collagen in the ECM in three dimensions, and specifically visualizing denatured collagen is a significant first step.

Collagen is the most abundant protein in the body, comprising 75% of the dry weight of the skin.⁷ Collagen provides the structural scaffold of the ECM upon which cellular epithelialization takes place. It also serves a significant role within the dermal layer of the skin for cell adhesion and migration.^{8,9} Our group has shown that this migration potential can be hijacked during metastasis in breast and ovarian cancer, such that collagen structure and alignment can be a prognostic indicator of cancer progression,^{10,11} implicating collagen as a significant cell behavior mediator. During wound repair, deposition of collagen occurs alongside regeneration of the ECM thereby playing an important role in determining morphological skin properties.⁹ Thermal injuries result in collagen denaturation and structural changes.¹² Despite efforts to visualize architectural changes of collagen in the ECM in wound models, the clinical impact of collagen assessment for determining burn severity or depth after burn injury is still relatively unexplored.¹³ Characterization of the wound healing response from thermal injury in a zebrafish model indicates a dramatic loss of collagen fibers associated with delayed healing when compared to a transection wound.¹⁴ With better visualization of collagen damage *in vivo* and an understanding of the relationship between this damage and cellular viability, therapeutic development could be directed toward collagen repair. The current inability to assess the role of collagen in thermal injury is due largely to the lack of available methods to visualize damaged collagen *in vivo* and a lack of human models of thermal injury.

Clinically, understanding the changes to collagen architecture due to thermal injury is important, as collagen plays a major role in the mechanical integrity of the skin and the resultant functional outcomes and scarring. Tanaka *et al.* performed second harmonic generation (SHG) imaging on *in vivo* samples to observe changes in dermal collagen fibers in living rat burn models.¹² SHG occurs when two incident photons interact with the non-centrosymmetric triple-helical structure of collagen and combine to form a single emitted photon of exactly half the wavelength or twice the energy, offering a label-free, quantitative measure of intact collagen with high spatial resolution.^{10,11,12} However, SHG has several issues that limit its clinical application including being expensive, bulky, utilizes non eye-safe class IV lasers and does not currently have US Food and Drug Administration (FDA) approval.¹¹ Therefore, exploring other imaging techniques to specifically identify denatured collagen may prove useful during burn excision.

Currently, approximately 80% of burn surgeons in the United States determine burn depth by visualization alone, an assessment method that is accurate in only 64–76% of cases.^{2,15} Furthermore, indeterminate burn depths—the burn depth at which decisions about surgical intervention necessity is least clear—are even less accurately identified.^{16,17} The next most common method to determine healing potential of a thermal injury is Laser Doppler imaging (LDI),^{18,19} which has a 95–100% positive predictive value for prognosing healing within three weeks, increasing over time as does visual interpretation.²⁰ Use of LDI is limited by day-to-day and intra-operator differences resulting from training variability.²⁰ Furthermore, LDI is not useful intraoperatively during excision when the wound is excised under tourniquet as the blood flow is artificially decreased. Optical Coherence Tomography (OCT) offers high-resolution, multi-sectional imaging but the long acquisition times limit the applicability.²¹ Fluorescence-guidance-based techniques can be used to assess wounds, but the availability of specifically targeted probes and instrumentation is limited. The clinically approved fluorescence agent Indocyanine green (ICG), is used for determining vascular flow during reconstructive surgery. However, it can be challenging to interpret when applied for burn cases.²² The use of perfusion as a readout for viability is particularly challenging where tourniquets or vasoconstriction injections are used, as is common intraoperatively. Furthermore, ICG does not provide any information regarding collagen structural damage to the ECM.^{3,23} Visualization of ICG requires injection and is more invasive than potentially non-intravenous alternatives that may identify thermal collagen damage more explicitly, such as a fluorescently tagged collagen-mimetic peptide.

Collagen-mimetic peptides (CMPs) are synthetic chains of amino acid residues that replicate the strands in natural collagen fibers.⁷ CMPs can bind to damaged collagen and have been used for several purposes, including identifying atherosclerotic plaques.²⁴ In previous work, we synthesized a CMP that contains three types of residues: (2*S*,4*S*)-4-fluoroproline (flp), (2*S*,4*R*)-4-hydroxyproline (Hyp), and glycine. This CMP, (flpHypGly)₇,²⁵ anneals to damaged collagen but not to itself.²⁶ By appending a cyanine 5 (Cy5) dye, we imaged damaged collagen in a burned rat tendon by excitation ($\lambda_{\text{ex}} = 652 \text{ nm}$) using either multiphoton or single-photon fluorescence microscopy, and capturing emission ($\lambda_{\text{em}} = 672 \text{ nm}$).²⁶ We validated the binding of the Cy5-CMP conjugate specifically to damaged collagen through SHG imaging, which generates a signal only for intact collagen, and observed that Cy5-CMP does not bind to adjacent intact collagen.²⁶

Here, we probe collagen damage following thermal injury in human skin by leveraging signals obtained from the fluorescence of Cy5-CMP adhered to denatured collagen, combined with SHG imaging of intact collagen. We show that Cy5-CMP binds specifically to regions of denatured collagen in human *in vivo* clinical burn biopsies and provides contrast from normal tissue. The images add insight alongside histologic staining with lactate dehydrogenase (LDH) for cellular viability in an *ex vivo* thermally injured skin model. Finally, we demonstrate the potential for Cy5-CMP to be visualized in the clinic through the use of single-photon fluorescence imaging.

METHODS

Cy5-Collagen mimetic peptide synthesis, preparation and validation

A Cy5-CMP conjugate, Cy5-Gly-(Ser-Gly)₂-(flpHypGly)₇, was prepared as described in Dones *et al.*²⁶ Briefly, the peptide was synthesized by microwave-assisted solid-phase peptide synthesis, purified by reversed-phase HPLC, and conjugated at its N terminus to Cy5. The specific binding of Cy5-CMP to damaged collagen was validated by using a Cy5-CI conjugate, Cy5-Gly-(Ser-Gly)₂-(HypflpGly)₇ (a compositional isomer), as a negative control. Cy5-CI contains the same three amino acid residues as does Cy5-CMP but arranged in an order that does not allow for specific annealing to damaged natural collagen.²⁶ Washing of residual unbound Cy5-CMP or Cy5-CI from human skin samples was performed with 1× PBS (Sigma, St. Louis, MO) (see Supplemental Data, Figure S1).

Clinical *in vivo* patient selection and tissue preparation

Clinical tissue samples were obtained through the University of Wisconsin Hospital as described previously.³ This study was approved by the University of Wisconsin Human Subjects Committee Institutional Review Board in compliance with the 1975 Declaration of Helsinki. Briefly, samples were obtained from patients with indeterminate depth burn injury at the time of surgery, 6–13 days after burn injury following a period of observation to assess healing trajectory. Intraoperatively, burn wound depth was determined by the burn surgeon using visual assessment prior to and during excision.³ Normal skin tissue used as a control was obtained from non-burn related elective reconstructive surgery. Control non-burned and burned tissue samples were stored for up to 1 hour in normal saline (Baxter) prior to further processing. A 6-mm biopsy punch was taken from the skin samples and 100 µL of 100 µM Cy5-CMP or 100 µL of 100 µM Cy5-CI was applied directly onto the superficial side of the biopsy, fully submerged in dye solution, and incubated in a petri dish covered with aluminum foil at room temperature for 1 hour. The samples (n=6 burn pairs) were washed three times in rapid succession with 1× PBS (Sigma) prior to imaging and kept covered in 1× PBS. Images were collected from both superficial and deep surfaces of the biopsies immediately following rinsing.

Tissue sample preparation

Normal skin tissue for use in the *ex vivo* model was obtained from patients receiving elective surgery at the University of Wisconsin Hospital (Madison, WI). Discarded de-identified tissue was transported in normal saline after surgical excision.

Thermal injury, tissue culture and Cy5-CMP labeling

A novel burn device was created in collaboration with the Morgridge Institute for Research FabLab group (Madison, WI) (unpublished results). Briefly, the device generated a burn on the *ex vivo* skin with an 8 mm circular face applied with uniform pressure of 0.12 kg/cm² at 150°C for 3 seconds to simulate a partial thickness burn. After thermal injury, the tissues (including non-burn control skin) were biopsied using a 6 mm punch and cultured in Dulbecco's Modified Eagle's Medium (Gibco) supplemented with 10% bovine calf serum (Hyclone), 0.625 µg/ml amphotericin B (Gemini Bio Products), and 100 µg/ml of Pen/Strep (Gibco by Life Technologies) at 37°C for 48 hours prior to Cy5-CMP application and imaging. After 48 hours, the necrotic epidermis on the 3s samples was mechanically debrided with a forceps to expose the dermis. All biopsies received 100 µL of 100 µM Cy5-CMP, were fully submerged and incubated for 1 hour, and washed 3 times with 1× PBS, as described above. After Cy5-CMP application, incubation and washing, the 6 mm *ex vivo* thermally injured and debrided biopsies were bisected. One-half of the biopsy was used for multiphoton imaging of Cy5 and SHG, the other half was cryopreserved and sectioned for lactate dehydrogenase (LDH) staining.

Lactate dehydrogenase staining on histological sections

Lactate dehydrogenase (LDH) staining was performed on cryosections of *ex vivo* tissue sections to assess cell viability. Staining was performed as described in Gibson *et al.*²⁷ Briefly, sections were air-dried for a minimum of 3 hours (range 3–18 hours) then washed twice with PBS for 5 minutes each time. Sections were incubated with freshly prepared LDH solution containing 5% Polypep (Sigma); 2 mM Gly-Gly (Sigma); 0.75% NaCl (Fisher, Hampton, NH); 60 mM lactic acid (Dot Scientific, Burton, MI); 1.75 mg/ml β-nicotinamide adenine (Sigma, St. Louis, MO); and 3 mg/ml Nitroblue Tetrazolium (Sigma) pH 8.0, for 3.5 hours at 37°C. Slides were washed twice for 2 minutes each with 50°C tap water, followed by two washes with PBS for 2 minutes each. Tissues were counterstained with aqueous eosin (Newcomer Supply, Middleton, WI) for 4 minutes. Slides were washed with PBS for 1 second, dehydrated with acetone for 30 seconds followed by acetone:xylene (1:1) for 1 minute, and finally with xylene alone for 1 minute. Slides were then cover slipped with Cytoseal 60 (Thermo Fisher Scientific).

Multiphoton excitation of Cy5 and second harmonic generation

All imaging was performed on the Bruker Ultima IV multiphoton microscope system described in Dones *et al.*, 2019.²⁶ A Ti:Sapphire pulsed laser (Spectra-Physics Insight DeepSee) was tuned to 820 nm to excite the Cy5 fluorophore of Cy5-CMP, using a 690 (50) nm emission bandpass filter (Chroma Technology Corporation, VT) to isolate the fluorescence emission. SHG of intact collagen was performed on all tissues on the same microscope by tuning the laser wavelength to 890 nm and using a 445 (50) nm bandpass filter (Semrock Inc., Rochester, NY). Samples were imaged using a 10× CFI Super Fluor objective lens (Nikon, NA = 0.50). Images generated were acquired as z-stacks from superficial to deep at 40 micron intervals spanning 800–1200 microns in depth. Slices were summed and stitched to capture the full field of view using FIJI.²⁸ A high pass Gaussian

filter was used to correct for gradients in the images in Figures 3 and 4, which were taken with a 4× Super Fluor objective (Nikon, NA = 0.20).

Wide field fluorescence with CMOS camera

Wide field fluorescence imaging was performed on Cy5-CMP applied to a 6-mm *ex vivo* human skin biopsy that was burned as described above. A PhotoFluor II mercury halogen lamp (Chroma) excited the Cy5-CMP fluorescence through a Cy5.5 dichroic cube (Chroma) consisting of a 685 nm long pass beam splitter, a 650 (50) nm bandpass excitation filter and a 720 nm (50) nm bandpass emission filter. The resulting signal was captured with a complementary metal-oxide semiconductor (CMOS) camera (Digital Sight DS-U3, Nikon) using a 4× Super Fluor objective (Nikon, NA = 0.20) on the same Bruker Ultima microscope as the multiphoton imaging was performed. A non-burned biopsy labeled with Cy5-CMP was used as a control. Images generated were stitched to capture the full field of view using FIJI.²⁸

RESULTS

Collagen mimetic peptides are specific to sites of collagen denaturation in excised human burn tissue

Tissues obtained from patients who sustained thermal injuries, diagnosed at the time of excision (6–13 days after burn injury) as deep partial thickness,³ were imaged after treatment with Cy5-CMP (n=6). Cy5-CI washes from the denatured collagen and the fluorescence intensity of Cy5-CMP was 25 times greater at the burn site compared to neighboring intact collagen as identified by SHG imaging. Multiphoton images taken of the deep dermal surfaces of human burn tissue reveal that Cy5-CMP binds specifically to regions of denatured collagen as validated by the inverse relationship of these images to those obtained using SHG imaging, which detects only intact collagen (Figure 1A–C). Burn depth is heterogeneous within a given wound; therefore, during surgical excision of burn wounds it is expected that the deep surface of an excision will have a mixture of intact and denatured collagen present. Figure 1B illustrates this phenomenon where the SHG imaging identifies the intact collagen, while the Cy5-CMP image correlates with the low-signal regions in the SHG image (Figure 1A and B, yellow circles), representing regions of denatured collagen that remained after surgical excision. In this case, Cy5-CMP was able to clearly identify and confirm that regions of collagen damage were residually present. These data show the potential for Cy5-CMP to be used to identify regions of damaged collagen resulting from thermal injury in human patients.

Cy5-CMP and SHG on *ex vivo* thermally injured human tissue

Using an *ex vivo* human burn injury model, we qualitatively evaluated Cy5-CMP penetration on thicker sections in experimentally generated burns by generating a histogram of pixel intensities from Cy5 fluorescence throughout the sample as measured in FIJI²⁸, as well as determined the relationship between collagen denaturation and cell viability with LDH-staining (Figure 2). Deep partial thickness thermal injuries were generated in *ex vivo* human skin. Burned tissue was cultured for 48 hours prior to Cy5-CMP application and imaging to simulate the clinically relevant period of time in which initial depth determination for

operative planning occurs. The non-burned control (Figure 2A) shows very little Cy5 signal, with no specific binding to the intact dermal collagen, as determined by the correlation with the SHG signal (Figure 2B). LDH-staining shows cell viability, that is blue color, throughout the depth of the tissue (Figure 2D). The 3s burns show higher Cy5-CMP uptake where SHG signal is low and, conversely, low Cy5-CMP binding where SHG is brighter (Figure 2E–G). This region of intact collagen appears to localize near the region with viable cells identified with LDH-staining (Figure 2H). This suggests that, following thermal injury, the location of viable cells correlates with the areas of intact collagen.

Wide field fluorescence excitation of Cy5-CMP can distinguish between normal skin and thermally injured skin

To demonstrate that visualization of Cy5-CMP would not be limited to multiphoton excitation, which would greatly curtail any clinical application given the cumbersome methods needed for image acquisition, we imaged the superficial surface of a control non-burned and *ex vivo* thermally injured human skin sample with Cy5-CMP using wide field fluorescence excitation (Figure 3). Using a mercury-arc light source, we found intensely bright Cy5 fluorescence on the superficial face of an *ex vivo* thermal injury post Cy5-CMP application when compared to the non-burned control (Figure 3A and B). Alongside the wide field images, we captured images on the same microscope platform with multiphoton microscopy for the same sample conditions of non-burned control and thermally injured skin (Figure 3C and D). Wide field fluorescence excitation provides sufficient contrast to substantially excite Cy5-CMP and discriminate burned from non-burned human skin.

DISCUSSION

Collagen plays a significant role in maintaining the structural integrity of human skin as it provides a scaffold for cells to migrate and regenerate in the wound healing process.^{7–9} Currently, there are few clinically viable imaging techniques to probe collagen, and fewer still that permit real-time visualization of regions of denatured collagen. SHG offers the ability to image intact collagen without exogenous contrast at a high spatial resolution. Tanaka *et al.* assessed whether a change in SHG signal could be associated with burn degree, where the vanishing of the signal corresponds to denatured collagen, and proposed a quantitative parameter for burn-assessment based on the depth profile of the mean SHG intensity across the entire image.¹² Pierce *et al.* used polarization-sensitive optical coherence tomography (OCT) to show that partial thickness burns had a loss of collagen birefringence due to denaturation.²⁹ These previous studies suggest that collagen denaturation can be quantified with SHG and OCT and correlated to burn severity. Still, both techniques can only image intact collagen rather than identifying regions of damage. Our results show that in clinical burn tissue, Cy5-CMP specifically labels damaged collagen. SHG coupled with multiphoton fluorescence excitation provide an opportunity to validate the Cy5 signal from Cy5-CMP on the same multiphoton microscope platform.

We aimed to test whether a thermal injury can be assessed using Cy5-CMP and whether the binding of Cy5-CMP correlated with injury using an *ex vivo* skin model of burn injury. The use of a dye to localize denatured and potentially non-viable tissue will allow more precision

with removal of necrotic tissue in real-time during surgical excision or bedside wound debridement compared to the current standard of care which is visual evaluation of tissue characteristics, thought further testing would be necessary to facilitate clinical implementation. Future studies would be directed to understand the mechanism of how the temperature of the wound affects collagen damage and how this damage correlates further with cellular viability in physiological models. It is likely that, at certain temperatures, cells are more sensitive to damage than collagen is to denaturation, but the basis for this observation requires future investigation. *Ex vivo* skin also lacks the vascular flow of blood and lymphatics present in the clinical *in vivo* patient samples. This perfusion undoubtedly plays a crucial role in wound repair and may affect collagen response to injury. Additionally, burns are a highly heterogeneous even within the adjacent regions on the same patient. Because of this, we reported on relative contrast in fluorescence signal between non-burned regions and burned regions, as a numerical rubric for image intensity may not directly correlate to clinical relevance. Further, it would be beneficial to explore other dye delivery solvents, such as DMSO, that might improve the ability for the superficial application of Cy5-CMP to penetrate the skin surface.

Widefield fluorescence imaging offers the capability to visualize subcellular structures and dynamic biological processes with high spatial resolution in the clinic.³⁰ While multiphoton excitation successfully allowed us to visualize Cy5-CMP-labeled damaged collagen, in conjunction with the intact collagen structures provided by SHG, neither modality is currently clinically available. Therefore, we showed that the far-red fluorescence of Cy5-CMP can be captured using the more readily available widefield fluorescence with a mercury arc lamp and appropriate filters with a CMOS camera system. Such a widefield imaging instrumentation setup is potentially suitable for fluorescence-guided surgery, whereas, as mentioned previously, SHG is not a clinically available modality. Cy5-CMP may provide a clinically viable labeling methodology for assessing the extent of collagen damage following thermal injury, and commercial fluorescence-guided imaging systems can potentially capture a Cy5-CMP signal.³⁰

Conclusion

Cy5-CMP anneals to the damaged collagen of thermally injured skin while washing away from skin with intact collagen. This peptide could be beneficial in conjunction with current histologic methods to assist in visualizing locations of collagen denaturation as a result of thermal injury. This integrative approach would allow the correlation of collagen and cellular damage, which could help to illuminate the molecular mechanism of injury progression and wound repair following thermal injury. Moreover, Cy5-CMP could serve as the basis for the intraoperative labeling of thermally damaged tissue to be removed during burn excision. The utility of seeing a signal corresponding directly to denatured collagen is that directed excision of necrotic tissue would be an improvement to the visual characteristics that currently lead to over-excision of burn wounds.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

We thank Dr. Jayne Squirrel for useful comments on the manuscript and Josiah Wolf for assistance with the burn device. We acknowledge funding from the Laboratory for Optical and Computational Instrumentation (LOCI), the Morgridge Institute for Research (KWE), the UW Department of Surgery (AG), and the National Institute of Arthritis and Musculoskeletal and Skin Diseases (RTR).

References

1. World Health Organization. Burns Fact Sheet. 3 6, 2018 (accessed July 29, 2019).
2. Paul DW, Ghassemi P, Ramella-Roman JC, Prindeze NJ, Moffatt LT, Alkhalil A, et al. Noninvasive imaging technologies for cutaneous wound assessment: a review. *Wound Repair Regen* 2015; 23(2): 149–62. [PubMed: 25832563]
3. Karim AS, Yan A., Ocotl E, Bennett DD, Wang Z, Kendzierski C, et al. Discordance between histologic and visual assessment of tissue viability in excised burn wound tissue. *Wound Repair Regen* 2018; 27(2): 150–61. [PubMed: 30585657]
4. Kowalske KJ, Burn wound care. *Phys Med Rehabil Clin N Am* 2011; 22: 213–27. [PubMed: 21624717]
5. Hernandez P, Buller D, Mitchell T, Wright J, Liang H, Manchanda K, et al. Severe Burn-Induced Inflammation and Remodeling of Achilles Tendon in a Rat Model. *Shock* 2018; 50(3): 346–50. [PubMed: 29065066]
6. Zhai Q, Zhou F, Ibrahim MM, Zhao J, Liu X, Wu J et al. An immune-competent rat split thickness skin graft model: useful tools to develop new therapies to improve skin graft survival. *Am J Transl Res* 2018;10(6): 1600–10. [PubMed: 30018703]
7. Shoulders MD, and Raines RT, Collagen structure and stability. *Annu Rev Biochem* 2009; 78: 929–58. [PubMed: 19344236]
8. Van Goethem E, Poincloux R, Gauffre F, Maridonneau-Parini I, Le Cabec V. Matrix architecture dictates three-dimensional migration modes of human macrophages: differential involvement of proteases and podosome-like structures. *J Immunol* 2010; 184: 1049–61. [PubMed: 20018633]
9. Morris JL, Cross SJ, Lu Y, Kadler KE, Lu Y, Dallas SL, et al. Zebrafish: Live imaging of collagen deposition during skin development and repair in a collagen I – GFP fusion transgenic zebrafish line. *Dev Biol* 2018; 441(1): 4–11. [PubMed: 29883658]
10. Conklin MW, Eickhoff JC, Riching KM, Pehlke CA, Eliceiri KW, Provenzano PP, et al. Aligned collagen is a prognostic signature for survival in human breast carcinoma. *Am J Pathol* 2011; 178(3): 1221–32. [PubMed: 21356373]
11. Tilbury K, Campagnola PJ. Applications of second-harmonic generation imaging microscopy in ovarian and breast cancer. *Perspect Medic Chem* 2015; 7: 21–32.
12. Tanaka R, Fukushima S, Sasaki K, Tanaka Y, Murota H, Matsumoto T, et al. *In vivo* visualization of dermal collagen fiber in skin burn by collagen-sensitive second-harmonic-generation microscopy. *J Biomed Optics* 2013; 18(6): 061231–1–8.
13. Chvapil M, Speer DP, Owen JA, Chvapil TA. Identification of the depth of burn injury by collagen stainability. *Plast Reconstr Surg* 1984; 73(3):438–441. [PubMed: 6199804]
14. Miskolci V, Squirrel J, Rindy J, Vincent S, Sauer JD, Gibson ALF, et al. Distinct inflammatory and wound healing responses to complex caudal fin injuries of larval zebrafish. *ELife* 2019; 8: e45976. [PubMed: 31259685]
15. Jaskille AD, Ramella-Roman JC, Shupp JW, Jordan MH, Jeng JC. Critical review of burn depth assessment techniques: part II. Review of laser Doppler technology. *J Burn Care Res* 2010; 31(10): 151–7. [PubMed: 20061851]
16. Heimbach D, Engrav L, Grube B, Marvin J. Burn depth: a review. *World J Surg* 1992; 16: 10–15. [PubMed: 1290249]
17. Monstrey S, Hoeksema H, Verbelen J, Pirayesh A, Blondeel P. Assessment of burn depth and burn wound healing potential. *Burns* 2008; 34: 761–769. [PubMed: 18511202]

18. Khatib M, Jabir S, Fitzgerald O'Connor E, Philp B. A systematic review of the evolution of laser Doppler techniques in burn depth assessment. *Plast Surg Int* 2014; 2014: 621792. [PubMed: 25180087]
19. Israel JS, Greenhalgh DG, Gibson AL. Variations in burn excision and grafting: a survey of the American burn association. *J Burn Care Res* 2017; 38: 125–132. [PubMed: 28253212]
20. Sharma VP, O'Boyle CP, Jeffery SL. Man or machine? The clinimetric properties of laser Doppler imaging in burn depth assessment. *J Burn Care Res* 2011; 32: 143–9. [PubMed: 21107272]
21. Thatcher JE, Li W, Rodriguez-Vaqueiro Y, Squiers JJ, Mo W, Lu Y, et al. Multispectral and Photoplethysmography optical imaging techniques identify important tissue characteristics in an animal model of tangential burn excision. *J Burn Care Res* 2016; 37: 38–52. [PubMed: 26594863]
22. McUmber H, Dabek RJ, Bojovic B, Driscoll DN. Burn depth analysis using indocyanine green fluorescence: a review. *J Burn Care Res* 2019; 40(4): 513–6. [PubMed: 31046089]
23. Kaiser M, Yafi A, Cinat M, Choi B, Durkin AJ. Noninvasive assessment of burn wound severity using optical technology: a review of current and future modalities. *Burns* 2011; 37: 377–86. [PubMed: 21185123]
24. Megens RTA, oude Egbrink MGA, Cleutjens JPM, Kuijpers MJE, Schiffrers PHM, Merckx M, Slaaf DW, et al. Imaging collagen in intact viable healthy and atherosclerotic arteries using fluorescently labeled CNA35 and two-photon laser scanning microscopy. *Mol Imaging* 2007; 6: 247–60. [PubMed: 17711780]
25. Barth D, Milbradt AG, Renner C, Moroder L. A (4R)- or a (4S)-fluoroproline residue in position Xaa of the (Xaa-Yaa-Gly) collagen repeat severely affects triple-helix formation. *ChemBioChem* 2004; 5: 79–86. [PubMed: 14695516]
26. Dones JM, Tanrikulu IC, Chacko JV, Schroeder AB, Hoang TT, Gibson ALF, et al. Optimization of interstrand interactions enables burn detection with a collagen-mimetic peptide. *Org Biomol Chem* 2019.
27. Gibson AL, Shatadal S. A simple and improved method to determine cell viability in burn-injured tissue. *J Surg Res* 2017; 215: 83–7. [PubMed: 28688666]
28. Schindelin J, Arganda-Carreras I, Frise E. et al. Fiji: an open-source platform for biological-image analysis. *Nat Methods* 2012; 9(7): 676–82. [PubMed: 22743772]
29. Pierce MC, Sheridan RL, Park BH, Cense B, de Boer JF. Collagen denaturation can be quantified in burned human skin using polarization-sensitive optical coherence tomography. *Burns* 2004; 30(4): 511–7. [PubMed: 15302415]
30. Zhang R, Schroeder AB, Grudzinski JJ, Rosenthal EL, Warram JM, Pinchuk AN, et al. Beyond the margins: real-time detection of cancer using targeted fluorophores. *Nat Rev Clin Oncol* 2017; 14(6): 347–64. [PubMed: 28094261]

Figure 1:

Specificity of Cy5-CMP to denatured collagen in patient burn tissue. Stitched z-projections of multiphoton micrographs of (A) Cy5 fluorescence image with regions of interest (ROIs) (yellow circles) showing Cy5-CMP labeling on the deep surface (en face view) of 6 mm biopsies of tangentially excised deep partial thickness burn tissue, and (B) absence of SHG signal in the same ROIs (yellow circles). (C) Composite image of Cy5 fluorescence and SHG. The region of brighter Cy5 fluorescence near the right side of the image corresponds to a portion of the cut edge of the biopsy where we see some fluorescence signal from the superficial side of the biopsy penetrating through the z stacked images. (D) Burn tissue sample labeled with Cy5-CI (negative control) imaged with Cy5 parameters and (E) SHG parameters. Yellow ROIs and (F) the composite image of tissue labeled with Cy5-CI indicating that the dark regions in panel E are not fluorescent in panel D. The Cy5 fluorescence on the circumference of the biopsy corresponds to non-specific binding around the superficial cut edge of the biopsy. Scale bars = 1 mm. Color version is available in the online copy.

Figure 2:

Cy5 excitation of Cy5-CMP reveals collagen damage with depth on cross-sectional views. Shown are the side profiles of a 6 mm biopsy of (A-D) normal non-burned skin and (E-H) skin burned for 3s at 150 °C with a custom burn device. (A) Multiphoton excitation of Cy5-CMP shows little fluorescence while (B) SHG image reveals intact collagen throughout the normal biopsy. In contrast, (E) Cy5-CMP extensively labels the burned sample while the (F) SHG signal is minimal. SHG signal exhibits an inversion of signal intensity to the Cy5 channel, highlighted by the composite images of (C) Cy5-CMP + SHG for normal skin and (G) 3s burned skin. LDH images show that viable cells (blue) are present throughout the (D) non-burned skin and in the (H) deep portion in the 3s burned skin, correlating with intact collagen. Scale bar = 1 mm. Color version is available in the online copy.

Figure 3:

Assessment of Cy5-CMP labeling using wide field fluorescence microscopy. Fluorescence of an *ex vivo* 6 mm biopsy from (A) normal non-burned skin and (B) burned skin relative to the same samples imaged with multiphoton microscopy in (C) non-burned and (D) burned skin. Scale bar = 1 mm. Color version is available in the online copy.