Published in final edited form as:

Handb Exp Pharmacol. 2020 ; 258: 147–165. doi:10.1007/164_2019_268.

THE RISE AND FALL OF KAPPA-OPIOID RECEPTORS IN DRUG ABUSE RESEARCH

Matthew L. Banks

Department of Pharmacology and Toxicology, Virginia Commonwealth University

Abstract

Substance use disorders represent a global public health issue. This mental health disorder is hypothesized to result from neurobiological changes as a result of chronic drug exposure and clinically manifests as inappropriate behavioral allocation towards the procurement and use of the abused substance and away from other behaviors maintained by more adaptive nondrug reinforcers (e.g. social relationships, work). The dynorphin/kappa-opioid receptor (KOR) is one receptor system that has been altered following chronic exposure to drugs of abuse (e.g. cocaine, opioids, alcohol) in both laboratory animals and humans; implicating the dynorphin/KOR system in the expression, mechanisms, and treatment of substance use disorders. KOR antagonists have reduced drug self-administration in laboratory animals under certain experimental conditions, but not others. Recently, several human laboratory and clinical trials have evaluated the effectiveness of KOR antagonists as candidate pharmacotherapies for cocaine or tobacco use disorder to test hypotheses generated from preclinical studies. KOR antagonists failed to significantly alter drug use metrics in humans suggesting translational discordance between some preclinical drug self-administration studies and consistent with other preclinical drug self-administration studies that provide concurrent access to an alternative nondrug reinforcer (e.g. food). The implications of this translational discordance and future directions for examining the therapeutic potential of KOR agonists or antagonists as candidate substance use disorder pharmacotherapies are discussed.

1. Preclinical Evaluation of Candidate Substance Use Disorder Treatments

Substance use disorders (SUDs; i.e. drug addiction) are an insidious and global public health issue. This complex and multi-faceted mental health disorder is most commonly modeled in the laboratory using a drug self-administration (SA) procedure to provide an opportunity to measure volitional drug intake. Both preclinical and human laboratory drug SA procedures have made significant contributions to improving our understanding of psychoactive compounds for more than 50 years. In general, preclinical drug SA procedures are used to address two main categories of scientific questions. One category is for abuse liability assessment of psychoactive compounds for potential scheduling as controlled substances by the Drug Enforcement Agency, and there are already excellent reviews on the utility of drug SA procedures (Ator and Griffiths 2003; Carter and Griffiths 2009). The other category is for understanding the expression, mechanisms, and treatment of drug-taking behavior as a

model of SUDs. This chapter will focus on the use of drug SA procedures to address this latter scientific category.

Although there are infinite iterations of drug SA procedures, all use the classic 3-term contingency of operant conditioning to investigate the stimulus properties of drugs (Skinner 1938). This 3-term contingency can be diagrammed as follows in (7.1):

$$S^D \rightarrow R \rightarrow S^C \quad (7.1)$$

where S^D designates a *discriminative stimulus*, R designates a *response* on the part of the organism, and S^C designates a *consequent stimulus*. The arrows specify the contingency that, in the presence of the discriminative stimulus S^D , performance of response R will deliver the consequent stimulus S^C . As a simple example, a rat implanted with a chronic indwelling venous catheter might be connected to an infusion pump containing a dose of a psychoactive drug and placed into an experimental chamber that contains a stimulus light and a response lever. Contingencies can be programmed such that, if the stimulus light is illuminated (the discriminative stimulus), then depression of the response lever (the response) will result in delivery of a drug injection (the consequent stimulus). Conversely, if the stimulus light is not illuminated, then responding does not result in the delivery of the drug injection. Under these conditions, subjects typically learn to respond when the discriminative stimulus is present. Consequent stimuli that increase responding leading to their delivery are operationally defined as *reinforcers*, whereas stimuli that decrease responding leading to their delivery are defined as *punishers*. The contingencies that relate discriminative stimuli, responses and consequent stimuli are defined by the schedule of reinforcement (Ferster and Skinner 1957).

Although there are Food and Drug Administration (FDA)-approved pharmacotherapies for some SUDs (e.g., opioid, nicotine, and ethanol), FDA-approved pharmacotherapies are absent for many other classes of abused drugs (e.g., cocaine, methamphetamine, and cannabis). Moreover, the development of safer and more efficacious medications to treat SUDs remains a priority for both preclinical and human laboratory/clinical drug abuse research. Preclinical evaluation of candidate medication treatment effects on drug SA has demonstrated good, but not perfect, concordance with both medication effects in human laboratory drug SA studies and metrics of drug abuse in clinical trials (Comer et al. 2008; Haney and Spealman 2008a; Mello and Negus 1996).

2. Rationale for Kappa-Opioid Receptors as Candidate SUD Treatments

This chapter will focus on kappa-opioid receptor (KOR) agonist and antagonist effects on preclinical drug SA endpoints. The KOR is a seven transmembrane $G_{i/o}$ -protein coupled receptor that is ubiquitously expressed in the central and peripheral nervous system and is hypothesized to be involved in mental health disorders including stress, anxiety, depression, and SUD (for review, see (Chavkin and Koob 2015; Crowley and Kash 2015; Tejeda and Bonci 2019; Wee and Koob 2010b)). In general, KOR activation results in inhibition of neuronal function and can occur following either endogenous release of the dynorphin peptide (Chavkin et al. 1982; Oka et al. 1982) or synthetic KOR agonist administration. For

example, administration of synthetic KOR agonists (e.g. U59,593, U50,488, or salvinorin A) decreases extracellular dopamine levels in the nucleus accumbens (Carlezon et al. 2006; Di Chiara and Imperato 1988; Leidl et al. 2014).

Table 1 lists the most prominent KOR agonists and antagonists used in preclinical and clinical research along with the relative selectivity for the KOR over other similar homology receptors such as the mu-opioid (MOR) and delta-opioid (DOR) receptors. In general, the greater the selectivity for KOR over MOR and DOR, the more confidence the result is due to KOR activation or inhibition and not due to an off-target receptor. In addition, the availability of both selective KOR agonists and antagonists allows for sufficient and necessary experimentation in the role of KOR for a specific pharmacological effect. For example, if you were interested in whether activation of KOR receptors by dynorphin was sufficient to decrease mesolimbic dopamine levels, then you could administer a KOR agonist (e.g. salvinorin A) and measure mesolimbic dopamine levels. As mentioned in above, this is a reported effect of KOR agonists. However, if you were interested in whether KOR were necessary for salvinorin A to decrease mesolimbic dopamine levels, then you would administer a KOR antagonist before salvinorin A to determine whether salvinorin A decreases mesolimbic dopamine levels. The availability of selective agonists and antagonists allows for rigorous experimentation into the mechanisms associated involved in SUDs.

Repeated exposure to several drugs of abuse (e.g. cocaine, methamphetamine, mu-opioid receptor agonists) has been shown to activate or sensitize the endogenous dynorphin/KOR system. This engagement of the dynorphin/KOR system by drugs of abuse has been theorized to contribute to the development of compulsive drug use and SUD (Chavkin and Koob 2015; Shippenberg et al. 2007). For example, polymorphisms of prodynorphin have been correlated with increased diagnosis of opioid use disorder (Clarke et al. 2012). In addition, both increased dynorphin expression and KOR density have been observed in cocaine overdose patients (Hurd and Herkenham 1993; Staley et al. 1997). Furthermore, cocaine SA leads to a reduction in KOR availability in humans as measured using positron emission tomography (Martinez et al. 2019). Consistent with these clinical data, a history of either heroin or cocaine SA under extended access conditions increases prodynorphin expression in the nucleus accumbens shell of rats or monkeys (Daunais et al. 1993; Fagergren et al. 2003; Schlosburg et al. 2013; Solecki et al. 2009). These results have been interpreted to suggest that chronic KOR activation is one key neurobiological system mediating the progression from drug abuse to severe SUD (Koob and Moal 2008; Wee and Koob 2010a). Table 2 summarizes the 34 publications that have examined either acute or chronic (i.e. repeated administration for at least 3 days) treatment effects of KOR agonists or antagonists in preclinical drug SA procedures. The predominant drugs of abuse examined have been cocaine (44%), ethanol (35%), and opioids (24%). The predominant research subjects have been rats (74%) and rhesus monkeys (26%); most studies only involved male subjects (~76%). The results and implication of this literature will be reviewed in more detail below.

3. KOR Agonists Effects on Preclinical Drug SA

KOR agonists and drugs of abuse appear to produce opposing effects on both abuse-related neurochemical and behavioral endpoints. For example, KOR agonists decrease mesolimbic dopamine levels in rats (Devine et al. 1993; Donzanti et al. 1992; Leidl et al. 2014; Spanagel et al. 1990), produce dysphoric subjective effects in humans (Pfeiffer et al. 1986; Walsh et al. 2001), and fail to function as positive reinforcers in rodent and nonhuman primate drug SA procedures (Marinelli et al. 1998; Negus et al. 2008; Tang and Collins 1985; Townsend et al. 2017; Woods and Gmerek 1985). This line of research led to the hypothesis that KOR agonists may have clinical utility as candidate SUD pharmacotherapies by either punishing drug-taking behavior if the KOR agonist and the drug of abuse were combined or antagonizing the abuse-related effects of central nervous system-active drugs if the KOR agonist was administered as an acute or repeated pretreatment to subsequent drug-taking behavior. Two general types of experiments have been conducted examining the effects of KOR agonists on drug SA.

3.1 Preclinical SA of KOR Agonists and Drug of Abuse Combinations

One type of experiment involves combining a KOR agonist and the drug of abuse in the same syringe for SA to determine whether the KOR agonist would function as a punisher (i.e., presentation of the stimulus (KOR agonist) decreases the probability of the preceding behavior). Thus, the research animal would self-administer a mixture of KOR agonist and drug of abuse. Based on the literature cited above, the hypothesis for these studies would be that the KOR agonist mixed with the drug of abuse would lead to a decrease in rates of drug SA. Three studies (#1–3 in Table 2) have examined combining a KOR agonist and cocaine (Freeman et al. 2014) or a MOR agonist (Negus et al. 2008; Townsend et al. 2017). All three studies reported that mixtures of the abused drug and KOR agonist were less reinforcing (i.e., maintained lower rates of operant responding) than the abused drug alone. These results have been interpreted to suggest that clinical combinations of MOR agonists and KOR agonists might retain the therapeutic desirable effects (e.g. analgesia) of MOR agonists with reduced undesirable effects (e.g. abuse liability). However, illicit drug manufacturers or dealers are unlikely to adulterate their product with something that would deter abuse liability.

3.2 KOR Agonists as Pretreatments to Preclinical Drug of Abuse SA

A second type of experiment has examined either acute KOR agonist pretreatment to a single drug SA or repeated KOR agonist treatment effects across multiple days of drug SA. The studies that employed a repeated dosing procedure were modeling aspects of repeated candidate medication administration utilized in human laboratory studies, clinical trials, and clinical prescribing patterns for current FDA-approved treatments for SUDs (e.g. buprenorphine, methadone, varenicline, and naltrexone). Thirteen studies (#4–16 in Table 2) have examined the effects of either acute or repeated KOR agonist treatment on drug SA in mice, rats, and rhesus monkeys. Acute administration of U50,488 (Glick et al. 1995), U69,593 (Schenk et al. 1999; Schenk et al. 2001), cyclazocine (Glick et al. 1998), spiradoline (Glick et al. 1995), and enadoline (Bowen et al. 2003; Höltner et al. 2000) has been reported to decrease rates of cocaine, morphine, and ethanol SA in both rats and rhesus

monkeys. These results were interpreted as evidence that KOR agonists may have clinical utility as candidate medications for SUD treatment.

However, when KOR agonists effects on drug SA were examined under repeated dosing conditions, a more complicated profile of effects emerged compared to the acute KOR agonist pretreatment studies described above. For example, repeated ethylketocyclazine, U50,488, enadoline, spiradoline, PD117302, bremazocine, and cyclazocine treatments all produced sustained decreases in rates of cocaine and ethanol SA in rhesus monkeys (Cosgrove and Carroll 2002; Mello and Negus 1998; Negus et al. 1997). Repeated bremazocine, but not U50,488, produced sustained decreases in rates of ethanol SA in rats (Nestby et al. 1999). Repeated U50,488 treatment significantly decreased rates of nicotine SA in rats on the third day of treatment and without altering food-maintained responding (Ismayilova and Shoaib 2010). However, the KOR agonist doses that decreased drug SA also decreased rates of food-maintained responding when food-maintained responding was assessed (Cosgrove and Carroll 2002; Mello and Negus 1998; Negus et al. 1997). Thus, repeated KOR agonist treatment effects were not behaviorally selective for drug vs. nondrug reinforcers and suggestive of overall depression of behavior. In contrast to these results, repeated enadoline treatment increased rates of ethanol SA in rats (Hölter et al. 2000) and repeated U50,488 treatment shifted both cocaine and morphine SA dose-effect functions to the left in rats (Kuzmin et al. 1997). Furthermore, when repeated U50,488 treatment effects were examined on cocaine SA in rhesus monkeys under conditions where there was an alternative nondrug food reinforcer available concurrently to intravenous cocaine injections, repeated U50,488 increased cocaine “choice” and decreased food “choice” (Negus 2004). Consistent with these later results, repeated enadoline treatment failed to attenuate cocaine-vs.-money choice in humans and there was a trend for increased cocaine choice at the largest enadoline dose examined (Walsh et al. 2001). No double-blind, placebo controlled clinical trials examining KOR agonists as candidate pharmacotherapies for SUD treatment have been published.

In summary, the effects of KOR agonists on preclinical drug SA studies and the single human laboratory drug SA study have thus far generated three main findings. First, when KOR agonists are mixed with the abuse drug and co-administered, KOR agonists can function as punishers and decrease SA of the abused drug. Second, acute pretreatment with a KOR agonist decreased rates of drug SA across a broad range of abused drug classes (e.g. cocaine, ethanol, morphine). Lastly, repeated KOR agonist treatment either decreased rates of abused drug SA typically at doses that also decreased rates of food-maintained responding or increases rates of drug SA including under a drug-vs.-food choice procedure. Overall, this body of preclinical literature does not support the clinical utility of KOR agonists as candidate SUD treatments.

4. KOR Antagonist Effects on Preclinical Drug Self-Administration

One prominent and emerging SUD theory is that chronic exposure to drugs of abuse and withdrawal produce a “motivational withdrawal syndrome” that increase the magnitude and alter the mechanisms of drug reinforcement and serves as “one of the driving factors of compulsivity in addiction” (Koob and Mason 2016). Chronic exposure to drugs of abuse is

hypothesized to alter the state of the patient or research subject and thereby alter the mechanisms of and increase the magnitude of drug reinforcement. For example, opioid abuse often leads to physical dependence, and opioid withdrawal in dependent subjects increases the reinforcing effects of opioids, decreases the reinforcing efficacy of nondrug reinforcers like food, and promotes a maladaptive allocation of behavior toward further drug use at the expense of behaviors maintained by more adaptive behaviors. Specifically, chronic drug exposure has been shown to 1) decrease basal activity of dopamine and/or opioid reward systems, and 2) recruit activation of other neural systems, sometimes described as “stress” or “anti-reward” systems, that involve neurotransmitters including corticotropin releasing factor (CRF) and dynorphin (Koob and Mason 2016; Koob and Moal 2008; Koob and Volkow 2009). Overall, consumption of any abused drug dose is hypothesized to produce an initial increase in dopamine/opioid signaling as drug levels rise and peak, followed by a longer period of depressed dopamine/opioid signaling and enhanced stress hormone signaling that together mediate negative affective states of drug withdrawal. Increasingly intensive regimens of abused drug exposure are hypothesized to produce a cumulative increase in the later effects and increasingly intense aversive subjective states. Under these circumstances, drug-induced reinforcing effects are often described as “negative,” because the drug is now hypothesized to produce its reinforcing effects not by increasing dopamine/opioid signaling from a normal basal level, but rather by alleviating the aversive state produced by a depressed reward system and activated stress system; however, this interpretation is not consistent with the operational definition of negative reinforcement (i.e. response requirement completion results in removal of S^C proposed by Skinner (1938)). The subject in a drug SA procedure is responding to receive the drug injection (S^C) which is positive reinforcement and not the removal of some hypothesized internal state (Negus and Banks 2017).

Evidence to support this hypothesized transition comes in part from preclinical rat drug SA studies that have used single-operant drug SA procedures in which the primary dependent measure is the rate of drug SA. Two observations have been seminal in support of this hypothesized transition. First, the recruitment of negative reinforcement processes that occurs with extended drug SA is hypothesized to not only modify the mechanisms of drug reinforcement, but also to increase its magnitude (i.e., by summing positive and negative reinforcement mechanisms). In support of this hypothesis, regimens of “extended access” (produced by increasing the number of hours per day that subjects can self-administer drug) have been shown to increase rates of SA, a phenomenon referred to as “escalation” (Koob and Kreek 2007). Second, the recruitment of these negative reinforcement processes is also hypothesized to render drug SA sensitive to experimental manipulations that attenuate the KOR/stress system signaling. For example, dynorphin acting at KOR is one stress-related neurotransmitter implicated in “negative reinforcement” processes, and KOR antagonists such as nor-binaltorphimine (nor-BNI) and CERC-501 have been reported to block escalated rates of cocaine, methamphetamine, heroin, and ethanol SA in rats (Domi et al. 2018; Schlosburg et al. 2013; Walker et al. 2011; Wee et al. 2009; Wee et al. 2012; Whitfield et al. 2015). Furthermore, KOR antagonists do not attenuate cocaine, methamphetamine, or heroin SA under more limited (~1–2 h) drug access conditions in both rats (Deehan et al. 2012; Doyon et al. 2006; Glick et al. 1995; Höltner et al. 2000; Liu and Jernigan 2011; Negus et al.

1993) and rhesus monkeys (Negus 2004; Negus et al. 1997) suggesting that the recruitment of the dynorphin/KOR system only occurs during extended drug access conditions. Although the effectiveness of KOR antagonists to alter drug SA appears to be dependent on the length of the drug SA session for MOR agonists and monoamine transporter ligands (cocaine and methamphetamine; Table 2), the literature suggests the same principle does not necessarily hold true for ethanol. For example, under limited ethanol access conditions, KOR antagonists have been shown to increase (Mitchell et al. 2005), have no effect (Deehan et al. 2012; Doyon et al. 2006), or decrease (Cashman and Azar 2014; Rorick-Kehn et al. 2014; Schank et al. 2012) ethanol SA in rats. Overall, one implication of this literature is that different classes of abused drugs might recruit the dynorphin/KOR system in different manners.

4.1 KOR Antagonist Effects on Preclinical Drug Choice SA

However, these KOR antagonist treatment effects on preclinical drug SA endpoints have not translated when evaluated in nonhuman primates or humans. For example, in opioid-dependent rhesus monkeys, the KOR antagonist 5'-guanidinaltrindole (GNTI) failed to attenuate opioid withdrawal-associated increases in opioid vs. food choice (Negus and Rice 2008). In addition, nor-BNI failed to attenuate both rates of cocaine SA during extended cocaine access sessions and cocaine vs. food choice in rhesus monkeys self-administering cocaine 22-h per day (Hutsell et al. 2016). Whether these differences in KOR antagonist effects are due to species differences between rats and rhesus monkeys or procedural differences in the drug SA schedule or reinforcement remain unexplored scientific space.

4.2 KOR Antagonist Effects on Human Drug SA Metrics

In humans, buprenorphine plus naloxone and naltrexone maintenance, combined to produce a KOR antagonist effect, failed to attenuate cocaine use in a double-blind, placebo-controlled, multi-centered clinical trial (Ling et al. 2016). Repeated LY2456302 treatment also failed to attenuate cocaine craving in humans (Reed et al. 2017). Furthermore, CERC-501 (i.e. LY2456301) failed to attenuate cigarette smoking, craving, or nicotine withdrawal in humans (Jones et al. 2019). Moreover, a recent positron emission tomography study examining KOR binding in healthy controls and cocaine abusers reported no significant differences in KOR binding in any brain region examined (Martinez et al. 2019). These results suggest that a history of repeated cocaine exposure and a diagnosis of cocaine use disorder was not sufficient to alter KOR binding in humans and are in contrast to previous results in cocaine overdose patients using autoradiography methods (Hurd and Herkenham 1993; Staley et al. 1997). Reasons for differences between the in vivo positron emission tomography study and the post-mortem mRNA and autoradiography study are not presently clear, but could be related to the affinity of the ligands for different KOR subtypes or states (i.e. high vs. low affinity). In summary, the promising results of KOR antagonists in preclinical rodent models of cocaine use disorder have so far failed to translate when evaluated in higher-order species such as nonhuman primates or humans, and on endpoints that focus on behavioral allocation between drug and nondrug reinforcers instead of rates of drug-taking behavior.

5. Conclusions and Future Directions

Over the past two decades, preclinical research has improved our understanding of the role of KOR in the expression, mechanisms, and treatment of SUD. Converging lines of evidence from both preclinical and human studies support the conclusion that chronic abused drug exposure changes the dynorphin/KOR system. Furthermore, mixtures of MOR and KOR agonists (Freeman et al. 2014; Negus et al. 2008; Townsend et al. 2017) or mixed-action MOR/KOR agonists such as pentazocine (Hoffmeister 1979) appear to have reduced abuse-related effects compared to MOR agonists. However, both KOR agonist and KOR antagonist treatments have thus far failed to significantly alter metrics of cocaine use in humans. Unfortunately, there are no published human laboratory drug SA studies or clinical trials evaluating KOR agonist or antagonist effects on other SUDs, such as methamphetamine, opioids, ethanol, or tobacco. CERC-501 is currently being evaluated in clinical trials as a candidate medication for tobacco smoking and smoking relapse (Helal et al. 2017). The evaluation of candidate medication effects in SUD patients provides critical reverse translational feedback to improve the predictive validity of preclinical SUD models. However, the results of KOR agonists and antagonists in cocaine use disorder and tobacco use disorder patients thus far fails to support the continued development and evaluation of novel chemical entities targeting the dynorphin/KOR system as candidate SUD pharmacotherapies. There are presently no published clinical data on the effectiveness of KOR antagonists for alcohol or opioid use disorder.

In the broader preclinical drug abuse literature, there are two experimental features that appear to promote accurate translation of preclinical-to-clinical results. First, repeated treatment with the candidate medication to match the subchronic-to-chronic treatment regimens commonly employed in clinical SUD treatment (Czoty et al. 2016; Haney and Spealman 2008b; Mello and Negus 1996). The effects of acute vs. repeated KOR agonists on preclinical drug SA endpoints reviewed above are consistent with this conclusion. However, the examination of repeated KOR antagonist effects on drug SA endpoints has been problematic because most currently available KOR antagonists are irreversible or receptor-inactivating antagonists (Butelman et al. 1998; Butelman et al. 1993; Schmid et al. 2013). The long duration of action of irreversible KOR antagonists complicates dose titration and potentially increases the risk of off-target undesirable effects. The development of short-acting KOR antagonists CERC-501 (i.e. LY2456302) (Helal et al. 2017; Reed et al. 2017) has facilitated clinical SUD research to provide necessary feedback regarding hypotheses developed using preclinical drug SA procedures. Second, assessment of candidate medication effects in preclinical drug SA procedures that use behavioral allocation between the target drug of abuse and an alternative nondrug reinforcer (e.g. food or social) rather than rates of drug SA behavior has shown strong translational concordance with clinical results (for review, see (Banks et al. 2015; Banks and Negus 2012; Banks and Negus 2017)). A critical step in efficiently evaluating candidate SUD medications is the utilization of preclinical testing procedures that are both sensitive to FDA-approved medications (if available) and selective for those positive controls in comparison to active negative controls known to be clinically ineffective (Banks et al. 2019). Given the human laboratory and clinical literature cited above in Sections 3 and 4, both KOR agonists and KOR antagonists

could function as active negative controls in the development and evaluation of either novel candidate SUD medications or novel preclinical drug SA procedures.

References

- Ator NA, Griffiths RR (2003) Principles of drug abuse liability assessment in laboratory animals. *Drug and Alcohol Dependence* 70: S55–S72. [PubMed: 12759197]
- Banks ML, Hutsell BA, Schwientek KL, Negus SS (2015) Use of Preclinical Drug Vs. Food Choice Procedures to Evaluate Candidate Medications for Cocaine Addiction. *Curr Treat Options Psychiatry* 2: 136–150. [PubMed: 26009706]
- Banks ML, Negus SS (2012) Preclinical Determinants of Drug Choice under Concurrent Schedules of Drug Self-Administration. *Advances in pharmacological sciences* 2012: 281768. [PubMed: 23243420]
- Banks ML, Negus SS (2017) Insights from Preclinical Choice Models on Treating Drug Addiction. *Trends Pharmacol Sci* 38: 181–194. [PubMed: 27916279]
- Banks ML, Townsend EA, Negus SS (2019) Testing the 10 most wanted: a preclinical algorithm to screen candidate opioid use disorder medications. *Neuropsychopharmacology*.
- Bowen CA, Stevens Negus S, Zong R, Neumeyer JL, Bidlack JM, Mello NK (2003) Effects of Mixed-Action κ/μ Opioids on Cocaine Self-Administration and Cocaine Discrimination by Rhesus Monkeys. *Neuropsychopharmacology* 28: 1125–1139. [PubMed: 12637953]
- Butelman ER, Ko M-C, Sobczyk-Kojiro K, Mosberg HI, Van Bommel B, Zernig G, Woods JH (1998) Kappa-Opioid Receptor Binding Populations in Rhesus Monkey Brain: Relationship to an Assay of Thermal Antinociception. *J Pharmacol Exp Ther* 285: 595–601. [PubMed: 9580603]
- Butelman ER, Negus SS, Ai Y, de Costa BR, Woods JH (1993) Kappa opioid antagonist effects of systemically administered nor-binaltorphimine in a thermal antinociception assay in rhesus monkeys. *J Pharmacol Exp Ther* 267: 1269–1276. [PubMed: 8263790]
- Carlezon WA, Béguin C, DiNieri JA, Baumann MH, Richards MR, Todtenkopf MS, Rothman RB, Ma Z, Lee DY-W, Cohen BM (2006) Depressive-Like Effects of the κ -Opioid Receptor Agonist Salvinorin A on Behavior and Neurochemistry in Rats. *J Pharmacol Exp Ther* 316: 440–447. [PubMed: 16223871]
- Carter LP, Griffiths RR (2009) Principles of laboratory assessment of drug abuse liability and implications for clinical development. *Drug Alcohol Depend* 105, Supplement 1: S14–S25. [PubMed: 19443137]
- Cashman JR, Azar MR (2014) Potent Inhibition of Alcohol Self-Administration in Alcohol-Preferring Rats by a κ -Opioid Receptor Antagonist. *Journal of Pharmacology and Experimental Therapeutics* 350: 171–180.
- Chavkin C, James I, Goldstein A (1982) Dynorphin is a specific endogenous ligand of the kappa opioid receptor. *Science* 215: 413–415. [PubMed: 6120570]
- Chavkin C, Koob GF (2015) Dynorphin, Dysphoria, and Dependence: the Stress of Addiction. *Neuropsychopharmacology* 41: 373.
- Clarke T-K, Ambrose-Lanci L, Ferraro TN, Berrettini WH, Kampman KM, Dackis CA, Pettinati HM, O'Brien CP, Oslin DW, Lohoff FW (2012) Genetic association analyses of PDYN polymorphisms with heroin and cocaine addiction. *Genes, Brain and Behavior* 11: 415–423.
- Comer SD, Ashworth JB, Foltin RW, Johanson CE, Zacny JP, Walsh SL (2008) The role of human drug self-administration procedures in the development of medications. *Drug Alcohol Depend* 96: 1–15. [PubMed: 18436394]
- Cosgrove KP, Carroll ME (2002) Effects of Bremazocine on Self-Administration of Smoked Cocaine Base and Orally Delivered Ethanol, Phencyclidine, Saccharin, and Food in Rhesus Monkeys: A Behavioral Economic Analysis. *Journal of Pharmacology and Experimental Therapeutics* 301: 993–1002.
- Crowley NA, Kash TL (2015) Kappa opioid receptor signaling in the brain: Circuitry and implications for treatment. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 62: 51–60. [PubMed: 25592680]

- Czoty PW, Stoops WW, Rush CR (2016) Evaluation of the “Pipeline” for Development of Medications for Cocaine Use Disorder: A Review of Translational Preclinical, Human Laboratory, and Clinical Trial Research. *Pharmacol Rev* 68: 533–562. [PubMed: 27255266]
- Daunais JB, Roberts DC, McGinty JF (1993) Cocaine self-administration increases preprodynorphin, but not c-fos, mRNA in rat striatum. *Neuroreport* 4: 543–6. [PubMed: 8099817]
- Deehan GA, McKinzie DL, Carroll FI, McBride WJ, Rodd ZA (2012) The long-lasting effects of JDTC, a kappa opioid receptor antagonist, on the expression of ethanol-seeking behavior and the relapse drinking of female alcohol-preferring (P) rats. *Pharmacology Biochemistry and Behavior* 101: 581–587.
- Devine DP, Leone P, Pocock D, Wise RA (1993) Differential involvement of ventral tegmental mu, delta and kappa opioid receptors in modulation of basal mesolimbic dopamine release: in vivo microdialysis studies. *Journal of Pharmacology and Experimental Therapeutics* 266: 1236–1246.
- Di Chiara G, Imperato A (1988) Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely moving rats. *Proc Natl Acad Sci USA* 85: 5274–5278. [PubMed: 2899326]
- Domi E, Barbier E, Augier E, Augier G, Gehlert D, Barchiesi R, Thorsell A, Holm L, Heilig M (2018) Preclinical evaluation of the kappa-opioid receptor antagonist CERC-501 as a candidate therapeutic for alcohol use disorders. *Neuropsychopharmacology* 43: 1805–1812. [PubMed: 29463912]
- Donzanti BA, Althaus JS, Payson MM, Von Voigtlander PF (1992) Kappa agonist-induced reduction in dopamine release: site of action and tolerance. *Res Commun Chem Pathol Pharmacol* 78: 193–210. [PubMed: 1282270]
- Doyon WM, Howard EC, Shippenberg TS, Gonzales RA (2006) κ -Opioid receptor modulation of accumbal dopamine concentration during operant ethanol self-administration. *Neuropharmacology* 51: 487–496. [PubMed: 16781738]
- Fagergren P, Smith HR, Daunais JB, Nader MA, Porrino LJ, Hurd YL (2003) Temporal upregulation of prodynorphin mRNA in the primate striatum after cocaine self-administration. *Eur J Neurosci* 17: 2212–2218. [PubMed: 12786988]
- Ferster C, Skinner B (1957) *Schedules of Reinforcement*. Appleton-Century-Croft, New York
- Freeman KB, Naylor JE, Prinszano TE, Woolverton WL (2014) Assessment of the kappa opioid agonist, salvinorin A, as a punisher of drug self-administration in monkeys. *Psychopharmacology* 231: 2751–2758. [PubMed: 24481567]
- Glick SD, Maisonneuve IM, Raucci J, Sydney A (1995) Kappa opioid inhibition of morphine and cocaine self-administration in rats. *Brain Res* 681: 147–152. [PubMed: 7552272]
- Glick SD, Visker KE, Maisonneuve IM (1998) Effects of cyclazocine on cocaine self-administration in rats. *European Journal of Pharmacology* 357: 9–14. [PubMed: 9788768]
- Haney M, Spealman R (2008a) Controversies in translational research: drug self-administration. *Psychopharmacology* 199: 403–19. [PubMed: 18283437]
- Haney M, Spealman R (2008b) Controversies in translational research: drug self-administration. *Psychopharmacology* 199: 403–419. [PubMed: 18283437]
- Helal MA, Habib ES, Chittiboyina AG (2017) Selective kappa opioid antagonists for treatment of addiction, are we there yet? *European Journal of Medicinal Chemistry* 141: 632–647. [PubMed: 29107424]
- Hoffmeister F (1979) Progressive-ratio performance in the rhesus monkey maintained by opiate infusions. *Psychopharmacology* 62: 181–186. [PubMed: 111282]
- Hölter SM, Henniger MSH, Lipkowski AW, Spanagel R (2000) Kappa-opioid receptors and relapse-like drinking in long-term ethanol-experienced rats. *Psychopharmacology* 153: 93–102. [PubMed: 11255932]
- Hunter JC, Leighton GE, Meecham KG, Boyle SJ, Horwell DC, Rees DC, Hughes J (1990) CI-977, a novel and selective agonist for the κ -opioid receptor. *Br J Pharmacol* 101: 183–189. [PubMed: 2178014]
- Hurd YL, Herkenham M (1993) Molecular alterations in the neostriatum of human cocaine addicts. *Synapse* 13: 357–369. [PubMed: 7683144]

- Hutsell BA, Cheng K, Rice KC, Negus SS, Banks ML (2016) Effects of the kappa opioid receptor antagonist nor-binaltorphimine (nor-BNI) on cocaine versus food choice and extended-access cocaine intake in rhesus monkeys. *Addict Biol* 00: 360–373.
- Ismayilova N, Shoaib M (2010) Alteration of intravenous nicotine self-administration by opioid receptor agonist and antagonists in rats. *Psychopharmacology* 210: 211–220. [PubMed: 20401605]
- Jones JD, Babalonis S, Marcus R, Vince B, Kelsh D, Lofwall MR, Fraser H, Paterson B, Martinez S, Martinez DM, Nunes EV, Walsh SL, Comer SD (2019) A randomized, double-blind, placebo-controlled study of the kappa opioid receptor antagonist, CERC-501, in a human laboratory model of smoking behavior. *Addiction Biology* 0: e12799.
- Koob GF, Kreek MJ (2007) Stress, Dysregulation of Drug Reward Pathways, and the Transition to Drug Dependence. *American Journal of Psychiatry* 164: 1149–1159.
- Koob GF, Mason BJ (2016) Existing and Future Drugs for the Treatment of the Dark Side of Addiction. *Ann Rev Pharmacol Toxicol* 56: 299–322. [PubMed: 26514207]
- Koob GF, Moal ML (2008) Addiction and the Brain Antireward System. *Annual Review of Psychology* 59: 29–53.
- Koob GF, Volkow ND (2009) Neurocircuitry of Addiction. *Neuropsychopharmacology* 35: 217.
- Kuzmin AV, Semenova S, Gerrits MAFM, Zvartau EE, Van Ree JM (1997) κ -Opioid receptor agonist U50,488H modulates cocaine and morphine self-administration in drug-naïve rats and mice. *European Journal of Pharmacology* 321: 265–271. [PubMed: 9085036]
- Leitl MD, Onvani S, Bowers MS, Cheng K, Rice KC, Carlezon WA Jr., Banks ML, Negus S (2014) Pain-Related Depression of the Mesolimbic Dopamine System in Rats: Expression, Blockade by Analgesics, and Role of Endogenous kappa-opioids. *Neuropsychopharmacology* 39: 614–624. [PubMed: 24008352]
- Ling W, Hillhouse MP, Saxon AJ, Mooney LJ, Thomas CM, Ang A, Matthews AG, Hasson A, Annon J, Sparenborg S, Liu DS, McCormack J, Church S, Swafford W, Drexler K, Schuman C, Ross S, Wiest K, Korthuis PT, Lawson W, Brigham GS, Knox PC, Dawes M, Rotrosen J (2016) Buprenorphine + naloxone plus naltrexone for the treatment of cocaine dependence: the Cocaine Use Reduction with Buprenorphine (CURB) study. *Addiction* 111: 1416–1427. [PubMed: 26948856]
- Liu X, Jernigan C (2011) Activation of the opioid μ 1, but not δ or κ , receptors is required for nicotine reinforcement in a rat model of drug self-administration. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 35: 146–153. [PubMed: 20965223]
- Marinelli M, Barrot M, Simon H, Oberlander C, Dekeyne A, Moal ML, Piazza PV (1998) Pharmacological stimuli decreasing nucleus accumbens dopamine can act as positive reinforcers but have a low addictive potential. *European Journal of Neuroscience* 10: 3269–3275.
- Martinez D, Slifstein M, Matuskey D, Nabulsi N, Zheng M-Q, Lin S-f, Ropchan J, Urban N, Grassetti A, Chang D, Salling M, Foltin R, Carson RE, Huang Y (2019) Kappa-opioid receptors, dynorphin, and cocaine addiction: a positron emission tomography study. *Neuropsychopharmacology*.
- Mello NK, Negus SS (1996) Preclinical evaluation of pharmacotherapies for treatment of cocaine and opioid abuse using drug self-administration procedures. *Neuropsychopharmacology* 14: 375–424. [PubMed: 8726752]
- Mello NK, Negus SS (1998) Effects of Kappa Opioid Agonists on Cocaine- and Food-Maintained Responding by Rhesus Monkeys. *J Pharmacol Exp Ther* 286: 812–824. [PubMed: 9694938]
- Mitchell JM, Liang MT, Fields HL (2005) A single injection of the kappa opioid antagonist norbinaltorphimine increases ethanol consumption in rats. *Psychopharmacology* 182: 384–392. [PubMed: 16001119]
- Munro TA, Huang X-P, Inglese C, Perrone MG, Van't Veer A, Carroll FI, Béguin C, Carlezon WA Jr., Colabufo NA, Cohen BM, Roth BL (2013) Selective κ Opioid Antagonists nor-BNI, GNTI and JDTic Have Low Affinities for Non-Opioid Receptors and Transporters. *PLOS ONE* 8: e70701. [PubMed: 23976952]
- Nagase H, Imaide S, Hirayama S, Nemoto T, Fujii H (2012) Essential structure of opioid κ receptor agonist nalfurafine for binding to the κ receptor 2: Synthesis of decahydro(iminoethano)phenanthrene derivatives and their pharmacologies. *Bioorg Med Chem Letts* 22: 5071–5074. [PubMed: 22742909]

- Negus SS (2004) Effects of the kappa opioid agonist U50,488 and the kappa opioid antagonist nor-binaltorphimine on choice between cocaine and food in rhesus monkeys. *Psychopharmacology* 176: 204–213. [PubMed: 15112031]
- Negus SS, Banks ML (2017) Modulation of drug choice by extended drug access and withdrawal in rhesus monkeys: Implications for negative reinforcement as a driver of addiction and target for medications development. *Pharmacology Biochemistry and Behavior*.
- Negus SS, Henriksen SJ, Mattox A, Pasternak GW, Portoghese PS, Takemori AE, Weinger MB, Koob GF (1993) Effect of antagonists selective for mu, delta and kappa opioid receptors on the reinforcing effects of heroin in rats. *Journal of Pharmacology and Experimental Therapeutics* 265: 1245–1252.
- Negus SS, Mello NK, Portoghese PS, Lin C-E (1997) Effects of Kappa Opioids on Cocaine Self-Administration by Rhesus Monkeys. *J Pharmacol Exp Ther* 282: 44–55. [PubMed: 9223538]
- Negus SS, Rice KC (2008) Mechanisms of Withdrawal-Associated Increases in Heroin Self-Administration: Pharmacologic Modulation of Heroin vs Food Choice in Heroin-Dependent Rhesus Monkeys. *Neuropsychopharmacology* 34: 899–911. [PubMed: 18704098]
- Negus SS, Schrode K, Stevenson GW (2008) Mu/kappa opioid interactions in rhesus monkeys: implications for analgesia and abuse liability. *Exp Clin Psychopharmacol* 16: 386–99. [PubMed: 18837635]
- Nestby P, Schoffelmeer ANM, Homberg JR, Wardeh G, De Vries TJ, Mulder AH, Vanderschuren LJMJ (1999) Bremazocine reduces unrestricted free-choice ethanol self-administration in rats without affecting sucrose preference. *Psychopharmacology* 142: 309–317. [PubMed: 10208324]
- Oka T, Negishi K, Suda M, Sawa A, Fujino M, Wakimasu M (1982) Evidence that dynorphin-(1–13) acts as an agonist on opioid κ -receptors. *European Journal of Pharmacology* 77: 137–141. [PubMed: 6277659]
- Pfeiffer A, Brantl V, Herz A, Emrich H (1986) Psychotomimesis mediated by kappa opiate receptors. *Science* 233: 774–776. [PubMed: 3016896]
- Raynor K, Kong H, Chen Y, Yasuda K, Yu L, Bell GI, Reisine T (1994) Pharmacological characterization of the cloned kappa-, delta-, and mu-opioid receptors. *Mol Pharmacol* 45: 330–334. [PubMed: 8114680]
- Reed B, Butelman ER, Fry RS, Kimani R, Kreek MJ (2017) Repeated Administration of Opra Kappa (LY2456302), a Novel, Short-Acting, Selective KOP-r Antagonist, in Persons with and without Cocaine Dependence. *Neuropsychopharmacology* 43: 739. [PubMed: 28857070]
- Rorick-Kehn LM, Witkin JM, Statnick MA, Eberle EL, McKinzie JH, Kahl SD, Forster BM, Wong CJ, Li X, Crile RS, Shaw DB, Sahr AE, Adams BL, Quimby SJ, Diaz N, Jimenez A, Pedregal C, Mitch CH, Knopp KL, Anderson WH, Cramer JW, McKinzie DL (2014) LY2456302 is a novel, potent, orally-bioavailable small molecule kappa-selective antagonist with activity in animal models predictive of efficacy in mood and addictive disorders. *Neuropharmacology* 77: 131–144. [PubMed: 24071566]
- Roth BL, Baner K, Westkaemper R, Siebert D, Rice KC, Steinberg S, Ernsberger P, Rothman RB (2002) Salvinorin A: A potent naturally occurring nonnitrogenous κ opioid selective agonist. *Proc Natl Acad Sci USA* 99: 11934–11939. [PubMed: 12192085]
- Schank JR, Goldstein AL, Rowe KE, King CE, Marusich JA, Wiley JL, Carroll FI, Thorsell A, Heilig M (2012) The kappa opioid receptor antagonist JDTC attenuates alcohol seeking and withdrawal anxiety. *Addiction Biology* 17: 634–647. [PubMed: 22515275]
- Schenk S, Partridge B, Shippenberg TS (1999) U69593, a kappa-opioid agonist, decreases cocaine self-administration and decreases cocaine-produced drug-seeking. *Psychopharmacology* 144: 339–346. [PubMed: 10435406]
- Schenk S, Partridge B, Shippenberg TS (2001) Effects of the Kappa-opioid Receptor Agonist, U69593, on the Development of Sensitization and on the Maintenance of Cocaine Self-administration. *Neuropsychopharmacology* 24: 441. [PubMed: 11182539]
- Schlosburg JE, Whitfield TW, Park PE, Crawford EF, George O, Vendruscolo LF, Koob GF (2013) Long-Term Antagonism of κ Opioid Receptors Prevents Escalation of and Increased Motivation for Heroin Intake. *J Neurosci* 33: 19384–19392. [PubMed: 24305833]

- Schmid CL, Streicher JM, Groer CE, Munro TA, Zhou L, Bohn LM (2013) Functional Selectivity of 6'-Guanidinonaltrindole (6'-GNTI) at κ -Opioid Receptors in Striatal Neurons. *Journal of Biological Chemistry* 288: 22387–22398.
- Shippenberg TS, Zapata A, Chefer VI (2007) Dynorphin and the pathophysiology of drug addiction. *Pharmacology & Therapeutics* 116: 306–321. [PubMed: 17868902]
- Skinner B (1938) *The behavior of organisms*. Appleton-Century-Crofts, New York
- Solecki W, Ziolkowska B, Krowka T, Gieryk A, Filip M, Przewlocki R (2009) Alterations of prodynorphin gene expression in the rat mesocorticolimbic system during heroin self-administration. *Brain Research* 1255: 113–121. [PubMed: 19100723]
- Spanagel R, Herz A, Shippenberg TS (1990) The Effects of Opioid Peptides on Dopamine Release in the Nucleus Accumbens: An In Vivo Microdialysis Study. *Journal of Neurochemistry* 55: 1734–1740. [PubMed: 1976759]
- Staley JK, Rothman RB, Rice KC, Partilla J, Mash DC (1997) κ 2 Opioid Receptors in Limbic Areas of the Human Brain Are Upregulated by Cocaine in Fatal Overdose Victims. *The Journal of Neuroscience* 17: 8225–8233. [PubMed: 9334398]
- Tang AH, Collins RJ (1985) Behavioral effects of a novel kappa opioid analgesic, U-50488, in rats and rhesus monkeys. *Psychopharmacology* 85: 309–314. [PubMed: 3923516]
- Tejeda HA, Bonci A (2019) Dynorphin/kappa-opioid receptor control of dopamine dynamics: Implications for negative affective states and psychiatric disorders. *Brain Research* 1713: 91–101. [PubMed: 30244022]
- Townsend EA, Naylor JE, Negus SS, Edwards SR, Qureshi HN, McLendon HW, McCurdy CR, Kapanda CN, do Carmo JM, da Silva FS, Hall JE, Sufka KJ, Freeman KB (2017) Effects of nalfurafine on the reinforcing, thermal antinociceptive, and respiratory-depressant effects of oxycodone: modeling an abuse-deterrent opioid analgesic in rats. *Psychopharmacology* 234: 2597–2605. [PubMed: 28567699]
- Walker BM, Zorrilla EP, Koob GF (2011) Systemic κ -opioid receptor antagonism by nor-binaltorphimine reduces dependence-induced excessive alcohol self-administration in rats. *Addiction Biology* 16: 116–119. [PubMed: 20579007]
- Walsh SL, Geter-Douglas B, Strain EC, Bigelow GE (2001) Enadoline and Butorphanol: Evaluation of κ -Agonists on Cocaine Pharmacodynamics and Cocaine Self-Administration in Humans. *J Pharmacol Exp Ther* 299: 147–158. [PubMed: 11561074]
- Wee S, Koob G (2010a) The role of the dynorphin– κ opioid system in the reinforcing effects of drugs of abuse. *Psychopharmacology* 210: 121–135. [PubMed: 20352414]
- Wee S, Koob GF (2010b) The role of the dynorphin– κ opioid system in the reinforcing effects of drugs of abuse. *Psychopharmacology* 210: 121–135. [PubMed: 20352414]
- Wee S, Orio L, Ghirmai S, Cashman J, Koob G (2009) Inhibition of kappa opioid receptors attenuated increased cocaine intake in rats with extended access to cocaine. *Psychopharmacology* 205: 565–575. [PubMed: 19484223]
- Wee S, Vendruscolo LF, Misra KK, Schlosburg JE, Koob GF (2012) A Combination of Buprenorphine and Naltrexone Blocks Compulsive Cocaine Intake in Rodents Without Producing Dependence. *Science Transl Med* 4: 146ra110.
- Whitfield TW, Schlosburg JE, Wee S, Gould A, George O, Grant Y, Zamora-Martinez ER, Edwards S, Crawford E, Vendruscolo LF, Koob GF (2015) κ Opioid Receptors in the Nucleus Accumbens Shell Mediate Escalation of Methamphetamine Intake. *J Neurosci* 35: 4296–4305. [PubMed: 25762676]
- Woods JH, Gmerek DE (1985) Substitution and primary dependence studies in animals. *Drug and Alcohol Dependence* 14: 233–247. [PubMed: 4039649]

Kappa-opioid receptor (KOR) agonists and antagonists that have been most commonly used in preclinical studies examining the potential of KOR ligands as candidate substance use disorder medications. Columns show the affinity (K_i nM) for the ligand at the KOR and three potential off-target receptor affinities mu-opioid receptor (MOR), delta-opioid receptor (DOR), and nociceptin opioid peptide receptor (NOPR). Together, these results provide one index of KOR selectivity.

Table 1:

Ligand	KOR (K_i nM)	MOR (K_i nM)	DOR (K_i nM)	NOPR (K_i nM)	Reference
<i>KOR Agonists</i>					
Dynorphin A	0.5	32	>1000	Not determined	(Raynor et al. 1994)
U-50,488	0.12	>1000	>1000	Not determined	(Raynor et al. 1994)
U-69,593	0.59	>1000	>1000	Not determined	(Raynor et al. 1994)
Salvinorin A	16	>10,000	>10,000	Not determined	(Roth et al. 2002)
Nalfurafine (TRK-820)	0.2	0.6	96.5	Not determined	(Nagase et al. 2012)
Enadoline (CI-977)	0.11	99	1	Not determined	(Hunter et al. 1990)
Spiradoline	0.036	21	>1000	Not determined	(Raynor et al. 1994)
Bremazocine	0.09	0.75	2.3	Not determined	(Raynor et al. 1994)
<i>KOR Antagonists</i>					
nor-binaltorphimine	4	41	20	4400	(Munro et al. 2013)
JD1c	1	3	44	12	(Munro et al. 2013)
6'-guanidonaltrindole	3	54	58	2500	(Munro et al. 2013)
CERC-501 (LY2456302)	0.8	24	155	Not determined	(Rorick-Kehn et al. 2014)

Table 2:

Summary of published preclinical manuscripts reporting effects of kappa-opioid receptor (KOR) agonists or antagonists on IV drug self-administration (SA). Columns show the self-administered drug, the specific KOR ligand, the sex and species in which the studies were conducted, the primary effect reported in the study, and the reference. Numbers in parentheses indicate unit drug doses in μg or mg/kg /injection for the self-administered drug and drug doses for the KOR ligand.

#	Self-administered Drug (dose)	KOR ligand (dose)	Sex and Species	Main Effect	Ref
		<i>KOR Agonists Studied</i>			
1	Fentanyl (0.01–1 $\mu\text{g/kg}$ /inj)	U69,593 (fixed proportion based on fentanyl dose)	Male Rhesus Monkey	Increasing proportions of U69,593 in the U69,593/fentanyl mixture decreased rates of fentanyl self-administration (SA)	(Negus et al. 2008)
2	Oxycodone (0.032–0.18 mg/kg /inj)	Nalfurafine (fixed proportion based on oxycodone dose)	Male Sprague-Dawley (S-D) Rat	Increasing proportions of nalfurafine in the nalfurafine/oxycodone mixture decreased rates of oxycodone SA	(Townsend et al. 2017)
3	Cocaine (0.1 mg/kg /inj) Remifentanyl (0.1 $\mu\text{g/kg}$ /inj)	Salvinorin A (0.0003–0.01 mg/kg /inj)	Male Rhesus Monkey	Increasing doses of Salvinorin A decreased both cocaine and remifentanyl SA in a drug vs. drug choice procedure	(Freeman et al. 2014)
4	Cocaine (0.4 mg/kg , inj) Morphine (0.04 mg/kg /inj)	U50,488 (2.5–10 mg/kg , IP) Spiradoline (1.25–5 mg/kg , IP) Nor-BNI (10 mg/kg , SC)	Female S-D Rat	Acute U50,488 and spiradoline decreased rates of cocaine and morphine SA. Nor-BNI had no effect alone, but blocked KOR agonist effects.	(Glick et al. 1995)
5	Cocaine (0.007–0.06 mg/kg /inj) Morphine (0.2–2.4 $\mu\text{g/kg}$ /inj)	U50,488 (2.5–10 mg/kg , IP)	Male Wistar Rat (cocaine) Male Mice (morphine)	Repeated U50,488 shifted cocaine and morphine SA dose-effect functions to the left	(Kuzmin et al. 1997)
6	Cocaine (0.0032–0.1 mg/kg /inj)	U50,488 (0.0032–0.1 mg/kg /h, IV) Nor-BNI (3.2 mg/kg , IV)	Male Rhesus Monkey	Chronic U50,488 increased cocaine choice. Acute nor-BNI did not alter cocaine choice, but blocked U50,488 effects.	(Negus 2004)
7	Cocaine (0.001–0.3 mg/kg /inj)	Enadoline (0.001–0.01 mg/kg /h, IV)	Male Rhesus Monkey	Acute enadoline decreased cocaine and food-maintained responding.	(Bowen et al. 2003)
8	Cocaine (0.01–0.032 mg/kg /inj)	Enadoline (0.00032–0.0032 mg/kg /h, IV) Spiradoline (0.0032–0.018 mg/kg /h, IV) PD117302 (0.032–0.32 mg/kg /h, IV) Bremazocine (0.0032–0.032 mg/kg /h, IV) Mr2033 (0.0032–0.032 mg/kg /h, IV) Cyclazocine (0.001–0.1 mg/kg /h, IV)	Male and Female Rhesus Monkey	Repeated KOR agonist decreased rates of cocaine SA at doses that decreased food SA with one exception. Mr2033 selectively attenuate cocaine SA compared to food SA.	(Mello and Negus 1998)
9	Cocaine (0.01–0.032 mg/kg /inj)	Ethylketocyclazocine (EKC; 0.0032–0.032 mg/kg /h, IV) U50,488 (0.032–0.1 mg/kg /h, IV) Nor-BNI (3.2 mg/kg , IV)	Male and Female Rhesus Monkey	Repeated EKC and U50,488 treatment decreased rates of cocaine SA at doses that also tended to decrease food SA. Nor-BNI did not alter cocaine or food SA.	(Negus et al. 1997)
10	Smoke Cocaine (1 mg/kg /delivery)	Bremazocine (0.00032, 0.001, 0.0025 mg/kg , IM)	Male Rhesus Monkey	Repeated bremazocine decreased rates of cocaine, EtOH, saccharin, PCP, and food SA.	(Cosgrove and Carroll 2002)

#	Self-administered Drug (dose)	KOR ligand (dose)	Sex and Species	Main Effect	Ref
	PCP (0.25 mg/mL) EtOH (not reported)				
11	Cocaine (0.4 mg/kg/inj)	Cyclazocine (0–1 8 mg/kg, PO)	Female Long-Evans Rat	Acute racemic, but not enantiomers, cyclazocine decreased rates of cocaine SA, but not water-maintained behavior	(Glick et al. 1998)
12	Cocaine (0.015–1 mg/kg/inj)	U69,593 (0, 0.32 mg/kg, SC)	Male S-D Rat	Acute U69,593 decreased rates of low dose, but not high dose cocaine SA	(Schenk et al. 1999)
13	Cocaine (0.5 mg/kg/inj)	U69,593 (0, 0.32 mg/kg, SC)	Male S-D Rat	Acute U69,593 decreased rates of cocaine SA when cocaine delivery was paired with a visual stimulus.	(Schenk et al. 2001)
14	Nicotine (0.03 mg/kg/inj)	U50,488 (0.3–3 mg/kg, IP)	Male Wistar Rat	Repeated U50,488 treatment decreased rates of nicotine self-administration at doses that did not alter food-maintained responding	(Ismayilova and Shoaib 2010)
15	EtOH (10% v/v)	Bremazocine (0.1 mg/kg, SC) U50,488 (10 mg/kg, SC)	Male Wistar Rat	Repeated bremazocine produced sustained decrease in EtOH SA. Repeated U50,488 decreased EtOH SA on day 1 only.	(Nestby et al. 1999)
16	EtOH (20% v/v)	Enadoline (0.01 mg/kg/h, SC) Nor-BNI (5 mg/kg, IP × 2)	Male Wistar Rat	Repeated enadoline increased EtOH SA. Acute enadoline decreased EtOH SA. Acute nor-BNI had no effect.	(Hölter et al. 2000)
		<i>KOR Antagonists Studied</i>			
17	Heroin (0–0.1 mg/kg/inj)	5'-guanidinaltrindole (GNTI; 1.0 mg/kg, IM)	Male Rhesus Monkey	Acute GNTI did not attenuate withdrawal-associated increases in heroin choice.	(Negus and Rice 2009)
18	Heroin (0.06 mg/kg/inj)	Nor-BNI (5 or 10 mg/kg, SC)	Male Wistar Rat	Acute nor-BNI did not alter heroin SA.	(Negus et al. 1993)
19	Heroin (60 µg/kg/inj)	Nor-BNI (30 mg/kg, SC or 4 µg/side)	Male Wistar Rat	Acute nor-BNI blocked increased rates "escalation" of heroin SA	(Schlosburg et al. 2013)
20	Cocaine (0–0.1 mg/kg/inj)	Nor-BNI (3.2–10 mg/kg, IM)	Male Rhesus Monkey	Nor-BNI did not block or reverse cocaine choice or rates of cocaine SA	(Hutsell et al. 2016)
21	Cocaine (0.03, 0.06 mg/kg/inj)	Nor-BNI (30 mg/kg, SC)	Male Wistar Rat	Acute nor-BNI decreased rates of cocaine SA at 0.03, but not 0.06 mg/kg/inj cocaine.	(Kuzmin et al. 1998)
22	Cocaine (0.5 mg/kg/inj)	Nor-BNI (15, 30 mg/kg, SC)	Male Wistar Rat	Nor-BNI blocked increased "escalated" rates of cocaine SA.	(Wee et al. 2009)
23	Cocaine (0.5 mg/kg/inj)	Buprenorphine (3 mg/kg alone or in combination with 0.3–10 mg/kg, naltrexone)	Male Wistar Rat	Repeated 2-day Buprenorphine/naltrexone blocked increased rates "escalation" of heroin SA and reduced cocaine PR responding.	(Wee et al. 2012)
24	Methamphetamine (0.05 mg/kg/inj)	Nor-BNI (30 mg/kg, SC)	Male Wistar Rat	Acute nor-BNI blocked increased rates "escalation" of methamphetamine SA	(Whitfield et al. 2015)
25	Nicotine (0.03 mg/kg/inj)	GNTI (0–1 mg/kg, IP)	Male S-D Rat	GNTI pretreatment failed to alter rates of nicotine self-administration	(Liu and Jernigan 2011)
26	EtOH (10% v/v)	Nor-BNI (10 mg/kg, SC)	Male Lewis Rat	Acute nor-BNI increased EtOH consumption.	(Mitchell et al. 2005)

#	Self-administered Drug (dose)	KOR ligand (dose)	Sex and Species	Main Effect	Ref
27	EtOH (10% v/v) w/sucrose	Nor-BNI (0 or 20 mg/kg, SC)	Male Long-Evans Rat	Acute nor-BNI did not alter EtOH SA.	(Doyon et al. 2006)
28	EtOH (10% v/v)	Nor-BNI (5 mg/kg, SC)	Male Wistar Rat	Nor-BNI blocked increased "escalated" rates of EtOH SA.	(Walker et al. 2011)
29	EtOH (10% v/v)	JDTic (0–10 mg/kg, IP) Nor-BNI (0, 30 mg/kg, SC)	Male Wistar Rat	JDTic and nor-BNI decreased rates of EtOH SA.	(Schank et al. 2012)
30	EtOH (15% v/v)	JDTic (0–10 mg/kg, SC)	Female alcohol-preferring (P) Rat	JDTic did not alter EtOH SA.	(Deehan et al. 2012)
31	EtOH (15% v/v)	LY2456302 (3–10 mg/kg, PO)	Female P Rat	Acute LY2456302 decreased the amount of EtOH consumed	(Rorick-Kehn et al. 2014)
32	EtOH/saccharin/sucrose (10%/0.125%/3%)	Nor-BNI (0–10 mg/kg, SC)	Male and Female S-D Rat	Nor-BNI increased ethanol/saccharin consumption in male rats, decreased in female rats. Nor-BNI had no effect on adolescent EtOH/saccharin consumption.	(Morales et al. 2014)
33	EtOH (10% v/v)	Compound 5 (0.0031–0.0125 mg/kg, SC)	Male P Rat	Acute compound 5 (KOR antagonist) decreased rates of EtOH SA	(Cashman and Azar 2014)
34	EtOH (20% v/v)	CERC-501 (0–10 mg/kg, PO)	Male Wistar Rat	Acute CERC-501 decreases rates of EtOH SA	(Domi et al. 2018)