environmental services workers who had passed the temperature and symptom screening evaluation on entry to the campus. Samples were obtained from both nares of consenting individuals and ran daily on a validated polymerase chain reaction (PCR) platform.

Results: A total of 1394 employees consented to participate by June 15, 2020 and none of them had a positive result for COVID-19. Thus, the prevalence rate among asymptomatic employees was zero during this period (95%CI 0%-0.26%). Thirteen employees developed symptoms after initial enrollment and testing of which 1 tested positive for COVID-19. Based on these data, we estimate a monthly incidence rate of 0.8 new COVID-19 cases per 1000 asymptomatic employees (95%CI 0.2 - 4.4 per 1000). Of note, at the time of submission the R_0 for our region was 1.4, and the prevalence of COVID-19 infection among symptomatic employees was 9.9% (68/686 tested in our system, 95%CI: 7.8%-12.3%).

Conclusion: We did not identify any COVID-19-positive asymptomatic hospital employees who passed screening measures. This suggests very low risk of nosocomial transmission to other employees, patients, and families. Along with low community prevalence and capture of COVID-19-positive symptomatic employees, we could confidently advise staff that universal surgical masking and eye protection were likely adequate to prevent significant exposure.

Disclosures: All Authors: No reported disclosures

464. Rapid whole genome sequence typing reveals multiple waves of SARS-CoV-2 spread

Ahmed M. Moustafa, PhD¹; Paul J. Planet, MD, PhD²; Paul J. Planet, MD, PhD²; ¹Sackler Institute for Comparative Genomics, American Museum of Natural History, New York, NY; The Children's Hospital of Philadelphia, Philadelphia, PA., Philadelphia, PA; ²Children's Hospital of Philadelphia/UPenn, Philadelphia, Pennsylvania

Session: P-14. COVID-19 Epidemiology and Screening

Background: As the pandemic SARS-CoV-2 virus has spread globally its genome has diversified and distinct clones can now be recognized, tracked, and traced. Identifying clonal groups allows for assessment of geographic spread, transmission events, and identification of more virulent or transmissible emerging strains.

Methods: All SARS-CoV-2 genomes (n=17,504) that are complete and high coverage were downloaded from GISAID on May 17th 2020. We developed a GNU-based Virus Identification (GNUVID) tool that implements a whole genome multilocus sequence typing (wgMLST) scheme composed of all ten ORFs in the SARS-CoV-2 genome. The 10,422 genomes that passed our quality check were fed to the GNUVID tool, which assigned a ST profile to each genome. Global optimum eBURST was then used to cluster the STs in clonal complexes (CCs).

Results: Our ST/CC analysis uncovered strong associations of ST/CCs with certain geographical regions but also dynamic local changes in ST/CC prevalence. We also identified several unexpected putative global transmission events (e.g., from the US to the Middle East and reintroduction to China later in the pandemic). We have made our tool (GNUVID) available so that new WG sequences can be rapidly assigned to an ST/CC (<https://github.com/ahmedmagds/GNUVID>).

Conclusion: Our sequence typing system uncovered previously unappreciated transmission events and waves of expansion and replacement of SARS-CoV-2 STs and CCs in different geographical locations, suggesting complex dynamics in viral populations that previously seemed monomorphic. Because, our tool can be rapidly updated with new sequencing data it can track emerging clones and identifying new hotspots.

Disclosures: All Authors: No reported disclosures

465. Risk Factors Associated with Critical COVID-19 Requiring Mechanical Ventilation

Robert B. Chris, MD MSc¹; Rabeeya Khalid, MD¹; Urmila Ravichandran, MS²; Mary Ellen Acree, MD; Jennifer Grant, MD²; Chethra Muthiah, MD²; Nirav Shah, MD, MPH²; ¹University of Chicago (NorthShore), Evanston, Illinois; ²NorthShore University HealthSystem, Evanston, Illinois

Session: P-14. COVID-19 Epidemiology and Screening

Background: Mechanical ventilation of patients with COVID-19 is associated with high mortality. Understanding risk factors for developing mechanical ventilation may allow for more targeted monitoring and therapeutics that may improve outcomes.

Methods: We performed a retrospective case series of all patients admitted within thirty days of a positive Sars CoV-2 test to an integrated health system near Chicago, Illinois between March 12 and May 31, 2020. Covariates evaluated included demographics, symptoms on admission, vital signs, medications, comorbidities, census tract data and social history. Univariable analysis was performed and variables with an alpha of 0.05 or less were included in multivariable regression modeling to identify factors associated with mechanical ventilation. The area under the curve (AUC) was used to assess performance of the model.

Results: Of 990 patients admitted with COVID-19 12.6% (125) were mechanically ventilated. The median age was 68 (interquartile range 55-82), 48.4% (479) were female and 49.6% (491) were Caucasian. Independent factors associated with mechanical ventilation included female sex (Adjusted OR [AOR] 0.621, Confidence Interval [CI] 0.427-0.903; p=0.0363), body mass index (BMI) (AOR 1.035, CI 1.011-1.060; p=0.0175), percent of english speaking population within patient's census tract (AOR 0.989, CI 0.979-0.998; p=0.0454), respiratory rate (AOR 1.054, CI 1.027-1.083; p=0.0011), oxygen saturation (AOR 0.922, CI 0.901-0.943; p< 0.0001),

cerebrovascular accident (CVA) (AOR 0.176, CI 0.051-0.605; p=0.0207) and hematologic malignancy (AOR 3.668, CI 1.403-9.590; p=0.0261). AUC of the model was 0.8 (0.75-0.84).

Characteristics of patients admitted with COVID-19

	Population N=990		Population N=990
Age (yr)	68.0 [55.0 - 82.0]	Past medical History	
Age <18	1 (0.1%)	Acute Myocardial Infarction	31 (3.1%)
Female	479 (48.4%)	Asthma	78 (7.9%)
Body Mass Index (Kg/m2)	28.3 [24.6 - 32.8]	Coronary Artery Disease	133 (13.4%)
Body Mass Index >=30	352 (35.6%)	Congestive Heart Failure	120 (12.1%)
Past or Current Smoker	255 (25.8%)	Chronic Kidney Disease	126 (12.7%)
Pregnant During Encounter	13 (1.3%)	Conduction Abnormalities	217 (21.9%)
Race		Chronic Obstructive Pulmonary Disease	119 (12.0%)
Caucasian	491 (49.6%)	Cerebrovascular Accident	93 (9.4%)
African American	92 (9.3%)	Diabetes	279 (28.2%)
Asian	75 (7.6%)	End Stage Renal Disease	32 (3.2%)
Other	332 (33.5%)	Hematologic Malignancy	22 (2.2%)
Ethnicity		Human Immunodeficiency Virus	2 (0.4%)
Hispanic/Latino	181 (18.3%)	Hypertension	482 (48.7%)
Non-Hispanic	809 (81.7%)	Inflammatory Bowel Disease	13 (1.3%)
Insurance		Liver Disease	56 (5.7%)
Private	277 (28.0%)	Metastatic Solid Tumor	14 (1.4%)
Medicare	503 (50.8%)	Systemic Lupus Erythematosus	4 (0.4%)
Medicaid	188 (19.0%)	Peripheral Vascular Disease	64 (6.5%)
Self-Pay/Unknown	22 (2.2%)	Solid Tumor	181 (18.3%)
Census Tract		Transplant	4 (0.4%)
Distance To Closest NorthShore Hospital (miles)	3.5 [2.0 - 5.8]	Medications Taken in the Past 6 Months	
Median Family Income in Thousands (\$)	76765.0 [57558.0 - 110666.0]	ACE/ARB	184 (18.6%)
Percent Of English Speaking Population	59.9 [43.5 - 74.5]	Chemotherapy	15 (1.5%)
Symptoms on Admission		Insulin	88 (8.9%)
Cough	133 (13.4%)	Statins	174 (17.6%)
Decreased Sense Of Taste Or Smell	0 (0.0%)	Steroid In Last Month	55 (5.6%)
Shortness Of Breath	411 (41.3%)	Relationship with NorthShore	
Fever	225 (22.7%)	Medical Group PCP	257 (26.0%)
Gastrointestinal Symptoms	60 (6.1%)	Affiliate PCP	20 (2.0%)
Fatigue	28 (2.8%)	NorthShore Encounter in Last 24 Months	457 (46.2%)
Myalgia	76 (7.7%)		
Upper Respiratory Symptoms	6 (0.6%)		
First Vital After Admission			
Temperature (°F)	99.4 [98.5 - 100.9]		
Systolic Blood Pressure (mmHg)	129.0 [116.0 - 144.0]		
Diastolic Blood Pressure (mmHg)	76.0 [66.0 - 85.0]		
Respiration (per minute)	22.0 [19.0 - 26.0]		
Pulse (per minute)	94.0 [81.2 - 108.0]		
Oxygen Saturation (percent)	94.0 [91.0 - 97.0]		

Multivariable logistic regression to identify risk factors associated with mechanical ventilation

	Adjusted Odds Ratio	P-Value
Age (yr)	0.997 [0.985-1.009]	0.7023
Female	0.621 [0.427-0.903]	0.0363
Body Mass Index (Kg/m2)	1.035 [1.011-1.060]	0.0175
Percent Of English Speaking Population	0.989 [0.979-0.998]	0.0454
Shortness Of Breath	1.244 [0.844-1.835]	0.3549
Respiration (per minute)	1.054 [1.027-1.083]	0.0011
Pulse (per minute)	1.005 [0.995-1.014]	0.4005
Oxygen Saturation (percent)	0.922 [0.901-0.943]	0.0000
Conduction Abnormalities	0.634 [0.374-1.077]	0.1569
Cerebrovascular Accident	0.176 [0.051-0.605]	0.0207
Hematologic Malignancy	3.668 [1.403-9.590]	0.0261
Non-Hispanic	1.125 [0.706-1.795]	0.6771

Conclusion: Risk factors associated with mechanical ventilation included male gender, elevated BMI, census tract with lower percentage of english speakers, increased respiratory rate, low oxygen saturation, hematologic malignancy and not having a CVA. We suspect that history of CVA may have been associated with overall patient debility in which aggressive measures such as intubation were not deemed appropriate. Identifying patients with risk factors associated with mechanical ventilation may allow for early and targeted interventions to improve outcomes.

Disclosures: All Authors: No reported disclosures

466. SARS-CoV-2 Diagnosis And Point Prevalence in a Non-Cohorted Tertiary Care Center

Spencer Schrank, DO¹; Katherine McAleese, MD¹; Amanda B. Spence, MD²; Madhuri Natarajan, MS³; Joseph Timponi, MD³; Joseph Chang, MD³; Gayle p. Balba, MD⁶; Princy Kumar, MD⁶; ¹MedStar Georgetown University Hospital, Washington, District of Columbia; ²Georgetown University Medical Center, Washington, District of Columbia; ³Medstar Georgetown University Hospital, Washington, DC; ⁴Georgetown University, Washington, DC; ⁵Fellow, Washington, District of Columbia; ⁶Georgetown University School of Medicine, Washington, District of Columbia

Session: P-14. COVID-19 Epidemiology and Screening

Background: The CDC recommends testing for SARS-CoV-2 in patients who present with symptoms consistent with COVID-19 and to cohort hospitalized patients diagnosed with COVID-19. Up to 35% of persons infected with SARS-CoV-2 are asymptomatic; however, no recommendations exist for universal testing in hospitalized patients. We assessed the point prevalence of SARS-CoV-2 infection amongst hospitalized patients at a tertiary care center during a time when there was a regional surge of cases.

Methods: Nasopharyngeal SARS-CoV-2 PCR testing was performed on inpatients at Georgetown University Hospital on 4/27/20, excluding those who were SARS-CoV-2 positive, tested within 72 hours or admitted to pediatric, psychiatric, labor & delivery or ICUs. Patients within the hospital were not cohorted based on COVID-19