Figure 3

Conclusion: Infectious disease providers are trained to consider antimicrobial stewardship. Antimicrobials directly impact therapeutic choices for patients. However, less frequently considered is the insertion of PICCs in patients with tenuous renal function. Given the prevalence of PICCs in patients with GFR < 45, contrary to guideline recommendations, as well as increasing evidence for use of shorter durations and oral antibiotics for treatment there is opportunity for quality improvement and inter-sectional synergy.

Disclosures: All Authors: No reported disclosures

123. Utilization of the BioFire® FilmArray® Pneumonia Panel in Critically Ill Patients with Influenza Pneumonia

Benjamin Ereshefsky, PharmD, BCIDP¹; Mina Raju, DO, American Board Of Internal Medicine - Internal Medicine, American Board Of Internal Medicine - Infectious Disease¹; Kaweah Delta Medical Center, Visalia, California

Session: P-4. Antimicrobial Stewardship: Diagnostics/Diagnostic Stewardship

Background: Severe influenza pneumonia is associated with poor clinical outcomes and may be complicated by bacterial co-infections. Traditional culture techniques may fail to identify organisms, especially if antibiotics were administered prior to culture collection. In critically ill patients, negative cultures may lead to prolonged use of empiric, broad-spectrum antibiotics.

Methods: This is a single-center descriptive study on the use of the BioFire® FilmArray® Pneumonia (PNA) panel in critically ill patients with influenza. All intubated patients diagnosed with influenza pneumonia admitted to an ICU between December 1, 2019 and March 31, 2020 were included for analysis. The purpose of this study was to compare the results from rapid diagnostic testing with traditional culture results, and to describe the impact on time to results and the time to de-escalation of antibiotics.

Results: A total of 21 patients with 22 PNA panels were included. The mean age was 64 years, and 43% were female. Diabetes and COPD were present in 29% and 33%, respectively. A majority of patients (91%) had influenza A. Bacterial co-infection was found in 82% of cases, with all having bacteria identified on the PNA panel. Polymicrobial results from the PNA panel occurred 32% of the time. The most frequently identified bacterial pathogens by PNA panel were *S. pneumoniae* (41%), MSSA (27%), *H. influenzae* (14%), and *P. aeruginosa* (9%). Respiratory cultures had no growth 50% of the time, with MSSA most commonly identified at 23%. *P. aeruginosa* and MRSA were the only organisms that were detected by both PNA panel and respiratory culture with 100% agreement. PNA panel results were available 5.1 (2.7–11.3) hours after respiratory culture collection, compared to 45.8 (31.5–55.6) hours for culture results. De-escalation of empiric antibiotics occurred 13.2 (1.7–24.0) hours after PNA panel results in the 18 evaluable patients.

Table 1: Pathogens Identified by BioFire Pneumonia Panel and Respiratory Culture

Table 1: Pathogens Identified by BioFire® Pneumonia Panel and Respiratory Culture

Organism	BioFire® Pneumonia (n=22)	Respiratory Culture (n=21)
Coronavirus	2 (9%)	N/A
Rhino/Enterovirus	2 (9%)	N/A
Influenza A	20 (91%)	N/A
Influenza B	2 (9%)	N/A
<i>S. pneumoniae</i>	9 (41%)	1 (5%)
MSSA	6 (27%)	5 (23%)
MRSA	1 (5%)	1 (5%)
<i>S. agalactiae</i>	2 (9%)	0
<i>H. influenzae</i>	3 (14%)	0
<i>M. catarrhalis</i>	2 (9%)	0
<i>P. aeruginosa</i>	2 (9%)	2 (9%)
<i>K. pneumoniae</i>	1 (5%)	0
<i>S. marcescens</i>	1 (5%)	0
<i>S. maltophilia</i>	N/A	1 (5%)

Conclusion: The PNA panel identified a high rate of bacterial pathogens in this cohort of critically ill patients with influenza. It also provided actionable results much quicker than traditional respiratory cultures. Further study is needed to evaluate the significance of positive molecular tests when cultures are negative.

Disclosures: All Authors: No reported disclosures

124. Acute Uncomplicated Cystitis in Urgent Care Centers: Changes in Antibiotic Prescribing Practices Before and After Provider Education

Rebecca S. Cofsky, PharmD, BCIDP¹; David Goodman, Bachelors of Science in Kinesiology²; Arianna Morton, MLS (ASCP)³; Debra L. Powell, MD¹; Edward Blanchard, MD¹; ¹Reading Hospital, Sinking Spring, Pennsylvania; ²Pottstown Hospital, Reading, Pennsylvania; ³PCOM, Reading, Pennsylvania

Session: P-5. Antimicrobial Stewardship: Non-Inpatient Settings

Background: Acute uncomplicated cystitis (AUC) is a frequent infection that requires antibiotic treatment that is seen in women who seek care at urgent care centers.¹ The Infectious Diseases Society of America (IDSA) has published guidelines for the management of AUC. They recommend nitrofurantoin, trimethoprim-sulfamethoxazole, or fosfomycin as the first-line agents for the treatment of AUC in healthy adult women.¹ Previous evaluation of prescribing practices at Tower Health Urgent Care Centers demonstrated a 16% compliance rate with treatment guidelines for AUC. Subsequently, provider education and an AUC medication order set in the electronic medical record (EMR) were implemented. The purpose of this study was to evaluate the effectiveness of these interventions on prescribing practices for AUC.

Methods: A retrospective electronic chart review was performed evaluating the antibiotic treatment of patients diagnosed with AUC between January 1, 2018, to June 30, 2019, excluding July 2018 to September 2018, the washout period during which education was provided. Inclusion criteria included age ≥ 18 years, female, and treated at one of three Tower Health Urgent Care Centers.

Results: 1,455 patients met inclusion criteria during our study period. Our three-month washout period included 273 of those patients. 421 patients were included in the pre-education analysis with 761 patients in the post-education analysis. When both first and second-line antibiotic prescriptions were evaluated in conjunction with considering patient allergies and concomitant infections, compliance was 59% in our post-education analysis (improved from 16% compliance at baseline).

Conclusion: Provider education and an AUC medication order set in the EMR were successful in increasing the compliance rate with antibiotic treatment guidelines for the management of AUC in adult females in the urgent care setting.

Disclosures: All Authors: No reported disclosures

125. Comparison of Diagnosis and Prescribing Practices between Virtual Visits and Office Visits for Adults Diagnosed with Uncomplicated Urinary Tract Infections within a Primary Care Network

Kaitlyn Johnson, PharmD¹; Lisa E. Dumkow, PharmD, BCPS, BCIDP¹; Lisa Salvati, PharmD²; Kristen Johnson, PharmD³; Megan Yee, MD³; Nnaemeka Egwuatu, MD, MPH¹; ¹Mercy Health Saint Mary's, Grand Rapids, Michigan; ²Ferris State University, Grand Rapids, Michigan; ³Mercy Health Physician Partners, Grand Rapids, MI

Session: P-5. Antimicrobial Stewardship: Non-Inpatient Settings

Background: Urinary tract infections (UTIs) are one of the most common infectious indications for antibiotic prescribing in the outpatient setting. With the exponential growth of virtual visits over the past decade, virtual visits represent an important ambulatory care target for antimicrobial stewardship programs outside of traditional office visits. This study aimed to compare the appropriateness of antimicrobial therapy between virtual visits and office visits for adult females diagnosed with uncomplicated UTIs within a primary care network.

Methods: This retrospective cohort study evaluated adult female patients diagnosed with a UTI within a primary care network comprised of 44 outpatient sites. The primary objective was to compare guideline-concordant antibiotic prescribing between virtual visits and office visits. Guideline-concordance was determined based on local antibiogram-based treatment recommendations. Secondary objectives included comparing appropriate treatment duration and use of diagnostic testing resources between groups. Additionally, patient outcomes were compared between groups including 48-hour, 7-day, and 30-day re-visits, or development of *Clostridioides difficile* infection within 30 days.

Results: A total of 350 patients were included in this study, with 175 patients in each group. Patients treated for a UTI via a virtual visit were more likely to be prescribed a first-line antibiotic (74.9% vs 59.4%; $P = 0.002$). Additionally, virtual visits were more likely to prescribe an appropriate duration (100% vs 53.1%; $P = < 0.0001$). Patients treated via office visits were more likely to have a urinalysis (0% vs 97.1%; $P < 0.001$) and urine culture (0% vs 73.1%; $P < 0.0001$) ordered. There was no difference between groups in 48-hour or 30-day revisits, however, patients completing office visits were more likely to have a revisit within 7 days (18.9% vs 5.1%; $P < 0.0001$). In multivariate logistic regression, UTI care via office visit was the only independent risk factor for 7-day revisit (OR 3.74, 95% CI 1.31–10.67).

Conclusion: In adult female patients presenting with uncomplicated UTIs, care at a virtual visit was associated with significantly improved antimicrobial prescribing compared to office visits and decreased utilization of diagnostic and follow-up resources.

Disclosures: All Authors: No reported disclosures