Brazilian Journal of Physical Therapy

<https://www.journals.elsevier.com/brazilian-journal-of-physical-therapy>

ORIGINAL RESEARCH

Can *Kinesio Taping*® influence the electromyographic signal intensity of trunk extensor muscles in patients with chronic low back pain? A randomized controlled trial

Leandro Garcia Pires, Rosimeire Simprini Padula, Maurício Antônio Da Luz Junior, Irlei Santos, Matheus Oliveira Almeida, Shaiane Silva Tomazoni, Lucíola Cunha Menezes Costa, Leonardo Oliveira Pena Costa*

Masters and Doctoral Programs in Physical Therapy, Universidade Cidade de São Paulo (UNICID), São Paulo, SP, Brazil

Received 3 July 2019; received in revised form 10 October 2019; accepted 6 December 2019

Available online 15 December 2019

KEYWORDS

Chronic low back pain;
Kinesio taping;
Electromyography;
Physical therapy

Abstract

Background: The evidence of the influence of *Kinesio Taping*® in changing electromyographic signal intensity of the lumbar musculature in patients with chronic non-specific low back pain (LBP) is very sparse.

Objectives: To evaluate if *Kinesio Taping*® changes the electromyographic signal intensity of the longissimus and iliocostalis muscles in patients with chronic non-specific LBP.

Methods: Prospectively registered, three-arm randomized controlled trial with a blinded assessor. Patients were randomly allocated to the following interventions: 1) *Kinesio Taping*® Group (n = 21), where patients received the tape according to the manufacturer's manual; 2) Placebo Group (i.e. normal surgical tape) (n = 21); and 3) Non-treatment control Group (n = 21). Assessments were performed at baseline, immediately after, and 30 min after the intervention. The primary outcome was muscle activity of the iliocostalis and longissimus muscles as measured by surface electromyography. The secondary outcome was pain intensity (measured with a 0–10 Numerical Rating Scale). The effects of treatment were calculated using linear mixed models.

Results: A total of 63 patients were recruited. Follow up rate was high (98.4%). Patients were mostly women with moderate levels of pain and disability. *Kinesio Taping*® was better than the control and placebo groups in only 4 of 96 statistical comparisons, likely reflective of type I error due to multiple comparisons. No statistically significant differences were identified for the immediate reduction in pain intensity between groups.

* Corresponding author at: Masters and Doctoral Programs in Physical Therapy, Universidade Cidade de São Paulo, Rua Cesário Galeno, 448 - Tatuapé, CEP: 03071-000, São Paulo, SP, Brazil.
E-mail: lc3060@gmail.com (L.O. Costa).

Conclusion: *Kinesio Taping*® did not change the electromyographic signal intensity of the longissimus and iliocostalis muscles or reduce pain intensity in patients with chronic low back pain.

Clinicaltrials.gov: NCT02759757 (<https://clinicaltrials.gov/ct2/show/NCT02759757>)
© 2019 Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia. Published by Elsevier Editora Ltda. All rights reserved.

Introduction

Low back pain (LBP) is one of the most prevalent public health issues worldwide.¹ According to the most recent Global Burden of Disease Study, LBP is the leading cause of years lived with disability.^{2,3} Episodes of acute LBP usually have a good prognosis, with significant improvement in the first few weeks.⁴ After this time, however, more than 40% of patients are likely to develop chronic LBP (pain that persists for more than 12 weeks).⁵ Studies have shown that the use of health services due to LBP has increased in recent years,⁶ and patients with chronic LBP tend to seek more treatment than patients with acute LBP.^{7,8}

Non-pharmacological interventions such as exercise, manual therapy, psychological therapies, and multidisciplinary biopsychosocial rehabilitation⁹ are the first options in the treatment of patients with chronic LBP.^{9–11} The latest guidelines published for the treatment of LBP have recommended that medication should only be used in patients who have not improved with non-pharmacological treatment due to the potential adverse effects and limited effectiveness of drugs.^{9,12}

Kinesio Taping (KT)[®] has become very popular, especially among athletes,¹³ and has also been proposed for the treatment of patients with LBP.^{14,15} KT is an intervention consisting of the application of a porous, adhesive elastic bandage, without chemicals or drugs.¹⁶ KT can extend up to 140% of its original length, allowing good range of motion when compared to other types of tape.^{16,17} The creators of the technique claim that KT reduces pain intensity and abnormal muscle tension, change joint position and movement perception, and reduces muscle spasms by reducing muscle activation, among other mechanisms of action.^{17,18} Patients with chronic LBP typically experience moderate levels of pain and disability. In addition, superficial trunk muscles (such as the longissimus and iliocostalis muscles) are usually over-activated in response to pain stimulus.^{19,20,21}

Although the creators of the technique propose that KT reduces muscle activation (ie, electromyographic (EMG) signal intensity), only three studies,^{15,22,23} to date, have investigated this possible mechanism. Because these trials presented methodological limitations related to study design, small sample size, and inadequate description of EMG parameters, further investigation is warranted. Therefore, we conducted a clinical trial aimed to determine the effects of *Kinesio Taping*® on the EMG signal intensity of the lumbar musculature of patients with chronic LBP.

Methods

Design

A three-arm, randomized controlled trial with a blinded assessor was conducted.

Ethics and clinical trial registration

The project was approved by the Research Ethics Committee of Universidade Cidade de São Paulo (UNICID), São Paulo, Brazil (#50548115.8.0000.0064) and was prospectively registered on Clinicaltrials.gov (NCT02759757). All eligible participants were informed about the objectives of the study and were asked to complete an informed consent form. The study was conducted at the Physical Therapy Clinic of the Universidade Cidade de São Paulo.

Study population

The present study included participants of both sexes, aged between 18 and 60 years, presenting with chronic non-specific LBP (pain or discomfort with duration of at least three months between the costal margins and the inferior gluteal folds, with or without referred pain to the lower limbs). Patients who had previously used KT for their LBP and/or any other adhesive tape and patients with any contraindication to their use were excluded from the study. Other exclusion criteria were: serious spinal pathologies (fracture, tumor, inflammatory or infectious disease), previous spinal surgery, patients with nerve root compromise, decompensated cardiorespiratory and metabolic diseases, pregnancy, and previous use of KT for any condition.

Procedure

Initially, an individual physical examination was conducted,⁶ including collection of sociodemographic (such as age, gender, marital status, and education levels) and anthropometric data (such as weight, height, and body mass index) and assessment of pain intensity using the 11-point Numeric Pain Rating Scale (NPRS).²⁴ Disability was assessed with the Roland-Morris Disability Questionnaire (RMDQ)²⁵ scored from 0 (no disability) to 24 (high disability).

Next, the area to be tested was shaved and cleansed with 70% alcohol. The surface EMG electrodes were applied with T350 double-sided tape, and the electrogoniometer sensors, with T10 double-sided tape. The participants were positioned in slight trunk flexion for the instrumentation: placement of the electrogoniometer and surface EMG electrodes made according to SENIAM^{26–28} recommendations and placement of the lumbar dynamometry sensors. The EMG electrodes were applied bilaterally on the longissimus, two fingers width from the L1 spinous process and over the iliocostalis at the L2 level (Fig. 1a).²⁹ In addition, an electrogoniometer was positioned between the T12 and S1 vertebrae. One arm of the electrogoniometer was directed caudally and the other cranially with the slightly tensioned spring aligned with the spine (Fig. 1a). The outputs of each channel of the electrogoniometer were previously calibrated according to manufacturers' instructions to capture the angle variations during data collection.³⁰ Then the participant was positioned in trunk extension, from the position of 30 degrees of trunk flexion measured by a common goniometer (Fig. 1b),³¹ as well as extension of the cervical spine, elbows, and knees.

Subsequently, the participants received instructions from the therapist to become familiar with the equipment through the trunk extension movement. From that position, the initial evaluation (pre-intervention) was conducted for a maximum voluntary isometric contraction (MVIC) test, in a sustained position for 10 s, with the EMG signal synchronized to the electrogoniometer and dynamometer data.³² The first two seconds and the final two seconds were disregarded to minimize signal contamination, therefore only the central six seconds of data collection were used for analysis.

After this pre-intervention assessment, participants were randomized into three treatment groups: Intervention/KT, Placebo, or Control.

EMG processing and data analysis

An electrogoniometer synchronized to the EMG was used to simultaneously capture the range of motion of the thoracolumbar region during the MVIC test on the lumbar dynamometer. The lumbar dynamometer also recorded the mean and maximum trunk muscle strength. The EMG signal was processed and analyzed using the Biometrics EMG software for DataLOG version 8.51. The raw EMG data were captured and the software provided the Root Mean Square (RMS) and Median Frequency (MF) values for the right/left iliocostalis and longissimus muscles. Data were collected for 10 s, with the initial and final seconds subsequently trimmed and eliminated. The remaining six-second window was analyzed. The raw EMG signal was used as reference in this study,³³ because in situations of the EMG set-up and controlled environment, the use of the non-normalization signal presents good reliability.³⁴

Randomization

The randomization process was generated by a software program (Excel Office 2010) and performed by a researcher not involved with data collection. Concealed allocation was achieved through the use of sequentially numbered, sealed, and opaque envelopes. After the initial evaluation, the

participants were randomly allocated to one of the three groups: *Kinesio Taping* group; Placebo group, and Control group.

Interventions

Kinesio Taping Group: Patients allocated to this group received the Tex Gold *Kinesio Taping*[®] bandage according to manufacturers' instructions.¹⁷ The bandage was applied in the shape of an 'I' in the direction of insertion (S1 region) to the origin (T12 region), with tension of 10–15% (Paper-off), in the trunk flexion position. This procedure was conducted minutes after instrumentation and pre-intervention assessment. KT was applied over the EMG electrodes monitoring the longissimus muscles at the L1 level (portion of the longissimus dorsi; Fig. 1c).

Placebo Group: Patients allocated to this group had a Missner[®] surgical tape (5 cm in width) placed on the longissimus muscle. The form of application adopted generated folds in the skin, and an 'I' shape in the trunk flexion position. The tape was applied over the EMG electrodes, bilaterally to the spine towards the insertion at S1 to the origin at T12, minutes after instrumentation (Fig. 1d).

Control Group: The patients allocated to this group did not receive any type of bandage.

Immediately after randomization and application of the intervention, the participants repeated the maximum voluntary isometric contraction (MVIC) test, in a sustained position for 10 s, as performed pre-intervention. The same test was again performed 30 min post-intervention with the tape remaining in place.

Outcome measures

The primary and secondary outcome measures of the study were obtained by an assessor who was unaware of the allocation of patients to their treatment groups by covering their back with a shirt. The primary outcome of the study was the EMG signal of the iliocostalis and longissimus muscles.

The EMG signal was acquired with an eight channel Biometrics DataLog MWX8 wireless data system via SX230–1000 EMG electrodes (Biometrics Ltd., Gwent, UK) with the following specifications: active (single differential), bipolar, reusable surface disk electrodes with eighth order preamplifier and elliptical filter (-60 dB at 550 Hz), 38 mm long, 20 mm wide, 0.6 mm thick, 10 mm diameter discs, 20–450 Hz band-pass filter, and third order high pass filter (18 dB/octave; 20 Hz), CMRR > 96 dB; noise ratio < 5 μ V Input impedance > 10¹⁵ Ω Ohms. The connection cables of each channel were previously tagged with the name of the evaluated muscles and the tags remained in place for all data collections. The EMG signal was amplified (K800) and sampled at 1000 Hz and data were reported according to the International Society of Electrophysiology and Kinesiology (ISEK) guidelines. A reference electrode (Biometrics R206) was placed on the right wrist.³⁵ In our study, crosstalk was minimized through wireless transmission and the use of surface sensors with amplifiers and internal filters. This choice was based on the small difference in crosstalk when compared to intramuscular electrodes,³⁶ as well as good reliability.³⁷ In addition, the MVIC angle was limited to 30°, minimiz-

Figure 1 Placement of the EMG electrodes and electrogoniometer (A) Initial position on the dynamometer (B), KT application (C) Missner[®] surgical tape[®] application (d).

ing sensor movement. A 20 Hz filter was also used, which is above the minimum recommended of 10 Hz,³⁸ thus minimizing contamination by up to 47%.³⁹ In the EMG literature, signal normalization is important to eliminate signal variability between evaluations⁴⁰ and this process depends very much on the type of activity in which signal was collected, e.g., static⁴¹ or dynamic condition,⁴² healthy or unhealthy participants,⁴³ by means of maximum voluntary contraction or not.⁴⁴ The EMG signal of the right and left iliocostalis and longissimus muscles was captured for signal amplitude analysis (RMS) and Medium Frequency (MF) in the initial evaluation (pre-intervention), after randomization (immediately post-intervention), and 30 min post-intervention.

Data collection was performed by three individuals (one therapist and two assessors). The therapist participated in 100% of the EMG data collections. Narayanaswami et al.⁴⁵ reported the importance of having the same therapist for good reliability with EMG equipment.

The secondary outcome of the study was pain intensity as measured by the 11-point NPRS scored which ranges from 0 (no pain) to 10 (worst possible pain).⁴⁶ Pain intensity was collected minutes before the initial evaluation (pre-intervention), shortly after randomization (immediately post-intervention), and 30 min after randomization (30 min post-intervention).

In addition to the collection of the primary and secondary outcomes, RMDQ^{25,47} was only administered pre-intervention to measure disability at baseline. The questionnaire consists of 24 items that describe activities that patients with LBP may have difficulty performing on a daily basis, where each

affirmative answer corresponds to one point on the scale. The final score of the RMDQ was determined by the sum of the values obtained: the higher the score, the greater the disability.^{25,46} Patients were instructed to complete the statements according to how they felt on the day of the evaluation.

Sample size calculation

The sample calculation of the study was performed to detect a difference of 2.8 units in RMS of the EMG signal with estimated standard deviation of 2.5 units of the EMG signal.⁴⁸ Assuming a statistical power of 95%, alpha of 5%, and a possible sample loss of up to 20%, a total of 63 participants (21 participants per group) were included.

Statistical analysis

The statistical analysis followed intention-to-treat principles (i.e. patients were analyzed into the groups they were originally allocated to).⁴⁹ The normality of the data was tested by visual inspection of histograms, and the characterization of the participants was presented using descriptive statistical tests. The primary analysis of interest was the difference in EMG signal between the 3 groups. The between-group differences (effects of treatment) and their respective 95% confidence intervals (CI) were calculated by constructing mixed linear models⁵⁰ using interaction terms (group versus time). Treatment effects were adjusted for

baseline values for all outcomes and for multiple comparisons. Missing data were handled using linear mixed models. The analyses were performed using the software SPSS 19.

Results

Recruitment and baseline evaluation

From a total of 100 potential participants identified between August and December 2016, 63 were considered eligible and were included in the study. Potential participants were excluded due to different ineligibility reasons: age over 60 years ($n = 25$), age under 18 years ($n = 1$), acute non-specific LBP ($n = 5$), previous spinal surgery ($n = 2$), knowledge of the *Kinesio Taping*[®] technique ($n = 2$), nerve root involvement ($n = 1$), and knee pain only ($n = 1$). All randomized patients were included in the statistical analysis (Fig. 2). One patient dropped out from the remaining assessments after been assessed at baseline due to personal issues. The characteristics of the included participants in the initial evaluation are described in Table 1. The baseline characteristics of all variables were similar in the three groups.

Primary and secondary outcomes

Table 2 presents the descriptive data of the primary and secondary variables at all time points. There was no significant change in muscle activity of the iliocostalis and longissimus muscles as measured by EMG amplitude (RMS) and frequency (MF) with the use of *Kinesio Taping*. Of the 96 statistical comparisons (Table 3), a statistically significant difference in EMG signal was only observed for the RMS max of the

Figure 2 Flow diagram of participants throughout the study.

right iliocostalis muscle and the medium frequency (MF) of the right longissimus muscle in favor of the *Kinesio Taping* group compared to the placebo (surgical tape) and control group.

Discussion

We aimed to determine whether KT would change the EMG signal intensity of the iliocostalis and longissimus muscles of patients with chronic non-specific LBP. Compared to control or placebo, KT reduced EMG signal in only 4 of 96 statistical comparisons, which suggests that these findings are spurious due to the high number of statistical comparisons. No between-group differences were identified for the immediate reduction in pain intensity.

Table 1 Characteristics of participants at baseline ($n = 63$).

Variables	Groups		
	<i>Kinesio Taping</i> [®] ($n = 21$)	Placebo ($n = 21$)	Control ($n = 21$)
Age (years)	46.0 $\pm$ 11.1	38.9 $\pm$ 10.3	45.3 $\pm$ 12.0
Male gender	4 (19.0)	6 (28.6)	6 (28.6)
Weight (kg)	72.9 $\pm$ 12.9	74.4 $\pm$ 16.3	72.9 $\pm$ 9.9
Height (m)	1.6 $\pm$ 0.1	1.6 $\pm$ 0.1	1.6 $\pm$ 0.1
Body mass index (kg/m ²)	27.2 $\pm$ 5.3	27.0 $\pm$ 4.1	27.2 $\pm$ 3.5
Marital status			
Single	7 (33.3)	9 (42.9)	4 (19.0)
Married	14 (66.7)	10 (47.6)	10 (47.6)
Divorced	0 (0.0)	2 (9.5)	7 (33.3)
Academic level education status			
Elementary school	5 (23.8)	1 (4.8)	3 (14.3)
Secondary school	9 (42.9)	18 (85.7)	11 (52.4)
University	7 (33.3)	2 (9.5)	7 (33.3)
Work compensation	0 (0)	0 (0)	0 (0)
Physically active	7 (33.3)	4 (19.0)	6 (28.6)
Use of medication	10 (47.6)	11 (52.4)	11 (52.4)
Smoker	3 (14.3)	1 (4.8)	2 (9.5)
Duration of symptoms (months)	96.9 $\pm$ 96.6	44.4 $\pm$ 52.6	58.2 $\pm$ 75.2
Pain intensity (0–10)	6.0 $\pm$ 2.4	7.2 $\pm$ 1.3	7.0 $\pm$ 2.5
Disability (0–24)	12.4 $\pm$ 6.3	13.2 $\pm$ 5.3	12.5 $\pm$ 6.7

Categorical variables are expressed in n (%) and continuous variables as mean $\pm$ standard deviation.

Table 2 Outcomes at baseline, after Intervention, and after 30 min.

Outcome	Baseline			Immediately after treatment			Assessment at 30 min		
	Kinesio Taping® Group	Placebo Group	Control Group	Kinesio Taping® Group	Placebo Group	Control Group	Kinesio Taping® Group	Placebo Group	Control Group
Pain (0–10)	6.00 ± 2.36	7.19 ± 1.32	7.00 ± 2.54	5.00 ± 2.55	6.00 ± 2.80	5.0 ± 3.20	4.00 ± 2.52	5.00 ± 3.20	5.0 ± 3.20
Disability (0–24)	12.38 ± 6.30	13.24 ± 5.29	12.52 ± 6.66	–	–	–	–	–	–
RMS Maximum Iliocostalis_R	0.48 ± 0.31	0.42 ± 0.20	0.54 ± 0.54	0.55 ± 0.45	0.42 ± 0.17	0.39 ± 0.24	0.52 ± 0.45	0.44 ± 0.22	0.42 ± 0.24
RMS Maximum Iliocostalis_L	0.58 ± 0.39	0.49 ± 0.32	0.46 ± 0.33	0.56 ± 0.50	0.46 ± 0.19	0.43 ± 0.23	0.54 ± 0.41	0.48 ± 0.21	0.45 ± 0.30
RMS Maximum Longi_R	0.67 ± 0.42	0.73 ± 0.37	0.61 ± 0.31	0.70 ± 0.45	0.67 ± 0.29	0.56 ± 0.40	0.79 ± 0.54	0.70 ± 0.27	0.60 ± 0.31
RMS Maximum Longi_L	0.87 ± 0.76	0.76 ± 0.46	0.63 ± 0.32	0.91 ± 0.80	0.75 ± 0.36	0.65 ± 0.30	0.75 ± 0.50	0.72 ± 0.32	0.60 ± 0.31
RMS Mean Erector_R	0.09 ± 0.05	0.07 ± 0.03	0.08 ± 0.05	0.10 ± 0.05	4.40 ± 19.82	0.11 ± 0.12	0.10 ± 0.06	0.08 ± 0.03	0.08 ± 0.05
RMS Mean Erector_L	0.13 ± 0.16	0.09 ± 0.04	0.07 ± 0.05	0.10 ± 0.06	4.64 ± 20.85	0.08 ± 0.05	0.10 ± 0.07	0.08 ± 0.03	0.08 ± 0.04
RMS Mean Longi_R	0.13 ± 0.07	0.14 ± 0.08	0.11 ± 0.06	0.12 ± 0.06	4.86 ± 21.69	0.11 ± 0.06	0.11 ± 0.06	0.12 ± 0.04	0.10 ± 0.06
RMS Average Longi_L	0.20 ± 0.41	0.17 ± 0.23	0.11 ± 0.06	0.20 ± 0.31	5.30 ± 23.75	0.10 ± 0.06	0.13 ± 0.10	0.12 ± 0.04	0.13 ± 0.10
MF Maximum Iliocostalis_R	88.12 ± 16.74	92.48 ± 18.11	90.86 ± 17.58	89.14 ± 17.50	91.41 ± 15.20	90.03 ± 15.84	88.83 ± 15.35	89.80 ± 11.24	88.35 ± 16.90
MF Maximum Iliocostalis_L	92.32 ± 19.85	96.03 ± 19.09	95.70 ± 20.31	90.84 ± 21.30	96.13 ± 17.70	94.88 ± 20.60	87.25 ± 22.85	96.04 ± 17.20	92.96 ± 16.94
MF Maximum Longi_R	95.15 ± 18.96	106.68 ± 18.91	102.45 ± 16.29	96.67 ± 19.96	106.78 ± 16.18	104.10 ± 12.57	96.02 ± 17.10	102.83 ± 15.95	102.58 ± 15.02
MF Maximum Longi_L	99.26 ± 19.57	106.70 ± 17.99	107.00 ± 21.87	99.39 ± 19.83	109.80 ± 15.16	104.16 ± 17.90	97.96 ± 17.38	104.03 ± 13.30	102.89 ± 18.18
MF Mean Iliocostalis_R	81.31 ± 16.62	84.37 ± 13.87	84.23 ± 15.53	82.07 ± 17.30	84.27 ± 13.45	83.70 ± 14.86	81.73 ± 15.95	80.92 ± 11.28	83.53 ± 15.87
MF Mean Iliocostalis_L	85.91 ± 19.60	88.96 ± 18.82	88.09 ± 14.73	84.50 ± 20.40	89.55 ± 17.36	87.42 ± 14.42	80.77 ± 22.92	86.03 ± 16.43	85.92 ± 14.11
MF Mean Longi_R	88.14 ± 17.28	97.84 ± 14.46	94.18 ± 12.20	88.80 ± 17.80	96.94 ± 12.80	96.79 ± 13.15	88.37 ± 16.80	92.62 ± 11.45	95.46 ± 11.53
MF Mean Longi_L	91.19 ± 17.86	97.74 ± 14.93	97.62 ± 13.71	91.31 ± 18.50	99.02 ± 16.26	95.91 ± 13.50	90.06 ± 16.76	95.00 ± 14.50	94.87 ± 11.76

RMS = Root Mean Square; MF = Medium Frequency ; R = Right Side ; L = Left Side; Longi = Longissimus. Data are mean ± standard deviation.

Table 3 Effects of interventions.

Outcome	Mean between-group differences adjusted (95% CI)											
	Reassessment immediately after treatment						Reassessment 30 min after treatment					
	Kinesio × Placebo	p	Kinesio × Control	p	Placebo × Control	p	Kinesio × Placebo	p	Kinesio × Control	p	Placebo × Control	p
Pain Intensity (0–10)	−0.19 (−1.49, 1.11)	0.77	0.43 (−0.88, 1.73)	0.52	0.24 (−1.06, 1.54)	0.72	−0.57 (−1.87, 0.73)	0.39	0.05 (−1.25, 1.35)	0.94	−0.52 (−1.83, 0.78)	0.43
RMS Maximum Iliocostalis_R_	−0.03 (−0.18, 0.10)	0.62	0.18 (0.03, 0.32)*	0.01	0.15 (−0.00, 0.30)	0.05	−0.01 (− 0.17, 0.12)	0.80	0.15 (0.01, 0.30)*	0.03	0.14 (−0.00, 0.28)	0.05
RMS Maximum Iliocostalis_L_	−0.01 (−0.13, 0.10)	0.76	0.01 (−0.10, 0.13)	0.75	0.00 (−0.11, 0.12)	0.98	0.01 (−0.11, 0.13)	0.85	−0.02 (−0.14, 0.10)	0.73	0.00 (−0.13, 0.11)	0.88
RMS Maximum Longi_R_	−0.07 (−0.20, 0.04)	0.20	0.07 (−0.05, 0.19)	0.25	−0.00 (−0.13, 0.11)	0.91	−0.01 (−0.14, 0.11)	0.81	0.01 (−0.10, 0.15)	0.77	0.00 (−0.12, 0.12)	0.95
RMS Maximum Longi_L_	−0.05 (−0.22, 0.11)	0.52	0.01 (−0.15, 0.18)	0.85	−0.04 (−0.22, 0.14)	0.64	0.09 (−0.07, 0.26)	0.30	−0.10 (−0.27, 0.07)	0.30	−0.00 (−0.20, 0.20)	0.99
RMS Mean Iliocostalis_R_	4.33 (−1.40, 10.06)	0.14	−0.02 (−5.75, 5.70)	0.99	4.30 (−1.42, 10.04)	0.15	0.00 (−5.75, 5.80)	0.99	−0.00 (−5.72, 5.72)	1.00	0.00 (−5.75, 5.80)	0.99
RMS Mean Iliocostalis_L_	4.60 (−1.45, 10.61)	0.14	−0.05 (−6.07, 6.00)	0.99	4.55 (−1.50, 10.60)	0.14	0.02 (−6.04, 6.10)	0.99	−0.4 (−6.05, 6.00)	0.99	−0.00 (−6.07, 6.05)	0.99
RMS Mean Longi_R_	4.75 (−1.55 to 11.00)	0.15	−0.00 (−6.30, 6.30)	1.00	4.75 (−1.55, 11.00)	0.15	−0.01 (−6.31, 6.30)	0.99	−0.00 (−6.30, 6.30)	1.00	−0.01 (−6.35, 6.30)	0.99
RMS Mean Longi_L_	5.14 (−1.74, 12.00)	0.15	0.00 (−6.87, 6.90)	1.00	5.13 (−1.74, 12.00)	0.15	0.00 (−6.90, 6.92)	0.99	−0.08 (−7.00, 6.80)	0.99	−0.07 (−7.00, 6.90)	0.99

Table 3 (Continued)

Outcome	Mean between-group differences adjusted (95% CI)											
	Reassessment immediately after treatment						Reassessment 30 min after treatment					
	Kinesio × Placebo	p	Kinesio × Control	p	Placebo × Control	p	Kinesio × Placebo	p	Kinesio × Control	p	Placebo × Control	p
MF Maximum Iliocostalis_R_	−2.45 (−7.40 to 2.50)	0.33	1.85 (−3.10, 6.80)	0.46	−0.60 (−5.55, 4.33)	0.80	−2.22 (−7.30, 2.80)	0.40	3.22 (−1.75, 8.15)	0.20	0.99 (−4.00, 6.00)	0.70
MF Maximum Iliocostalis_L_	1.60 (−5.65, 8.83)	0.70	−0.66 (−7.92, 6.60)	0.85	0.91 (−6.33, 8.17)	0.80	6.24 (−1.06, 13.55)	0.09	−2.35 (−9.60, 4.90)	0.53	3.90 (−3.40, 11.20)	0.30
MF Maximum Longi_R_	−1.45 (−7.10, 4.25)	0.65	−0.12 (−5.80, 5.55)	0.99	−1.55 (−7.20, 4.15)	0.60	−4.25 (−9.95, 1.50)	0.15	0.75 (−4.92, 6.45)	0.80	−3.50 (−9.20, 2.22)	0.30
MF Maximum Longi_L_	2.97 (−2.87, 8.81)	0.31	3.00 (−2.90, 8.85)	0.35	5.95 (−0.09, 11.80)	0.04	−0.70 (−6.60, 5.20)	0.85	2.85 (−3.02, 8.70)	0.35	2.11 (−3.80, 8.01)	0.50
MF Mean Iliocostalis_R_	−0.86 (−4.05, 2.30)	0.60	1.30 (−1.90, 4.50)	0.45	0.45 (−2.75, 3.60)	0.80	−2.35 (−5.55, 0.88)	0.20	1.15 (−2.05, 4.30)	0.50	−1.20 (−4.40, 2.00)	0.50
MF Mean Iliocostalis_L_	2.02 (−4.01, 8.10)	0.50	−0.80 (−6.80, 5.30)	0.80	1.30 (−4.80, 7.30)	0.70	3.50 (−2.60, 9.60)	0.30	−3.00 (−9.00, 3.10)	0.35	0.55 (−5.60, 6.65)	0.90
MF Mean Longi_R_	−1.55 (−6.15, 3.05)	0.50	−2.00 (−6.55, 2.60)	0.40	−3.55 (−8.10, 1.05)	0.15	−4.90 (−9.50, −0.30)*	0.03	−1.05 (−5.65, 3.55)	0.65	−5.94 (−10.55, −1.35)*	0.01
MF Mean Longi_L_	1.15 (−3.15, 5.45)	0.60	1.85 (−2.50, 6.15)	0.40	3.00 (−1.30, 7.30)	0.20	−0.85 (−5.15, 3.55)	0.75	1.65 (−2.70, 5.95)	0.45	0.85 (−3.55, 5.15)	0.75

RMS = Root Mean Square; MF = Medium Frequency; R = Right Side; L = Left side; Longi = Longissimus; *Significant difference ($p < 0.05$).

To our knowledge, this is the first trial that aimed to identify the effects of *Kinesio Taping*[®], compared to a placebo treatment, on the EMG signal intensity of lumbar muscles of patients with LBP. The study's strengths were 1) control of the main sources of bias in clinical trials, such as concealed allocation, assessor blinding, and intention-to-treat analysis,⁵¹ 2) rigorous use of all SENIAM guidelines,⁵² and 3) strict adherence to ISEK (International Society of Electrophysiology and Kinesiology)³⁸ guidelines.

However, our study has some limitations. First of all, we did not include only patients with high levels of muscle activity as we assumed that this is very common in patients with LBP. Due to the nature of the interventions, it was not possible to blind the therapist and patients. Another limitation was that signal contamination could not be completely eliminated. Also, the tapes were placed on top of the EMG electrodes, however these electrodes were very small and it is unlikely to influence the outcomes. Although randomization generated similar groups in most variables, the duration of symptoms among the three treatment arms were different. Nevertheless all participants from this trial were patients with long-lasting chronic LBP. There is no evidence that the prognosis of a patient with 97 months (e.g. a patient from the KT group) would be different compared to a patient from the placebo group with 44 months of symptoms. Clinically speaking, it is unlikely that a physical therapist would consider these patients as separate subgroups. The most important systematic review⁵³ on the prognosis of patients with chronic LBP only provides data up to 12 months. Therefore, we do not have data to support or refute the hypothesis that these differences in duration of symptoms could influence the outcomes of the trial. Finally, we conducted quite a large number of statistical comparisons. Therefore the chances of type 1 errors (i.e. false positive findings) were very likely. We used linear mixed models to test between group differences adjusted for baseline estimates even knowing that these differences at baseline were not clinically relevant. Although we did not use corrections such as Bonferroni corrections, we ended up getting most regression coefficients not statistically significant. Only four of 96 estimates (4.2%) were statistically significant, very likely reflecting spurious findings.

Among all clinical trials on the efficacy of *Kinesio Taping*[®] performed in patients with chronic LBP^{14,54–56} three used EMG as an outcome.^{15,22,23,57} In these trials, no clinically important results were found for *Kinesio Taping*[®], which is consistent with our findings. Regarding the primary outcome measure (EMG signal), the difference between the treatment, control, and placebo groups was not statistically significant for most of the comparisons. Fong et al.²³ also did not observe a significant change in the EMG signal intensity of the lumbar multifidus muscles after the use of *Kinesio Taping*[®] compared to the control group that received no intervention. Similar results were observed from Paoloni et al.¹⁵, who measured EMG signals on a three-arm trial comparing exercises versus KT versus KT plus exercises in patients with LBP. These authors¹⁵ also did not find correlations among changes in EMG signals and changes in pain intensity. Conversely, a very small (n=20) and low quality trial conducted by Bae et al.²² concluded that KT changed EMG findings compared to "ordinary physical therapy". From a purely statistical point of view, the few statistically

significant findings may have been random and caused by type I statistical errors.^{58,59} Therefore, the prescription of KT with the goal to reduce muscle activation does not seem justified.

There is a large amount of clinical evidence, including two meta-analyses,^{13,60} four systematic reviews^{56,61–63} in musculoskeletal conditions, and one for LBP⁶⁴ that consistently conclude that the effects of *Kinesio Taping*[®] for these patients are not better than placebo.⁶⁵ Therefore, the current evidence does not support the use of KT in patients with chronic non-specific LBP. On the other hand, the number of studies on the mechanisms of action of KT in patients with musculoskeletal conditions is still very small. Therefore, more research is needed in this area.

We conclude that KT did not change the EMG signal intensity of the longissimus and iliocostalis muscles or reduced pain intensity in patients with chronic LBP. This specific proposed mechanism of action of this intervention was not supported by the findings of this trial.

Acknowledgement

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. The first author of this manuscript received a scholarship from CAPES – Brazil (finance code 001).

Conflict of interest

The authors declare that there is no conflict of interest.

References

1. Ferreira G, Costa LM, Stein A, et al. Tackling low back pain in Brazil: a wake-up call. *Braz J Phys Ther.* 2019;23(3):189–195.
2. Disease GBD, Injury I, Prevalence C. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet.* 2017;390:1211–1259.
3. Maher C, Underwood M, Buchbinder R. Non-specific low back pain. *Lancet.* 2017;389:736–747.
4. da Costa LMC, Maher CG, Hancock MJ, et al. The prognosis of acute and persistent low-back pain: a meta-analysis. *CMAJ.* 2012;184:E613–624.
5. da Costa LMC, Maher CG, McAuley JH, et al. Prognosis for patients with chronic low back pain: inception cohort study. *BMJ.* 2009;339:b3829. Multicenter Study Research Support, Non-U.S. Gov't 2009/10/08.
6. Delitto A, George SZ, Van Dillen LR, et al. Low back pain. *J Orthop Sports Phys Ther.* 2012;42:A1–57.
7. Molano SM, Burdorf A, Elders LA. Factors associated with medical care-seeking due to low-back pain in scaffolders. *Am J Ind Med.* 2001;40:275–281.
8. Mortimer M, Ahlberg G, Group MU-NS. To seek or not to seek? Care-seeking behaviour among people with low-back pain. *Scand J Public Health.* 2003;31:194–203.
9. Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive treatments for acute, subacute, and chronic low back pain: a clinical practice guideline from the American College of Physicians. *Ann Intern Med.* 2017;166:514–530.

10. Foster NE, Anema JR, Cherkin D, et al. Prevention and treatment of low back pain: evidence, challenges, and promising directions. *Lancet*. 2018;391(10137):2368–2383.
11. Sahrmann S, Azevedo DC, Dillen LV. Diagnosis and treatment of movement system impairment syndromes. *Braz J Phys Ther*. 2017;21:391–399.
12. National Institute for Health and Care Excellence. *Low back pain and sciatica in over 16s: assessment and management*. NICE guideline [NG59]; 2017 (2016, accessed November 15, 2017) <https://www.nice.org.uk/guidance/ng59>.
13. Williams S, Whatman C, Hume PA, et al. Kinesio taping in treatment and prevention of sports injuries: a meta-analysis of the evidence for its effectiveness. *Sports Med*. 2012;42:153–164.
14. Castro-Sanchez AM, Lara-Palomo IC, Mataran-Penarrocha GA, et al. Kinesio Taping reduces disability and pain slightly in chronic non-specific low back pain: a randomised trial. *J Physiother*. 2012;58:89–95.
15. Paoloni M, Bernetti A, Fracocchi G, et al. Kinesio Taping applied to lumbar muscles influences clinical and electromyographic characteristics in chronic low back pain patients. *Eur J Phys Rehabil Med*. 2011;47:237–244.
16. Chang WD, Chen FC, Lee CL, et al. Effects of Kinesio Taping versus McConnell taping for patellofemoral pain syndrome: a systematic review and meta-analysis. *Evid Based Complement Alternat Med*. 2015;2015:471208.
17. Kase KWJ, Kase T. *Clinical therapeutic applications of the Kinesio Taping method Tokyo*. Japan: Ken Ikai Co. Ltd; 2003.
18. Kase KTH, Tomoki O. Development of Kinesiotape. KinesioTaping perfect manual. *Kinesio Taping Association*. 1996;6:117–118.
19. Saragiotto BT, Maher CG, Yamato TP, et al. Motor control exercise for chronic non-specific low-back pain. *Cochrane Database Syst Rev*. 2016;CD012004.
20. Costa LO, Maher CG, Latimer J, et al. Motor control exercise for chronic low back pain: a randomized placebo-controlled trial. *Phys Ther*. 2009;89:1275–1286.
21. van Dieën JH, Selen LP, Cholewicki J. Trunk muscle activation in low-back pain patients, an analysis of the literature. *J Electromyogr Kinesiol*. 2003;13:333–351.
22. Bae SH, Lee JH, Oh KA, et al. The effects of kinesio taping on potential in chronic low back pain patients anticipatory postural control and cerebral cortex. *J Phys Ther Sci*. 2013;25:1367–1371.
23. Fong SS, Tam YT, Macfarlane DJ, et al. Core muscle activity during TRX suspension exercises with and without kinesiology taping in adults with chronic low back pain: implications for rehabilitation. *Evid Based Complement Alternat Med*. 2015;2015:910168.
24. Turk DC, Melzack R. *Handbook of pain assessment*. Guilford Press; 2011.
25. Costa LO, Maher CG, Latimer J, et al. Psychometric characteristics of the Brazilian-Portuguese versions of the functional rating index and the roland morris disability questionnaire. *Spine (Phila Pa 1976)*. 2007;32:1902–1907.
26. You YL, Su TK, Liaw LJ, et al. The effect of six weeks of sling exercise training on trunk muscular strength and endurance for clients with low back pain. *J Phys Ther Sci*. 2015;27:2591.
27. Hermans HJFB. *European recommendations for surface electromyography: Results of the SENIAM project*; 2015.
28. Merletti R. Standards for reporting EMG data. *J Electromyogr Kinesiol*. 1999;9:3–4.
29. Ltd B. *EMG sensor operating manual*; 2007.
30. LB. *Goniometer and tensiometer operating manual*; 2007.
31. Esola MA, McClure PW, Fitzgerald GK, et al. Analysis of lumbar spine and hip motion during forward bending in subjects with and without a history of low back pain. *Spine (Phila Pa 1976)*. 1996;21:71–78.
32. Mendes A, de Freitas S, Amorin CF, et al. Electromyographic activity of the erector spinae: the short-effect of one workday for welders with nonspecific chronic low back pain, an observational study. *J Back Musculoskelet Rehabil*. 2018;31:147–154.
33. Halaki M, Ginn K. *Normalization of EMG signals: To normalize or not to normalize and what to normalize to?*; 2012.
34. Zakaria D, Kramer JF, Harburn KL. Reliability of non-normalized and normalized integrated EMG during maximal isometric contractions in females. *J Electromyogr Kinesiol*. 1996;6:129.
35. Ball N, Scurr J. An assessment of the reliability and standardisation of tests used to elicit reference muscular actions for electromyographical normalisation. *J Electromyogr Kinesiol*. 2010;20:81–88, <http://dx.doi.org/10.1016/j.jelekin.2008.09.004>.
36. McGill S, Jucker D, Kropf P. Appropriately placed surface EMG electrodes reflect deep muscle activity (psoas, quadratus lumborum, abdominal wall) in the lumbar spine. *J Biomech*. 1996;29:1503–1507.
37. Arena JG, Bruno GM, Brucks AG, et al. Reliability of an ambulatory electromyographic activity device for musculoskeletal pain disorders. *Int J Psychophysiol*. 1994;17:153–157.
38. ISEK International Society, et al (accessed 30/05/2016 2016) <http://www.isek.org/wp-content/uploads/2015/05/Standards-for-Reporting-EMG-Data.pdf>.
39. De Luca CJ, Donald Gilmore L, Kuznetsov M, et al. Filtering the surface EMG signal: movement artifact and baseline noise contamination. *J Biomech*. 2010;43:1573–1579.
40. Gabriel DA. Changes in kinematic and EMG variability while practicing a maximal performance task. *J Electromyogr Kinesiol*. 2002;12:407–412.
41. Roy AL, Keller TS, Colloca CJ. Posture-dependent trunk extensor EMG activity during maximum isometrics exertions in normal male and female subjects. *J Electromyogr Kinesiol*. 2003;13:469–476.
42. Burden AM, Trew M, Baltzopoulos V. Normalisation of gait EMGs: a re-examination. *J Electromyogr Kinesiol*. 2003;13:519–532.
43. Burden A. How should we normalize electromyograms obtained from healthy participants? What we have learned from over 25 years of research. *J Electromyogr Kinesiol*. 2010;20:1023–1035.
44. Cholewicki J, van Dieën J, Lee AS, et al. A comparison of a maximum exertion method and a model-based, sub-maximum exertion method for normalizing trunk EMG. *J Electromyogr Kinesiol*. 2011;21:767–773.
45. Narayanaswami P, Geisbush T, Jones L, et al. Critically re-evaluating a common technique: accuracy, reliability, and confirmation bias of EMG. *Neurology*. 2016;86:218–223.
46. Costa LO, Maher CG, Latimer J, et al. Clinimetric testing of three self-report outcome measures for low back pain patients in Brazil: which one is the best? *Spine (Phila Pa 1976)*. 2008;33:2459–2463.
47. Nusbaum L, Natour J, Ferraz MB, et al. Translation, adaptation and validation of the Roland-Morris questionnaire–Brazil Roland-Morris. *Braz J Med Biol Res*. 2001;34:203–210.
48. Bicalho E, Setti JA, Macagnan J, et al. Immediate effects of a high-velocity spine manipulation in paraspinal muscles activity of nonspecific chronic low-back pain subjects. *Man Ther*. 2010;15:469–475.
49. Elkins MR, Moseley AM. Intention-to-treat analysis. *J Physiother*. 2015;61:165–167, <http://dx.doi.org/10.1016/j.jphys.2015.05.013>.
50. Twisk J. *Applied longitudinal data analysis for epidemiology: a practical guide*; 2003.

51. de Morton NA. The PEDro scale is a valid measure of the methodological quality of clinical trials: a demographic study. *Aust J Physiother.* 2009;55:129–133.
52. SENIAM. *European recommendations for surface electromyography*; 2015.
53. da CMCL, Maher CG, Hancock MJ, et al. The prognosis of acute and persistent low-back pain: a meta-analysis. *CMAJ.* 2012;184:E613–624.
54. Lim EC, Tay MG. Kinesio taping in musculoskeletal pain and disability that lasts for more than 4 weeks: is it time to peel off the tape and throw it out with the sweat? A systematic review with meta-analysis focused on pain and also methods of tape application. *Br J Sports Med.* 2015;49:1558–1566.
55. Nelson NL. Kinesio taping for chronic low back pain: a systematic review. *J Bodyw Mov Ther.* 2016;20:672–681.
56. Parreira Pdo C, Costa Lda C, Hespanhol Junior LC, et al. Current evidence does not support the use of Kinesio Taping in clinical practice: a systematic review. *J Physiother.* 2014;60:31–39.
57. Hung CC, Shen TW, Liang CC, et al. Using surface electromyography (SEMG) to classify low back pain based on lifting capacity evaluation with principal component analysis neural network method. *Conf Proc IEEE Eng Med Biol Soc.* 2014;2014:18–21.
58. Timmesfeld N, Schafer H, Muller HH. Increasing the sample size during clinical trials with t-distributed test statistics without inflating the type I error rate. *Stat Med.* 2007;26:2449–2464.
59. Wang B, Ting N. Sample size determination with familywise control of both type I and type II errors in clinical trials. *J Biopharm Stat.* 2016;26:951–965.
60. Csapo R, Alegre LM. Effects of Kinesio((R)) taping on skeletal muscle strength—a meta-analysis of current evidence. *J Sci Med Sport.* 2015;18:450–456.
61. Kalron A, Bar-Sela S. A systematic review of the effectiveness of Kinesio Taping—fact or fashion? *Eur J Phys Rehabil Med.* 2013;49:699–709.
62. Morris D, Jones D, Ryan H, et al. The clinical effects of Kinesio(R) Tex taping: a systematic review. *Physiother Theory Pract.* 2013;29:259–270.
63. Mostafavifar M, Wertz J, Borchers J. A systematic review of the effectiveness of kinesio taping for musculoskeletal injury. *Phys Sportsmed.* 2012;40:33–40.
64. Grzeskowiak M, Szulc P, Szwedziak M, et al. The effect of the Kinesio Taping method on spinal motion and physiological spinal curvatures. Literature review. *Ortop Traumatol Rehabil.* 2014;16:221–226.
65. Luz Junior MAD, Almeida MO, Santos RS, et al. Effectiveness of Kinesio Taping in patients with chronic nonspecific low back pain: a systematic review with meta-analysis. *Spine (Phila Pa 1976).* 2019;44:68–78.