Published in final edited form as:

Psychophysiology. 2020 October ; 57(10): e13632. doi:10.1111/psyp.13632.

Progesterone modulates theta oscillations in the frontal-parietal network

Justin Riddle^{1,2,3}, Sangtae Ahn^{1,2}, Trevor McPherson^{1,2}, Susan Girdler^{1,3,*}, Flavio Frohlich^{1,2,4,5,6,7,*}

¹Department of Psychiatry, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA 27599

²Carolina Center for Neurostimulation, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA 27599

³Center for Women's Mood Disorders, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA 27599

⁴Department of Neurology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA 27599

⁵Department of Cell Biology and Physiology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA 27599

⁶Department of Biomedical Engineering, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA 27599

⁷Neuroscience Center, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA 27599

Abstract

The neuroactive metabolites of the steroid hormones progesterone (P4) and testosterone (T) are GABAergic modulators that influence cognition, yet the specific effect of P4 and T on brain network activity remains poorly understood. Here, we investigated if a fundamental oscillatory network activity pattern, often related to cognitive control, frontal midline theta (FMT) oscillations, are modulated by steroid hormones, P4 and T. We measured the concentration of P4 and T using salivary enzyme immunoassay and FMT oscillations using high-density electroencephalography (EEG) during eyes-open resting state in fifty-five healthy women and men. Electrical brain activity was analyzed using Fourier analysis, aperiodic signal fitting, and beamformer source localization. Steroid hormone concentrations and biological sex were used as predictors for scalp and source-estimated amplitude of theta oscillations. Elevated concentrations of P4 predicted increased amplitude of FMT oscillations across both sexes, and no relationship was found with T. The positive correlation with P4 was specific to the frontal-midline electrodes and survived correction for the background aperiodic signal of the brain. Using source localization, FMT oscillations were localized to the frontal-parietal network. Additionally, theta amplitude within the frontal-parietal network, but not the default mode network, positively

Corresponding Author: Flavio Frohlich, flavio_frohlich@med.unc.edu, (919) 966-4584.

*Senior authors contributed equally

correlated with P4 concentration. Our results suggest that P4 concentration modulates brain activity via upregulation of theta oscillations in the frontal-parietal network.

1. Introduction

The metabolites of the sex steroids, progesterone (P4) and testosterone (T), are allosteric modulators of the GABA_A receptor and impact cognition (Zurkovsky et al. 2007; Schumacher et al. 2014; Celec, Ostatníková, and Hodosy 2015), as well as other biological and affective systems. The progestogens (e.g. P4) and androgens (e.g. T) are classes of sex steroids that differentially engage neural activity patterns in functional neuroimaging (Peper et al. 2011) and differentially influence cognition (Pletzer, Petasis, and Cahill 2014). In particular, a recent study found that elevated concentrations of progesterone (P4) correlate with better performance in the N-back, working memory task (Hidalgo-Lopez and Pletzer 2017) suggesting a role for P4 in boosting cognitive control. Consistent with a role of P4 in cognitive control, increased resting-state connectivity within the frontal-parietal network in functional magnetic resonance imaging positively correlated with P4 concentrations (Syan et al. 2017). Despite emerging evidence for a role of P4 in cognitive control, there has yet to be an investigation into the electrical brain activity patterns that might underlie this influence. Frontal midline theta (FMT) oscillations (4–7Hz) are an ideal candidate for how P4 may influence brain activity (see (Cavanagh and Frank 2014) for review).

T concentrations are associated with alterations in social cognition: social decision-making (Eisenegger and Naef 2011), aggression (Montoya et al. 2012), and social status (Eisenegger, Haushofer, and Fehr 2011). Consistent with the role of T in social cognition, greater concentrations of T are correlated with decreased functional connectivity between the orbito-frontal cortex (critical region for social cognition) and the amygdala (critical region for processing emotionally salient perception) (Ackermann et al. 2012). Frontal theta oscillations also have been shown to play a role in social cognition (Tendler and Wagner 2015; Narayanan et al. 2011). Therefore, T concentration may also be predictive of the amplitude of FMT oscillations.

We investigated the relationship between the steroid hormones, P4 and T, and the amplitude of FMT oscillations using salivary enzyme immunoassay and recorded high-density electroencephalography (EEG) during the resting-state in men and women. Using source localization, we investigated the neural origin of FMT oscillations and correlated theta activity in the frontal-parietal network to steroid hormone concentrations.

2. Method

Data from 55 participants was pooled across three experiments (22 from National Clinical Trial 0324450 (Sheffield et al. 2019), 14 from [NCT03243084](#) (Ahn et al. 2019), and 19 from [NCT03178344](#)). 23 participants were female and 32 participants were male. Participants were ages 18 to 70 years with an average age of 28.8 years and standard deviation of 12.9 years (Table 1). Seven participants were 50 years of age or older. In each of these studies, we sampled steroid hormone concentrations via saliva sample and collected resting-state electrical brain activity via high-density EEG. All experiments were approved by the

institutional review board at the University of North Carolina at Chapel Hill and data were collected at the Carolina Center for Neurostimulation. Participants were screened prior to enrollment with the following inclusion criteria: no personal or immediate family history of neurological or psychiatric illness, no ongoing psychotherapy treatment or use of medication to treat a neurological or psychiatric illness, no major head injury or brain surgery, no brain implants (including cochlear implants), no history of cardiovascular disease. All participants were required to pass a urinary drug test and women were screened with a pregnancy test. Participants were asked to maintain a regular sleep schedule prior to the experimental sessions, as well as abstain from alcohol and caffeine for the 24 hours prior to the visit.

2.1 Statistical Analysis

Our primary analysis was to test for a correlation (Pearson) between the amplitude of FMT oscillations with P4 or T concentration. To correct for multiple comparisons, we used the Bonferroni method and consider a correlation significant at alpha of .025. Upon finding a significant correlation, we tested for the specificity of this relationship by comparing correlations between steroid hormones. For this analysis, we used Pearson and Filon's comparison of correlations with an overlapping variable (Pearson and Filon 1898) implemented by the "cocor" toolbox (Diedenhofen and Musch 2015) in R (The R Foundation).

Steroid hormone concentration of P4 and T is known to decrease with age (Ukkola et al. 2001). Therefore, to control for age, we investigated if the difference in correlations between steroid hormones was still significant after removing the variance that is explained by age (Pearson partial correlation). The physiological impact of steroid hormones is also known to be mediated by sex. Therefore, we also investigated if the relationship was sex-specific by comparing correlations between women and men for each steroid hormone to FMT amplitude (Pearson and Filon). If no difference was found, then follow-up analyses were run on the full group. However, given the relevance of sex to the function of steroid hormones, we report sex-specific effects as an exploratory analysis.

We ran three control analyses that tested for the frequency specificity of the relationship (Method 2.3), scalp specificity of the relationship (Method 2.3), and addressed a confound to interpreting neural oscillations (Method 2.4). Frequency specificity was evaluated using a difference in correlation with neighboring oscillations (Pearson and Filon). Spatial specificity was quantitatively investigated in a control site and qualitatively assessed via an exhaustive topographic correlation analysis (Pearson). A confound that theta oscillations are an artifact of the aperiodic signal (described in Method 2.4) was assessed by correlation (Pearson) after aperiodic signal removal. Finally, we ran a follow-up analysis that investigated the network origin of the theta oscillation using source localization (Method 2.5). The relationship of source localized theta amplitude to steroid hormone concentration was investigated by comparing correlations between steroid hormones (Pearson and Filon) and using post-hoc correlation analysis (Pearson).

2.2 Steroid Hormone Enzyme Immunoassay

Steroid hormone concentrations were assessed via four saliva samples taken throughout a single day: before breakfast, before lunch, before dinner and before bed. Participants were instructed not to eat for an hour before giving the sample. By sampling across four timepoints in a single day, we controlled for the known diurnal fluctuation in salivary hormone concentration (Konishi et al. 2012). In each experiment, participants were provided with the saliva sample kit in their first session and returned the kit in their second session that was scheduled for a week later. EEG data were used from the first session of each experiment. Thus, the saliva samples reflect hormone concentrations in the week following the EEG. Samples were sent to Labrix for assay (Clackamas, OR, USA). Labrix pooled the four samples into a single sample estimate and conducted enzyme immunoassays for P4 and T. The P4 (4-pregnen-3,20-dione) enzyme immunoassay kit was provided by Salimetrics Inc (State College, PA, USA) with a limit of detection of 5 pg/mL, a strong correlation between salivary P4 and serum P4 ($r(35) = .80$), and no detectable ($<0.004\%$) cross-reactivity with T (Salimetrics Salivary Progesterone Enzyme Immunoassay Kit Item No. 1–2502). The kit measured free P4 in the saliva that is not bound to serum proteins and is considered biologically active. The T enzyme immunoassay kit was the Pantex Salivary Direct Testosterone EIA Kit by PANTEX Division of Bio-Analysis (Santa Monica, CA, USA) with a limit of detection of 2.1 pg/mL and low cross-reactivity with P4 (0.28%) (Pantex Salivary Direct Testosterone EIA Kit Catalog No. 635). One to two percent of T is free in plasma, and this concentration is highly correlated with free T in saliva. Like P4, the kit measures only biologically active T. The immunoassays have an exponential distribution. Therefore, raw values were log-transformed prior to all statistical analysis. Range and descriptive metrics are provided in Table 1.

2.3 High Density Electroencephalography

EEG data were collected with a high-density 128-channel electrode net at 1000 Hz (HydroCel Geodesic Sensor Net) and EGI system (NetAmps 410, Electrical Geodesics Inc., OR, USA). The impedance of each electrode was ensured to be below 50 k Ω at the start of each session. Four-minutes (five-minutes in 19 participants from [NCT03178344](#)) of eyes-open resting-state EEG was collected.

The resting-state EEG data were preprocessed using custom scripts in MatLab and the EEGLAB toolbox (Delorme and Makeig 2004). All EEG data were downsampled to 250Hz with anti-aliasing filtering and band-pass filtered from 1 to 50 Hz. The data were then preprocessed using an artifact subspace reconstruction algorithm to remove high-variance signal and reconstruct missing data (Mullen et al. 2013). Channels that were found in the previous step to contain above threshold noise were interpolated. All data was common average referenced. An infomax independent component analysis (ICA) was performed to separate plausible neural activity from eye blinks, eye movement, muscle activity, heartbeats, and channel noise (Jung et al. 2000). All ICA components were visually inspected and components corresponding to noise were manually rejected. Resting-state EEG data were epoched into two second windows.

Our analysis was restricted to the 90 channels on the scalp, referred to as data channels. The fast Fourier transform was applied to each two-second epoch and the average spectral amplitude was calculated by median to reduce spurious amplitude fluctuations from outlier epochs. Frontal midline theta (FMT) amplitude was calculated as the average amplitude from 4–7 Hz, the canonical theta band, for the FCz channel. The range and descriptive metrics for the amplitude of FMT oscillations is provided in Table 1. To reduce multiple comparisons, we restricted our primary analysis to this single channel. We were interested in the frequency specificity of our effects. Thus, as a control analysis, we calculated the amplitude of alpha oscillations (8–12 Hz) and beta (15–30 Hz) oscillations and compared the correlation of these bands with steroid hormone concentrations with that of FMT amplitude. As an additional control analysis, we also estimate spectral amplitude in an occipital-midline electrode, Oz, that is known to be dominated by alpha, but not theta, oscillations, and correlated these estimates with steroid hormone concentration.

Furthermore, we hypothesized that theta amplitude, and thus its correlation with steroid hormone concentration would be spatially localized to the frontal-midline. To estimate the spatial distribution of theta amplitude across the scalp, we log-transformed theta amplitude for each of the 90 data channels and applied a spatial normalization using the z -transformation for each participant. We hypothesized that a genuine correlation with FMT oscillations would be confined to a cluster of electrodes surrounding FCz.

2.4 Correction for Aperiodic Signal

Recent evidence suggests that the aperiodic signal in the background of the brain may play a meaningful role in modulating cognition and predicting healthy aging (Voytek et al. 2015). As the aperiodic signal of the brain, a $1/f$ power distribution, steepens, there will be an increase in theta and delta power (2–7 Hz). The change in slope could lead to the erroneous conclusion that theta oscillations are increasing, when the more parsimonious explanation is a steepening of the aperiodic signal slope (Gao, Peterson, and Voytek 2017). A true oscillation under this framework is defined as a band-limited increase in spectral power over and above the aperiodic background spectra (He 2014). Therefore, we calculated the aperiodic signal for each participant and subtracted it in order to confirm that the measured theta oscillations represent true oscillations. We calculated the aperiodic signal as a linear fit to the $\log(\text{amplitude})$ and $\log(\text{frequency})$ of each participant. As the alpha oscillation is prominent in EEG during the resting state, we calculated the aperiodic signal as the linear fit to amplitude values from 2–4 Hz and 40–50 Hz (see (Voytek et al. 2015) for a similar method). After fitting the data for each participant, we subtracted the fit amplitude across all frequencies in order to calculate theta amplitude after aperiodic signal correction.

2.5 Network Analysis

Our large sample size and use of a 128-channel EEG system provided a methodological foundation for source localization analysis to investigate the spatial origination of FMT oscillations. Source localization was run using the Dynamical Imaging of Coherent Sources (DICS) beamformer algorithm implemented in the FieldTrip toolbox (Oostenveld et al. 2011). For each two second epoch of resting-state data, we calculated the spectral amplitude for each channel via 5-cycle Morlet wavelet convolution at 6 Hz. Next, we calculated the

cross-spectral density matrix to estimate the phase difference and shared amplitude of each channel pair for source localization. The beamformer algorithm was run using a lead field calculated in the Montreal Neurological Institute (MNI) space using standard skull, skin, and brain tissue estimates provided in the FieldTrip toolbox. Theta amplitude was source localized into MNI space for each two second epoch and then averaged across epochs. The theta amplitude estimates from source localization were not normally distributed. Therefore, the data were log-transformed. We applied spatial normalization using the z-transformation for all voxels within a grey-matter mask in MNI space derived from tissue segmentation estimates provided by the SPM12 toolbox (Penny et al. 2011). For display purposes, an independent t-test between participants with high and low FMT amplitude in sensor-space (median split) was run on theta-amplitude for every voxel in source-space.

Investigation into the temporal and spatial engagement of the human brain during cognitive control has found two primary signals: one spatial and one temporal. Electrophysiology researchers find increased amplitude of FMT oscillations and hypothesized that these signals originate from the anterior cingulate cortex or medial prefrontal cortex directly beneath the scalp sensors with peak FMT amplitude (see (Cavanagh and Frank 2014) for review), whereas functional neuroimaging researchers find increased functional connectivity and activation within the FPN as a function of increased cognitive control as driven by task demands (see (Badre and Nee 2018) for review). When dipoles in homologous brain regions are aligned in opposite directions (e.g. medial to lateral), the resulting electric field is found in the space between the two dipoles where the field inverts. This principle of electric field modeling is found in task-evoked electrical activity. For example, in auditory tasks, peak auditory-evoked electrophysiological signal is found in central midline electrode, Cz; but via source localization auditory activity is found to be driven by bilateral auditory cortex (Stropahl et al. 2018). Therefore, we hypothesized that source localization of FMT oscillations would reveal greatest theta amplitude in bilateral frontal cortex (Sasaki et al. 1994; Sasaki et al. 1996) and not in superior frontal gyrus or medial prefrontal cortex.

Bilateral middle frontal gyri have network membership in the task-positive frontal-parietal network (FPN), whereas anterior superior frontal gyrus and medial prefrontal cortex have network membership in the default mode network (DMN) (Yeo et al. 2011). To quantify the degree to which theta oscillations were source localized to the FPN as opposed to the DMN, we averaged theta amplitude in grey matter voxels of the FPN and DMN. FPN and DMN were defined as the 6th and 7th network from the seven-network Yeo atlas derived from resting-state functional connectivity analysis (Yeo et al. 2011). This network analysis provides a critical dimensionality reduction by which 10,000s of voxels are reduced to two networks; and, therefore, addresses the multiple comparisons problem. Analyses were run using custom code written in MatLab using functions from SPM12 and Fieldtrip.

2.6 Code accessibility

Scripts used for these analyses can be found in our code repository for this study on the Open Science Framework (<https://osf.io/6j253/>).

3. Results

The goal of the study was to test for a possible relationship between P4 or T concentration and the amplitude of FMT oscillations. We applied the Bonferroni method for multiple comparisons and considered an alpha of .025 to be significant. P4 concentration was positively correlated with FMT amplitude for all participants ($r(54) = .370, p = .005$) (Figure 1A), and T concentration was not correlated with FMT amplitude ($r(54) = -.003, p = .981$; Figure 1B). The correlation between FMT amplitude and P4 concentration was significantly greater than the correlation with T concentration (Pearson and Filon, $z(54) = 2.462, p = .014$). Thus, the relationship of FMT amplitude to P4 concentration was hormone specific. Furthermore, the correlation between FMT amplitude and P4 concentration explained unique variance when controlling for T concentration (partial-correlation, $r(54) = .386, p = .004$) despite a positive relationship between P4 and T concentration ($r(54) = .279, p = .039$). P4 and T concentration are known to decrease with age (Ukkola et al. 2001), and indeed this was the case within our dataset for both P4 ($r(54) = -.352$) and T ($r(54) = -.355$). When controlling for age, the correlation between P4 concentration and FMT amplitude was still greater than the correlation with T concentration (partial-correlation Pearson and Filon, $z(54) = 3.301, p = .001$).

As a follow-up analysis, we investigated a potential sex difference between these findings. Critically, the difference in correlation of P4 concentration to FMT amplitude between women and men was not significant (Fisher, $z(53) = 1.240, p = .215$), nor was the difference significant for T concentration (Fisher, $z(53) = 0.824, p = .410$). As there was no difference in correlation between women and men, all follow-up and control analyses were run ignoring sex as a factor. However, given the relevance of sex as a biological factor in the action of steroid hormones, we include the separate analyses for women and men as uncorrected exploratory analyses in Table 4, and qualitative differences between the sexes are reviewed in the discussion section.

To assess the frequency specificity of the relationship between P4 and peak theta amplitude, we analyzed the difference in correlation from peak alpha (8–12 Hz) and peak beta (15–30 Hz) amplitude (Figure 2A and Table 2). The increased correlation from FMT amplitude to P4 concentration was greater than alpha amplitude at a trend-level (Pearson and Filon, $z(54) = 1.812, p = .070$), and the difference in correlation from beta amplitude was significant (Pearson and Filon, $z(54) = 2.059, p = .040$). These findings provide evidence that the correlation of FMT amplitude with P4 concentration is not widespread across all frequencies, but is band-limited to canonical theta frequency.

The aperiodic signal, or background noise, of the human brain is a biologically relevant signal independent of neuronal oscillations (Voytek et al. 2015). With a sharper slope in the background noise, low frequency power is increased, despite no band-specific increase in power. In order to confirm that our results are specific to the amplitude of theta frequency neuronal oscillations and not a result of a difference in aperiodic signal, we fit and removed the aperiodic signal (Figure 2B). We found that our initial correlation between FMT amplitude and P4 concentration across all participants was still significant after correction for aperiodic signal ($r(54) = .284, p = .035$). The correlation of FMT amplitude after

aperiodic signal correction with P4 concentration was greater than the correlation with T concentration (Pearson and Filon, $r(54) = 2.118$, $p = .034$). Therefore, the relationship between P4 concentration and the amplitude of FMT oscillations in our data is likely a genuine neuronal oscillation.

As a control analysis, we analyzed peak theta amplitude in a scalp location that displays elevated alpha oscillations, electrode Oz at the occipital-midline (Table 2). Theta amplitude in Oz was neither significantly correlated with P4 concentration in the raw spectra ($r(54) = .198$, $p = .148$) (Figure 2C) nor after aperiodic signal correction ($r(54) = .057$, $p = .667$) (Figure 2D). There was no difference in correlation with P4 concentration for alpha or beta amplitude in Oz in the raw spectra or after aperiodic signal correction. This control analysis demonstrates that the frequency specific relationship of theta amplitude to P4 concentration was limited to the frontal midline.

To confirm that the correlation between FMT amplitude and P4 concentration is spatially specific to the frontal-midline electrodes, we correlated P4 concentration with the spatial-normalized theta amplitude across the scalp (Figure 3A). Consistent with the canonical distribution of FMT oscillations, the positive correlation with P4 was specific to FCz and its surrounding electrodes. To better understand the neural origin of theta oscillations, we ran source localization on theta oscillations and performed a median split of our participants based on FMT amplitude in sensor-space to analyze the spatial distribution of theta amplitude in source-space (Figure 3B). As hypothesized, participants with the greatest FMT amplitude showed increased theta amplitude in bilateral prefrontal cortex and not in medial prefrontal cortex or superior frontal gyrus (Figure 4A and 4B). We hypothesized that FMT oscillations originated in the frontal-parietal network (FPN) and not the default mode network (DMN). The FPN and DMN are visualized in Figure 4C and 4D to illustrate the spatial consistency between the spatial distribution of theta amplitude in source-space and the independently-derived FPN network from the seven-network Yeo atlas (Yeo et al. 2011).

As hypothesized, a one-tailed t-test between participants with high and low FMT amplitude found that theta amplitude was greater in the FPN relative to the DMN (one-tailed, $t(53) = 1.798$, $p = .039$, $d = .485$). Post-hoc t-tests between the high and low FMT groups revealed that this effect was driven by an increase in theta amplitude within the FPN (one-tailed, $t(53) = 1.697$, $p = .048$, $d = .460$), but not the DMN (one-tailed, $t(53) = -0.501$, $p = .691$, $d = .136$). Finally, we ran a correlation between theta amplitude derived from source localization and steroid hormone concentrations. As expected, we found a significant positive correlation between P4 concentration and theta amplitude in the FPN relative to DMN (Table 3). Post-hoc correlation analysis revealed that this effect was driven by a significant correlation between theta amplitude in the FPN and P4 concentration, but there was no relationship between theta amplitude in the DMN and P4 concentration (Table 3). Furthermore, P4 concentration showed a significantly greater correlation to source localized FMT amplitude in the FPN relative to DMN than T concentration (Pearson and Filon, $r(54) = 2.546$, $p = .011$). Post-hoc analysis revealed that this effect was driven by a significant increase in correlation for P4 concentration to theta amplitude in the FPN relative to T concentration (Pearson and Filon, $r(54) = 2.373$, $p = .018$), and there was no difference in correlation between steroid hormones to theta amplitude in the DMN (Pearson and Filon,

$z(54) = -0.745, p = .456$). These findings provide evidence that P4 is positively related to theta amplitude within the frontal-parietal network, and that this relationship is hormone and network specific.

4. Discussion

We investigated if the amplitude of frontal midline theta (FMT) oscillations were related to the concentration of steroid hormones progesterone (P4) or testosterone (T). We found that P4, but not T, concentrations correlated positively with FMT oscillatory amplitude. The relationship with P4 concentration was frequency specific to the theta band and spatially specific to the frontal midline scalp electrodes. We found supporting evidence that theta frequency oscillatory amplitude in frontal midline electrodes represented a genuine neuronal oscillation by removing the aperiodic signal, or background spectra, of the brain for each participant. Source localization analysis found that FMT oscillations originate in the frontal-parietal network (FPN), not the default-mode network (DMN), and source localized theta amplitude in the FPN correlated with P4 concentrations. Together, these findings suggest a role for P4 to specifically increase the amplitude of theta oscillations in the FPN that is measured from the scalp at the frontal midline.

P4 has been found to play a neuroprotective role in both women and men and readily passes through the blood-brain barrier (see (Schumacher et al. 2014) for review). P4, particularly its neuroactive metabolite allopregnanolone (ALLO), is critical to downregulation of hypothalamic-pituitary-adrenal (HPA) activation following stress by facilitating GABAergic inhibitory signaling; and this compensatory response to stress is impaired in depression (Schumacher et al. 2014) (see (Girdler and Klatzkin 2007) for review). ALLO is a potent, positive allosteric modulator of the gamma-aminobutyric acid (GABA_A) receptor, and it is through this mechanism that P4 exerts antidepressant, antianxiety, and HPA-axis modulatory effects. This research culminated in FDA approval for ALLO as a treatment of post-partum depression (Osborne et al. 2017; Kanes et al. 2017). Although we did not directly measure ALLO in this study, concentrations of P4 and ALLO are highly correlated in blood and in the CNS (e.g., correlation strength in serum ranging from .80 to .95) (Freeman et al. 1993; Andréen et al. 2005; Palliser et al. 2015). In some cases, such as postpartum, P4 shows a reduced correlation with ALLO and a lower concentration of ALLO predicted increased depressive symptoms (Nappi et al. 2001). This finding suggests an additional mechanism by which P4, a steroid hormone, is converted into its neuroactive metabolite ALLO, despite their strong correlation in most studies with healthy participants.

The neural target of ALLO, GABA_A receptors, are found throughout the brain, but are found in the greatest concentration in the striatum (Brittain and Brown 2014). The striatum plays a critical role in cognitive control from operant-conditioning (Pagnoni et al. 2002) to decision-making (Balleine, Delgado, and Hikosaka 2007) due to its primary function in inhibiting motor output (Grillner et al. 2005). The FPN and the striatum have reciprocal anatomical projections that contribute to a dynamic hierarchically-organized system for cognitive control (Jarbo and Verstynen 2015; Badre and Nee 2018). Although the current study did not directly measure cognitive control processes, facilitation of GABAergic signaling by P4 may facilitate the frontal-parietal network in exerting inhibitory control.

Furthermore, FMT oscillations are an emerging biomarker relevant to the treatment of depression. Previous observational studies have found decreased amplitude of theta oscillations (Saletu, Anderer, and Saletu-Zyhlarz 2010; Pizzagalli, Oakes, and Davidson 2003) and decreased theta frequency functional connectivity (Linkenkaer-Hansen et al. 2005) during the resting-state in patients with major depressive disorder. A recent experiment found that depressed patients with increased amplitude of FMT oscillations during a working memory task were more likely to respond to non-invasive brain stimulation treatment (Bailey et al. 2018). Variability in recruitment of FMT oscillations during rest also serves as a predictor for pharmaceutical intervention for major depressive disorder (Mulert et al. 2007). Therefore, elucidating contributing factors to FMT oscillations may be of critical importance to future interventions.

As in any scientific study, the work presented here has limitations. We only measured peripheral concentrations of P4 and T. Therefore, we cannot draw inferences about concentrations in the brain. However, P4 readily crosses the blood-brain barrier and significantly contributes to central P4 and ALLO concentrations (Hu et al. 2009). Another important factor that mediates steroid hormone concentration is age (Ukkola et al. 2001). Our effects were robust when accounting for variance explained by the age-related decrease in P4 and T concentration. Sex differences were not captured in the current dataset, although our exploratory analysis found that P4 concentration shows a qualitatively greater correlation with FMT amplitude in women relative to men. Exogenous administration of P4 results in a sedative effect in both men and women (Söderpalm et al. 2004) suggesting that there may be a common mechanism for P4 action on neural activity across sexes. It should be noted that although women had a numerically higher concentration of P4, this difference was not significant. Therefore, the women in our sample must have been by majority pre-ovulatory (follicular phase), using oral contraception (low P4 concentration across the menstrual cycle), or in menopause, although these data were not collected. None of the women in our study were pregnant. Under these conditions, men display comparable P4 concentration to women (Oettel and Mukhopadhyay 2004). The particular reasons for a similar concentration between women and men does not invalidate the correlation of P4 concentration to FMT amplitude. Future research needs to be conducted to remove any non-linear effects introduced by reproductive events, such as the menopause transition or the postpartum period, where there may be a breakdown in relationship between P4 concentration and FMT amplitude.

In our cross-sectional study, we were unable to conclude that within a participant FMT amplitude tracks fluctuations in P4 concentration over time. Recent evidence suggests that change in hormone concentration is most critical for predicting its impact on mood (Schmidt et al. 2017; Gordon et al. 2016). Therefore, future research should employ a within-participant design that investigates the relative change in P4 concentration and FMT amplitude. Additionally, emerging experimental evidence suggests that some women are more sensitive to changes in hormone concentration than others and these women are more likely to develop reproductive mood disorders (Schiller, Meltzer-Brody, and Rubinow 2015). For example, the late luteal phase is characterized by a sharp decrease in P4 concentration and women with hormone sensitivity experience this withdrawal with mood disturbances (Schmidt et al. 1998; Farage, Osborn, and MacLean 2008). Future research is required to

investigate whether the relationship between P4 concentration and FMT amplitude breaks down in women with hormone sensitivity.

A recent review on the effect of progesterone in the menstrual cycle found that P4 concentration was a key predictor of emotional reactivity in women with premenstrual dysphoric disorder (PMDD) (Sundström-Poromaa 2018). P4 concentration during the late luteal phase positively correlated with increased activity in the amygdala and increased symptoms of anxiety (Gingnell et al. 2014; Gingnell et al. 2012). Consistent with our prediction, previous research using functional neuroimaging found that connectivity in the FPN was modulated in the late luteal phase as a function of P4 concentration (Syau et al. 2017) and treatment with ALLO during the luteal phase reduced symptoms of PMDD (Bixo et al. 2017). While speculative, the decrease in P4 during the late luteal phase may downregulate theta oscillations in the FPN resulting in hyperactivity of the amygdala in PMDD (Gingnell et al. 2012). Causal investigation using double-blinded administration of P4 or ALLO with EEG is required to establish a causal link between P4 concentration and FMT amplitude. Furthermore, the use of a task that drives FMT oscillations as a function of cognitive control demands would further elucidate the functional role of FMT oscillations with respect to P4.

Finally, non-invasive brain stimulation is a promising new therapeutic that has been used to target the lateral prefrontal cortex in depression (Perera et al. 2016), and can be tailored to target specific brain networks in a frequency specific manner (Alexander et al. 2019; Ahn et al. 2019). In cognitive neuroscience, transcranial alternating current stimulation (tACS) that synchronizes theta oscillations in frontal and parietal regions increases performance during cognitive control tasks (Reinhart and Nguyen 2019; Jaušovec and Jaušovec 2014; Polanía et al. 2012; Vosskuhl, Huster, and Herrmann 2015). Future research should investigate the application of theta frequency tACS targeted to the FPN as a potential treatment for reproductive mood disorders typified by P4 withdrawal.

Acknowledgments

Special thanks to David Rubinow who provided useful feedback and discussion in the interpretation of our findings. Thank you to Jhana Parikh, Julianna Prim, and Alec Sheffield for collecting the data, and to Alec Sheffield for preprocessing a subset of the EEG data. This study was funded by the Department of Psychiatry at the University of North Carolina at Chapel Hill and supported in part by the National Institute of Mental Health of the National Institutes of Health under Award Numbers R01MH101547 and R01MH111889 and the postdoctoral training program in reproductive mood disorders T32MH09331502. FF is the lead inventor of IP filed by UNC. FF is the founder, CSO, and majority owner of Pulvinar Neuro LLC. Pulvinar Neuro had nothing to do with this study.

5. References

- Ackermann Sandra, Spalek Klara, Rasch Björn, Gschwind Leo, Coynel David, Fastenrath Matthias, Papassotiropoulos Andreas, and Dominique J-F. 2012 ‘Testosterone levels in healthy men are related to amygdala reactivity and memory performance’, *Psychoneuroendocrinology*, 37: 1417–24. 10.1016/j.psyneuen.2012.01.008 [PubMed: 22341731]
- Ahn Sangtae, Prim Julianna H, Alexander Morgan L, McCulloch Karen L, and Fröhlich Flavio. 2019 ‘Identifying and engaging neuronal oscillations by transcranial alternating current stimulation in patients with chronic low back pain: a randomized, crossover, double-blind, sham-controlled pilot study’, *The Journal of Pain*, 20: 277 e1–77. e11 10.1016/j.jpain.2018.09.004 [PubMed: 30268803]

- Alexander Morgan L, Alagapan Sankaraleengam, Lugo Courtney E, Mellin Juliann M, Lustenberger Caroline, Rubinow David R, and Fröhlich Flavio. 2019 'Double-blind, randomized pilot clinical trial targeting alpha oscillations with transcranial alternating current stimulation (tACS) for the treatment of major depressive disorder (MDD)', *Translational psychiatry*, 9: 106 10.1038/s41398-019-0439-0 [PubMed: 30837453]
- Andréen Lotta, Sundström-Poromaa Inger, Bixo Marie, Andersson Agneta, Nyberg Sigrid, and Bäckström Torbjörn. 2005 'Relationship between allopregnanolone and negative mood in postmenopausal women taking sequential hormone replacement therapy with vaginal progesterone', *Psychoneuroendocrinology*, 30: 212–24. 10.1016/j.psyneuen.2004.07.003 [PubMed: 15471618]
- Badre David, and Nee Derek Evan. 2018 'Frontal cortex and the hierarchical control of behavior', *Trends in cognitive sciences*, 22: 170–88. 10.1016/j.tics.2017.11.005 [PubMed: 29229206]
- Bailey NW, Hoy KE, Rogasch NC, Thomson RH, McQueen S, Elliot D, Sullivan CM, Fulcher BD, Daskalakis ZJ, and Fitzgerald PB. 2018 'Responders to rTMS for depression show increased fronto-midline theta and theta connectivity compared to non-responders', *Brain stimulation*, 11: 190–203. 10.1016/j.brs.2017.10.015 [PubMed: 29128490]
- Balleine Bernard W, Delgado Mauricio R, and Hikosaka Okihide. 2007 'The role of the dorsal striatum in reward and decision-making', *Journal of Neuroscience*, 27: 8161–65. 10.1523/JNEUROSCI.1554-07.2007 [PubMed: 17670959]
- Bixo Marie, Ekberg Karin, Poromaa Inger Sundström, Hirschberg Angelica Lindén, Jonasson Aino Fianu, Andréen Lotta, Timby Erika, Wulff Marianne, Ehrenborg Agneta, and Bäckström Torbjörn. 2017 'Treatment of premenstrual dysphoric disorder with the GABAA receptor modulating steroid antagonist Sepranolone (UC1010)—A randomized controlled trial', *Psychoneuroendocrinology*, 80: 46–55. 10.1016/j.psyneuen.2017.02.031 [PubMed: 28319848]
- Brittain John-Stuart, and Brown Peter. 2014 'Oscillations and the basal ganglia: motor control and beyond', *Neuroimage*, 85: 637–47. 10.1016/j.neuroimage.2013.05.084 [PubMed: 23711535]
- Cavanagh James F, and Frank Michael J. 2014 'Frontal theta as a mechanism for cognitive control', *Trends in cognitive sciences*, 18: 414–21. 10.1016/j.tics.2014.04.012 [PubMed: 24835663]
- Celec Peter, Ostatníková Daniela, and Hodosy Július. 2015 'On the effects of testosterone on brain behavioral functions', *Frontiers in neuroscience*, 9: 12 10.3389/fnins.2015.00012 [PubMed: 25741229]
- Delorme A, and Makeig S. 2004 'EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis', *Journal of neuroscience methods*, 134: 9–21. 10.1016/j.jneumeth.2003.10.009 [PubMed: 15102499]
- Diedenhofen Birk, and Musch Jochen. 2015 'cocor: A comprehensive solution for the statistical comparison of correlations', *PloS one*, 10 10.1371/journal.pone.0121945
- Eisenegger Christoph, Haushofer Johannes, and Fehr Ernst. 2011 'The role of testosterone in social interaction', *Trends in cognitive sciences*, 15: 263–71. 10.1016/j.tics.2011.04.008 [PubMed: 21616702]
- Eisenegger Christoph, and Naef Michael. 2011 'Combining behavioral endocrinology and experimental economics: testosterone and social decision making', *JoVE (Journal of Visualized Experiments)*: e2065 10.3791/2065
- Farage Miranda A, Osborn Thomas W, and MacLean Allan B. 2008 'Cognitive, sensory, and emotional changes associated with the menstrual cycle: a review', *Archives of gynecology and obstetrics*, 278: 299 10.1007/s00404-008-0708-2 [PubMed: 18592262]
- Freeman Ellen W, Purdy Robert H, Coutifaris Christos, Rickels Karl, and Paul Steven M. 1993 'Anxiolytic metabolites of progesterone: correlation with mood and performance measures following oral progesterone administration to healthy female volunteers', *Neuroendocrinology*, 58: 478–84. 10.1159/000126579 [PubMed: 7904330]
- Gao Richard, Peterson Erik J, and Voytek Bradley. 2017 'Inferring synaptic excitation/inhibition balance from field potentials', *Neuroimage*, 158: 70–78. 10.1016/j.neuroimage.2017.06.078 [PubMed: 28676297]
- Gingnell Malin, Ahlstedt Victoria, Bannbers Elin, Johan Wikström Inger Sundström-Poromaa, and Fredrikson Mats. 2014 'Social stimulation and corticolimbic reactivity in premenstrual dysphoric

disorder: a preliminary study', *Biology of mood & anxiety disorders*, 4: 3 10.1186/2045-5380-4-3 [PubMed: 24572042]

- Gingnell Malin, Morell Arvid, Bannbers Elin, Wikström Johan, and Poromaa Inger Sundström. 2012 'Menstrual cycle effects on amygdala reactivity to emotional stimulation in premenstrual dysphoric disorder', *Hormones and behavior*, 62: 400–06. 10.1016/j.yhbeh.2012.07.005 [PubMed: 22814368]
- Girdler Susan S, and Klatzkin Rebecca. 2007 'Neurosteroids in the context of stress: implications for depressive disorders', *Pharmacology & therapeutics*, 116: 125–39. 10.1016/j.pharmthera.2007.05.006 [PubMed: 17597217]
- Gordon Jennifer L, Rubinow David R, Eisenlohr-Moul Tory A, Leserman Jane, and Girdler Susan S. 2016 'Estradiol variability, stressful life events and the emergence of depressive symptomatology during the Menopause Transition', *Menopause (New York, NY)*, 23: 257 10.1097/GME.0000000000000528
- Grillner Sten, Hellgren Jeanette, Menard Ariane, Saitoh Kazuya, and Wikström Martin A. 2005 'Mechanisms for selection of basic motor programs—roles for the striatum and pallidum', *Trends in neurosciences*, 28: 364–70. 10.1016/j.tins.2005.05.004 [PubMed: 15935487]
- He Biyu J. 2014 'Scale-free brain activity: past, present, and future', *Trends in cognitive sciences*, 18: 480–87. 10.1016/j.tics.2014.04.003 [PubMed: 24788139]
- Hidalgo-Lopez Esmeralda, and Pletzer Belinda. 2017 'Interactive effects of dopamine baseline levels and cycle phase on executive functions: the role of progesterone', *Frontiers in neuroscience*, 11: 403 10.3389/fnins.2017.00403 [PubMed: 28751855]
- Hu Zhiying, Li Yan, Fang Marong, Wai Maria SM, and Yew David T. 2009 'Exogenous progesterone: a potential therapeutic candidate in CNS injury and neurodegeneration', *Current medicinal chemistry*, 16: 1418–25. 10.2174/092986709787846523 [PubMed: 19355896]
- Jarbo Kevin, and Verstynen Timothy D. 2015 'Converging structural and functional connectivity of orbitofrontal, dorsolateral prefrontal, and posterior parietal cortex in the human striatum', *Journal of Neuroscience*, 35: 3865–78. 10.1523/JNEUROSCI.2636-14.2015 [PubMed: 25740516]
- Jaušovec Norbert, and Jaušovec Ksenija. 2014 'Increasing working memory capacity with theta transcranial alternating current stimulation (tACS)', *Biological psychology*, 96: 42–47. 10.1016/j.biopsycho.2013.11.006 [PubMed: 24291565]
- Jung TP, Makeig S, Humphries C, Lee TW, McKeown MJ, Iragui V, and Sejnowski TJ. 2000 'Removing electroencephalographic artifacts by blind source separation', *Psychophysiology*, 37: 163–78. 10.1111/1469-8986.3720163 [PubMed: 10731767]
- Kanes Stephen, Colquhoun Helen, Gunduz-Bruce Handan, Raines Shane, Arnold Ryan, Schacterle Amy, Doherty James, Epperson C Neill, Deligiannidis Kristina M, and Riesenber Robert. 2017 'Brexanolone (SAGE-547 injection) in post-partum depression: a randomised controlled trial', *The Lancet*, 390: 480–89. 10.1016/S0140-6736(17)31264-3
- Konishi Shoko, Brindle Eleanor, Guyton Amanda, and O'Connor Kathleen A. 2012 'Salivary concentration of progesterone and cortisol significantly differs across individuals after correcting for blood hormone values', *American journal of physical anthropology*, 149: 231–41. 10.1002/ajpa.22114 [PubMed: 22826025]
- Linkenkaer-Hansen Klaus, Monto Simo, Rytsälä Heikki, Suominen Kirsii, Isometsä Erkki, and Kähkönen Seppo. 2005 'Breakdown of long-range temporal correlations in theta oscillations in patients with major depressive disorder', *Journal of Neuroscience*, 25: 10131–37. 10.1523/JNEUROSCI.3244-05.2005 [PubMed: 16267220]
- Montoya Estrella R, Terburg David, Bos Peter A, and Van Honk Jack. 2012 'Testosterone, cortisol, and serotonin as key regulators of social aggression: A review and theoretical perspective', *Motivation and emotion*, 36: 65–73. 10.1007/s11031-011-9264-3 [PubMed: 22448079]
- Mulert Christoph, Juckel Georg, Brunnermeier Michael, Karch Susanne, Leicht Gregor, Mergl Roland, Möller Hans-Jürgen, Hegerl Ulrich, and Pogarell Oliver. 2007 'Rostral anterior cingulate cortex activity in the theta band predicts response to antidepressive medication', *Clinical EEG and neuroscience*, 38: 78–81. 10.1177/155005940703800209 [PubMed: 17515172]
- Mullen T, Kothe C, Chi YM, Ojeda A, Kerth T, Makeig S, Cauwenberghs G, and Jung TP. 2013 'Real-time modeling and 3D visualization of source dynamics and connectivity using wearable EEG',

- Conference proceedings IEEE Engineering in Medicine and Biology Society 2013: 2184–7. 10.1109/EMBC.2013.6609968
- Nappi Rossella E, Petraglia Felice, Luisi Stefano, Polatti Franco, Farina Claudio, and Genazzani Andrea R. 2001 ‘Serum allopregnanolone in women with postpartum “blues”’, *Obstetrics & Gynecology*, 97: 77–80. 10.1016/s0029-7844(00)01112-1 [PubMed: 11152912]
- Narayanan Venu, Rebecca S Heiming Friederike Jansen, Lesting Jörg, Sachser Norbert, Pape Hans-Christian, and Seidenbecher Thomas. 2011 ‘Social defeat: impact on fear extinction and amygdala-prefrontal cortical theta synchrony in 5-HTT deficient mice’, *PloS one*, 6: e22600 10.1371/journal.pone.0022600 [PubMed: 21818344]
- Oettel M, and Mukhopadhyay AK. 2004 ‘Progesterone: the forgotten hormone in men?’, *The Aging Male*, 7: 236–57. 10.1080/13685530400004199 [PubMed: 15669543]
- Oostenveld Robert, Fries Pascal, Maris Eric, and Schoffelen Jan-Mathijs. 2011 ‘FieldTrip: open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data’, *Computational intelligence and neuroscience*, 2011: 1 10.1155/2011/156869 [PubMed: 21837235]
- Osborne Lauren M, Gispén Fiona, Sanyal Abanti, Yenokyan Gayane, Meilman Samantha, and Payne Jennifer L. 2017 ‘Lower allopregnanolone during pregnancy predicts postpartum depression: an exploratory study’, *Psychoneuroendocrinology*, 79: 116–21. 10.1016/j.psyneuen.2017.02.012 [PubMed: 28278440]
- Pagnoni Giuseppe, Zink Caroline F, Montague P Read, and Berns Gregory S. 2002 ‘Activity in human ventral striatum locked to errors of reward prediction’, *Nature neuroscience*, 5: 97 10.1038/nn802 [PubMed: 11802175]
- Palliser Hannah Kate, Kelleher Meredith A, Tolcos Mary, Walker David William, and Hirst JJ. 2015 ‘Effect of postnatal progesterone therapy following preterm birth on neurosteroid concentrations and cerebellar myelination in guinea pigs’, *Journal of developmental origins of health and disease*, 6: 350–61. 10.1017/S2040174415001075 [PubMed: 25907069]
- Pearson Karl, and Filon Louis Napoleon George. 1898 ‘Mathematical contributions to the theory of evolution. IV. On the probable errors of frequency constants and on the influence of random selection on variation and correlation’, *Proceedings of the Royal Society of London*, 62: 173–76. 10.1098/rspl.1897.0091
- Penny William D, Friston Karl J, Ashburner John T, Kiebel Stefan J, and Nichols Thomas E. 2011 *Statistical parametric mapping: the analysis of functional brain images* (Elsevier).
- Peper Jiska S, van den Heuvel Martijn P, Mandl René CW, Hulshoff Pol Hilleke E, and van Honk Jack. 2011 ‘Sex steroids and connectivity in the human brain: a review of neuroimaging studies’, *Psychoneuroendocrinology*, 36: 1101–13. 10.1016/j.psyneuen.2011.05.004 [PubMed: 21641727]
- Perera Tarique, George Mark S, Geoffrey Grammer, Janicak Philip G, Pascual-Leone Alvaro, and Wirecki Theodore S. 2016 ‘The clinical TMS society consensus review and treatment recommendations for TMS therapy for major depressive disorder’, *Brain stimulation*, 9: 336–46. 10.1016/j.brs.2016.03.010 [PubMed: 27090022]
- Pizzagalli Diego A, Oakes Terrence R, and Davidson Richard J. 2003 ‘Coupling of theta activity and glucose metabolism in the human rostral anterior cingulate cortex: an EEG/PET study of normal and depressed subjects’, *Psychophysiology*, 40: 939–49. 10.1111/1469-8986.00112 [PubMed: 14986847]
- Pletzer Belinda, Petasis Ourania, and Cahill Larry. 2014 ‘Switching between forest and trees: Opposite relationship of progesterone and testosterone to global–local processing’, *Hormones and behavior*, 66: 257–66. 10.1016/j.yhbeh.2014.05.004 [PubMed: 24874173]
- Polanía Rafael, Nitsche Michael A, Korman Carolin, Batsikadze Giorgi, and Paulus Walter. 2012 ‘The importance of timing in segregated theta phase-coupling for cognitive performance’, *Current Biology*, 22: 1314–18. 10.1016/j.cub.2012.05.021 [PubMed: 22683259]
- Reinhart Robert MG, and Nguyen John A. 2019 ‘Working memory revived in older adults by synchronizing rhythmic brain circuits’, *Nature neuroscience*, 22: 820 10.1038/s41593-019-0371-x [PubMed: 30962628]
- Saletu B, Anderer P, and Saletu-Zyhlarz GM. 2010 ‘EEG topography and tomography (LORETA) in diagnosis and pharmacotherapy of depression’, *Clinical EEG and neuroscience*, 41: 203–10. 10.1177/155005941004100407 [PubMed: 21077572]

- Sasaki K, Tsujimoto T, Nambu A, Matsuzaki R, and Kyuhou S. 1994 'Dynamic activities of the frontal association cortex in calculating and thinking', *Neuroscience research*, 19: 229–33. 10.1016/0168-0102(94)90147-3 [PubMed: 8008251]
- Sasaki Kazuo, Tsujimoto Toru, Nishikawa Satoru, Nishitani Nobuyuki, and Ishihara Tsutomu. 1996 'Frontal mental theta wave recorded simultaneously with magnetoencephalography and electroencephalography', *Neuroscience research*, 26: 79–81. 10.1016/0168-0102(96)01082-6 [PubMed: 8895895]
- Schiller Crystal Edler, Meltzer-Brody Samantha, and Rubinow David R. 2015 'The role of reproductive hormones in postpartum depression', *CNS spectrums*, 20: 48–59. 10.1017/S1092852914000480 [PubMed: 25263255]
- Schmidt Peter J, Martinez Pedro E, Nieman Lynnette K, Koziol Deloris E, Thompson Karla D, Schenkel Linda, Wakim Paul G, and Rubinow David R. 2017 'Exposure to a change in ovarian steroid levels but not continuous stable levels triggers PMDD symptoms following ovarian suppression', *The American journal of psychiatry*, 174: 980 10.1176/appi.ajp.2017.16101113 [PubMed: 28427285]
- Schmidt Peter J, Nieman Lynnette K, Danaceau Merry A, Adams Linda F, and Rubinow David R. 1998 'Differential behavioral effects of gonadal steroids in women with and in those without premenstrual syndrome', *New England Journal of Medicine*, 338: 209–16. 10.1056/NEJM199801223380401
- Schumacher M, C Mattern, Ghomari A, Oudinet JP, Liere P, Labombarda F4, Sitruk-Ware Régine, De Nicola Alejandro Federico, and Guennoun R. 2014 'Revisiting the roles of progesterone and allopregnanolone in the nervous system: resurgence of the progesterone receptors', *Progress in neurobiology*, 113: 6–39. 10.1016/j.pneurobio.2013.09.004 [PubMed: 24172649]
- Sheffield Alec, Ahn Sangtae, Alagapan Sankaraleengam, and Fröhlich Flavio. 2019 'Modulating neural oscillations by transcranial static magnetic field stimulation of the dorsolateral prefrontal cortex: A crossover, double-blind, sham-controlled pilot study', *European Journal of Neuroscience*, 49: 250–62. 10.1111/ejn.14232
- Söderpalm Anna HV, Lindsey Sommer, Purdy Robert H, Hauger Richard, and De Wit Harriet. 2004 'Administration of progesterone produces mild sedative-like effects in men and women', *Psychoneuroendocrinology*, 29: 339–54. 10.1016/s0306-4530(03)00033-7 [PubMed: 14644065]
- Stropahl Maren, Bauer Anna-Katharina Raphaela, Debener Stefan, and Bleichner Martin Georg. 2018 'Source-modelling auditory processes of EEG data using EEGLAB and Brainstorm', *Frontiers in neuroscience*, 12: 309 10.3389/fnins.2018.00309 [PubMed: 29867321]
- Sundström-Poromaa Inger. 2018 'The menstrual cycle influences emotion but has limited effect on cognitive function' in, *Vitamins and hormones* (Elsevier). 10.1016/bs.vh.2018.01.016
- Syan Sabrina K, Minuzzi Luciano, Costescu Dustin, Smith Mara, Allegra Olivia R, Coote Marg, Hall Geoffrey BC, and Frey Benicio N. 2017 'Influence of endogenous estradiol, progesterone, allopregnanolone, and dehydroepiandrosterone sulfate on brain resting state functional connectivity across the menstrual cycle', *Fertility and sterility*, 107: 1246–55. e4 10.1016/j.fertnstert.2017.03.021 [PubMed: 28476183]
- Tendler Alex, and Wagner Shlomo. 2015 'Different types of theta rhythmicity are induced by social and fearful stimuli in a network associated with social memory', *Elife*, 4: e03614 10.7554/eLife.03614
- Ukkola Olavi, Gagnon J, Rankinen T, Thompson PA, Hong Y, Leon AS, Rao DC, Skinner JS, Wilmore JH, and Bouchard C. 2001 'Age, body mass index, race and other determinants of steroid hormone variability: the HERITAGE Family Study', *European journal of endocrinology*, 145: 1–9. 10.1530/eje.0.1450001 [PubMed: 11415846]
- Vosskuhl Johannes, Huster René J, and Herrmann Christoph S. 2015 'Increase in short-term memory capacity induced by down-regulating individual theta frequency via transcranial alternating current stimulation', *Frontiers in human neuroscience*, 9: 257 10.3389/fnhum.2015.00257 [PubMed: 26005411]
- Voytek Bradley, Kramer Mark A, Case John, Lepage Kyle Q, Tempesta Zechari R, Knight Robert T, and Gazzaley Adam. 2015 'Age-related changes in 1/f neural electrophysiological noise', *Journal of Neuroscience*, 35: 13257–65. 10.1523/JNEUROSCI.2332-14.2015 [PubMed: 26400953]

- Yeo BT, Krienen Fenna M, Sepulcre Jorge, Sabuncu Mert R, Lashkari Danial, Hollinshead Marisa, Roffman Joshua L, Smoller Jordan W, Zöllei Lilla, and Polimeni Jonathan R. 2011 'The organization of the human cerebral cortex estimated by intrinsic functional connectivity', *Journal of neurophysiology*, 106: 1125–65. 10.1152/jn.00338.2011 [PubMed: 21653723]
- Zurkovsky L, Brown SL, Boyd SE, Fell JA, and Korol Donna L. 2007 'Estrogen modulates learning in female rats by acting directly at distinct memory systems', *Neuroscience*, 144: 26–37. 10.1016/j.neuroscience.2006.09.002 [PubMed: 17052857]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1. Frontal midline theta amplitude to steroid hormone correlation analysis

Correlation analysis (Pearson; $N=55$) of progesterone (A) and testosterone (B) concentration to frontal midline theta oscillatory amplitude. Solid line depicts significant relationship, dotted line is non-significant. Shaded area is 95% confidence interval. $** p < 0.01$. F = female, M = male.

Figure 2. Spectral analysis and aperiodic signal correction

Frontal midline theta oscillations survive aperiodic signal correction. Raw spectral amplitude for (A) FCz and (C) Oz electrode. Aperiodic signal was estimated by linear regression of the log(amplitude) and log(frequency) values for each participant. Then aperiodic signal was removed from the data for (B) FCz and (D) Oz electrode. Individual participants are displayed in light grey and the group average is displayed in black. Theta band is highlighted in a dark grey box. Alpha and beta band are highlighted in a light grey box. FCz and Oz are depicted on the scalp. Correlation to P4: ** $p < 0.01$, * $p < 0.05$, no asterisks $p > 0.05$, w/o = without.

Figure 3. Spatial distribution of frontal midline theta (FMT)

oscillations (A) Sensor-space correlation (Pearson; $N = 55$) between theta amplitude and P4 across women and men revealed the canonical distribution of FMT. FCz is outlined with a black square. A black circle depicts a correlation with significance of $p < 0.05$. (B) Source localization of theta amplitude. For display purposes, voxel-wise independent t -values ($df = 53$) between participants median split on sensor-space FMT amplitude. Theta oscillations localized to bilateral prefrontal cortex. Anterior-to-posterior axial view of prefrontal cortex. L = left, R = right. The red ellipses highlight increased theta oscillations in the lateral prefrontal cortices, and the blue circle highlights decreased theta oscillations in the superior frontal gyri.

Figure 4. Network analysis of source localized theta amplitude

Source localization analysis of frontal midline theta amplitude reveals activation in bilateral middle frontal gyrus (MFG), and not in superior frontal gyrus (SFG) or medial prefrontal cortex (mPFC). Lateral (A) and medial (B) view of the spatial distribution of increased theta amplitude (source-space) with increased frontal-midline theta amplitude (sensor-space) in left hemisphere. Lateral (C) and medial (D) view of the frontal-parietal (red) and default mode (blue) networks from an atlas that was independently derived from resting-state functional connectivity in fMRI (Yeo et al. 2011).

Table 1.**Sex differences**

Descriptive estimates, mean and standard deviation (SD), for all variables are displayed by sex. Data were normalized prior to t-test according to the description in Methods.

MEAN (SD)	Women (N = 23)	Men (N = 32)	Women & Men (N = 55)	Sex Difference (w – m)	
				t(53)	d
<i>Age [18–70 years]</i>	31.1 (13.7)	27.2 (12.3)	28.8 (12.9)	1.10	.30
<i>Progesterone (P4) [pg/mL]</i>	142.3 (103.7)	112.4 (76.9)	124.9 (89.5)	0.16	.04
<i>Testosterone (T) [pg/mL]</i>	20.2 (8.51)	94.3 (38.5)	63.3 (47.3)	–13.4***	3.62
<i>Frontal Midline Theta [μV]</i>	0.994 (0.31)	0.928 (0.24)	0.956 (0.27)	0.13	.04

p < 0.001.

Table 2.
Spectral analysis correlated to P4 concentration

Correlation analysis (Pearson; N = 55) between steroid hormone concentrations and peak spectral amplitude in three frequency bands (theta, alpha, and beta) in the frontal-midline (FCz) and the occipital-midline (Oz) in the raw power spectra and after correction for the aperiodic signal.

R-VALUES	FCz		Oz	
	Raw	w/o Aperiodic	Raw	w/o Aperiodic
Theta band [4–8Hz]	0.377 **	0.284 *	0.198	0.057
Alpha band [8–12Hz]	0.165	0.115	0.080	0.062
Beta band [15–30Hz]	–0.004	–0.131	–0.031	–0.124

**
p < 0.01,

*
p < 0.05, w/o = without.

Table 3.
Network analysis correlated with steroid hormones

Correlation analysis (Pearson; $N = 55$) between steroid hormone concentrations and theta amplitude within the frontal parietal network (FPN) and default mode network (DMN). FPN and DMN were independently derived from fMRI analysis.

	Progesterone (P4)	Testosterone (T)
<i>FPN - DMN</i>	0.310[*]	-0.079
<i>FPN</i>	0.343[*]	-0.019
<i>DMN</i>	-0.024	0.096

*
 $p < 0.05$

Table 4.**Exploratory analysis by sex**

Exploratory correlation analysis (Pearson) between steroid hormone concentrations and the amplitude of theta oscillations for each primary analysis (Methods section 2.3, 2.4, and 2.5) independently for both sexes.

R-VALUES	Women (N=23)		Men (N=32)	
	<i>P4</i>	<i>T</i>	<i>P4</i>	<i>T</i>
<i>Raw FMT Amplitude</i>	0.498 *	0.322	0.184	0.094
<i>FMT w/o Aperiodic</i>	0.383~	0.201	0.104	-0.018
<i>Theta [FPN – DMN]</i>	0.380~	0.284	0.217	0.127

* $p < 0.05$,

~ $p < 0.1$,

FMT = frontal midline theta, w/o = without, FPN = frontal-parietal network, DMN = default-mode network.