Bulging fontanel during a measles infection

Kübra Şahin^{1*}, Gülsüm İclal Bayhan², Başak Alan³

Abstract

Introduction Causes of a bulging fontanel in children include viral/bacterial meningitis, encephalitis, hydrocephaly, intracranial bleeding, psedotumor cerebri and central nervous system tumors. A less well-known cause of acutely-developing fontanel bulging is transient intracranial hypertension.

Case report A 7-month-old female was brought to our hospital with fever, cough, watery nasal secretion and bulging fontanel for the last 3 days. The cranial ultrasound examination and cerebrospinal fluid examinations were normal. The patient's irritability and fontanel bulging improved the day after admission but a maculopapular rash developed. The measles Ig M was positive. Measles PCR was positive from urine and nasopharyngeal swab samples. The signs of intracranial hypertension of the patient recovered completely the day after admission and did not recur during the follow-up. The patient was discharged on the 5th day of admission.

Discussion Transient intracranial hypertension is thought to result from a transient increase in the quantity of the circulating cerebrospinal fluid but the pathogenesis is not clear.

Conclusions Based on this case, we report that transient intracranial hypertension could develop during a measles infection.

Keywords Transient intracranial hypertension, measles, intracranial hypertension, fontanel bulging, fever, meningitis

Introduction

Causes of a bulging fontanel in children include viral/bacterial meningitis, encephalitis, hydrocephaly, intracranial bleeding, psedotumor cerebri and central nervous system tumors. A less well-known cause of acutely-developing fontanel

bulging is transient intracranial hypertension. Transient intracranial hypertension is a benign cause of increased intracranial pressure and is thought to develop due to a temporary increase in the amount of cerebrospinal fluid (CSF).¹ We report a seven-month-old infant who presented with fever, irritability and a bulging fontanel, and subsequently developed a measles rash.

Case report

A 7-month-old female was brought to our hospital with fever, cough, watery nasal secretion and bulging fontanel for the last 3 days. The patient had no history of an upper respiratory tract infection or contact with a child who had a rash. The patient was irritable with a hyperemic oropharynx and vesicles in the posterior pharynx on admission to our hospital. The fontanel was 2×1 cm in size and markedly bulging when evaluated with the patient calm and in the upright position. The neurological examination results were normal. The patient was admitted with a preliminary diagnosis of meningitis. The laboratory tests revealed the following: hemoglobin level 11.2 g/dL, MCV 82.5 fL, RDW 40.7 fL, white blood cell count 7770/μL

Received: 07 January 2020; revised: 09 April 2020; accepted: 13 June 2020.

¹MD, Yildirim Beyazit University, Faculty of Medicine, Department of General Pediatrics, Ankara City Hospital, Children's Hospital, Üniversiteler Mahallesi 1604., Cadde no: 9, Çankaya, Ankara, Turkey; ²MD, Yildirim Beyazit University, Faculty of Medicine, Ankara City Hospital, Children's Hospital, Üniversiteler Mahallesi 1604., Cadde no: 9, Çankaya, Ankara, Turkey; ³MD, Yildirim Beyazit University, Faculty of Medicine, Department of General Pediatrics, Ankara City Hospital, Children's Hospital, Üniversiteler Mahallesi 1604., Cadde no: 9, Çankaya, Ankara, Turkey.

*Corresponding author: Kübra Şahin, kubra.guneri.94@gmail.com

Article downloaded from www.germs.ro

Published September 2020

© GERMS 2020

ISSN 2248 - 2997

ISSN - L = 2248 - 2997

(neutrophils 45%, lymphocytes 49%, monocytes 5% on peripheral smear), and thrombocyte count 409 000/mm³; C-reactive protein (CRP) 9.7 mg/L (normal range 0-5 mg/L); normal biochemical parameters. The cranial ultrasound examination demonstrated normal cerebral and cerebellar hemispheres, corpus callosum and ventricular system. Lumbar puncture was performed, but the opening cerebrospinal fluid (CSF) pressure could not be measured. Direct microscopic examination of the CSF revealed no cells. CSF glucose was 66 mg/dL (simultaneous blood sugar 101 mg/dL) and CSF proteins were 17.5 mg/dL, thus excluding meningitis. The patient's irritability and fontanel bulging improved the day after admission without any treatment except for lumbar puncture which addressed intracranial hypertension but a maculopapular rash that started from the face and spread to the whole body developed. Koplik's spots were seen on oral mucosa evaluation. The measles Ig M level was 20.77 U/mL (positive). Measles PCR was positive from urine and nasopharyngeal swab samples. The patient was administered perorally 100 000 IU once a day vitamin A for two consecutive days with a diagnosis of measles. The cough increased on follow-up and the respiratory sounds became coarse. Sulbactam-ampicillin (intravenous 200 mg/kg/day q6 h) was started on the 3rd day of admission, when bilateral interstitial infiltration was observed on the chest X-ray. There was no growth on CSF, blood or urine cultures. The patient was discharged on the 5th day of admission when the rash and the respiratory system findings recovered. The patient continued the sulbactam-ampicillin treatment for five days perorally (50 mg/kg/day). The neurologic examination was normal at the 6th month follow-up.

Discussion

Measles morbillivirus, which is the causative agent of measles, is an RNA virus from the *Paramyxoviridae* family. The measles virus enters the body by inhalation of respiratory droplets followed by the invasion of and multiplication in nasopharyngeal epithelial cells. The next step is spread to the regional lymph nodes and

secondary viral replication there. The virus then infects lymphocytes and monocytes to cause viremia and to spread to almost all organs. The skin, conjunctivae and respiratory tract are typically involved, leading to the clinical signs of cough, nasal secretion, conjunctivitis, fever and a rash.² In addition to this typical clinical picture, the virus can access many organs and systems through viremia and cause other complications.³

The central nervous system (CNS) complications of measles are rare but quite serious and the main complications consist of primary measles encephalitis, measles inclusion body encephalitis, postinfectious encephalomyelitis and subacute sclerosing panencephalitis (SSPE). Primary measles encephalitis and measles inclusion body encephalitis are differentiated from the other forms by their development during acute infection.⁴ Measles inclusion body encephalitis is seen in immunocompromised individuals. In primary measles encephalitis, the virus invades the CNS and replicates here. The CNS infection typically develops simultaneously with the rash although it can also rarely develop before the rash.^{4,5} CSF evaluation reveals lymphocytic pleocytosis and mildly elevated protein levels.^{4,6} The presence of fever, irritability and a bulging fontanel in our patient with no immune deficiency resulted in a preliminary diagnosis of CNS infection. However, the completely normal cranial imaging and CSF findings and the rapid recovery of the irritability excluded this diagnosis.

Transient intracranial hypertension is a benign cause of increased intracranial pressure. It has been defined as consisting of a normal level of consciousness and normal neurological findings, no pathology on cranial imaging, normal CSF findings, the lack of another cause and normal neurological development on follow-up in patients with a bulging fontanel.⁷ It has been reported that the irritability of the patient improves and the fontanel bulging disappears after lumbar puncture.^{8,9} The normal CSF and cranial imaging results, the immediate improvement in the irritability and fontanel bulging following lumbar puncture, the normal neurological findings, and the normal neurological development at follow-up were

consistent with transient intracranial hypertension in our patient.

Transient intracranial hypertension is thought to result from a transient increase in the quantity of the circulating CSF but the pathogenesis is not clear. One hypothesis is that the proinflammatory cytokines secreted into the circulation during viral infections influence the choroid plexus which has a high degree of vascularity and results in increased CSF production here.¹ Transient intracranial hypertension has been reported to develop following human herpes virus 6 infection, diphtheria-tetanus-pertussis vaccination and vitamin A supplementation.^{7,9,10} We could not study the changes in plasma cytokine levels. Temporary fontanel bulging has been reported to develop after measles vaccination, however measles infection-related transient intracranial hypertension has not been previously reported.⁸

Conclusions

Many pediatricians have observed that the irritability of an infant with preliminary diagnosis of meningitis is relieved immediately after performing lumbar puncture. We therefore believe that transient intracranial hypertension could develop without the presence of meningitis during various infections, but the diagnosis could be missed as the condition is not commonly known. By presenting this case, we want to remind clinicians of the transient intracranial hypertension diagnosis and also to report that it could develop during a measles infection.

Consent: Written informed consent was obtained from the patient for publication of this case report and images. KŞ, GİB, BA

Authors' contributions statement: GİB, KŞ and BA contributed to concept, design and drafting of the manuscript. GİB, KŞ and BA contributed to analysis and

critical revision. All authors read and approved the final version of manuscript.

Conflicts of interest: All authors – none to declare.

Funding: None to declare.

References

1. Cristoforo T, Le NK, Rye-Buckingham S, Hudson WB, Carroll LF. The not-so-soft spot: pathophysiology of the bulging fontanelle in association with roseola. *Pediatr Emerg Care.* 2018. <https://doi.org/10.1097/PEC.0000000000001447>
2. Moench TR, Griffin DE, Obrecht CR, Vaisberg AJ, Johnson RT. Acute measles patients with and without neurological involvement: distribution of measles virus antigen and RNA. *J Infect Dis.* 1988;158:433-42. <https://doi.org/10.1093/infdis/158.2.433>
3. Makhene MK, Diaz PS. Clinical presentations and complications of suspected measles in hospitalized children. *Pediatr Infect Dis J.* 1993;12:836-40. <https://doi.org/10.1097/00006454-199310000-00008>
4. Buchanan R, Bonthius DJ. Measles virus and associated central nervous system sequelae. *Semin Pediatr Neurol.* 2012;19:107-14. <https://doi.org/10.1016/j.spen.2012.02.003>
5. Fox A, Hung TM, Wertheim H, et al. Acute measles encephalitis in partially vaccinated adults. *PLoS One.* 2013;8:e71671. <https://doi.org/10.1371/journal.pone.0071671>
6. Hänninen P, Arstila P, Lang H, Salmi A, Panelius M. Involvement of the central nervous system in acute, uncomplicated measles virus infection. *J Clin Microbiol.* 1980;11:610-3. <https://doi.org/10.1128/JCM.11.6.610-613.1980>
7. Freedman SB, Reed J, Burwen DR, Wise RP, Weiss A, Ball R. Transient bulging fontanelle after vaccination: case report and review of the vaccine adverse event reporting system. *J Pediatr.* 2005;147:640-4. <https://doi.org/10.1016/j.jpeds.2005.06.009>
8. Raju SS, Chacko B. Transient bulging fontanelle after measles vaccination. *Indian Pediatr.* 2014;51:673. <https://doi.org/10.1007/s13312-014-0478-9>
9. Mann NP, McLellan NJ, Cartledge PH. Transient intracranial hypertension of infancy. *Arch Dis Child.* 1988;63:966-8. <https://doi.org/10.1136/adc.63.8.966>
10. Goldberg EM. Fever and bulging fontanelle mimicking meningitis in an infant diagnosed with benign intracranial hypertension. *Pediatr Emerg Care.* 2013;29:513-4. <https://doi.org/10.1097/PEC.0b013e31828a3a28>

Please cite this article as:

Şahin K, Bayhan Gİ, Alan B. Bulging fontanel during a measles infection. *GERMS.* 2020;10(3):385-387. doi: 10.18683/germs.2020.1231