Published in final edited form as:

Clin Breast Cancer. 2018 April ; 18(2): e179–e185. doi:10.1016/j.clbc.2017.08.001.

Trends in Treatment Patterns and Clinical Outcomes in Young Women Diagnosed With Ductal Carcinoma In Situ

Hannah Lui Park, Jenny Chang, Gagandeep Lal, Krustina Lal, Argyrios Ziogas, Hoda Anton-Culver

Department of Epidemiology, University of California, Irvine, School of Medicine, Irvine, CA

Abstract

Younger women diagnosed with ductal carcinoma in situ (DCIS) are more likely to receive a more aggressive treatment compared with older women. Our analysis of Surveillance, Epidemiology, and End Results data (n = 3648) showed that young women who received mastectomy with or without contralateral prophylactic mastectomy did not have improved survival compared with breast-conserving surgery with radiation therapy. Thoughtful consideration should be given to treatment selection for DCIS in young women.

Background: Although it is known that the risk of a second breast cancer event among young women diagnosed with ductal carcinoma in situ (DCIS) is higher than in older women, the effect of current treatment options on long-term outcomes in this subgroup of women remains poorly defined. We aimed to evaluate national treatment trends and determine their effect on second breast cancer risk and overall survival among young women diagnosed with DCIS.

Materials and Methods: Surveillance, Epidemiology, and End Results data from 1998 to 2011 were used to analyze 3648 DCIS patients younger than age 40 years.

Results: Among all treatment options, breast-conserving surgery (BCS) with radiation therapy (BCS + RT) was the most prevalent (36.1%) followed by mastectomy (MTX) without contralateral prophylactic MTX (CPM; 25.8%), BCS alone (22.2%), and MTX with CPM (15.8%). Risk of a second ipsilateral event was > 5-fold and > 2-fold lower within 2 years and 5 years of initial DCIS diagnosis, respectively, in women who received BCS + RT compared with BCS alone; and overall survival was 3-fold higher in women who received BCS + RT. However, MTX with or without CPM did not show an increase in overall survival compared with BCS + RT. In addition, although the percentage of young women who receive MTX with CPM has increased in recent years, MTX with CPM did not show an increased benefit in survival compared with MTX without CPM.

Conclusion: The results of our study suggest that more aggressive treatments do not offer survival benefits over BCS + RT; thus, clinical treatment options in young women with DCIS should be carefully considered.

Address for correspondence: Hannah Lui Park, PhD, Department of Epidemiology, University of California, Irvine, School of Medicine, 222 Irvine Hall, Irvine, CA 92697 hlpark@uci.edu.

Disclosure

The authors have stated that they have no conflicts of interest.

Keywords

Breast-conserving surgery; Contralateral prophylactic mastectomy; Epidemiology; Mastectomy; Radiation therapy

Introduction

Ductal carcinoma in situ, also known as DCIS, is a noninvasive form of breast cancer, which might be a potential precursor to invasive breast cancer. In women between the ages of 20 and 49 years, DCIS accounted for approximately 24% of breast cancers diagnosed in 2013.¹ Women younger than 45 years are at a higher risk of a second event after diagnosis with DCIS compared with older women (27% vs. 11%),^{2,3} and up to half of these second events after DCIS are invasive.^{4,5} Treatment options for DCIS include breast-conserving surgery (BCS), BCS and radiation therapy (BCS + RT), and mastectomy (MTX). In addition, some women opt to undergo MTX with prophylactic removal of the contralateral breast, or contralateral prophylactic MTX (CPM).⁶

The goal of treatment for DCIS is not only to minimize the development of a second breast cancer event but also reduce its effect on the patient's quality of life.^{7,8} Among women diagnosed with DCIS, mental quality of life was significantly lower in younger women compared with older women diagnosed with DCIS.⁹ Because overtreatment might have potential effects on a patients' quality of life that are especially relevant in young women, to reach a balanced decision about which treatment option a young woman diagnosed with DCIS should take, it is important to understand the clinical benefits of the different treatment options in terms of developing a second breast cancer and survival in young women.

Studies comparing BCS + RT versus BCS alone have shown a survival benefit for DCIS patients overall who received BCS + RT,^{10–12} and numerous clinical trials have demonstrated that radiation therapy (RT) lowers the incidence for a second ipsilateral event.^{13,14} Studies have also shown that there is no difference in survival between women overall treated with BCS + RT versus MTX for women with early stage breast cancer.^{15–20} However, DCIS in young women specifically has been minimally studied in terms of second ipsilateral events and survival in the United States. In addition, there has been an increase in the use of CPM in women diagnosed with DCIS, especially among young women in the United States^{21–23}; however, the clinical benefits of CPM are not well understood in young women with DCIS and there have been controversial findings for young women diagnosed with low-grade invasive breast cancer.^{24,25} The current analysis focuses on recent trends in treatment options and clinical outcomes in young women with DCIS.

Materials and Methods

This is a retrospective population-based study using Surveillance, Epidemiology, and End Results (SEER) data. The SEER program of the National Cancer Institute contains approximately 97% of all incident cancer cases from population-based cancer registries covering approximately 28% of the US population. The SEER Program Registries routinely

collect data on patient demographic characteristics, primary tumor site, tumor morphology and stage, first course of treatment, and follow-up for vital status in 18 registries.

Our analysis included data on young women (age at diagnosis between 18 and 40 years old) diagnosed between January 1, 1998 and December 31, 2011 with primary DCIS breast cancer as their first or only tumor in their lifetime.²⁶ Breast cancer was identified using SEER primary site recode 26,000 with the median follow-up time of 7 years. International Classification of Disease for Oncology morphology codes categorized as DCIS included 85,002, 85,012, 85,032, 85,042, 85,222, 85,432, 80,502, and 82,012. Patients who were treated with MTX as well as radiation comprised 0.7% of data and were excluded from analysis because of small sample size. There was 1 case identified according to autopsy or death certificate only, which was excluded. The final analytic data set included 3648 subjects.

Variables and Statistical Analysis

Treatment for DCIS was identified for each patient using the “surgery for primary site” and “radiation therapy” variables provided in SEER data. The 4 types of treatment were BCS alone, BCS + RT, MTX with CPM, and MTX without CPM.

Other covariates included demographic and tumor characteristics. Race/ethnicity of the patient was categorized into 5 groups: White, African American, Hispanic, Asian/Pacific Islander, or other/unknown. Patient marital status was categorized into married, not married, or unknown. Registry region had 4 categories: central (Detroit, Iowa, Utah, Kentucky, and Louisiana), eastern (Connecticut, Georgia, and New Jersey), western (Hawaii, New Mexico, Seattle, and Alaska), and California. Tumor characteristics included tumor grade (low grade, high grade, or unknown), tumor size (≤ 15 mm, > 15 mm, or unknown size), tumor histology (comedo or other histology), estrogen receptor status, and progesterone receptor status, all obtained from the SEER data.

Frequency distributions of patients’ demographic and clinical characteristics were analyzed using the χ^2 test for categorical variables or analysis of variance test for continuous variables in bivariate analysis. Multivariate logistic regression models were used to test for associations between receiving BCS + RT and the characteristics. The Kaplan—Meier method was used to determine the cumulative risk of a second ipsilateral breast tumor from the date of first DCIS diagnosis to the date of second ipsilateral tumor diagnosis. Incidence rates of a second ipsilateral breast tumor were calculated in BCS alone and BCS + RT treatment groups for follow-up times of 2 years, 5 years, and the entire study period. After verifying the proportionality assumption, Cox proportional hazards model was fitted to evaluate the effects of additional RT on the risk of developing second ipsilateral breast cancer controlling for other characteristics. Patients’ follow-up time was defined as the time between diagnosis date and death from all causes or last follow-up date. After verifying proportionality assumption, a proportional hazards model was fitted to evaluate the effects of treatment option and other predictor on overall survival, and possible interaction terms of main effects were also tested by comparing a reduced model with the full model. All statistical tests were 2-sided. *P* values < .05 were considered statistically significant. All statistical analyses were performed using SAS version 9.4 (SAS Institute, Cary, NC).

Results

Trends in Treatment Options

We identified 3648 DCIS patients younger than the age of 40 years who met the eligibility criteria. The greatest proportion of patients underwent BCS + RT (36.1%) followed by MTX without CPM (25.8%), then BCS alone (22.2%), and MTX with CPM (15.8%). Treatment trends have significantly changed in the recent years going from 1998 through 1999 to 2005 through 2011. The percentage of women who received MTX with CPM increased (7% to 21.5%) whereas the percentages of women getting BCS + RT (38.8% to 35.3%), BCS alone (25.1% to 20%), and MTX without CPM (29.1% to 23.1%) decreased (Table 1).

Women who were diagnosed with DCIS and got MTX in 2005 to 2011 were 4.18-fold more likely to get CPM than women diagnosed in 1998 to 1999 (Table 2). Among the same women, African American women were significantly less likely to get CPM (odds ratio [OR], 0.29; 95% confidence interval [CI], 0.20–0.44), followed by Asian women (OR, 0.48; 95% CI, 0.31–0.73) and Hispanic women (OR, 0.57; 95% CI, 0.38–0.84) compared with white women (Table 2).

Second Ipsilateral Breast Tumor

Young women who got BCS alone were more likely to have a second ipsilateral breast tumor compared with women who received BCS + RT, at 2 years as well as at 5 years after DCIS diagnosis (hazard ratio [HR], 5.36; 94% CI, 2.02–14.3; $P = .0008$ and HR, 2.46; 95% CI, 1.47–4.11; $P = .0006$, respectively), after adjusting for year of diagnosis, race, registry region, tumor grade, histology, size, and hormone receptor status (Table 3). The small numbers of second ipsilateral events limited our stratified analysis according to second DCIS events, but the trend remained statistically significant for second invasive ipsilateral events within 5 years (HR, 2.06; 95% CI, 1.12–3.77; $P = .019$).

Survival

Overall survival was 3-fold lower in young women who received BCS alone (HR, 3.26; 95% CI, 1.58–6.73; $P = .0014$) compared with BCS + RT, after adjusting for potential confounders (Table 4). Overall survival in young women was not significantly different between those who received MTX with (HR, 1.16; 95% CI, 0.38–3.56; $P = .80$) or without (HR, 1.65; 95% CI, 0.79–3.46; $P = .19$) CPM compared with BCS + RT. The results were consistent after propensity weight scoring for adjusting for confounded and inverse probability weighting. There was also no significant difference in overall survival in young women who received MTX with CPM compared with without CPM (data not shown).

Discussion

With advancements in mammographic techniques, the incidence and detection of DCIS is on the rise.²⁷ Because only a limited number of studies on DCIS focused on women younger than 40 years, there is little clarity as to which treatment option is best for this age group of women. The trends in treatment patterns among young women diagnosed with DCIS observed in our study are different than in a previous analysis of women of all age groups,

which reported that the greatest proportion opted for BCS + RT (43%), followed by BCS alone (26.5%), unilateral MTX (23.8%), and MTX with CPM (4.5%).²⁸ The rise in use of MTX with CPM among young women diagnosed with DCIS observed in our study is in agreement with previous studies, which have also shown that MTX is increasingly becoming a popular choice for women over the past decade.^{21,22} Because young women are more likely to undergo MTX compared with women overall and the percentage of young women who choose MTX with CPM is increasing, it is imperative to study not only treatment trends but also clinical outcomes in this specific age group of women.

Our observation that young white women were more likely to get MTX with CPM compared with women of other race/ethnicities is consistent with a previous study done in women of all age groups diagnosed with DCIS, which suggested that the trend toward getting a more aggressive treatment option most prominent among white women compared with other ethnic groups.²⁸ An unexpected finding was that women with tumors > 15 mm were less likely to get MTX with CPM than MTX without CPM compared with women with tumors up to 15 mm. This might be because a large proportion of tumors classified with “unknown” size actually are smaller tumors that cannot be measured. Because we do not have data that this was actually the case, we allowed tumors of unknown size to remain as a separate category; however, if we combined women with tumors of unknown size and women with tumors up to 15 mm, there would have been no statistically significant difference in likelihood of undergoing the different treatments.

Consistent with previous studies, which showed that women of all ages diagnosed with DCIS who got RT had a 60% decreased risk of second ipsilateral event compared with women who got BCS alone,²⁹ our data also showed that young women with DCIS who got BCS + RT also had a > 5-fold and > 2-fold lower risk of a second ipsilateral event within 2 years and 5 years of initial DCIS diagnosis, respectively. These are important parameters because previous long-term studies have shown that approximately half of the recurrences after DCIS diagnosis occurred within the first 5 years.^{30,31} However, considering all second ipsilateral events that occurred in our study cohort, which had up to 13 years of follow-up, there was no difference in risk, suggesting that the protective effect of radiation decreases over time, which is consistent with recent findings in women with DCIS in the Netherlands.³²

Although our data indicate that, after adjusting for potential confounders, young women who received BCS alone were more than 3-fold likely to die compared with women who received BCS + RT, there does not appear to be a survival benefit associated with undergoing MTX with or without CPM over BCS + RT. There are other reasons women might opt for MTX with or without CPM. For example, a woman might choose MTX because of physician advice, personal fear of developing a second breast cancer event, high levels of stress and anxiety related to future breast cancer screening of the remaining breast(s), family history, and others reasons.^{33,34} Some women might opt for MTX with CPM in part because of a desire for better cosmetic symmetry.³⁴ In addition, the use of magnetic resonance imaging (MRI) preoperatively allows for a better interpretation of the contralateral breast, and women who receive MRI are more likely to choose CPM.³⁵ In young women, these issues are particularly relevant because they have a longer expected

number of years to live and also tend to overestimate their risk of having a contralateral breast cancer in the future.³⁶ Thus, although a survival benefit is not evident, MTX with or without CPM might offer young women diagnosed with DCIS better outcomes in terms of alleviating anxiety of a second breast cancer in the future and general peace of mind. Because the rate of young women diagnosed with DCIS who choose to undergo MTX with CPM has recently increased, future studies are needed to better understand the long-term effects of this treatment in relation to overall and breast cancer-free survival. In addition, the increasing trend warrants studies on other aspects of women's health including quality of life and well-being.

Limitations of this study include the lack of family history, mutation status of the BRCA1 and BRCA2, genetic testing, and hormone therapy results available in the SEER database. Because our study focused on DCIS in young women, our results are also limited by sample size and follow-up time. However, despite these limitations, SEER data account for a large diverse set of patients from all parts of the United States, thus making our findings broadly applicable to overall breast cancer trends in the country. To our knowledge, this study is unique in that it uses the most recent patient data from the SEER database (diagnosed between 1998 and 2011) to study risk of second ipsilateral events as well as survival between young women diagnosed with DCIS who received different clinical treatments. Further studies on large populations with longer follow-up time and risk factor and treatment information are warranted to better understand the most clinically beneficial treatment option for young women with DCIS.

Conclusion

Young women with DCIS and their physicians should give thoughtful consideration to all treatment options because although young women choose MTX more so than do older women, there appears to be no significant difference in overall survival between young women who received MTX compared with BCS + RT. In addition, there was no significant difference in overall survival between young women who received MTX with CPM compared with without CPM. Young women with DCIS should carefully consider all treatment options before making an informed decision, because more aggressive treatments (MTX with or without CPM) do not appear to confer survival benefit over BCS + RT.

Acknowledgments

The authors thank the UC Irvine Undergraduate Research Opportunities Program for their support of G.L. and K.L.

References

1. Howlader N, Noone AM, Krapcho M, et al., eds. SEER Cancer Statistics Review, 1975–2013. National Cancer Institute, Bethesda, MD, 2016, Available at: https://seer.cancer.gov/csr/1975_2013/.
2. Kong I, Narod SA, Taylor C, et al. Age at diagnosis predicts local recurrence in women treated with breast-conserving surgery and postoperative radiation therapy for ductal carcinoma in situ: a population-based outcomes analysis. *Curr Oncol* 2014; 21:e96–104. [PubMed: 24523627]
3. Alvarado R, Lari SA, Roses RE, et al. Biology, treatment, and outcome in very young and older women with DCIS. *Ann Surg Oncol* 2012; 19:3777–84. [PubMed: 22622473]

4. Romero L, Klein L, Ye W, et al. Outcome after invasive recurrence in patients with ductal carcinoma in situ of the breast. *Am J Surg* 2004; 188:371–6. [PubMed: 15474428]
5. Lee LA, Silverstein MJ, Chung CT, et al. Breast cancer-specific mortality after invasive local recurrence in patients with ductal carcinoma-in-situ of the breast. *Am J Surg* 2006; 192:416–9. [PubMed: 16978940]
6. Brown D, Shao S, Jatoi I, Shriver CD, Zhu K. Trends in use of contralateral prophylactic mastectomy by racial/ethnic group and ER/PR status among patients with breast cancer: a SEER population-based study. *Cancer Epidemiol* 2016; 42: 24–31. [PubMed: 26999400]
7. Smith BD. When is good enough really good enough? Defining the role of radiation in low-risk ductal carcinoma in situ. *J Clin Oncol* 2015; 33:686–91. [PubMed: 25605858]
8. Smith BD, Smith GL, Buchholz TA. Controversies over the role of radiation therapy for ductal carcinoma in situ. *Expert Rev Anticancer Ther* 2008; 8:433–41. [PubMed: 18366290]
9. Hart V, Sprague BL, Lakoski SG, et al. Trends in health-related quality of life after a diagnosis of ductal carcinoma in situ. *J Clin Oncol* 2016; 34:1323–9. [PubMed: 26884560]
10. Sagara Y, Freedman RA, Vaz-Luis I, et al. Patient prognostic score and associations with survival improvement offered by radiotherapy after breast-conserving surgery for ductal carcinoma in situ: a population-based longitudinal cohort study. *J Clin Oncol* 2016; 34:1190–6. [PubMed: 26834064]
11. Qian GW, Ni XJ, Wang Z, Jiang YZ, Yu KD, Shao ZM. Effect of radiotherapy on survival of women with locally excised ductal carcinoma in situ of the breast: a Surveillance, Epidemiology, and End Results population-based analysis. *Onco Targets Ther* 2015; 8:1407–18. [PubMed: 26089689]
12. Donker M, Litiere S, Werutsky G, et al. Breast-conserving treatment with or without radiotherapy in ductal carcinoma in situ: 15-year recurrence rates and outcome after a recurrence, from the EORTC 10853 randomized phase III trial. *J Clin Oncol* 2013; 31:4054–9. [PubMed: 24043739]
13. Julien JP, Bijker N, Fentiman IS, et al. Radiotherapy in breast-conserving treatment for ductal carcinoma in situ: first results of the EORTC randomised phase III trial 10853. EORTC Breast Cancer Cooperative Group and EORTC Radiotherapy Group. *Lancet* 2000; 355:528–33. [PubMed: 10683002]
14. Warnberg F, Garmo H, Emdin S, et al. Effect of radiotherapy after breast-conserving surgery for ductal carcinoma in situ: 20 years follow-up in the randomized SweDCIS trial. *J Clin Oncol* 2014; 32:3613–8. [PubMed: 25311220]
15. Blichert-Toft M, Nielsen M, Düring M, et al. Long-term results of breast conserving surgery vs. mastectomy for early stage invasive breast cancer: 20-year follow-up of the Danish randomized DBCG-82TM protocol. *Acta Oncol* 2008; 47:672–81. [PubMed: 18465335]
16. Fisher B, Anderson S, Bryant J, et al. Twenty-year follow-up of a randomized trial comparing total mastectomy, lumpectomy, and lumpectomy plus irradiation for the treatment of invasive breast cancer. *N Engl J Med* 2002; 347:1233–41. [PubMed: 12393820]
17. Litiere S, Werutsky G, Fentiman IS, et al. Breast conserving therapy versus mastectomy for stage I-II breast cancer: 20 year follow-up of the EORTC 10801 phase 3 randomised trial. *Lancet Oncol* 2012; 13:412–9. [PubMed: 22373563]
18. Simone NL, Dan T, Shih J, et al. Twenty-five year results of the national cancer institute randomized breast conservation trial. *Breast Cancer Res Treat* 2012; 132: 197–203. [PubMed: 22113254]
19. Veronesi U, Cascinelli N, Mariani L, et al. Twenty-year follow-up of a randomized study comparing breast-conserving surgery with radical mastectomy for early breast cancer. *N Engl J Med* 2002; 347:1227–32. [PubMed: 12393819]
20. Wilkinson JB, Vicini FA, Shah C, et al. Twenty-year outcomes after breast-conserving surgery and definitive radiotherapy for mammographically detected ductal carcinoma in situ. *Ann Surg Oncol* 2012; 19:3785–91. [PubMed: 22644510]
21. Tuttle TM, Jarosek S, Habermann EB, et al. Increasing rates of contralateral prophylactic mastectomy among patients with ductal carcinoma in situ. *J Clin Oncol* 2009; 27:1362–7. [PubMed: 19224844]

22. Rutter CE, Park HS, Killelea BK, Evans SB. Growing use of mastectomy for ductal carcinoma-in situ of the breast among young women in the United States. *Ann Surg Oncol* 2015; 22:2378–86. [PubMed: 25564175]
23. Güth U, Myrick ME, Viehl CT, Weber WP, Lardi AM, Schmid SM. Increasing rates of contralateral prophylactic mastectomy — a trend made in USA? *Eur J Surg Oncol* 2012; 38:296–301. [PubMed: 22305274]
24. Pesce C, Liederbach E, Wang C, Lapin B, Winchester DJ, Yao K. Contralateral prophylactic mastectomy provides no survival benefit in young women with estrogen receptor-negative breast cancer. *Ann Surg Oncol* 2014; 21:3231–9. [PubMed: 25081341]
25. Hwang ES. The impact of surgery on ductal carcinoma in situ outcomes: the use of mastectomy. *J Natl Cancer Inst Monogr* 2010; 2010:197–9. [PubMed: 20956829]
26. SEER Research Data 1998–2011 — ASCII Text Data: Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) Research Data (1998–2011). National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch.
27. Chuwa EW, Tan VH, Tan PH, Yong WS, Ho GH, Wong CY. Treatment for ductal carcinoma in situ in an Asian population: outcome and prognostic factors. *ANZ J Surg* 2008; 78:42–8. [PubMed: 18199204]
28. Worni M, Akushevich I, Greenup R, et al. Trends in treatment patterns and outcomes for ductal carcinoma in situ. *J Natl Cancer Inst* 2015; 107:djv263. [PubMed: 26424776]
29. Viani GA, Stefano EJ, Afonso SL, et al. Breast-conserving surgery with or without radiotherapy in women with ductal carcinoma in situ: a meta-analysis of randomized trials. *Radiat Oncol* 2007; 2:28. [PubMed: 17683529]
30. Habel LA, Daling JR, Newcomb PA, et al. Risk of recurrence after ductal carcinoma in situ of the breast. *Cancer Epidemiol Biomarkers Prev* 1998; 7:689–96. [PubMed: 9718221]
31. Wallis MG, Clements K, Kearins O, Ball G, Macartney J, Lawrence GM. The effect of DCIS grade on rate, type and time to recurrence after 15 years of follow-up of screen-detected DCIS. *Br J Cancer* 2012; 106:1611–7. [PubMed: 22516949]
32. Elshof LE, Schaapveld M, Schmidt MK, Rutgers EJ, van Leeuwen FE, Wesseling J. Subsequent risk of ipsilateral and contralateral invasive breast cancer after treatment for ductal carcinoma in situ: incidence and the effect of radiotherapy in a population-based cohort of 10,090 women. *Breast Cancer Res Treat* 2016; 159:553–63. [PubMed: 27624164]
33. Lostumbo L, Carbine NE, Wallace J. Prophylactic mastectomy for the prevention of breast cancer. *Cochrane Database Syst Rev* 2010:CD002748. [PubMed: 21069671]
34. Montgomery LL, Tran KN, Heelan MC, et al. Issues of regret in women with contralateral prophylactic mastectomies. *Ann Surg Oncol* 1999; 6:546–52. [PubMed: 10493622]
35. Sorbero ME, Dick AW, Beckjord EB, Ahrendt G. Diagnostic breast magnetic resonance imaging and contralateral prophylactic mastectomy. *Ann Surg Oncol* 2009; 16:1597–605. [PubMed: 19330381]
36. Rosenberg SM, Tracy MS, Meyer ME, et al. Perceptions, knowledge, and satisfaction with contralateral prophylactic mastectomy among young women with breast cancer: a cross-sectional survey. *Ann Intern Med* 2013; 159:373–81. [PubMed: 24042365]

Clinical Practice Points

- In 2013, DCIS accounted for approximately 24% of breast cancers in women between the ages of 20 and 49 years.
- Studies that have compared BCS + RT versus BCS alone have shown that RT lowers the incidence for a second ipsilateral event and increases survival in DCIS patients overall.
- Previous studies have also shown that there are no differences in survival between women treated with BCS + RT versus MTX for women overall with early stage breast cancer.
- In the present study we found that treatment trends for young women (younger than age 40 years) diagnosed with DCIS have significantly changed in the recent years from 1998 through 1999 to 2005 through 2011, with a significant increase in the percentage of young women who received MTX with CPM in recent years.
- Overall survival in young women with DCIS who received BCS + RT was significantly higher compared with BCS alone.
- There was no difference in overall survival between young women with DCIS who received MTX with or without CPM compared with BCS + RT.

Table 1
Patient Demographic and Tumor Characteristics According to Treatment in the Study Population, 1998–2011

	Total		BCS Alone		BCS and RT		MTX With CPM		MTX Without CPM	
	n	%	n	%	n	%	n	%	n	%
Total	3648	100.0	811	22.2	1318	36.1	577	15.8	942	25.8
Age at First DCIS Diagnosis										
Mean age $\pm$ SD	37.2 $\pm$ 3.4		37.1 $\pm$ 3.8		37.7 $\pm$ 3.0		36.6 $\pm$ 3.6		37.1 $\pm$ 3.3	
Year of Diagnosis										
1998–1999	402	11.0	101	25.1	156	38.8	28	7.0	117	29.1
2000–2004	1528	41.9	366	24.0	555	36.3	179	11.7	428	28.0
2005–2011	1718	47.1	344	20.0	607	35.3	370	21.5	397	23.1
Race										
White	2350	64.4	485	20.6	835	35.5	442	18.8	588	25.0
African American	462	12.7	110	23.8	164	35.5	39	8.4	149	32.3
Hispanic	379	10.4	94	24.8	131	34.6	50	13.2	104	27.4
Asian/Pacific Islander	415	11.4	111	26.7	172	41.4	39	9.4	93	22.4
Marital Status										
Married	2449	67.1	508	20.7	857	35.0	427	17.4	657	26.8
Not married	1061	29.1	261	24.6	419	39.5	130	12.3	251	23.7
Unknown	138	3.8	42	30.4	42	30.4	20	14.5	34	24.6
Registry Region										
California	1303	35.7	343	26.3	420	32.2	204	15.7	336	25.8
Central	782	21.4	120	15.3	297	38.0	142	18.2	223	28.5
Eastern	1230	33.7	288	23.4	464	37.7	194	15.8	284	23.1
Western	333	9.1	60	18.0	137	41.1	37	11.1	99	29.7
Grade										
Low grade (I or II)	1428	39.1	390	27.3	555	38.9	201	14.1	282	19.7
High grade (III or higher)	1428	39.1	201	14.1	485	34.0	270	18.9	472	33.1
Unknown	792	21.7	220	27.8	278	35.1	106	13.4	188	23.7
Histology										

	Total		BCS Alone		BCS and RT		MTX With CPM		MTX Without CPM	
	n	%	n	%	n	%	n	%	n	%
Comedo	654	17.9	85	13.0	247	37.8	114	17.4	208	31.8
All others	2994	82.1	726	24.2	1071	35.8	463	15.5	734	24.5
Size										
15 mm	1455	39.9	364	25.0	693	47.6	177	12.2	221	15.2
>15 mm	948	26.0	146	15.4	237	25.0	188	19.8	377	39.8
Unknown	1245	34.1	301	24.2	388	31.2	212	17.0	344	27.6
Laterality										
Right side	1808	49.6	407	22.5	641	35.5	296	16.4	464	25.7
Left side	1836	50.4	402	21.9	677	36.9	281	15.3	476	25.9
Hormone Status										
Either ER or PR is positive	1568	43.0	284	18.1	611	39.0	303	19.3	370	23.6
ER as well as PR are negative	254	7.0	35	13.8	78	30.7	52	20.5	89	35.0
Unknown	1826	50.1	492	26.9	629	34.4	222	12.2	483	26.5

Data set excludes MTX not otherwise specified; *P* values for χ^2 tests and analysis of variance test were all < .05, except for laterality (*P* = .72).

Abbreviations: BCS = breast-conserving surgery; CPM = contralateral prophylactic mastectomy; DCIS = ductal carcinoma in situ; ER = estrogen receptor; MTX = mastectomy; PR = progesterone receptor; RT = radiation therapy.

Table 2
Logistic Regression Models of Factors Associated With Treatment in Subsets of Young Women With DCIS

	Factors Associated With Receiving BCS and RT, Among the Subset of Patients Who Had BCS and RT or BCS Alone		Factors Associated With Receiving BCS and RT, Among the Subset of Patients Who Had BCS and RT or MTX (With or Without CPM)		Factors Associated With Receiving MTX and CPM, Among the Subset of Patients Who Had MTX With or Without CPM	
	OR (95% CI)	P	OR (95% CI)	P	OR (95% CI)	P
Age at Diagnosis						
	1.05 (1.02–1.08)	.0003	1.08 (1.05–1.11)	<.0001	0.94 (0.91–0.97)	.0003
Year of Diagnosis						
1998–1999	1.00	–	1.00	–	1.00	–
2000–2004	1.00 (0.74–1.34)	.98	0.85 (0.65–1.12)	.25	1.68 (1.06–2.65)	.027
2005–2011	0.85 (0.60–1.19)	.34	0.67 (0.50–0.91)	.011	4.18 (2.57–6.79)	<.0001
Race						
White	1.00	–	1.00	–	1.00	–
African American	0.87 (0.65–1.16)	.34	1.10 (0.85–1.41)	.47	0.29 (0.20–0.44)	<.0001
Hispanic	1.04 (0.76–1.42)	.80	1.27 (0.97–1.68)	.088	0.57 (0.38–0.84)	.0051
Asian/Pacific Islander	1.08 (0.81–1.44)	.61	1.75 (1.33–2.30)	<.0001	0.48 (0.31–0.73)	.0006
Other	1.32 (0.56–3.10)	.53	1.70 (0.80–3.63)	.17	1.22 (0.41–3.62)	.73
Marital Status						
Married	1.00	–	1.00	–	1.00	–
Not married	1.04 (0.85–1.27)	.74	1.47 (1.22–1.76)	<.0001	0.91 (0.70–1.19)	.51
Unknown	0.60 (0.37–0.95)	.03	0.90 (0.58–1.39)	.63	0.85 (0.46–1.56)	.60
Registry Region						
California	1.00	–	1.00	–	1.00	–
Central	2.20 (1.66–2.91)	<.0001	1.12 (0.89–1.40)	.34	0.88 (0.65–1.20)	.43
Eastern	1.51 (1.20–1.89)	.0005	1.28 (1.04–1.58)	.019	1.00	.99
Western	1.89 (1.33–2.69)	.0004	1.21 (0.90–1.63)	.20	0.60 (0.39–0.94)	.024
Grade						
Low grade (I or II)	1.00	–	1.00	–	1.00	–
High grade (III or higher)	1.47 (1.16–1.84)	.0012	0.70 (0.58–0.85)	.0002	0.85 (0.66–1.11)	.23
Unknown	0.95 (0.75–1.20)	.66	0.93 (0.75–1.16)	.52	0.93 (0.67–1.29)	.67
Histology						

	Factors Associated With Receiving BCS and RT, Among the Subset of Patients Who Had BCS and RT or BCS Alone		Factors Associated With Receiving BCS and RT, Among the Subset of Patients Who Had BCS and RT or MTX (With or Without CPM)		Factors Associated With Receiving MTX and CPM, Among the Subset of Patients Who Had MTX With or Without CPM	
	OR (95% CI)	P	OR (95% CI)	P	OR (95% CI)	P
Conedo	1.00	–	1.00	–	1.00	–
All others	0.59 (0.44–0.78)	.0002	0.94 (0.76–1.15)	.52	1.03 (0.78–1.36)	.85
Size						
15 mm	1.00	–	1.00	–	1.00	–
>15 mm	0.81 (0.63–1.05)	.11	0.27 (0.22–0.33)	<.0001	0.63 (0.47–0.83)	.0012
Unknown	0.68 (0.55–0.84)	.0003	0.41 (0.34–0.49)	<.0001	0.86 (0.65–1.15)	.31
Hormone Receptor Status						
Either ER or PR is positive	1.00	–	1.00	–	1.00	–
ER and PR are negative	0.87 (0.56–1.36)	.55	0.80 (0.58–1.11)	.19	0.75 (0.50–1.12)	.16
Unknown	0.56 (0.45–0.71)	<.0001	0.87 (0.71–1.07)	.19	0.92 (0.69–1.23)	.57

Abbreviations: BCS = breast-conserving surgery; CPM = contralateral prophylactic mastectomy; DCIS = ductal carcinoma in situ; ER = estrogen receptor; MTX = mastectomy; PR = progesterone receptor; RT = radiation therapy.

Table 3

Comparison of Number of Second Ipsilateral Events, Incidence Rates and Hazard Ratios During Follow-up in BCS and BCS With RT Treatment Groups

	BCS Alone			BCS and RT			Hazard Ratio (95% CI) and P Value for BCS and RT Compared With the BCS Alone Group From the Multivariable Cox Model ^a
	Person-Years	Cases	Incidence Rate (%) and 95% CI	Person-Years	Cases	Incidence Rate (%) and 95% CI	
2-Year Ipsilateral Breast Tumor	1486	19	1.28 (0.71–1.85)	2470	6	0.24 (0.05–0.44)	.0008
2-Year Ipsilateral In Situ		8	–		1	–	–
2-Year Ipsilateral Invasive		11	–		5	–	–
5-Year Ipsilateral Breast Tumor	3275	45	1.37 (0.98–1.77)	5479	29	0.53 (0.3–0.72)	.0006
5-Year Ipsilateral In Situ		17	–		5	–	–
5-Year Ipsilateral Invasive		28	0.86 (0.54–1.17)		24	0.44 (0.26–0.61)	.019
All Ipsilateral Breast Tumor	5444	72	1.32 (1.02–1.63)	8984	88	0.98 (0.78–1.18)	.79
All Ipsilateral In Situ		23	0.42 (0.25–0.59)		26	0.29 (0.18–0.40)	.48
All Ipsilateral Invasive		49	0.90 (0.65–1.15)		62	0.69 (0.52–0.86)	.85

Abbreviations: BCS = breast-conserving surgery; RT = radiation therapy.

^aHazard ratios are for the BCS with RT group compared with the BCS-alone group from multivariable proportional hazards models on time to second ipsilateral breast cancer controlling for age, year of diagnosis, race, registry region, tumor grade, histology, size, and hormone receptor status.

Table 4

Results From Survival Analysis on Overall Survival Using 3 Methods

Treatment	Multivariable Proportional Hazards Cox Model ^a			With Propensity Score Covariate Adjustment			With Inverse Probability Weighting		
	HR	95% CI	P	HR	95% CI	P	HR	95% CI	P
BCS Alone	3.26	1.58–6.73	.0014	3.24	1.553–6.763	.0017	3.43	1.667–7.041	.0008
BCS and RT	1.00	–	–	1.00	–	–	1.00	–	–
MTX With CPM	1.16	0.38–3.56	.7995	1.17	0.38–3.65	.7865	1.04	0.40–2.90	.9676
MTX Without CPM	1.65	0.79–3.46	.1855	1.68	0.79–3.54	.1765	1.77	0.84–3.75	.1351

Abbreviations: BCS = breast-conserving surgery; CPM = contralateral prophylactic mastectomy; MTX = mastectomy; RT = radiation therapy.

^aMultivariable Cox model also controlled for race, registry region, tumor grade, histology, size, and hormone receptor status. Age at diagnosis and diagnosis year were included in the model as strata.