Dysbaric osteonecrosis in technical divers: The new 'at-risk' group?

Brendan Coleman¹, F Michael Davis²

¹ Auckland Orthopaedic Practice, Auckland 1543, New Zealand

² Department of Anaesthesiology, Faculty of Medicine and Health Sciences, The University of Auckland, Auckland, New Zealand

Corresponding author: Dr Brendan Coleman, Auckland Orthopaedic Practice, PO Box 74 446, Auckland 1543, New Zealand
brendan@aoppractice.co.nz

Key words

Dysbaric osteonecrosis; Technical diving; Decompression sickness; Deep diving; Radiological imaging; Orthopaedics; Case reports

Abstract

(Coleman B, Davis FM. Dysbaric osteonecrosis in technical divers: The new 'at-risk' group? Diving and Hyperbaric Medicine. 2020 September 30;50(3):295–299. doi: 10.28920/dhm50.3.295-299. PMID: 32957134.)

Introduction: Dysbaric osteonecrosis (DON) in people working under increased atmospheric pressure is well documented. It is generally less common in military and commercial divers than in caisson workers, except in some high-risk groups, such as in many indigenous diving industries where workers have little or no understanding of decompression principles. With the increasing popularity within the recreational diving community of deep air and mixed-gas decompression diving ('technical diving'), it is likely that diving physicians may see an increase in the prevalence of DON in this group in the future.

Methods: The case report is presented of a technical diving instructor, with a 30-year history of deep diving, who developed bilateral humeral head DON and required a right shoulder hemi-arthroplasty. A focused literature search was also undertaken to identify published cases of DON in recreational divers.

Results: The frequency, duration and depth of exposure to pressure, inadequate decompression, the occurrence of DCS and increasing age have been common features associated with DON in both divers and caisson workers. Many of these features were present in this technical diver.

Conclusions: Whilst DON is uncommon in recreational air scuba divers, all the above risk factors are present to a greater degree in technical diving. It is suggested that medical review for DON is merited from time to time in this potentially high-risk group of recreational divers.

Introduction

Aseptic bone necrosis (AVN) is the final common pathway of various conditions leading to bone death,¹ most commonly long-term, high-dose steroid use and alcoholism. Other contributing conditions include pancreatitis, lupus, sickle cell disease, Gaucher's disease, radiotherapy, trauma, pregnancy and thrombotic conditions.^{1,2} A recent review paper discusses the diagnosis and staging of AVN of the femoral head.³ AVN in caisson workers and divers, termed dysbaric osteonecrosis (DON), is well documented. In the 1970s, the Medical Research Council (MRC) Decompression Sickness Panel reported an incidence of 19% in caisson workers and reported a link to decompression sickness (DCS).^{4,5} It is reported to be less common in military⁶ and commercial divers^{7,8} except in some high-risk groups⁹ and those with little or no understanding of decompression principles.¹⁰ Readers are referred to a useful review of DON in professional divers.¹¹ Lesions may be either juxta-articular (Type A, Table 1) or in the shaft of long bones (Type B), the former being far more likely to produce symptoms.

DON is considered rare in recreational divers, though individual cases have been reported.^{12–15} With the increasing popularity within the recreational diving community of deep mixed-gas decompression diving ('technical diving'), it is likely that diving physicians may see an increase in the prevalence of DON in this group in the future because of the nature of their diving activities. For this reason, we present an illustrative case of a technical diving instructor with DON and briefly review the relevant literature.

Literature review

PubMed, the Rubicon Foundation Research Repository (<http://archive.rubicon-foundation.org/xmlui/>), the complete collection of the *South Pacific Underwater Medicine Society Journal* and *Diving and Hyperbaric Medicine* journal and major textbooks on diving medicine were searched using the terms avascular necrosis, dysbaric osteonecrosis, and caisson disease and in combination with the terms diving, decompression sickness, scuba and technical diving. Further articles of potential interest from reference lists were also reviewed. The intention was not to perform a comprehensive

Table 1

Classification of dysbaric osteonecrosis lesions (after the United Kingdom Medical Research Council). * = classification of the present case

Lesion	Subtype	Comments
A type lesions Juxta-articular	A1 Dense area with intact articular cortex	Prevalence of A lesions: Tunnellers and saturation divers Femur > Humerus Other divers Humerus > Femur
	A2 Spherical opacities	
	A3 Linear opacities	
	A4 Structural failures - Translucent subcortical bands - Collapse of articular cortex - Sequestration of cortex*	
	A5 Secondary degenerative osteoarthritis	
B type lesions Shaft	B1 Dense areas	n/a
	B2 Irregular calcified areas	
	B3 Translucent and cystic areas	

literature search, but to focus on published evidence relevant to recreational and technical diving.

Case report

A man in his 40s had been actively involved in recreational diving since 1991. He qualified as a dive instructor in 1994, teaching a mix of recreational and technical scuba diving around the world, doing 200 or more hours a year in the water. During this period, technical mixed gas diving took up about a quarter of his diving hours, but no detailed records of his diving were kept. Between 2010 and 2018, he recorded on his annual medical questionnaires over 1,800 dives of which 80% were to depths greater than 30 metres' sea water (msw) and over 1,500 were on mixed gas/trimix using closed-circuit rebreathers.

In 2002, he did a wreck dive to 115 msw using trimix, followed by a gas switch to air at 30 msw during the ascent. At the 9 msw decompression stop, he developed severe vertigo which persisted post dive. Audiometry was normal. A diagnosis of inner-ear decompression sickness (DCS) was made and he underwent a RNZN heliox Table 1A (similar to a Comex 30 treatment table) with a good response. Following two further short hyperbaric oxygen treatments he was symptom-free. Two years later in 2004, he dived on the same wreck to a depth of 120 msw. Again, a gas switch to air was made, this time at 40 msw. On reaching the 15 msw stop, he developed severe vertigo, vomiting and tinnitus. Because symptoms had improved at 6 msw after switching to 100% oxygen, he did not present for assessment until the following day when examination revealed a fine right nystagmus, very poor sharpened Romberg test and a 70 dB hearing loss at 8 KHz in the right ear. The diagnostic difficulty of differentiating between inner-ear DCS and barotrauma necessitated a cautious approach to recompression on a Royal Navy Treatment Table 62; however, his symptoms responded well. Again, after two further short hyperbaric oxygen treatments, he was symptom-free.

In 2011, aged 42 years, he developed pain in the left shoulder. Given his diving history, he underwent a long-bone X-ray survey which showed *"a localised lucency in the left humeral head with a surrounding sclerotic rim, the right shoulder was normal in appearance and there were faint, slightly serpiginous-appearing sclerotic lesions in the proximal left and distal right femoral shafts."* In 2013, he was seen by an orthopaedic surgeon (BC) for assessment of the left shoulder lesion, which was managed conservatively at that time. No other risk factors for DON other than diving were identified.

In late 2018, now 49-years old, he presented following minor trauma to his right shoulder, with significant pain and limitation of movement associated with crepitus. Plain X-ray (Figure 1) and magnetic resonance imaging (MRI) (Figure 2) of the right shoulder showed a juxta-articular DON lesion of the humeral head measuring 8 mm by 24 mm with fragmentation of the articular surface and disruption of the articular cartilage. Given his symptoms were impeding his ability to work, he proceeded to surgery, at which time the articular surface of the right humerus had a 25 mm by 20 mm unstable osteochondral fragment in the central head. A right shoulder pyrocarbon hemiarthroplasty of the humeral head was inserted.

Recovery was uneventful with steady return of strength and range of motion such that he was cleared to return to work as a diving instructor four months post-operatively. For the hemiarthroplasty, there is little increased risk of aseptic loosening above what is already expected for patients under the age of 55. These patients have approximately double the failure rate of over 55-year olds.¹⁶ The prognosis for the opposite shoulder is for gradual deterioration in symptoms over time due to the presence of DON in the humeral head with subchondral fracturing and collapse of the joint surface. This will eventually lead to arthritis or displacement of the fragmentation of the osteonecrotic segment, resulting in arthroplasty when the symptoms warrant.

Figure 1

Plain X-ray of the right shoulder of a recreational and technical diving instructor showing an avascular lesion of the humeral head

Figure 2

Coronal t2-weighted magnetic resonance image of the right shoulder of a recreational and technical diving instructor showing an avascular lesion of the humeral head

Discussion

AVN secondary to DCS was first noted in the late nineteenth century, culminating in work such as that of the MRC Decompression Sickness Panel in the 1960s and 1970s,^{4,5,7} and later studies in China¹⁷ and elsewhere. A high incidence of DON has been confirmed in high-risk diving groups such as long-term Japanese commercial divers,⁹ and Hawaii's diving fishermen.¹⁰ The United Kingdom DCS Central Registry and Radiological Panel determined that the risk of DON in caisson workers and commercial divers was directly related to the degree of pressure and the number of exposures,^{5,7} whilst in Japanese commercial divers, diving exposure over 17–25 years was associated with DON, particularly in those diving to 35 msw or deeper and where there was a history of DCS.⁹ In the majority of reports, juxta-articular lesions appear to be more common in the humeral head than in the femoral head of divers,^{7,8} though this was not so in one study of 450 'hard-hat' Japanese divers in which the distribution was similar for both sites.⁹ The prevalence of DON in divers in these various studies shows widely differing rates.

Thus, the frequency, duration and (deeper) depth of pressure exposure, inadequate decompression, the occurrence of DCS (and possibly delayed recompression) and increasing age have been common features associated with DON in both divers and caisson workers.^{4–9,17} Whether age is, in fact, an independent risk factor or secondary to the length of the pressurisation/diving career is debatable. Although inert gas embolism is thought to be a mechanism linked to DON,¹⁸ there is no clear evidence of a cause-and-effect relationship,^{19,20} this more likely being a gradual process over many dives. Other mechanisms, such as a hypercoagulable

state and fat emboli, also may be involved.²¹ No certain aetiology for DON has been established.

DON is uncommon in typical recreational diving. The first report of dysbaric osteonecrosis lesions in sports divers was in an Australian study of 110 navy, professional and sport divers.²² There were three lesions (one juxta-articular) in the 19 sport divers in the study. In the single cases reported,^{12–15} in one,¹⁵ subsequent enquiry from the author (Laden G, personal communication, 2019) revealed that this diver was a technical diver with several hundred deep and mixed-gas dives; in another,¹³ the diver had performed 190 dives over six years, over 100 having been to depths greater than 30 msw. By contrast, in a third,¹² only no-decompression air diving had been undertaken, mostly to less than 18 msw. None had had symptoms suggesting DCS. Nevertheless, single events of DCS, especially if there was a delay to recompression treatment, appear to be associated with an increased risk of DON even in recreational divers.^{23,24}

In a long-bone survey of 56 Turkish dive instructors each of whom had performed in excess of 500 dives, DON lesions were detected in 14, with only one juxta-articular lesion in the humeral head.²⁵ On univariate analysis of a wide range of factors, only increasing age was associated with DON and this, in itself, was correlated with diving experience and the total numbers of dives performed.

For occupational divers, long bone surveys are “*optional and as medically indicated*” in most jurisdictions.²⁶ In the Australian/New Zealand Standard,²⁷ juxta-articular DON is not considered to be a contraindication to continued

diving. Under these regulations, this dive instructor with known DON was cleared to dive following the condition and risks associated with continued diving having been fully discussed and understood, and a document signed by the diver to that effect.

Several features of the present case – many years of deep and decompression diving resulting in a higher likelihood of DCS and the possible role of gas switches in inducing a gas phase in sensitive tissues such as the inner ear²⁸ or bone – illustrate why technical divers are more likely to be at risk of DON than the average recreational diver. Therefore, we believe that recreational technical divers need greater regular medical screening than open-circuit air scuba divers. Early radiological referral in the presence of joint symptoms which might suggest a juxta-articular DON lesion is warranted, as in this case.

Reporting an amateur scuba diver who developed DON, the authors state “*avascular bone necrosis ... may lead to joint dysfunction and lifelong disability*”.¹⁵ Modern orthopaedic surgery for shoulder and hip joint pathology now allows a range of treatment options, including hemi- and total arthroplasty for patients with juxta-articular DON.²⁹ Joint preserving surgery for osteonecrosis of the shoulder includes core decompression, bone grafting or autologous bone marrow grafting. Core decompression can be effective in early stages of the disease when there is no collapse of the humeral head but is less effective once the humeral head shows signs of collapse.³⁰ Bone grafting, either simple autologous grafting, strut bone graft or vascularised bone graft are complex procedures with increased morbidity and variable results.^{31,32} It remains unclear as to the most effective joint sparing treatment of osteonecrosis. The natural history of conservatively treated advanced osteonecrosis of the humeral head is poor. Once the condition results in structural damage (equivalent of stage A4, Table 1), almost half will require shoulder arthroplasty.³³

Hyperbaric oxygen treatment (HBOT) has been reported in several studies to show long-term benefit for early (equivalent to stages A1–A3; Table 1) femoral head AVN. In one double-blind, randomised study of 20 patients, the HBOT group showed a significant reduction in pain at the end of 30 treatments ($P < 0.001$) compared with the sham air group who were then offered HBOT, which they all accepted.³⁴ At seven years’ follow up of 17 of the 20, “*all patients remained substantially pain-free ... with none requiring hip arthroplasty. Substantial radiographic healing ... was observed in seven of nine hips [on MRI]*.”³⁴

The medicolegal aspects of DON in working divers in the Tasmanian fish farming and abalone industries were discussed at a conference in Hobart in 1988 with reference to a new Workers’ Compensation Act that came into force that year in Australia.³⁵ This Act recognised DON in Schedule 4 as “*compressed air illness including avascular necrosis*

caused by any work involving exposure to increased or reduced atmospheric pressure from working underground or underwater or from working at high altitudes”. Similarly, this diving instructor’s DON was approved for treatment under the Accident Compensation Corporation (ACC) in New Zealand. Whether the ACC would recognise DON in a recreational diver who was not an ‘employed’ diver remains a moot point.

References

- 1 Fondi C, Franchi A. Definition of bone necrosis by the pathologist. Clin Cases Miner Bone Metab. 2007;4:21–26. PMID: 22460748. PMCID: PMC2781178.
- 2 Goodman SG. Osteonecrosis (ON). MSD Manual Professional Version; 2019. Available at: <https://www.msdmanuals.com/en-in/home/bone,-joint,-and-muscle-disorders/osteonecrosis/osteonecrosis>. [cited 2019 December 12].
- 3 Choi H-R, Steinberg ME, Cheng EY. Osteonecrosis of the femoral head: diagnosis and classification systems. Curr Rev Muskuloskelet Med. 2015;8:210–20. doi: 10.1007/s12178-015-9278-7. PMID: 26088795. PMCID: PMC4596207.
- 4 McCallum RI, Walder DN. Bone lesions in compressed air workers, with special reference to men who worked on the Clyde Tunnel 1958 to 1963. Report of Decompression Sickness Panel Medical Research Council. J Bone Joint Surg. 1966;48B:207–35.
- 5 Decompression Sickness Panel (Medical Research Council). Decompression sickness and aseptic necrosis of bone. Investigation carried out during and after the construction of the Tyne Road Tunnel (1962–1966). Br J Indust Med. 1971;28:1–21.
- 6 Hunter WL, Biersner RJ, Sphar RL, Harvey CA. Aseptic bone necrosis among U.S. Navy divers: survey of 934 nonrandomly selected personnel. NSMRL Report number 854. Groton (CT): Submarine Medical Research Laboratory; 1978. Available from: <https://apps.dtic.mil/dtic/tr/fulltext/u2/a058915.pdf>. [cited 2019 November 11].
- 7 Aseptic bone necrosis in commercial divers. Lancet. 1981;2:384–8. PMID: 6115158.
- 8 Yangsheng T, Anquan Y, Weimin L, Jingxi Q. Investigation and analysis of dysbaric osteonecrosis in 171 divers. J Hyperb Med. 1992;7:123–6. Available from: <http://archive.rubicon-foundation.org/4480>. [cited 2019 December 16].
- 9 Kawashima M. Aseptic bone necrosis in Japanese divers. Bull Tokyo Med Dent Univ. 1976;23:71–92.
- 10 Wade CE, Hayashi EM, Cashman TM Jr, Beckman EL. Incidence of dysbaric osteonecrosis in Hawaii’s diving fishermen. Undersea Biomed Res. 1978;5:137–47. PMID: 675879. Available from: <http://archive.rubicon-foundation.org/2705>. [cited 2019 December 16].
- 11 Uguen M, Pougnet R, Uguen A, Loddé B, Dewitte JD. Dysbaric osteonecrosis among professional divers: A literature review. Undersea Hyperb Med. 2014;41:581–9. PMID: 25562949.
- 12 Edmonds C, Harvey R, Randle R. Dysbaric osteonecrosis divers bone rot: A case report. SPUMS Journal. 1997;27:122–5. Available from: <http://archive.rubicon-foundation.org/6071>. [cited 2019 December 16].
- 13 Wilmshurst P, Ross K. Dysbaric osteonecrosis of the shoulder in a sport scuba diver. Br J Sports Med. 1998;32:344–5. doi: 10.1136/bjsm.32.4.344. PMID: 9865412. PMCID: PMC1756113.

- 14 Wong R, Wright D. Aseptic bone necrosis as a diagnostic puzzle. SPUMS Journal. 2001;31:135–8. Available from: <http://archive.rubicon-foundation.org/7732>. [cited 2019 December 16].
- 15 Laden GDM, Grout P. Aseptic bone necrosis in an amateur scuba diver. Br J Sports Med. 2004;38:E19. doi: 10.1136/bsjm.2002.003129. PMID: 15388563. PMCID: PMC1724939.
- 16 NZ Joint Registry Annual Report 2019. Available from: https://nzoa.org.nz/system/files/DH8328_NZJR_2019_Report_v4_7Nov19.pdf. [cited 2019 December 16].
- 17 Zhang L-D, Kang J-F, Xue H-L. Distribution of lesions in the head and neck of the humerus and the femur in dysbaric osteonecrosis. Undersea Biomed Res. 1990;17:353–8. PMID: 2396333. Available from: <http://archive.rubicon-foundation.org/2570>. [cited 2019 December 16].
- 18 Chryssanthou CP. Dysbaric osteonecrosis. Etiological and pathogenetic concepts. Clin Orthop Relat Res. 1978;(130):94–106. PMID: 639412.
- 19 Coulthard A, Pooley J, Reed J, Walder D. Pathophysiology of dysbaric osteonecrosis: A magnetic resonance imaging study. Undersea Hyperb Med. 1996;119:119–20. PMID: 8840481.
- 20 Kenney IJ, Sonksen C. Dysbaric osteonecrosis in recreational divers: A study using magnetic resonance imaging. Undersea Hyperb Med. 2010;37:281–8. PMID: 20929185.
- 21 Miyaniishi K, Kamo Y, Ihara H, Naka T, Hirakawa M, Sugioka Y. Risk factors for dysbaric osteonecrosis. Rheumatology (Oxford). 2006;45:855–8. doi: 10.1093/rheumatology/kei013. PMID: 16436490.
- 22 Williams B, Unsworth I. Skeletal changes in divers. Aust Radiol. 1976;20:83–94. PMID: 962757.
- 23 Gempp E, Blatteau JE, Simon O, Stephant E. Musculoskeletal decompression sickness and risk of dysbaric osteonecrosis in recreational divers. Diving Hyperb Med. 2009;39:200–4. Available from: https://www.dhmjournal.com/images/Journals/39/DHM_Vol39_No4.pdf. [cited 2019 December 16].
- 24 Gempp E, Louge P, de Maistre S. Predictive factors of dysbaric osteonecrosis following musculoskeletal decompression sickness in recreational SCUBA divers. Joint Bone Spine. 2016;83:357–8. doi: 10.1016/j.jbspin.2015.03.010. PMID: 26454506.
- 25 Cimsit M, Ilgezdi S, Cimsit C, Uzun G. Dysbaric osteonecrosis in experienced dive masters and instructors. Aviat Space Environ Med. 2007;78:1150–4. doi: 10.3357/ase.2109.2007. PMID: 18064920.
- 26 Association of Diving Contractors International. International consensus standards for commercial diving and underwater operations. 6.2 ed. Houston TX: ADC International; 2016. Available from: <https://www.adc-int.org/files/C12181International%20Consensus%20Standards.pdf>. [cited 2020 March 26].
- 27 Joint Technical Committee SF-017, Occupational Diving. Australian/New Zealand Standard. Occupational diving operations Part 1: Standard operational practice AS/NZS 2299.1:2015. Purchasable from: [https://shop.standards.govt.nz/catalog/2299.1:2015\(AS%7CNZS\)/scope](https://shop.standards.govt.nz/catalog/2299.1:2015(AS%7CNZS)/scope). [cited 2020 March 27].
- 28 Doolette DJ, Mitchell SJ. Biophysical basis for inner ear decompression sickness. J Appl Physiol (1985). 2003;94:2145–50. doi: 10.1152/japplphysiol.01090.2002. PMID: 12562679.
- 29 Sarris I, Weiser R, Sotereanos DG. Pathogenesis and treatment of osteonecrosis of the shoulder. Orthop Clin North Am 2004;35:397–404. doi: 10.1016/j.ocl.2004.03.004. PMID: 15271548.
- 30 Harreld KL, Marulanda GA, Ulrich SD, Marker DR, Seyler TM, Mont MA. Small-diameter percutaneous decompression for osteonecrosis of the shoulder. Am J Orthop. 2009;38:348–54. PMID: 19714276.
- 31 Galloway MR, Horodyski M, Wright TW. Arthroscopically assisted fibular strut allograft for treatment of osteonecrosis of proximal humerus. J Surg Orthop Adv. 2013;22:277–82. doi: 10.3113/jsoa.2013.0277. PMID: 24393185.
- 32 Makihara T, Yoshioka T, Sugaya H, Yamazaki M, Mishima H. Autologous concentrated bone marrow grafting for the treatment of osteonecrosis of the humeral head: A report of five shoulders in four cases. Case Rep Orthop. 2017;2017:4898057. doi: 10.1155/2017/4898057. PMID: 28713606. PMCID: PMC5496114.
- 33 Hatstrup SJ, Cofield RH. Osteonecrosis of the humeral head: relationship of disease stage, extent, and cause to natural history. J Shoulder Elbow Surg. 1999;8:559–64. doi: 10.1016/s1058-2746(99)90089-7. PMID: 10633888.
- 34 Camporesi EM, Vezzani G, Bosco G, Mangar D, Bernasek TL. Hyperbaric oxygen therapy in femoral head necrosis. J Arthroplasty. 2010;25(6 Suppl):118–23. doi: 10.1016/j.arth.2010.05.005. PMID: 20637561.
- 35 Mills A. Medicolegal aspects of avascular necrosis in divers. SPUMS Journal. 1989;19:181–4.

Acknowledgements

We thank the patient for permission to report his professional and clinical histories and radiological images.

Conflicts of interest and funding: nil

Submitted: 11 February 2020

Accepted after revision: 09 May 2020

Copyright: This article is the copyright of the authors who grant *Diving and Hyperbaric Medicine* a non-exclusive licence to publish the article in electronic and other forms.