CORRIGENDUM

A neuronal enhancer network upstream of MEF2C is compromised in patients with Rett like characteristics

Eva D'haene^{1,†}, Reut Bar-Yaacov^{2,3}, Inbar Bariah^{2,3}, Lies Vantomme¹, Sien Van Loo¹, Francisco Avila Cobos^{1,4,5}, Karen Verboom^{1,4}, Reut Eshel^{2,3}, Rawan Alatawna^{2,3}, Björn Menten¹, Ramon Y Birnbaum^{2,3,*} and Sarah Vergult^{1,*}

¹Center for Medical Genetics, Ghent University, 9000 Ghent, Belgium ²Department of Life Sciences, Faculty of Natural Sciences, The Ben-Gurion University of the Negev, 84105 Beersheba, Israel ³Center of Evolutionary Genomics and Medicine, The Ben-Gurion University of the Negev, 84105 Beersheba, Israel ⁴Cancer Research Institute Ghent (CRIG), 9000 Ghent, Belgium ⁵Bioinformatics Institute Ghent from Nucleotides to Networks (BIG N2N), 9000 Ghent, Belgium

*To whom correspondence should be addressed. Ramon Y. Birnbaum, Tel: +97286479178; Email: ramonb@bgu.ac.il and Sarah Vergult, Tel: 3293325189; Email: Sarah.Vergult@ugent.be

Human Molecular Genetics, 2019, 28(5), 818–827.
doi: 10.1093/hmg/ddy393.

In the published article, Figure 1 did not include all 11 deletions (red bars) that were described in Table 1. This has now been corrected in both the PDF and HTML version of the article.

[†]Eva D'haene, <http://orcid.org/0000-0002-5936-8294>

[‡]Ramon Y Birnbaum, <http://orcid.org/0000-0002-3792-1095>

© The Author(s) 2020. Published by Oxford University Press. All rights reserved. For Permissions, please email: journals.permissions@oup.com