Safety and diagnostic accuracy of percutaneous CT-guided transthoracic biopsy of small lung nodules (≤ 20 mm) adjacent to the pericardium or great vessels

Chao Chen* Lichao Xu* Xiaofei Sun* Xiaoxia Liu* Zhi Han Wentao Li

PURPOSE

We aimed to evaluate the safety and diagnostic accuracy of computed tomography (CT)-guided transthoracic biopsy of small lung nodules (≤ 20 mm) adjacent to the pericardium or great vessels.

METHODS

This retrospective study examined the safety and diagnostic accuracy of percutaneous CT-guided biopsy for small lung nodules (≤ 20 mm) located within 10 mm of the pericardium or great vessels. Technical aspects and factors influencing complications were assessed, and diagnostic accuracy was calculated.

RESULTS

A total of 168 biopsies were performed in 168 patients. The complications were mainly pneumothorax (34.5%; 58 of 168 patients), chest tube insertion (5.3%; 9 of 168 patients), and pulmonary hemorrhage (61.3%; 103 of 168 procedures), with no patient mortality. One patient (0.6%) was admitted because of hemorrhage complications. Significant independent risk factors for pneumothorax were nodules resided in upper or middle lobes and lateral patient position, and for hemorrhage, longer distance from structures and longer needle trajectory through the lung parenchyma. Overall, the sensitivity, accuracy, and specificity were 91.0%, 92.2%, and 100%, respectively.

CONCLUSION

Percutaneous CT-guided transthoracic biopsy was highly accurate in small lung nodules (≤ 20 mm) adjacent to the pericardium or great vessels. Complications are common, but most were minor and self-limited.

In recent years, with wide availability of computed tomography (CT) and the development of low-dose CT screening techniques, more small lung nodules (≤ 20 mm) have been detected (1–3). Lee et al. (4) reported that small lung nodules were detected at a rate of 44.5%, and that 39.7% of lung cancers were found in small nodules. Accurate histopathological diagnosis is of great importance in management of these nodules, but biopsy via fluoroscopy or ultrasound guidance is technically more difficult, and CT guidance is usually required (5).

Percutaneous CT-guided transthoracic lung biopsy is a reasonably safe and accurate approach to the histological diagnosis of primary and secondary lung nodules (4, 6, 7). However, nodule size is the major determinant of diagnostic accuracy, which reached 96.9% for larger nodules (7), but ranged from 52% to 78.8% for small nodules (8, 9).

Biopsy of small lung nodules close to proximity to the pericardium or great vessels is particularly challenging, and precise needle placement is essential to avoid complications resulting from non-target tissue injury near vital mediastinal or vascular structures. Hsu et al. (10) suggested that CT-guided lung biopsy should be avoided if the adjacent nodule is within 10 mm of the pericardium.

The authors know of no related studies of biopsy of small lung nodules close to the pericardium or great vessels. This study, therefore, was conducted to examine the safety and diagnostic accuracy of CT-guided transthoracic biopsy of small lung nodules (≤ 20 mm) within 10 mm of the pericardium or great vessels.

From the Department of Interventional Radiology (C.C., L.X., X.S., W.L. ✉ wentaoli@fudan.edu.cn), and Department of Nursing (X.L.), Fudan University Shanghai Cancer Center, Shanghai, China; Department of Spine Surgery (Z.H.), Luoyang Orthopedic Hospital of Henan Province, Luoyang, Henan, China.

*Chao Chen, Lichao Xu, Xiaofei Sun, and Xiaoxia Liu contributed equally to this work.

Received 5 February 2020; revision requested 9 March 2020; last revision received 12 August 2020; accepted 24 August 2020.

Published online 16 October 2020.

DOI 10.5152/dir.2020.20051

You may cite this article as: Chen C, Xu L, Sun X, Liu X, Han Z, Li W. Safety and diagnostic accuracy of percutaneous CT-guided transthoracic biopsy of small lung nodules (≤ 20 mm) adjacent to the pericardium or great vessels. *Diagn Interv Radiol* 2021; 27: 94–101.

Figure 1. Flow diagram of patient management, interventions, and sample size for analyses.

Methods

Data were collected retrospectively from the institutional departmental database, and the study protocol and waiver of informed consent were approved by the Institutional Review Board (decision/protocol number 1904159-3-1905&1904159-3-1906). Patient management and intervention are summarized by the flow diagram shown in Fig. 1.

Main points

- Nodule in the upper or middle lobe was a significant risk factor for pneumothorax.
- Lateral position was a significant risk factor for pneumothorax.
- Longer transparenchymal needle trajectory increased the risk of hemorrhage.
- CT-guided biopsy was highly accurate for diagnosing small lung nodules near the pericardium or great vessels.

Patients

Between March, 2010, and March, 2018, 8217 patients underwent CT-guided transthoracic biopsy at our hospital. Patients were included if they had lung nodules measuring ≤ 20 mm in standard lung windows and adjacent to the pericardium or great vessels. Adjacent nodules were defined as index tumors seen at any distance within 10 mm of the pericardium or great vessels on axial CT images.

Biopsy procedures

CT-guided transthoracic lung biopsies were performed using a 64-slice spiral CT scanner (120 kV, 250 mA; Philips Healthcare) with 3 mm slice thickness by one of five interventional radiologists with more than 5-year experience.

Previous diagnostic images were reviewed to confirm the general location of the target lung nodule, and a planning CT scan was then performed to determine

the best patient position for the procedure: prone, supine, or lateral (biopsy side up; contralateral dependent), to optimize guided needle placement while ensuring the patient's safety and comfort. The most effective needle track was selected to approach the nodule, while avoiding crossing the adjacent large pulmonary vessels, visible airways, or fissures if possible. If feasible, a needle entry site as close as possible to the superior aspect of the rib was chosen to avoid intercostal vasculature. When possible, the needle track was kept parallel to the surface of the pericardium or great vessels, avoiding puncture of the pericardium or great vessel. The patients received local anesthesia, and the biopsies were performed using either a coaxial or non-coaxial technique, depending on a) whether or not immunohistochemical or molecular analysis was planned, and b) operator preference following risk/benefit analysis. For the coaxial technique, either a 17-gauge coaxial

Table 1. Final diagnoses			
Core needle biopsy histological results	Final diagnosis		Total
	Malignant	Benign	
Malignant	122	0	122
Benign	12	20	32
Total	134	20	154

needle with an 18-gauge semi-automated core biopsy needle or a 19-gauge coaxial needle with a 20-gauge semi-automated core biopsy needle was used (SuperCore; Argon Medical Devices). The needles were pushed into the peripheries of the lung nodules using a step-by-step technique under CT guidance. CT images were used to confirm the correct position of the needle tip. Great care was taken to puncture pleura only once during procedure. When the nodule was along the projected course of the needle, the procedure was continued. Otherwise, the needle was repositioned or the puncture site was changed. If the needle tip was located correctly within the target lesion, the biopsy was performed.

CT images were acquired immediately after the biopsy for checking any procedure-related complications. Patients were then positioned with the puncture site down and admitted for 4-hour continuous observation. The biopsies were always performed after 2:00 PM on Mondays to Thursdays, and all patients had erect anteroposterior chest radiographs 16 to 20 hours postbiopsy and were evaluated by the interventional radiologists. Our standard threshold for chest tube placement for pneumothorax was symptomatic pneumothorax, or at least 30% to 40% involvement of the hemithorax. Chest tubes were kept in place until air leakage stopped.

Data review

We recorded nodule size (0–15 mm, and 15.1–20 mm), nodule location (upper and middle lobes, lower lobes), pericardium vs. great vessel adjacent to the target nodule, and distance from pericardium or great vessel (0 mm, and 0.1–10 mm). Nodule size was determined by the maximum diameter measured on CT image.

Procedure data included position of patient (supine or prone vs. lateral), angle of pleural puncture, number of pleural punctures, and the length of needle trajectory through lung parenchyma (0–30 mm vs. >30 mm). Axial CT-images of needle placement

into the nodule were used to measure the angle of pleural puncture, that is, the lesser of the two angles between them, and to measure the length of the needle trajectory.

Complications were graded according to the Cardiovascular and Interventional Radiology Society of Europe (CIRSE) Classification System (11). Postprocedure pneumothorax was based on the immediate postbiopsy CT images or the chest X-ray at 16 to 20 hours postbiopsy. Pulmonary hemorrhage was defined by any new ground-glass opacity or consolidation on the immediate postbiopsy CT scan, and its severity was graded according to published methods (12, 13): 0, no hemorrhage; 1, hemorrhage along the needle tract less than 2 cm wide; 2, hemorrhage along the needle tract more than 2 cm wide but sublobar; 3, lobar hemorrhage or greater; or 4, hemothorax with hemodynamic instability.

We reviewed postbiopsy and post-resection pathologic reports and serial CT scans monitoring nodules over time. The CT-guided biopsy diagnoses of malignancy were classified as true positive (i.e., surgical confirmation of malignancy or same results of the known primary malignancy or postprocedural clinical evidence of malignancy) and false positive (i.e., no evidence of malignancy in surgical specimens). Postprocedural clinical evidence of malignancy was defined as fulfilling at least one of three criteria during the follow-up period: 1) increase in nodule size; 2) other proven metastases; 3) decrease in nodule size after radiotherapy, chemotherapy, or targeted therapy (14, 15). Benign CT-guided biopsy diagnoses were classified as true negative (i.e., surgical confirmation of benign nodule, subsequent disappearance or significant decrease in size of the nodule, or no change for at least 12 months postbiopsy) and false negative (i.e., subsequent proof of malignancy).

Statistical analysis

The data were grouped according to presence or absence of complications (pneumo-

thorax, pulmonary hemorrhage) after the biopsy procedure, and continuous and categorical variables were respectively compared between groups with and without complications by independent sample t test and Pearson chi-square test. Significant differences identified by univariate analysis were then analyzed with multivariate analysis by the use of logistic regression with forward selection. Based on the histopathologic diagnoses of the biopsy specimens with the final surgical, clinical, or follow-up radiologic findings, the sensitivity, specificity, and diagnostic accuracy of CT-guided biopsy for benign vs. malignant diagnoses were calculated by using 2×2 tables (Table 1). Statistical analysis using a commercially available statistical program was performed (IBM SPSS 20, IBM). Values of $p < 0.05$ were statistically significant.

Results

There were 168 biopsies in 168 patients (67 men, 101 women; mean age, 59±11 years). The characteristics of patients, lung nodules, and biopsy procedures are presented in Table 2. Cardiac chambers adjacent to the nodules were: left atrium (n=6; Fig. 2a), left ventricle (n=36; Fig. 2b), right atrium (n=26; Fig. 2c), and right ventricle (n=4; Fig. 2d). Great vessels adjacent to the nodules were: aorta (n=59; Fig. 3a), innominate artery (n=4; Fig. 3b), pulmonary artery shunt (n=12; Fig. 3c), subclavian artery (n=1; Fig. 3d), superior vena cava (n=13; Fig. 3e), inferior vena cava (n=2; Fig. 3f), innominate vein (n=1; Fig. 3g), and subclavian vein (n=4; Fig. 3h). No procedure-related mortality was recorded. One patient had CIRSE grade 2 pulmonary hemorrhage with hemoptysis (Fig. 4). The remaining complications, including pneumothorax in 58 cases and pulmonary hemorrhage in 102 cases, were categorized as grade 1.

The overall pneumothorax rate was 34.5% (58/168); pneumothorax was detected on the immediate postpuncture CT in 28.6% (10/58) and on chest X-ray the following morning in 6% (10/168) (Fig. 1). The placement of chest tube was required in 15.5% (9/58) of the pneumothorax cases, representing 5.4% (9/168) of the total study population. On univariate analysis, the significant risk factors for pneumothorax were nodule location ($p = 0.020$) and patient position ($p = 0.017$). Patient age, sex, nodule size, nodule adjacent to the pericardium or great vessels, distance from pericardium or great vessels to target nod-

Characteristics		Data
Age (years), mean±SD (range)		59±11 (15–79)
Sex, n (%)	Male	67 (39.9)
	Female	101 (60.1)
Nodule size, n (%)	0–15 mm	60 (35.7)
	15.1–20 mm	108 (64.3)
Nodule location, n (%)	Upper or middle lobe	117 (69.6)
	Lower lobe	51 (30.4)
Pericardium adjacent to the target nodule, n (%)	Left atrium	6 (8.3)
	Left ventricle	36 (50.0)
	Right atrium	26 (36.1)
	Right ventricle	4 (5.6)
Great vessel adjacent to the target nodule, n (%)	Aorta	59 (61.5)
	Innominate artery	4 (4.2)
	Pulmonary artery	12 (12.5)
	Subclavian artery	1 (1.0)
	Superior vena cava	13 (13.5)
	Inferior vena cava	2 (2.1)
	Innominate vein	1 (1.0)
	Subclavian vein	4 (4.2)
Distance from pericardium or great vessels, n (%)	0 mm	53 (31.5)
	0.1–10 mm	115 (68.5)
Patient position, n (%)	Supine and prone	145 (86.3)
	Lateral	23 (13.7)
Angle of pleural puncture, n (%)	≤50°	51 (30.4)
	>50°	117 (69.6)
Number of pleural punctures, n (%)	1	166 (98.8)
	2	2 (1.2)
Length of needle path through the lung parenchyma, n (%)	0–30 mm	63 (37.5)
	>30 mm	105 (62.5)

analysis of diagnostic performance included 154 of 168 patients (Fig. 1). Malignancy was finally diagnosed according to the gold standard in 134 patients. Among these, the diagnosis by CT-guided biopsy was true-positive in 122 patients and false-negative in 12 patients. Revisions in diagnosis of malignancy were based on surgical confirmation (n=13), histological findings that were similar to those of the known primary malignancy (n=46), and a postprocedural clinical course consistent with an obvious malignant process (n=75). Twenty nodules were identified as benign masses (true negative, n=20; false negative, n=0). Benign nodules were confirmed by surgical specimen (n=8), subsequent disappearance of the nodule (n=3), significant reduction in nodule size (n=7) in follow-up, or by stable status for more than 12 months postbiopsy (n=2). Overall, the sensitivity, accuracy, and specificity of CT-guided biopsy for diagnosis in small lung nodules were 91% (122 of 134), 92.2% (142 of 154), and 100% (20 of 20) (Table 1). The 95% confidence intervals were calculated as 0.846–0.951 for sensitivity and 0.800–1 for specificity.

Discussion

Percutaneous transthoracic needle biopsy with CT and CT-fluoroscopy guidance has proved to be an accurate and safe diagnostic procedure for small lung nodules (5, 16, 17). Beyond conventional CT, CT-fluoroscopy allows real-time visualization as the needle is advanced (18), and more recently, C-arm cone beam CT provides both real-time fluoroscopy and three-dimensional (3D) CT reconstruction images (19). However, both of these techniques impose significant radiation burden for patients and operators (20), and although navigation systems are now being incorporated to assist with image-guided percutaneous biopsy (21), those that have been tested *in vivo* thus far are few, and most of the trials have been conducted in small, heterogeneous cohorts (18, 19, 21). As such, even when real-time control is not available, conventional CT is still universally adopted in clinical practice for imaging-guided percutaneous needle biopsy of masses in lung.

The potential complications of CT-guided biopsy in close proximity to the pericardium or great vessels include cardiac tamponade, hemopericardium, pneumopericardium, right ventricular perforation, and great vessel laceration (22–24). These complications

ule, angle of pleural puncture, and length of needle trajectory through the lung parenchyma were not significantly different between groups (Table 3). Multivariate analysis revealed nodules residing in the upper or middle lobes ($p = 0.012$) and lateral position ($p = 0.011$) as significant independent risk factors for pneumothorax (Table 4).

Pulmonary hemorrhage occurred in 103 procedures (61.3%) among 168 patients (Fig. 1). Two (1.2%) of the 168 biopsies caused grade 3 hemorrhage, 24 (14.3%) caused grade 2 hemorrhage, and 77 (45.8%) caused grade 1 hemorrhage. Univariate analysis demonstrated that closer proximity to great vessels, longer distance (0.1–10 mm) from pericardium or great ves-

sels, and longer transparenchymal needle trajectory possessed significant risk of pulmonary hemorrhage ($p = 0.022$, $p = 0.027$, and $p < 0.001$, respectively), while patient age, sex, nodule size, nodule location, patient position, and angle of pleural puncture were not significant (Table 3). On multivariate analysis, longer distance (0.1–10 mm) from pericardium or great vessels and longer needle trajectory (>30 mm) through the lung parenchyma remained as significant independent risk factors for pulmonary hemorrhage ($p = 0.049$ and $p < 0.001$, respectively) (Table 5).

The biopsy results were considered nondiagnostic if the sample was not sufficient or if patient follow-up information was not available (n=14). Therefore, the statistical

are rare but potentially fatal and of primary concern to the interventional radiologist. It is important that a step-by-step technique

is used to monitor the relative position of the needle tip to the pericardium or great vessel. Once the correct needle trajectory

is confirmed, the needle is advanced incrementally until the tip lies within the nodule. If possible, the needle is positioned parallel to the surface of the pericardium or great vessels. Notwithstanding the recommendation that transcutaneous CT-guided biopsy should be avoided when the adjacent nodule is within 10 mm of the pericardium (10), there were no fatal complications recorded in our study of 168 patients with small lung nodules (≤ 20 mm) adjacent to the pericardium or great vessels.

In our cohort, 10 patients who did not have noticeable pneumothorax immediately postoperatively had pneumothorax detected by chest X-ray the following morning, and these accounted for 17% of 54 patients with pneumothorax. In other studies, chest X-rays have been obtained at 1 to 4 hours after biopsy (5, 25, 26). To our knowledge, no consensus exists regarding the most appropriate timing for post-biopsy chest X-rays. The overall rate of pneumothorax in our study was 34.5% and is well within the range of 12.8% to 62.0% reported in similar studies (3–5, 25–28). Nodule location in the upper or middle lobe is an

Figure 2. a–d. Representative CT images show lung nodules located within 10 mm of left atrium (a), left ventricle (b), right atrium (c), and right ventricle (d).

Table 3. Univariate analysis for risk factors associated with pneumothorax and pulmonary hemorrhage

Characteristics		Pneumothorax		<i>p</i>	Pulmonary hemorrhage		<i>p</i>
		Yes	No		Yes	No	
Age (years)	Mean $\pm$ SD	57 $\pm$ 10	58 $\pm$ 11	0.448 ^a	58 $\pm$ 11	57 $\pm$ 11	0.496 ^a
Sex, n (%)	Male	28 (41.8)	39 (58.2)	0.107 ^b	42 (62.7)	25 (37.3)	0.765 ^b
	Female	30 (29.7)	71 (70.3)		61 (60.4)	40 (39.6)	
Nodule size, n (%)	0–15 mm	20 (33.3)	40 (66.7)	0.809 ^b	37 (61.7)	23 (38.3)	0.944 ^b
	15.1–20 mm	38 (35.2)	70 (64.8)		66 (61.1)	42 (38.9)	
Nodule location, n (%)	Upper or middle lobe	47 (40.2)	70 (59.8)	0.020 ^b	71 (60.7)	46 (39.3)	0.801 ^b
	Lower lobe	11 (21.6)	40 (78.4)		32 (62.7)	19 (37.3)	
Nodule adjacent to, n (%)	Pericardium	27 (37.5)	45 (62.5)	0.482 ^b	37 (51.4)	35 (48.6)	0.022 ^b
	Great vessel	31 (32.3)	65 (67.7)		66 (68.8)	30 (31.2)	
Distance from pericardium or great vessels to target nodule, n (%)	0 mm	20 (37.7)	33 (62.3)	0.552 ^b	26 (49.1)	27 (50.9)	0.027 ^b
	0.1–10 mm	38 (33.0)	77 (67.0)		77 (67.0)	38 (33.0)	
Patient position, n (%)	Supine or prone	45 (31.0)	100 (69.0)	0.017 [†]	89 (61.4)	56 (38.6)	0.963 ^b
	Lateral	13 (56.5)	10 (43.5)		14 (60.9)	9 (39.1)	
Angle of pleural puncture, n (%)	$\leq 50^\circ$	14 (27.5)	37 (72.5)	0.286 ^b	29 (56.9)	22 (43.1)	0.435 ^b
	$> 50^\circ$	42 (35.9)	75 (64.1)		74 (63.2)	43 (36.8)	
Length of needle path through the lung parenchyma, n (%)	0–30 mm	18 (28.6)	45 (71.4)	0.209 ^b	25 (39.7)	38 (60.3)	$< 0.001^b$
	> 30 mm	40 (38.1)	65 (61.9)		78 (74.3)	27 (25.7)	

^aIndependent sample t test.

^bPearson chi-square test.

independent risk factor for pneumothorax, as previous data attest (5). Reported findings of the influence of nodule location on the pneumothorax rate are conflicting. In

Table 4. Multivariate analysis to identify independent risk factors for pneumothorax				
Characteristics	B coefficient	p	OR	95% CI
Nodule location	-1.016	0.012	0.362	0.164–0.801
Patient position	1.220	0.011	3.387	1.327–8.648

OR, odds ratio; CI, confidence interval.
p value for Hosmer-Lemeshow test was 0.738.

Table 5. Multivariate analysis to identify independent risk factors for pulmonary hemorrhage				
Characteristics	B coefficient	p	OR	95% CI
Nodule adjacent to pericardium or great vessels	0.593	0.089	1.809	0.914–3.578
Distance from pericardium or great vessels	0.724	0.049	2.063	1.003–4.240
Length of needle path through the lung parenchyma	1.476	<0.001	4.375	2.200–8.703

OR, odds ratio; CI, confidence interval.
p value for Hosmer-Lemeshow test was 0.913.

some studies, a nodule in the lower lobe increased the pneumothorax rate (4, 29, 30), whereas in others this was not an independent risk factor for pneumothorax (3, 8, 16, 31–33). The difference between our study and prior similar studies may be explained by the fact that a small nodule in upper and middle lobe is difficult to sample because of closely spaced ribs and the frequent proximity of such small nodules to the pericardium or great vessels. These conditions theoretically require more redirection and adjustment of the needle trajectory, causing longer operative time and higher pneumothorax rate.

The finding of a higher rate of pneumothorax in association with lateral dependent position does not contradict prior studies (25, 29). Rozenblit et al. (25) and Mills et al. (29) found that using the ipsilateral dependent position could reduce the rate of pneumothorax. For biopsies

Figure 3. a–h. Representative CT images show lung nodules located within 10 mm of aorta (a), innominate artery (b), pulmonary artery shunt (c), subclavian artery (d), superior vena cava (e), inferior vena cava (f), innominate vein (g), and subclavian vein (h).

Figure 4. a–c. CT-guided core biopsy of a small nodule in a 58-year-old woman. CT image (a) obtained before biopsy with patient in supine position shows a 17 mm lung nodule in the right lower lobe. The nodule is within 10 mm of the right atrium. CT image (b) obtained after needle placement shows the precise localization of the tip of the biopsy needle in the nodule. The length of needle trajectory through the lung parenchyma is 100 mm. CT image (c) obtained after needle removal shows grade 2 hemorrhage along the needle tract. The core biopsy diagnosis was adenocarcinoma, which was the final diagnosis after surgical resection.

performed in the contralateral dependent position, the patient is positioned with the nodule in the nondependent portion of the chest. Therefore, the relative hyperinflation and increased rib excursion of the nondependent lung may contribute to the development of pneumothorax (25).

The rate of pulmonary hemorrhage of 61.3% in our study is higher than the 30% to 42.4% range reported in other studies (12, 13, 34). This is probably because all of the CT-guided biopsies were performed in small lung nodules. Previous studies involving lung nodules ≤ 20 mm have registered higher rates of hemorrhage (12, 35, 36). Yeow et al. (12) had a rate of pulmonary hemorrhage of 66% for small (≤ 20 mm) lung nodules. The independent predictor of higher risk of pulmonary hemorrhage was a longer (> 30 mm) needle trajectory through the lung parenchyma, which compliments previous reports (36) maintaining that longer transparenchymal needle paths increase the risk of vessel puncture and hemorrhage. In addition, the pulmonary vessels enlarge when lung nodules are close to pericardium or great vessels, further increasing the incidence of major pulmonary hemorrhage. Nonetheless, the majority of the postprocedure pulmonary hemorrhages in our group were relatively minor and self-limited, requiring hospital admission in only one case (0.5%), similar to the result of Tai et al. (13), who had an admission rate of 0.4%.

It appears that the smaller the lung nodule, the more difficult inserting the needle

into the nodule using CT-guidance. Several studies reported decreased biopsy accuracy associated with decreasing nodule size (37, 38), whereas others have reported similar success rates in small and large nodules (35, 39). The diagnostic accuracy of biopsy of small lung nodules (≤ 20 mm) has been reported as 77.2%–98.2% (3, 8, 16, 38, 40), and the overall accuracy of diagnosis in our study was 92.2%. These variable results may be attributable to many study differences, including imaging guiding system, biopsy approach, nodule location, surgeon's skill, and patient characteristics.

Our present study had important limitations. First, the study was retrospective. A randomized controlled trial would clearly be preferable. Second, the definition of small lung nodules (≤ 20 mm) adjacent to the pericardium or great vessels was arbitrary. Third, the likely correlation between risk factors was not analyzed. Fourth, the 12-month follow-up period may be not sufficiently long, and there is still controversy about whether a 12- or 24-month follow-up period is optimal (3, 4, 40). For now, this study and others (3, 40) rely on a 12-month follow-up period for confirming the nature of nodules, while the 24-month follow-up period has prevailed elsewhere (4).

In conclusion, we found that percutaneous CT-guided transthoracic biopsy was highly accurate for the diagnosis of small lung nodules (≤ 20 mm) adjacent to the pericardium or great vessels. Complications were common, but mostly self-limited.

Financial disclosure

This work was supported by National Key Research and Development Program of China (No. 2016YFC0106203) and Program of Shanghai Hospital Development Center (No. SHDC22017102).

Conflict of interest disclosure

The authors declared no conflicts of interest.

References

1. Henschke CI. Early lung cancer action project: Overall design and findings from baseline screening. *Cancer* 2000; 89:2474–2482.
2. Diederich S, Wormanns D, Semik M, et al. Screening for early lung cancer with low-dose spiral CT: Prevalence in 817 asymptomatic smokers. *Radiology* 2002; 222:773–781. [\[Crossref\]](#)
3. Li Y, Du Y, Yang HF, Yu JH, Xu XX. CT-guided percutaneous core needle biopsy for small (≤ 20 mm) pulmonary lesions. *Clin Radiol* 2013; 68:e43–48. [\[Crossref\]](#)
4. Lee SM, Park CM, Lee KH, Bahn YE, Kim JI, Goo JM. C-arm cone-beam CT-guided percutaneous transthoracic needle biopsy of lung nodules: Clinical experience in 1108 patients. *Radiology* 2014; 271:291–300. [\[Crossref\]](#)
5. Wallace MJ, Krishnamurthy S, Broemeling LD, et al. CT-guided percutaneous fine-needle aspiration biopsy of small ($< \text{or} = 1\text{-cm}$) pulmonary lesions. *Radiology* 2002; 225:823–828. [\[Crossref\]](#)
6. Manhire A, Charig M, Clelland C, et al. Guidelines for radiologically guided lung biopsy. *Thorax* 2003; 58:920–936. [\[Crossref\]](#)
7. Li C, Liu B, Meng H, Lv W, Jia H. Efficacy and radiation exposure of ultra-low-dose chest CT at 100 kVp with tin filtration in CT-guided percutaneous core needle biopsy for small pulmonary lesions using a third-generation dual-source CT scanner. *J Vasc Interv Radiol* 2019; 30:95–102. [\[Crossref\]](#)
8. Ohno Y, Hatabu H, Takenaka D, et al. CT-guided transthoracic needle aspiration biopsy of small ($< \text{or} = 20$ mm) solitary pulmonary nodules. *AJR Am J Roentgenol* 2003; 180:1665–1669. [\[Crossref\]](#)

9. Ng YL, Patsios D, Roberts H, et al. CT-guided percutaneous fine-needle aspiration biopsy of pulmonary nodules measuring 10 mm or less. *Clin Radiol* 2008; 63:272–277. [\[Crossref\]](#)
10. Hsu LS, Chen CW, Chang CH, Liao CH, Hsu SL, Lin WM. Influences of cardiac motion on computed tomography-guided biopsy of lung nodules located near the heart. *Medicine* 2017; 96:e8558. [\[Crossref\]](#)
11. Filippiadis DK, Binkert C, Pellerin O, Hoffmann RT, Krajina A, Pereira PL. Cirs quality assurance document and standards for classification of complications: The cirse classification system. *Cardiovasc Intervent Radiol* 2017; 40:1141–1146. [\[Crossref\]](#)
12. Yeow KM, Su IH, Pan KT, et al. Risk factors of pneumothorax and bleeding: Multivariate analysis of 660 CT-guided coaxial cutting needle lung biopsies. *Chest* 2004; 126:748–754. [\[Crossref\]](#)
13. Tai R, Dunne RM, Trotman-Dickenson B, et al. Frequency and severity of pulmonary hemorrhage in patients undergoing percutaneous CT-guided transthoracic lung biopsy: Single-institution experience of 1175 cases. *Radiology* 2016; 279:287–296. [\[Crossref\]](#)
14. Yang W, Sun W, Li Q, et al. Diagnostic accuracy of CT-guided transthoracic needle biopsy for solitary pulmonary nodules. *PloS one* 2015; 10:e0131373. [\[Crossref\]](#)
15. Lee KH, Lim KY, Suh YJ, et al. Nondiagnostic percutaneous transthoracic needle biopsy of lung lesions: A multicenter study of malignancy risk. *Radiology* 2019; 290:814–823. [\[Crossref\]](#)
16. Choi JW, Park CM, Goo JM, et al. C-arm cone-beam CT-guided percutaneous transthoracic needle biopsy of small (≤ 20 mm) lung nodules: Diagnostic accuracy and complications in 161 patients. *AJR Am J Roentgenol* 2012; 199:W322–330. [\[Crossref\]](#)
17. Prosch H, Stadler A, Schilling M, et al. CT fluoroscopy-guided vs. Multislice CT biopsy mode-guided lung biopsies: Accuracy, complications and radiation dose. *Eur J Radiol* 2012; 81:1029–1033. [\[Crossref\]](#)
18. Santos RS, Gupta A, Ebright MI, et al. Electromagnetic navigation to aid radiofrequency ablation and biopsy of lung tumors. *Ann Thorac Surg* 2010; 89:265–268. [\[Crossref\]](#)
19. Choo JY, Park CM, Lee NK, Lee SM, Lee HJ, Goo JM. Percutaneous transthoracic needle biopsy of small (≤ 1 cm) lung nodules under c-arm cone-beam CT virtual navigation guidance. *Eur Radiol* 2013; 23:712–719. [\[Crossref\]](#)
20. Jin KN, Park CM, Goo JM, et al. Initial experience of percutaneous transthoracic needle biopsy of lung nodules using c-arm cone-beam CT systems. *Eur Radiol* 2010; 20:2108–2115. [\[Crossref\]](#)
21. Grasso RF, Faiella E, Luppi G, et al. Percutaneous lung biopsy: Comparison between an augmented reality CT navigation system and standard CT-guided technique. *Int J Comput Assist Radiol Surg* 2013; 8:837–848. [\[Crossref\]](#)
22. Kucharczyk W, Weisbrod GL, Cooper JD, Todd T. Cardiac tamponade as a complication of thin-needle aspiration lung biopsy. *Chest* 1982; 82:120–121. [\[Crossref\]](#)
23. Shevlan JE. Right ventricular perforation: A rare complication of percutaneous lung biopsy. *J Thorac Imaging* 1991; 6:85–86. [\[Crossref\]](#)
24. Mitchell MJ, Montgomery M, Reiter CG, Culp WC, Jr. Pericardial tamponade following CT-guided lung biopsy. *Cardiovasc Intervent Radiol* 2008; 31 Suppl 2:S227–230. [\[Crossref\]](#)
25. Rozenblit AM, Tuvia J, Rozenblit GN, Klink A. CT-guided transthoracic needle biopsy using an ipsilateral dependent position. *AJR Am J Roentgenol* 2000; 174:1759–1764. [\[Crossref\]](#)
26. Leger T, Jerjir N, Gregory J, et al. Does ipsilateral-dependent positioning during percutaneous lung biopsy decrease the risk of pneumothorax. *AJR Am J Roentgenol* 2019; 212:461–466. [\[Crossref\]](#)
27. Cheung YC, Chang JW, Hsieh JJ, Lin G, Tsai YH. Adequacy and complications of computed tomography-guided core needle biopsy on non-small cell lung cancers for epidermal growth factor receptor mutations demonstration: 18-gauge or 20-gauge biopsy needle. *Lung Cancer* 2010; 67:166–169. [\[Crossref\]](#)
28. Ozturk K, Soyulu E, Gokalp G, Topal U. Risk factors of pneumothorax and chest tube placement after computed tomography-guided core needle biopsy of lung lesions: A single-centre experience with 822 biopsies. *Pol J Radiol* 2018; 83:e407–e414. [\[Crossref\]](#)
29. Mills M, Choi J, El-Haddad G, et al. Retrospective analysis of technical success rate and procedure-related complications of 867 percutaneous CT-guided needle biopsies of lung lesions. *Clin Radiol* 2017; 72:1038–1046. [\[Crossref\]](#)
30. Hiraki T, Mimura H, Gobara H, et al. Incidence of and risk factors for pneumothorax and chest tube placement after CT fluoroscopy-guided percutaneous lung biopsy: Retrospective analysis of the procedures conducted over a 9-year period. *AJR Am J Roentgenol* 2010; 194:809–814. [\[Crossref\]](#)
31. Ko JP, Shepard JO, Drucker EA, et al. Factors influencing pneumothorax rate at lung biopsy: Are dwell time and angle of pleural puncture contributing factors? *Radiology* 2001; 218:491–496. [\[Crossref\]](#)
32. Rui Y, Han M, Zhou W, et al. Non-malignant pathological results on transthoracic CT guided core-needle biopsy: When is benign really benign? *Clin Radiol* 2018; 73:757 e751–757 e757. [\[Crossref\]](#)
33. Saji H, Nakamura H, Tsuchida T, et al. The incidence and the risk of pneumothorax and chest tube placement after percutaneous CT-guided lung biopsy: The angle of the needle trajectory is a novel predictor. *Chest* 2002; 121:1521–1526. [\[Crossref\]](#)
34. Anzidei M, Sacconi B, Fraioli F, et al. Development of a prediction model and risk score for procedure-related complications in patients undergoing percutaneous computed tomography-guided lung biopsy. *Eur J Cardiothorac Surg* 2015; 48:e1–6. [\[Crossref\]](#)
35. Laurent F, Latrabe V, Vergier B, Montaudon M, Vernejoux JM, Dubrez J. CT-guided transthoracic needle biopsy of pulmonary nodules smaller than 20 mm: Results with an automated 20-gauge coaxial cutting needle. *Clin Radiol* 2000; 55:281–287. [\[Crossref\]](#)
36. Nour-Eldin NE, Alsubhi M, Naguib NN, et al. Risk factor analysis of pulmonary hemorrhage complicating CT-guided lung biopsy in coaxial and non-coaxial core biopsy techniques in 650 patients. *Eur J Radiol* 2014; 83:1945–1952. [\[Crossref\]](#)
37. Li H, Boisselle PM, Shepard JO, Trotman-Dickenson B, McLoud TC. Diagnostic accuracy and safety of CT-guided percutaneous needle aspiration biopsy of the lung: Comparison of small and large pulmonary nodules. *AJR Am J Roentgenol* 1996; 167:105–109. [\[Crossref\]](#)
38. Shimizu K, Ikeda N, Tsuboi M, Hirano T, Kato H. Percutaneous CT-guided fine needle aspiration for lung cancer smaller than 2 cm and revealed by ground-glass opacity at CT. *Lung Cancer* 2006; 51:173–179. [\[Crossref\]](#)
39. Westcott JL, Rao N, Colley DP. Transthoracic needle biopsy of small pulmonary nodules. *Radiology* 1997; 202:97–103. [\[Crossref\]](#)
40. Gupta S, Krishnamurthy S, Broemeling LD, et al. Small (≤ 2 -cm) subpleural pulmonary lesions: Short- versus long-needle-path CT-guided biopsy—comparison of diagnostic yields and complications. *Radiology* 2005; 234:631–637. [\[Crossref\]](#)