Published in final edited form as:

Nat Rev Gastroenterol Hepatol. 2020 November ; 17(11): 655–672. doi:10.1038/s41575-020-0341-5.

Brain–gut–microbiome interactions in obesity and food addiction

Arpana Gupta^{1,2,3}, Vadim Osadchiy^{1,2}, Emeran A. Mayer^{1,2,3,4}

¹G. Oppenheimer Family Center for Neurobiology of Stress and Resilience, Ingestive Behavior and Obesity Program, University of California Los Angeles (UCLA)

²David Geffen School of Medicine, University of California Los Angeles (UCLA)

³Vatche and Tamar Manoukian Division of Digestive Diseases, University of California Los Angeles (UCLA)

⁴Ahmanson-Lovelace Brain Mapping Center at University of California Los Angeles (UCLA)

Abstract

Normal eating behavior is coordinated by the tightly regulated balance between intestinal and extra-intestinal homeostatic and hedonic mechanisms. By contrast, food addiction represents a complex, maladaptive eating behavior that reflects alterations in brain–gut–microbiome (BGM) interactions and a shift of this balance towards hedonic mechanisms. Each component of the BGM axis has been implicated in the development of food addiction, with both brain to gut and gut to brain signaling playing a role. Early life influences can prime the infant gut microbiome and brain for food addiction, which might be further reinforced by increased antibiotic usage and dietary patterns throughout adulthood. The ubiquitous availability and marketing of inexpensive, highly palatable and calorie dense food can further shift this balance towards hedonic eating through both central (disruptions in dopaminergic signaling) and intestinal (vagal afferent function, metabolic toxemia, systemic immune activation, changes to gut microbiome and metabolome) mechanisms. In this Review, we propose a systems biological model of BGM interactions, which incorporates published reports on food addiction, and provides novel insights into treatment targets aimed at each level of the BGM axis.

Introduction

The obesity epidemic continues to be a major public health problem both in the USA and globally.^{1,2} Obesity is defined as a body mass index (BMI) ≥ 30 kg/m² and a BMI ≥ 40 kg/m² is considered extreme obesity, with overweight classified as 25–29.9 kg/m².^{1,2} The prevalence of obesity worldwide has tripled since 1975, with ~39% of the world's adult population being overweight and 13% being obese in 2016.² In the USA alone, the number of individuals with obesity continues to dramatically increase, with >35% of individuals

* emayer@ucla.edu.

Competing interests

E.A.M. serves on the scientific advisory boards of Amare, Axial Biotherapeutics, Bloom Science, Danone, Viome, UBiome, Pendulum, Mahana Therapeutics and APC Microbiome Ireland. A.G. and V.O. declare no competing interests.

being overweight, >37% obese and 8% morbidly obese.^{2,3} Obesity is the biggest driver of preventable chronic diseases and healthcare costs in the USA, with current cost estimates ranging from \$147–210 billion per year.⁴ Despite the magnitude of the problem and the associated healthcare costs, drug development efforts have largely failed and proposed treatments have had disappointing outcomes, with only modest reductions and frequent weight regain after successful weight loss.^{4,5}

Obesity has a complex and multifactorial aetiology and the limited progress in obesity treatments can in large part be attributed to the failure to apply a systems biology-based approach to understand its pathophysiology and to develop individualized strategies to achieve sustained weight loss and prevention.^{6,7} A growing body of largely preclinical studies support the concept of bidirectional signaling within the brain–gut–microbiome (BGM) axis in the pathophysiology of obesity, mediated by metabolic, endocrine, neural and immune system mediated mechanisms.⁸ Signaling from the brain through the autonomic nervous system (ANS) and the hypothalamic–pituitary–adrenal (HPA) axis influences many gastrointestinal processes, including motility and transit,⁹ fluid and mucus secretion,⁹ immune activation, intestinal permeability,¹⁰ relative gut microbial abundances,¹¹ as well as gene expression patterns in certain pathogenic gut microorganisms.¹² Changes in the gut luminal environment can affect gut microbial community composition and function.^{13,14} Conversely, the gut microbiota can communicate with the brain via hundreds of metabolites,^{15–17} which are sensed by specialized cells in the gut, including enteroendocrine cells, enterochromaffin cells and primary or secondary afferent nerve endings. Sensing of bacterial metabolites by these cells results in neural signaling to the brain, interactions with gut-based immune cells leading to local and systemic immune activation, or the metabolite might achieve sufficient concentrations in the circulation to directly access brain circuits by crossing the blood–brain barrier.¹⁸ Short-chain fatty acids (SCFAs), the main byproduct of microbial fermentation of dietary fiber, have emerged as key mediators of BGM signaling.¹⁹ These saturated fatty acids can influence the central nervous system (CNS) through immune-,²⁰ endocrine-,²¹ and vagal- pathways.²²

The gut microbiome and bidirectional BGM interactions are programmed through influences during pregnancy and the first 1,000 days of life,²³ and are subject to multiple perturbations from within the body (including from metabolism, gut microbiota interactions and energy expenditure) and from the environment (for example, via food, stress and medications) throughout life (Figure. 1). Perturbations at any level of the BGM system, resulting in compromised inhibitory mechanisms that normally regulate food intake, can bias ingestive behaviors towards predominantly hedonic-driven eating behaviours, cravings and overeating.^{24–27} An extensive literature exists on the homeostatic regulation of food intake and maintenance of body weight via interactions between hypothalamic nuclei and orexogenic and anorexogenic gut hormones, in addition to chemical signals derived from adipose tissue, in particular leptin.^{28–30} However, it is ultimately the complex balance between gut-derived orexogenic (ghrelin, insulin) and anorexogenic signals (including cholecystokinin, neuropeptide Y (NPY) and glucagon-like peptide-1 (GLP1)), gut microbial metabolites (SCFAs and amino acid metabolites), stress mediators (corticotropin releasing factor (CRF)), and the motivational drive generated by the central reward system

(dopaminergic reward system) and prefrontal cortical inhibitory mechanisms that determine how much we eat.³¹⁻³³

A particular type of eating behavior which has been referred to as food addiction, plays an important part in the pathophysiology of obesity.^{34,35} Food addiction is the continued consumption of highly palatable foods even after energy requirements have been met and despite known negative physical and psychological consequences in response to uncontrolled food ingestion.^{34,36} Similar to other forms of substance abuse, food addiction represents an addiction-like response to food (especially foods rich in sugar and fat) or the process of eating itself in susceptible individuals.^{37,38}

Since it was first proposed by T. Randolph in 1956,³⁹ there has been an ongoing controversy over the term of food addiction,^{40,41} despite strong arguments supporting shared underlying pathophysiology between drug and food addiction.^{33,42} On a behavioral level, individuals with food addiction as identified by the Yale Food Addiction Scale (YFAS)⁴³ meet the diagnostic criteria for substance abuse disorder found in the Diagnostic and Statistical Manual of Mental Disorders⁴⁴, which involves loss of control over eating, excessive time or focus on food, neglect of other activities and continuation of the behavior despite known negative consequences.⁴⁵⁻⁴⁷ An increasing number of research reports on the biological alterations in the extended reward network in both humans and rodents also point towards strong similarities in the mechanisms underlying substance use disorders and food addiction, including substantial commonalities between the neural substrates underlying the substance abuse and at least some forms of obesity that involve food addiction.³¹⁻³³ For example, the biological similarities between individuals with obesity with food addition and individuals with drug addiction include, but are not limited to, changes in the dopaminergic pathways within the reward system, and in cortical performance monitoring, both of which are involved in processes associated with reward sensitivity, motivation, interoceptive awareness, stress reactivity and self-control.⁴⁸⁻⁵⁰ Despite these similarities, there are also clear differences. Although the development of predominantly hedonic driven eating behaviors involves food-induced alterations in multiple peripheral and central mechanisms of the BGM axis, drug addiction results from a direct effect of the drug on the brain.^{51,52} In addition, in contrast to the nearly universal development of addiction upon exposure to a drug, food addiction as assessed by the YFAS score is present in only a subgroup of individuals with obesity.^{46,53} On the basis of questionnaire-based surveys and by other methods of assessment, food addiction is present in 25–37% of individuals with obesity, reaching rates of up to 60% in those who are morbidly obese or in patients who undergo bariatric surgery.^{36,38,54-60} Food addiction is also highly associated with eating disorders such as bulimia nervosa and binge eating disorder.^{61,62}

Previous work on obesity and food addiction crosses over multiple fields of research: neuroscience, gastroenterology, microbiology, endocrinology, immunology and many others. For example, the gut microbiome, intestinal signaling, extraintestinal signaling (visual, olfactory, food memories), early life programming of food preferences and many other factors can contribute to food addiction. Here, we review and build on past work to create a systems-based BGM model of obesity and food addiction. Systems biology is an interdisciplinary field of study that focuses on complex interactions within multiple

biological systems, rather than focusing on individual mechanisms. One of the aims of systems biology is to model and discover emergent properties of cells, tissues and organisms functioning as a system rather than as individual parts. We believe that such an interdisciplinary, system-based approach is able to create a more nuanced understanding of food addiction, as shown in Figure 1. In this Review, we summarize the physiology of food addiction in obesity as it relates to alterations within the brain and the gut microbiome. We will introduce key factors that influence the BGM axis such as diet, antibiotics, early life adversity, food cues and psychological stress, during the prenatal and postnatal period, and during adulthood. This Review also discusses several therapies aimed at food addiction in individuals with obesity, including those targeted at the gut, the microbiome and the brain, and highlights limitations and areas for future research in the field.

Ingestive behaviour physiology

The role of the gut microbiome

Ingestive behavior represents a delicate balance between homeostatic and hedonic regulatory mechanisms in the CNS, orchestrated by a number of gut peptides, neuronal impulses, endocrine signals and countless other influences, including signals generated by the gut microbiota (Figure 1).

BGM interactions involving gut peptides that regulate ingestive behavior have been the most extensively studied. The gastric hormone ghrelin has an important role in producing hunger and craving,^{63,64} perhaps through amplification of dopaminergic signaling mechanisms,⁶⁵ whereas the intestinal hormones GLP1⁶⁶ and peptide YY⁶⁷ trigger satiety and associated behavioral changes. Increased production of microbiota-derived SCFAs can stimulate enteroendocrine cells to release GLP1⁶⁸ and peptide YY⁶⁹, while decreasing the secretion of ghrelin.⁷⁰

Insulin is another orexogenic hormone, as hyperinsulinaemia, regardless of plasma glucose levels, contributes to increased sensations of hunger and results in a heightened palatability of sucrose.⁷¹ Animal evidence suggests that disruptions of microbial SCFA metabolism can promote insulin resistance and hyperinsulinaemia,⁷² thereby potentially shifting the balance towards hedonic behaviors. For example, studies in mice have shown that increased production of acetate by an altered gut microbiota can lead to activation of the parasympathetic nervous system which in turn promotes increased glucose-stimulated insulin secretion and increased ghrelin secretion resulting in hyperphagia.⁷³

Additionally, gut microbiota-derived secondary bile acids can regulate insulin sensitivity through signaling involving the nuclear farnesoid X receptor (FXR) and the G protein-coupled receptor TGR5 (also known as G-protein coupled bile acid receptor 1 GBAR1).⁷⁴ Activation of intestinal FXR in a mouse model induces microbial production of the secondary bile acid lithocholic acid, driving TGR5 signaling and triggering GLP1 secretion from enteroendocrine L-cells.⁷⁵ Exposure to oral broad-spectrum antibiotics (combination of ampicillin, vancomycin, neomycin sulfate, and metronidazole) successfully inhibited microbial lithocholic acid production and completely reversed improvements in insulin sensitivity.⁷⁵ In a single blind randomized controlled trial of 20 individuals with obesity,

administration of an oral antibiotic (vancomycin) for just 1 week resulted in reduced microbial diversity (mainly affecting Firmicutes), with an associated drop in secondary bile acids as well as in insulin sensitivity.⁷⁶

An example of microbial regulation of food preferences has been shown in a study demonstrating that fruit flies fed a diet deficient in essential amino acids show preferential intake of amino acid-rich foods.⁷⁷ These preferences are, however, blunted by the presence of both *Acetobacter pomorum* and lactobacilli.⁷⁷ Of note, neither *Acetobacter pomorum* nor lactobacilli were capable of modulating food intake individually, suggesting that these microorganisms must work together to influence host behavior.⁷⁷ Although the mechanisms driving food seeking behaviors in this model remain unclear, microbial modulation of neuronal TOR signaling has been previously proposed as an important mediator of nutrient balance and growth in *Drosophila*.^{78,79} In fruit flies exposed to a nutrient scarce environment, *Lactobacillus plantarum* can promote protein assimilation from the diet, resulting in increased production of branched-chain amino acids.⁷⁹ These amino acids activate central nervous system (CNS) TOR kinase activity, resulting in the release of insulin-like peptides.⁷⁹

The role of the brain

Neuroimaging studies have improved our understanding of the role of the brain in ingestive behavior in both animals and more recently in humans, especially the interplay of various brain networks involved in homeostatic mechanisms versus food addiction (non-homeostatic).^{50,80-82} The homeostatic component of food intake is comprised of hormonal regulators of hunger, satiety and adiposity levels.^{83,84} The hypothalamus is the primary brain area within the homeostatic system that regulates food ingestion and energy balance, and hence is often referred to as the 'satiety centre' or 'feeding centre' of the brain.⁸⁵⁻⁸⁷

Normal ingestive behavior is under the control of the extended reward network, which includes brain regions from the core reward network such as the nucleus accumbens, ventral tagmental area or substantia nigra, and are regulated by cognitive network regions such as the prefrontal cortex.^{88,89} The extended reward network is involved in the processing of rewarding stimuli and modulation of food-seeking behavior,^{90,91} inhibitory control,⁹² cognitive performance monitoring,^{93,94} interoceptive and sensory awareness,^{88,95,96} and integrating salient information to make decisions regarding food intake.⁹⁷⁻¹⁰⁰ This processing includes brain regions concerned with interconnecting brain networks such as reward, salience, emotional regulation, the somatosensory system and cortical inhibition (prefrontal control) networks (Figure 2).^{31,33,90,99} The salience network is responsible for monitoring the homeostatic state of the body to make adaptive adjustments to real or expected disturbances in homeostasis through autonomic nervous system, as well as behavioral responses.^{98,101} In food addiction (as in substance abuse), the saliency of a specific type of reward (food or drug) becomes greater at the expense of other rewards.^{32,48} The emotional regulation network is activated by stimuli threatening the homeostasis of the organism, and provides a rapid feedback inhibition of such activation via its connections with the salience network.^{89,102,103} Advanced analytical techniques such as brain network metrics based on graph theory, which measure the underlying architecture and flow of

communication between brain regions and networks, and multivariate machine learning methods that predict obesity have been applied to phenotype hedonic ingestion-related brain signatures, with a focus on alterations in the extended reward network.^{88,89,104-107}

Homeostatic versus hedonic systems

Homeostatic system.—The hypothalamus acts as a hub integrating information from the external environment, such as food availability and stress, and from the internal milieu of the host to meet real or perceived nutrient needs.^{108,109} Lesions in the hypothalamus, in both humans and animals can lead to increases in appetite, food ingestion and weight gain.^{110,111} Numerous studies have been directed at identifying the molecular mechanisms within the hypothalamus underlying these processes.^{85,108,112} For example, animal models have demonstrated that brief electrical stimulation of nuclei within the hypothalamus can cause the increased expression of genes related to Agouti-related protein (AGRP)–NPY– γ -aminobutyric acid (GABA) signaling, which in turn can cause voracious food ingestion.^{113,114} These targeted cells have, therefore, been referred to as ‘hunger neurons’.^{86,115} The hypothalamus has close interactions with corticolimbic and medullary pontine regions integrating sensory information mediated by vagal afferents, affective state and cognitive modulation to generate appropriate motor responses and adaptive eating behaviors.⁸⁵

Hedonic system.—Individuals with obesity and with food addiction are more likely than individuals who are lean or obese without food addiction to display a heightened motivation to eat highly palatable foods, consume more calories from fat and protein, and have at least 20% prevalence rates with comorbid conditions, such as depression, binge eating, and decreased quality of life functioning, beyond that observed with obesity alone.¹¹⁶⁻¹¹⁹ The closely regulated balance between hedonic and homeostatic aspects of ingestive behavior can be altered when normal inhibitory regulation of the reward system is compromised via decreased modulation or connectivity, resulting in overconsumption of food. There are similarities between food addiction and other addictive behaviors, as both reflect an imbalance in responses within the brain’s extended reward system to stimuli from the environment.^{33,120} In food addiction, such uncoupling can be the result of central as well as peripheral disturbances in brain–gut interactions, including diet-induced neuroplastic changes in the sensitivity of vagal afferent nerve terminals and of hypothalamic nuclei to satiety hormones,¹²¹⁻¹²³ emotional state, and easy access to highly processed and palatable foods that all modify the rewarding properties of food, thereby leading to overconsumption.^{32,48} Both human and animal studies have shown that increased cravings and food addiction behaviors result in increased conditioning and motivation to seek these highly palatable foods, and these behaviors are based on alterations of regions within the extended reward network.^{31,33,124,125}

Evidence also exists suggesting that, similar to other addiction like behaviors, food addiction is associated with a reduced response of the reward network.¹²⁶ For example, ingestion of rewarding foods or drugs leads to reduced dopamine signaling within the reward system in both individuals with obesity and those with drug addiction,¹²⁷ suggesting that a reduction in dopamine signaling (by both the reduced release of dopamine and the downregulation of dopamine receptors) might contribute to the overconsumption and increased cravings of the

drug of choice.¹²⁸⁻¹³⁰ According to the dopamine deficiency hypothesis, reduced dopamine release in the striatum alters corticostriatal communication between the basal ganglia (core reward) and the extended reward system, resulting in compromised inhibition of connectivity in the reward regions.¹³¹ Reduced cortical inhibition of reward regions is also associated with greater cravings and reduced disinhibition scores.¹³²⁻¹³⁴ According to this theory, hypo-dopaminergic function also leads to reduced levels of subjective well-being, as it is linked to dysregulation of other neurotransmitters such as 5-HT, enkephalins and GABA.¹³¹ To compensate for this dopamine deficiency, it has been suggested that affected individuals will engage in behaviors that stimulate the brain's production and utilization of dopamine, such as by the overconsumption of highly palatable and rewarding foods,^{131,135,136} increasing the risk of developing food addiction and obesity.^{137,138} Thus, a stronger stimulus (for example, increased food intake) is required to overcome the reduced responsiveness of the dopamine system, similar to mechanisms identified in disorders of addiction, and failure to achieve this goal is associated with food cravings and the engagement of the stress response.^{31,32,48,139,140} Stress induced eating usually depends on a number of factors such as the length of the stress, type of stressor, type of foods available, especially if calorie dense and highly palatable, length of time exposed to the food, and satiety and hunger levels.¹²⁴ Studies have shown that during stress, increased cortisol levels could contribute to increased gluconeogenesis, an upregulation of corticotrophin releasing factor (CRF) in the amygdala and other limbic regions, and consequently the blunting of HPA axis function, which in turn leads to low dopamine and reward functioning commonly associated with food addiction.¹²⁵ Stress has also been associated with increases in ghrelin and cortisol and related increases in craving and intake of highly palatable foods, which were higher in those with obesity and overweight than lean individuals.¹⁴¹ A study performed in obese rats demonstrates downregulation of striatal dopamine D2 receptors compared to lean rats, similar to what has been shown in previous studies of humans addicted to drugs.^{142,143} Furthermore, D2 receptor knock down mice rapidly develop compulsive-like food seeking behaviors when high-fat food is readily available.¹⁴²

Even though dopamine has been the most thoroughly investigated signaling system in addictive behaviors, several neurotransmitters other than dopamine and neuropeptides are involved both in the homeostatic regulation of food intake (including orexin, leptin and ghrelin, corticotrophin releasing factor) and have been implicated in the rewarding effects of food, cannabinoids, opioids and serotonin.⁴⁸ Moreover, neurons in ventral tegmental area and/or nucleus accumbens express GLP1, ghrelin, leptin, insulin, orexin and melanocortin receptors, suggesting that these hormones or peptides can influence the rewarding responses to food.⁴⁸ Rats that are fed a diet high in saturated fats and refined sugars for two months demonstrate reduced hippocampal brain-derived neurotrophic factor, with a concomitant reduction in synaptic plasticity.¹⁴⁴

Food addiction pathological mechanisms

The first 1,000 days of life represent a crucial developmental period for the gut-associated immune system and the BGM axis.^{23,145,146} Preclinical models of myelination and brain development suggest that the early life microbiome regulates myelination of the prefrontal cortex¹⁴⁷ and facilitates proper striatal synapse function.¹⁴⁸ Additionally, commensal

microorganisms might also have a role in programming the HPA axis for stress responses,¹⁴⁹ a system implicated in obesity and food addiction behavior,¹⁵⁰⁻¹⁵³ whereby excess cortisol and related steroids, such as those from a disrupted of HPA axis, can drive adipogenesis¹⁵⁴ and increase food cravings.¹⁵² Early life exposure to different microorganisms, antibiotics, dietary factors and stress shape the relative abundances and richness of the gut microbiota, influence immune system and brain development, modulate microbial communication with the CNS and program maladaptive BGM interactions.^{8,155,156} Although these interactions and their roles in the development of obesity have been well described (Figure 3),^{8,155,156} their links with maladaptive eating behaviors is incompletely understood.

Prenatal developmental influences

Maternal prenatal factors have been shown to influence development of the infant BGM axis, with evidence suggesting an important role of the prenatal maternal diet in influencing the neonate gut microbiome. For example, greater maternal consumption of dairy during pregnancy was positively associated with a greater relative abundance of members of the genus *Clostridium* in the faecal microbiome of 145 infants, adjusted for maternal BMI, feeding method and parity.¹⁵⁷ Similarly, a maternal high-fat diet was associated with depletion of the genus *Bacteroides* in the neonatal gut microbiome, which persisted through 4–6 weeks of age.¹⁵⁸ These changes might be mediated indirectly by maternal dietary influences on the composition of breast milk, or directly from effects of the maternal diet on the fetal gut microbiome.^{157,158,159} Maternal psychosocial stress has been implicated in the development of an obese phenotype. For example, severe maternal stress due to bereavement in the prenatal period was associated with increased BMI in the male adult offspring in a study of 120,000 men, regardless of trimester that the bereavement occurred in.¹⁶⁰ In mice, moderate maternal stress during pregnancy was found to influence postnatal brain development and gene expression in the paraventricular nucleus of the hypothalamus, the hypothalamic regulator of the HPA axis, ultimately resulting in deficits of neuroplasticity and central stress responsiveness. This effect was partially mediated by stress-induced alterations in the maternal vaginal microbiome.¹⁶¹ As maternal antibiotic use during the second or third trimester is associated with an increased offspring risk of obesity regardless of pre-gravid BMI in a study of 436 mother–child dyads,¹⁶² and the vaginal microbiota play an important part in shaping the infant microbiome, as demonstrated in a very well-characterized group of 9 mothers and their 10 newborns,¹⁶³ it is possible that changes in maternal vaginal microbial abundance associated with antibiotic exposure or diet during pregnancy might also increase the risk of obesity in the offspring.

Postnatal influences

Early diet.—The infant gut microbiota is sensitive to early life nutrition. Human breast milk contains >200 different human milk oligosaccharides (HMOs), a type of prebiotic that cannot be degraded by gut glycoside hydrolases or absorbed via intestinal membrane transporters, suggesting that their primary target is the infant's gut microbiota.¹⁶⁴ The limited small intestinal bioavailability of HMOs enables efficient delivery to the developing infant gut microbiota, most notably *Bifidobacterium*, which can degrade these sugars.¹⁶⁴ Exclusively breastfed infants show a more diverse *Bifidobacterium* microbiota (175 faecal samples from 7 infants) compared with exclusively formula-fed infants (154 faecal samples

from 7 infants), which might set the stage for future, beneficial BGM interactions¹⁶⁵ given that a robust *Bifidobacterium* community has been shown to be protective against intestinal infections¹⁶⁶ and associated with appropriate development of the infant immune system.¹⁶⁷ Limited data exist on the effect of combination feeding (breastmilk and formula) on the gut microbiome, despite how common it is.¹⁶⁸ Bogen et al.,¹⁶⁹ in an observational study of over 70,000 infants, suggest that a longer duration of partial breastfeeding is necessary to yield a similar protective effect against obesity compared with exclusive breastfeeding; combination feeding for >26 weeks yielded a similar protective effect (odds ratio (OR) of developing obesity: 0.70; 95% CI: 0.61–0.81) as exclusive breastfeeding for 16–26 weeks (OR: 0.71; 95% CI: 0.56–0.92).

The relative abundances of the infant gut microbiota and its associated microbial transcriptome change substantially once solid food is incorporated at around 9 months of age, including increased abundances of Bacteroidetes and elevated SCFA levels, as suggested by a high-quality, 2.5-year case study of 60 faecal samples from a single infant¹⁷⁰. However, evidence has shown that the faecal microbiota profile assessed at 3 months of age (composed of primarily Bacteroidaceae, Bifidobacteriaceae, Enterobacteriaceae, Lachnospiraceae, Ruminococcaceae, and Veillonellaceae) is a more reliable predictor of future risk of being overweight compared to querying the microbiota profile at 12 months, as suggested by a study of 1,087 infants¹⁷¹. These findings are supported by a meta-analysis of >200,000 participants that found that breastfeeding was associated with a statistically significant reduced risk of obesity in children (pooled adjusted OR: 0.78; 95% CI: 0.74–0.81), with a subset of studies even revealing a dose–response relationship between breastfeeding duration and a reduction in obesity risk.¹⁷²

Highly processed foods, which are filled with large amounts of salt, sugar, fat and additives have become increasingly available in the developed world.^{173,174} A pattern of increased exposure and ingestion of such foods in childhood might program food preferences and increase the risk of the development of food addiction into adulthood.¹⁷⁵ Additionally, the relentless marketing of such foods, starting in childhood, has contributed to the increased uncontrollable consumption and cravings of unhealthy foods, especially in children.^{176,177}

Antibiotics.—An analysis of outpatient antibiotic prescription rates in 2010 found that >70% of prescriptions in the USA were written for antibiotics, with the highest prescribing rates for children under 10 years of age and with an average of three doses of antibiotics by the age of 2 years.¹⁷⁸ In a USA cohort study of 333,353 children, antibiotic prescriptions were significantly associated with a diagnosis of childhood obesity (HR: 1.26; 95% CI: 1.23–1.28).¹⁷⁹ In a longitudinal study of 39 healthy children, the gut microbiota of antibiotic-treated children was found to be less diverse at multiple phylogenetic levels, with some species even dominated by a single strain. However, the gut microbiome, largely appeared to return to baseline within 1 month of antibiotic exposure.¹⁸⁰ In a Danish study examining over 28,000 mother–child dyads, early administration of antibiotics – within the first 6 months of life – was associated with an increased risk of being overweight at a 7-year follow-up in children from normal weight, but not overweight, mothers; the gut microbiota of study participants were not examined in this study.¹⁸¹ Though it is difficult to draw definitive conclusions from natural history and cross sectional epidemiological studies,

numerous preclinical studies¹⁸²⁻¹⁸⁵ also support a negative effect of early life antibiotics on energy metabolism, the immune system and obesity. Mice that received a single dose of low-dose penicillin at birth showed enhanced high-fat diet-induced obesity as adults; this phenotype could be successfully transferred to germ-free mice by the penicillin-selected microbiota.¹⁸² Another mouse study showed that a single early-life macrolide antibiotic course resulted in persistent perturbations to the gut microbiome (increased *Akkermansia muciniphila* attributed to reductions in competitor mucin-degrading bacteria) and the immune system (reduction in small intestinal CD4⁺ IL-17A⁺ lymphocytes, decreased intestinal IgA secretion).¹⁸⁶ Collectively, these studies suggest that the antibiotic-altered microbiota, and not the antibiotic itself, has a causal role in driving obesogenic metabolic and immunological changes in mice. It remains to be determined if and how the antibiotic induced microbiome alterations can influence the brain and alter ingestive behaviors.

Early adversity.—A history of early adverse life events (EALs) such as natural disaster, parents divorcing, emotional or physical abuse, sickness or death of a family member, predisposes individuals to develop obesity and food addiction in adulthood, mainly through mechanisms associated with stress, inflammation, emotional perturbations, maladaptive coping responses and metabolic disturbances.^{187,188} Studies including a meta-analysis have demonstrated that trauma and abuse during the developmental period is significantly associated with greater odds of adulthood obesity and substance abuse (OR: 1.34, 95% CI: 1.24–1.45, $P < .001$).¹⁸⁹⁻¹⁹¹ Preclinical models of early adversity (maternal separation model) have observed that addictive behaviors can develop later in life, but these animal models require further investigation in humans, especially as the underlying mechanisms are unknown and these animal studies are poor models of human behaviour.¹⁹² For example, rats exposed to limited nesting stress in the post-natal period had an immature HPA axis, which was associated with reduced gut microbiota diversity, with an especially notable reduction in bacteria capable of degrading fiber.¹⁹³ Although the causal relationship between adversity during childhood and adult obesity is incompletely understood, it has been suggested that overconsumption of highly palatable foods even when hunger and satiation have been met are a possible coping mechanisms to deal with the increased stress responsiveness seen in individuals with a history of EALs.¹⁹⁴⁻¹⁹⁷ In a study of 186 men and women comparing healthy individuals with those with obesity, a history of EALs was associated with alterations in resting state functional connectivity, shown using MRI of brain regions in the extended reward network.¹⁹⁸ These EAL-related alterations probably contributed to an increased probability of developing food addiction and obesity later in life. In a network analysis, sex-differences were also noted in the interactions between early life adversity, brain connectivity and food addiction, suggesting that the development of food addiction might be driven by different factors in men versus women.¹⁹⁸ For example, compared to men, women show increased post-prandial activations in the brain's reward regions, which might increase susceptibility to cravings for highly palatable foods and result in hyperphagia.^{199,200} Although the exact molecular mechanisms by which women, compared to men, experience this increased susceptibility remain unclear, results from a pilot study of 63 individuals with varying BMI levels suggest that EALs might contribute to the development of food addiction by interfering with interactions between the brain and gut

microbiome, specifically microbial tryptophan metabolism and reward brain regions such as the amygdala, anterior insula and nucleus accumbens.²⁰¹

Adult environmental influences.

Several environmental factors might contribute to the pathophysiology of obesity by influencing BGM interactions in the adult. Below we discuss some of these environmental factors in detail.

Diet.—Cheap, highly processed and easily accessible high caloric and palatable foods are abundant in the developed world.²⁰² Studies have shown that foods that are enhanced for taste and salience not only increase cravings and ingestion of these foods, but contextual factors such as stress can serve as conditional cues for future food intake and long-term weight gain.^{125,203,204} In fact, overconsumption of highly palatable foods, in particular those containing high levels of fat and sugar, reduces the rewarding thresholds of such foods when ingested, in relationship to reduced levels of dopamine and dopamine receptors in the brain, therefore requiring an increased intake of such foods to generate the same satisfaction.^{205,206} Although long-term ingestion of such highly palatable foods has been shown to alter gut microbial diversity and relative abundances in humans, it is important to note that the adult microbiome is relatively resistant to short-term changes in diet, as suggested by a high-quality study of 98 individuals.^{207,208}

Food cues.—Studies have shown that portion sizes are directly related to a compromised ability to control food intake, an important feature of food addiction.²⁰⁹ Food labels and plate and utensil sizes can moderate the portion control effect by increasing food intake.²¹⁰ Although the exact mechanisms are unknown, these development and marketing driven food-related cues, which are ubiquitous in Western media and marketing, influence individuals with obesity to consume a greater number of calories than lean individuals.^{211,212,213,214} Individuals with obesity have also reported an increased preference and craving for foods rich in fat and sugar, in a study of over 46,000 adults.²¹⁵

Psychosocial stress.—Psychosocial stress can also stimulate ingestive behavior in the adult by increasing appetite, cravings and motivation to consume highly palatable foods, thereby contributing to stress-related weight gain in obesity.^{125,216,217} In individuals with obesity, strong associations have been shown between perceived stress and food addiction, snacking, cravings and abnormal eating patterns.^{124,218} For example, a study of 339 adults (mean BMI = 26.7 ± 5.4 kg/m²) demonstrated that chronic stress can influence levels of the orexogenic hormone insulin, as well as glucose and cortisol responses, which in turn can lead to increased food intake and weight gain.²¹⁹ Paradoxically, although the ingestion of ‘comfort foods’ high in fat and sugar can reduce subjectively perceived stress, various studies have shown that ingestion of such foods can also lead to increased autonomic responses, disrupt the HPA axis and increase cortisol and ghrelin levels, which have been associated with increased cravings and ingestive behaviors.^{132,220-222}

In a mouse model, chronic psychosocial stress resulted in a global reduction in gut microbiota richness and diversity, including a reduction of the relative abundance of

*Akkermansia*²²³, which has been suggested to have beneficial effects within the context of human obesity and metabolic syndrome by previous investigations.²²⁴ These stress-induced perturbations were also associated with changes in the functional profile of the gut microbiome, with decreased synthesis and metabolism of SCFAs, tryptophan and tyrosine.²²³ These changes might have been mediated, at least in part, by alterations in immunoregulatory signals, as these mice showed transient elevations in number of IL-10⁺ T regulatory (T_{reg}) cells, which were suppressed over time.²²³

Amino acid metabolites.—Tryptophan and its metabolites – serotonin (5-HT), kynurenine and indole – have been implicated as important mediators of BGM interactions within the context of obesity and food addiction.^{225,226} 5-HT, due in part to its diverse roles as a neurotransmitter in both the gastrointestinal tract (that is, in processes such as peristalsis, secretion and absorption) and the CNS (that is, in regulation of pain modulation, sleep and mood), is the most extensively studied of these metabolites.²²⁷ 95% of the body's 5-HT is stored in gastrointestinal enterochromaffin cells (ECCs) and the gut microbiota, through the production of SCFAs and secondary bile acids, can regulate 5-HT synthesis and its release from ECCs.²²⁸⁻²³⁰ Human studies incorporating acute tryptophan depletion (typically involving providing individuals with a large protein load containing non-tryptophan large neutral amino acids to saturate amino acid blood-brain barrier transporters, thereby limiting the transport of endogenous tryptophan), a validated method to transiently reduce peripheral and central 5-HT synthesis, underscore the effect of changes in 5-HT release on food preference.²³¹ In a study of 55 women following acute tryptophan depletion, participants with overweight showed a statistically significant increase in sweet calorie intake and preference for sweet foods compared with a placebo intervention.²³² By contrast, the lean group showed no differences, suggesting that individuals with overweight might be more susceptible to changes in tryptophan metabolism and 5-HT availability.²³² Host and microorganisms participate in different aspects of tryptophan metabolism: although host cells have the major role in the kynurenine pathway, microbial cells are primarily involved in the indole pathway.²³³

Although 5-HT has been the most extensively studied tryptophan metabolite, the majority of tryptophan is converted by host cells to kynurenine.²³⁴ In the gastrointestinal tract, kynurenine is synthesized from tryptophan by the rate limiting enzyme indoleamine-2,3-dioxygenase (IDO), which can be upregulated by inflammatory cytokines or downregulated by reactive oxygen species, such as hydrogen peroxide produced by intestinal *Lactobacillus*.^{235,236} As both kynurenine and tryptophan compete to cross the blood–brain barrier through the same, easily saturated transporter, inflammation-associated or microbiota-associated changes in peripheral kynurenine concentrations might also influence central 5-HT levels.²³⁷ Alternatively, increased flux through the kynurenine pathway can influence the brain through neuroactive downstream metabolites such as kynurenate and quinolinate, which function as an N-methyl-D-aspartate (NMDA) antagonist and an NMDA excitotoxin or neurotoxin, respectively.²³⁵ The balance of tryptophan metabolism might be preferentially shunted towards the kynurenine pathway in individuals with obesity, as serum kynurenine, kynurenate and quinolinate levels show positive associations with BMI.²³⁸

Another important group of tryptophan metabolites are the indoles. Most undigested dietary tryptophan in the gut lumen is converted exclusively by gut microorganisms to indole.²³⁹ In animal and human studies, indole has been shown to play an important part in modulating kynurenine synthesis⁷, strengthening the mucosal intestinal barrier,¹⁴ attenuating CNS inflammation,¹⁵ and modulating GLP1 secretion,²⁴⁰ all of which have been shown to be disrupted in states of obesity.^{238,241-243} One study investigating the role of stool indole metabolites in 63 healthy individuals found positive associations between indole, skatole and indoleacetic acid and food addiction behaviors, with regions of the extended reward network (nucleus accumbens, amygdala, and anterior insula) playing an important part in this interaction.²⁰¹

Metabolic toxemia.—In mouse models of obesity, a diet high in fat (60% lard) and low in dietary fibre has been implicated in the longterm disruptions in gut microbiota diversity,²⁴⁴ while diets high in fibre result in positive alterations in ingestive behavior (decreased food intake, increased satiety)²⁴⁵. When dietary fiber is reduced or unavailable, certain gut microorganisms such as *Akkermansia mucinophilia* consume the glycans making up mucins in the mucus layer of the gut, thereby compromising intestinal barrier function.²⁴⁶ This phenomenon is referred to as increased ‘leakiness’ of the gut (Figure 4). Sonnenburg et al.²⁴⁴ showed in mice that a low fibre diet results in a substantial loss of microbiota diversity and abundance, which was magnified in each successive generation, up to the fourth and final generation studied.²⁴⁴ Remarkably, supplementation with a high fibre diet alone was insufficient to normalize microbial diversity.²⁴⁴ Reduced intestinal barrier function results in increased access of membrane bound lipopolysaccharide (LPS) from Gram-positive microorganisms to TLR4 receptors on host epithelial and immune cells, contributing to inappropriate immune activation.²⁴⁷⁻²⁵⁰

The combination of a leaky gut and an overabundance of inflammatory bacterial products is thought to result in elevated plasma levels of LPS and proinflammatory cytokines, including IL-1 β , IL-6 and TNF.²⁵¹ Increased systemic immune activation can shift tryptophan metabolism towards the kynurenine pathway and away from 5-HT or indole synthesis, as described previously.²⁵² This state of metabolic endotoxaemia has been shown to reduce central satiety mechanisms by influencing enteroendocrine secretion of the satiety hormones PYY, cholecystokinin and 5-HT²⁵³⁻²⁵⁵ and by reducing the expression of anorexigenic peptide receptors and leptin receptors on vagal afferents²⁵⁶ and in the hypothalamus, respectively.²⁵⁷ In this way, vagal afferent neurons in the presence of a high-fat diet remain in an orexigenic state, regardless of whether food was consumed, driving hyperphagia and obesity.²⁵⁸ In addition to changes in ingestive behavior, there are likely numerous other mechanisms contributing to high-fat diet-induced obesity, such as gut microbiota-driven remodeling of the intestinal transcriptome to favor an obesogenic signaling cascade.²⁵⁹

Although the gut microbiota has an important role in the generation of inflammatory mediators, it might also be protective against the development of metabolic endotoxaemia. For example, mice fed a high-fat diet that was supplemented with oligofructose, a prebiotic fiber that preferentially increases gut *Bifidobacterium* abundance, showed reduced endotoxaemia, decreased levels of plasma and adipose tissue proinflammatory cytokines, and improved glucose tolerance.²⁶⁰ In this study, no relationship was seen between

endotoxaemia and any other bacterial group except for *Bifidobacterium*.²⁶⁰ The translatability of these preclinical findings to human metabolic disease remains to be determined.

In summary, disruptions during both early life (prenatal influences, including maternal diet, antibiotic exposure and early adversity) and adulthood (diet and/or psychosocial stressors) can have a profound effect on the gut microbiome and the brain, setting the stage for food addiction. The associated changes in amino acid metabolism and metabolic toxaemia perpetuate these maladaptive changes at all levels of the BGM axis (Figure 1).

Clinical implications of food addiction

The proposed system biological model of altered BGM interactions resulting in maladaptive changes in ingestive behavior provides not only a plausible explanation for the refractory nature of obesity to many traditional therapeutic strategies, but also presents a rationale for several new therapeutic strategies (Figure 5, Box 1).

Gut-directed therapies

As the gut is the primary source of hunger and satiety signals that regulate homeostatic feeding behaviors, it is not surprising that several obesity treatments, including bariatric surgery, have aimed to modify these gut mechanisms. Most of the bariatric procedures result in satisfactory and sustained weight loss and prompt resolution of the metabolic syndrome, a substantial improvement over the transient and more modest effects seen with medical therapies.^{261,262} Bariatric surgery-related weight loss is multifactorial, with the most common procedures, Roux-en-Y gastric bypass (RYGB) and laparoscopic sleeve gastrectomy (LSG), resulting in hypophagia. This hypophagia is not only a consequence of a reduced gastric capacity but also of marked reductions in appetite, food preferences and food addiction,²⁶³⁻²⁶⁵ which predicted long-term weight loss outcomes.²⁶⁶⁻²⁶⁸ Evidence also exists for the role of bariatric surgery-induced weight loss driving remission of food addiction (as assessed by YFAS); one study of 14 patients with obesity and with food addiction prior to surgery demonstrated remission of food addiction in 93% following surgery-induced weight loss ($P<0.001$).²⁶⁹ The mechanisms behind the post-bariatric surgery reductions in food addiction scores and brain responses to highly palatable food cues, as suggested by the aforementioned small, pilot studies, are incompletely understood.^{263,265} Both, RYGB and LSG result in increases in blood levels of anorexigenic gut peptides (GLP1, PYY) that, in part, mediate changes in appetite and food addiction after bariatric surgery.^{263,270-272} However, several other BGM pathways have also been implicated from both preclinical (mouse models) and clinical studies as possible explanations for these changes, including enhanced microbial production of polyamines and GABA, changes in bile acid profiles and FXR pathway signaling, and increased production of SCFAs.²⁷³⁻²⁷⁷ Preliminary work in individuals with obesity undergoing bariatric surgery suggest that changes in gut microbiome composition and microbial metabolism of aromatic amino acids and glutamate were associated with reductions in appetite, food addiction and changes in food preferences, suggesting a possible causal role of these metabolites in behavioral responses.²⁷⁸⁻²⁸³ However, future studies are needed to confirm causality between gut

microbial metabolites and food addiction in humans. Another important consideration is the well-known reduction in systemic low-grade inflammation and endotoxaemia seen after bariatric surgery.²⁸⁴ As discussed earlier, this anti-inflammatory effect could increase hypothalamic sensitivity to satiety signals and to insulin, resulting in a shift towards more homeostatic regulation of ingestive behaviors.²⁸⁵⁻²⁹⁰ Some studies have suggested that because of the involvement of the brain's reward system in both food addiction and addiction of other substances, bariatric surgery might increase alcohol use.²⁹¹

Microbiome-directed therapies

Microbiome-directed therapies, including faecal microbiota transplantation (FMT), represent a novel therapeutic option for obesity and metabolic syndrome. Small clinical studies (one study had 18 individuals, 9 receiving autologous FMT and 9 receiving allogenic FMT from a lean donor, and a second study had 38 individuals, 12 receiving autologous FMT, 26 receiving allogenic FMT from a lean donor) have shown that FMT from a lean donor resulted in an increase of butyrate-producing bacteria and improved insulin sensitivity in recipients with metabolic syndrome.^{292,293} The improved insulin sensitivity and associated changes in the faecal microbiome were not sustained at 18-week follow-up, suggesting a resilient faecal core microbiome.²⁹³ It remains unclear if FMT, when combined with lifestyle modification and brain-directed therapies, might result in longer-term success. Notably, lower recipient baseline faecal microbiota diversity was predictive of success of FMT.²⁹³ Ingestive behaviors were not assessed in those studies. It is well known that microbial products such as SCFAs modulate feeding behavior via central mechanisms.¹⁹ For example, the intake of a type of fiber that selectively increases gut microbial propionate production was associated with lowering the subjective appeal and the brain reward activation (as measured by brain MRI) to highly-palatable-food pictures in a study of 20 healthy men without obesity.²¹ It is important to note, however, that FMT is not without risks, including the rare but well-documented risks of bacteraemia or sepsis, ileus, perforation and aspiration, in addition to the more common, transient gastrointestinal complaints such as abdominal pain or changes in bowel habit.²⁹⁴

Time restricted eating

Increased interest exists in the potential health benefits of different types of time-restricted feeding (TRF) on obesity, cardiovascular health and ageing, including intermittent feeding and the fasting mimicking diet.²⁹⁵⁻²⁹⁷ In mice, ad libitum exposure to a high-fat diet resulted in changes in circadian rhythms and feeding behaviors that led to increased energy intake and weight gain.²⁹⁸ These changes could be reversed by TRF.²⁹⁹ One study showed that individuals with overweight, but otherwise healthy, adhering to TRF with the assistance of a smartphone application significantly reduced their daily-energy intake, in part by reducing late-night intake of alcohol and snacks. This behavioral change resulted in sustained weight loss up to 1 year after the intervention.³⁰⁰ The role that the microbiome plays in mediating the effects of TRF in humans is not known. However, in animal and in humans, gut microbiota composition and function display oscillations during the day that are associated with the circadian rhythm of the host and are dictated by the host's food intake patterns.³⁰¹ These oscillations in microbial metabolite production might have a major effect on the circadian epigenetic and transcriptional landscape of the gut and the liver, and eating

behaviors leading to compromised oscillations have been associated with obesity and metabolic syndrome.³⁰² There are other potential factors that might contribute to the benefits of TRF, including reduction of intake of snacks and calories, and changes in the gut microbial environment due to an increase in fasting associated patterns of motility and secretion.

Brain-directed therapies

Based on the premise that obesity is the result of an imbalance between energy intake and expenditure, for many years the dominant pharmacological approach to obesity was based on molecules that decrease appetite and/or stimulate energy expenditure.³⁰³⁻³⁰⁵ Of the medications now available for weight loss in the USA, some are aimed to decrease appetite by directly affecting the hypothalamus (phentermine, bupropion, naltrexone, or lorcaserin). Others are aimed to modulate reward circuits in the brain (naltrexone, bupropion, or topiramate), reducing the subjective pleasantness of palatable foods and compulsive food cravings, as well as decreasing the response to food cues at reward regions in the brain.³⁰⁶⁻³⁰⁹ A dual effect, reducing appetite and reward-based eating, is achieved through the use of hormonal satiety signals, like GLP1 agonists (liraglutide or exenatide).^{278,310} Very little is known about the effect of these anti-obesity medications on the gut microbiota. A small study in 19 individuals with type 2 diabetes mellitus have shown that treatment with liraglutide for 42 days resulted in a statistically significant increase in relative abundance of the genus *Akkermansia*.³¹¹

Cognitive behavioral therapy (CBT) in individuals with obesity with food addiction aims to change specific thoughts, beliefs, and cognition directly related to the feelings and behaviors attributed to uncontrollable ingestive behaviors and cravings.^{312,313} Since CBT strengthens prefrontal control mechanisms,³¹⁴⁻³¹⁹ cognitive reappraisal and attention strategies through CBT are thought to strengthen the inhibitory control of the prefrontal regions on the extended reward networks by influencing appetitive motivation and reducing food addiction in individuals with obesity.³²⁰⁻³²³

Key open research questions and future directions

Considerable progress has been made in our understanding of changes in BGM interactions in food addiction and obesity; yet, the majority of studies have been performed in rodents and there are few longitudinal, mechanistic studies in humans that support the translational relevance of these findings, which would guide more effective therapies. There is currently no evidence in humans that food addiction is the result of an altered gut microbiome, or that food addiction is driven by particular gut microbial metabolites. Furthermore, given the complexity and bidirectional signaling within the BGM axis, it is unlikely that a single microbial metabolite could causally explain the behavioral changes. Considering the influence of early life experiences, environmental factors, stress, emotions, genetic factors, and dietary influences in humans, microbial influences might only explain a small component of the variance in the development of food addiction. Several approaches are necessary to move this field beyond the current reliance on largely associative studies in small populations. Microbiome characterization by shotgun metagenomics,

metatranscriptomics and proteomics studies in well phenotyped human populations combined with big data analysis will be required to identify a microbial signature of food addiction. At the same time, mechanistic studies using targeted interventions, which have been shown to be effective in reducing predominantly hedonic driven eating behaviors in humans, such as bariatric surgery, TRF or cognitive behavioral therapy are needed to probe for a causal role of the gut microbiome. Such human studies should be combined with reverse translational studies, evaluating the effect of faecal microbial transfer on rodent feeding behaviors and body weight.

Conclusions

Altered BGM interactions manifesting as dysregulated eating behavior and resulting in obesity, can best be understood as a complex, circular system that is stable and highly resistant to change (Figure 4). The close interactions between diet and gut microbial signals, the effect of these signals on satiety and inflammatory mediators from the gut, and their disruptive effect on homeostatic mechanisms in the brain, leads to a shift towards a greater influence of hedonic reward mechanisms and a reduction in inhibitory control. These changes in turn drive the preferred intake of high caloric foods reinforcing the gut dysbiosis (Figure 4). As traditional therapies aimed at individual aspects of this system, including most traditional dieting strategies, have failed, novel therapies must be based on a new understanding of the systems properties of BGM interactions (Figure 1). A combination of therapeutic approaches targeting different nodes of this system, and individualization of treatments based on differences in gut microbial composition and function are required to provide greater clinical benefits.

Acknowledgements

We would like to acknowledge Claudia P. Sanmiguel for her contributions in making editorial suggestions to the gut-directed therapies section of this review to Cathy Liu for invaluable editorial services.

Acknowledgements/Funding Support:

EA Mayer has been supported by grants from the National Institute of Diabetes and Digestive and Kidney Diseases (DK048351, DK064539 and DK096606). AG has been supported by the following grants: DK106528 (NIDDK), ULTR001881/DK041301 (CURE/CTSI)

Glossary

Systems biology:

An interdisciplinary field of study that focuses on complex interactions within multiple biological systems, rather than focusing on individual mechanisms.

Hedonic-driven eating behaviors:

Also known as ‘food addiction’, these behaviours correspond to the continued consumption of highly palatable foods even after energy requirements have been met.

Dopaminergic reward system:

Refers to the extensive network of neurons in the extended reward network that depend on dopamine as the primary neurotransmitter for reward-related processing.

Extended reward network:

Used interchangeably with 'greater reward system', this term describes brain regions concerned with interconnecting brain networks such as reward and salience networks, and is associated with processing of rewarding stimuli and modulation of food-seeking behaviors.

Neural substrates:

A brain region or network that is associated with a specific behavior.

Cortical performance monitoring:

Refers to processes associated with reward sensitivity, motivation, interoceptive awareness, stress reactivity and self-control.

Salience:

The salience brain network is responsible for monitoring the homeostatic state of the body to make adaptive adjustments to real or expected disturbances in homeostasis through the autonomic nervous system and behavioral responses.

Corticostriatal communication:

Refers to the extensive communication network between the brain's cortex, which houses the extended reward network (including the frontal cortex and insula), and the striatum, which houses the core reward network (nucleus accumbens, basal ganglia).

Ventral tegmental area:

Key region of the midbrain that houses the dopaminergic cell bodies that project to all regions of the core and extended reward network.

Nucleus accumbens:

Region of the basal ganglia and a key hub for the core reward system, responsible for many dopaminergic processes, especially those related to pleasure, motivation, and aversion.

Prebiotic:

Refers to dietary fibre or other substrates that can only be digested by commensal gut microorganisms, thereby promoting gut microbiota diversity and health.

Maladaptive coping:

Behaviors that are used to cope with situations to alleviate stress or symptoms, but are not necessarily healthy and do not address the core cause of the stress.

Psychological stress:

Sufficient levels of stress originating from the environment that can cause dysregulation of homeostatic responses to cause physical or psychological symptoms.

Perceived stress:

A measure of the degree to which events in an individual's life are assessed as stressful. The most widely used scale for perceived stress is the Perceived Stress Scale.

References

1. CDC, C. f. D. C. a. P. Overweight and Obesity, <<http://www.cdc.gov/obesity/data/adult.html>> (2014).
2. <http://www.who.int/mediacentre/factsheets/fs311/en/>. Obesity and overweight. Fact sheet, (2016).
3. <http://stateofobesity.org/rates/>. Obesity Rates & Trends, (2016).
4. Biener A, Cawley J & Meyerhoefer C The High and Rising Costs of Obesity to the US Health Care System. *Journal of General Internal Medicine* 32, S6–S8, doi:10.1007/s11606-016-3968-8 (2017).
5. Mancini MC & de Melo ME The burden of obesity in the current world and the new treatments available: focus on liraglutide 3.0 mg. *Diabetol Metab Syndr* 9, 44, doi:10.1186/s13098-017-0242-0 (2017). [PubMed: 28580018]
6. Zhang Y et al. Obesity: pathophysiology and intervention. *Nutrients* 6, 5153–5183, doi:10.3390/nu6115153 (2014). [PubMed: 25412152]
7. Heymsfield SB & Wadden TA Mechanisms, Pathophysiology, and Management of Obesity. *N Engl J Med* 376, 254–266, doi:10.1056/NEJMra1514009 (2017). [PubMed: 28099824]
8. Osadchiy V, Martin CR & Mayer EA The Gut-Brain Axis and the Microbiome: Mechanisms and Clinical Implications. *Clin Gastroenterol Hepatol* 17, 322–332, doi:10.1016/j.cgh.2018.10.002 (2019). [PubMed: 30292888]
9. Mayer EA et al. Functional GI disorders: from animal models to drug development. *Gut* 57, 384–404, doi:10.1136/gut.2006.101675 [pii] 10.1136/gut.2006.101675 (2008). [PubMed: 17965064]
10. Keita AV & Soderholm JD The intestinal barrier and its regulation by neuroimmune factors. *Neurogastroenterol Motil* 22, 718–733, doi:10.1111/j.1365-2982.2010.01498.x (2010). [PubMed: 20377785]
11. Yu M et al. Variations in gut microbiota and fecal metabolic phenotype associated with depression by 16S rRNA gene sequencing and LC/MS-based metabolomics. *J Pharm Biomed Anal* 138, 231–239, doi:10.1016/j.jpba.2017.02.008 (2017). [PubMed: 28219800]
12. Moreira CG et al. Bacterial Adrenergic Sensors Regulate Virulence of Enteric Pathogens in the Gut. *MBio* 7, doi:10.1128/mBio.00826-16 (2016).
13. Houlden A et al. Brain injury induces specific changes in the caecal microbiota of mice via altered autonomic activity and mucoprotein production. *Brain Behav Immun* 57, 10–20, doi:10.1016/j.bbi.2016.04.003 (2016). [PubMed: 27060191]
14. Sovran B et al. Age-associated Impairment of the Mucus Barrier Function is Associated with Profound Changes in Microbiota and Immunity. *Sci Rep* 9, 1437, doi:10.1038/s41598-018-35228-3 (2019). [PubMed: 30723224]
15. Barrett E, Ross RP, O'Toole PW, Fitzgerald GF & Stanton C gamma-Aminobutyric acid production by culturable bacteria from the human intestine. *J Appl Microbiol* 113, 411–417, doi:10.1111/j.1365-2672.2012.05344.x (2012). [PubMed: 22612585]
16. Shishov VA, Kirovskaia TA, Kudrin VS & Oleskin AV [Amine neuromediators, their precursors, and oxidation products in the culture of *Escherichia coli* K-12]. *Prikl Biokhim Mikrobiol* 45, 550–554 (2009). [PubMed: 19845286]
17. Asano Y et al. Critical role of gut microbiota in the production of biologically active, free catecholamines in the gut lumen of mice. *Am J Physiol Gastrointest Liver Physiol* 303, G1288–1295, doi:10.1152/ajpgi.00341.2012 (2012). [PubMed: 23064760]
18. Maslowski KM et al. Regulation of inflammatory responses by gut microbiota and chemoattractant receptor GPR43. *Nature* 461, 1282–1286, doi:10.1038/nature08530 (2009). [PubMed: 19865172]
19. Dalile B, Van Oudenhove L, Vervliet B & Verbeke K The role of short-chain fatty acids in microbiota-gut-brain communication. *Nat Rev Gastroenterol Hepatol* 16, 461–478, doi:10.1038/s41575-019-0157-3 (2019). [PubMed: 31123355]
20. McLoughlin RF, Berthon BS, Jensen ME, Baines KJ & Wood LG Short-chain fatty acids, prebiotics, synbiotics, and systemic inflammation: a systematic review and meta-analysis. *Am J Clin Nutr* 106, 930–945, doi:10.3945/ajcn.117.156265 (2017). [PubMed: 28793992]

21. Byrne CS et al. Increased colonic propionate reduces anticipatory reward responses in the human striatum to high-energy foods. *Am J Clin Nutr* 104, 5–14, doi:10.3945/ajcn.115.126706 (2016). [PubMed: 27169834]
22. Lal S, Kirkup AJ, Brunsden AM, Thompson DG & Grundy D Vagal afferent responses to fatty acids of different chain length in the rat. *Am J Physiol Gastrointest Liver Physiol* 281, G907–915, doi:10.1152/ajpgi.2001.281.4.G907 (2001). [PubMed: 11557510]
23. Diaz Heijtz R Fetal, neonatal, and infant microbiome: Perturbations and subsequent effects on brain development and behavior. *Semin Fetal Neonatal Med* 21, 410–417, doi:10.1016/j.siny.2016.04.012 (2016). [PubMed: 27255860]
24. Bliss ES & Whiteside E The Gut-Brain Axis, the Human Gut Microbiota and Their Integration in the Development of Obesity. *Front Physiol* 9, 900, doi:10.3389/fphys.2018.00900 (2018). [PubMed: 30050464]
25. Torres-Fuentes C, Schellekens H, Dinan TG & Cryan JF The microbiota-gut-brain axis in obesity. *Lancet Gastroenterol Hepatol* 2, 747–756, doi:10.1016/S2468-1253(17)30147-4 (2017). [PubMed: 28844808]
26. Ochoa-Reparaz J & Kasper LH The Second Brain: Is the Gut Microbiota a Link Between Obesity and Central Nervous System Disorders? *Curr Obes Rep* 5, 51–64, doi:10.1007/s13679-016-0191-1 (2016). [PubMed: 26865085]
27. Buhmann H, le Roux CW & Bueter M The gut-brain axis in obesity. *Best Pract Res Clin Gastroenterol* 28, 559–571, doi:10.1016/j.bpg.2014.07.003 (2014). [PubMed: 25194175]
28. Myers MG Jr., Leibel RL, Seeley RJ & Schwartz MW Obesity and leptin resistance: distinguishing cause from effect. *Trends Endocrinol Metab* 21, 643–651, doi:10.1016/j.tem.2010.08.002 (2010). [PubMed: 20846876]
29. Guyenet SJ & Schwartz MW Clinical review: Regulation of food intake, energy balance, and body fat mass: implications for the pathogenesis and treatment of obesity. *J Clin Endocrinol Metab* 97, 745–755, doi:10.1210/jc.2011-2525 (2012). [PubMed: 22238401]
30. Rossi MA & Stuber GD Overlapping Brain Circuits for Homeostatic and Hedonic Feeding. *Cell Metab* 27, 42–56, doi:10.1016/j.cmet.2017.09.021 (2018). [PubMed: 29107504]
31. Volkow ND, Wang GJ, Fowler JS, Tomasi D & Baler R Food and drug reward: overlapping circuits in human obesity and addiction. *Curr Top Behav Neurosci* 11, 1–24, doi:10.1007/7854_2011_169 (2012). [PubMed: 22016109]
32. Volkow ND, Wang GJ, Tomasi D & Baler RD Obesity and addiction: neurobiological overlaps. *Obesity reviews : an official journal of the International Association for the Study of Obesity* 14, 2–18, doi:10.1111/j.1467-789X.2012.01031.x (2013). [PubMed: 23016694]
33. Lindgren E et al. Food addiction: A common neurobiological mechanism with drug abuse. *Front Biosci (Landmark Ed)* 23, 811–836 (2018). [PubMed: 28930574]
34. Gearhardt AN, Corbin WR & Brownell KD Food addiction: an examination of the diagnostic criteria for dependence. *J Addict Med* 3, 1–7, doi:10.1097/ADM.0b013e318193c993 (2009). [PubMed: 21768996]
35. Gearhardt AN, Grilo CM, DiLeone RJ, Brownell KD & Potenza MN Can food be addictive? Public health and policy implications. *Addiction* 106, 1208–1212, doi:10.1111/j.1360-0443.2010.03301.x (2011). [PubMed: 21635588]
36. Schulte EM & Gearhardt AN Associations of Food Addiction in a Sample Recruited to Be Nationally Representative of the United States. *Eur Eat Disord Rev* 26, 112–119, doi:10.1002/erv.2575 (2018). [PubMed: 29266583]
37. Schulte EM, Potenza MN & Gearhardt AN A commentary on the “eating addiction” versus “food addiction” perspectives on addictive-like food consumption. *Appetite* 115, 9–15, doi:10.1016/j.appet.2016.10.033 (2017). [PubMed: 27984189]
38. Gearhardt AN, Davis C, Kuschner R & Brownell KD The addiction potential of hyperpalatable foods. *Curr Drug Abuse Rev* 4, 140–145 (2011). [PubMed: 21999688]
39. Randolph TG The descriptive features of food addiction; addictive eating and drinking. *Q J Stud Alcohol* 17, 198–224 (1956). [PubMed: 13336254]
40. Meule A & Gearhardt AN Food addiction in the light of DSM-5. *Nutrients* 6, 3653–3671, doi:10.3390/nu6093653 (2014). [PubMed: 25230209]

41. Corsica JA & Pelchat ML Food addiction: true or false? *Curr Opin Gastroenterol* 26, 165–169, doi:10.1097/MOG.0b013e328336528d (2010). [PubMed: 20042860]
42. Hone-Blanchet A & Fecteau S Overlap of food addiction and substance use disorders definitions: analysis of animal and human studies. *Neuropharmacology* 85, 81–90, doi:10.1016/j.neuropharm.2014.05.019 (2014). [PubMed: 24863044]
43. Gearhardt AN, Corbin WR & Brownell KD Development of the Yale Food Addiction Scale Version 2.0. *Psychol Addict Behav* 30, 113–121, doi:10.1037/adb0000136 (2016). [PubMed: 26866783]
44. American Psychiatric Association. & American Psychiatric Association. *DSM-5 Task Force Diagnostic and statistical manual of mental disorders : DSM-5. 5th edn*, (American Psychiatric Association, 2013).
45. Sengor G & Gezer C Food addiction and its relationship with disordered eating behaviours and obesity. *Eat Weight Disord*, doi:10.1007/s40519-019-00662-3 (2019).
46. Penzenstadler L, Soares C, Karila L & Khazaal Y Systematic Review of Food Addiction as Measured with the Yale Food Addiction Scale: Implications for the Food Addiction Construct. *Curr Neuropharmacol* 17, 526–538, doi:10.2174/1570159X16666181108093520 (2019). [PubMed: 30406740]
47. Burrows T, Kay-Lambkin F, Pursey K, Skinner J & Dayas C Food addiction and associations with mental health symptoms: a systematic review with meta-analysis. *J Hum Nutr Diet* 31, 544–572, doi:10.1111/jhn.12532 (2018). [PubMed: 29368800]
48. Volkow ND, Wang GJ, Tomasi D & Baler RD The addictive dimensionality of obesity. *Biol Psychiatry* 73, 811–818, doi:10.1016/j.biopsych.2012.12.020 (2013). [PubMed: 23374642]
49. Chen M, Sun Y, Lu L & Shi J Similarities and Differences in Neurobiology. *Adv Exp Med Biol* 1010, 45–58, doi:10.1007/978-981-10-5562-1_3 (2017). [PubMed: 29098667]
50. Kalon E, Hong JY, Tobin C & Schulte T Psychological and Neurobiological Correlates of Food Addiction. *International review of neurobiology* 129, 85–110, doi:10.1016/bs.irn.2016.06.003 (2016). [PubMed: 27503449]
51. DiLeone RJ, Taylor JR & Picciotto MR The drive to eat: comparisons and distinctions between mechanisms of food reward and drug addiction. *Nat Neurosci* 15, 1330–1335, doi:10.1038/nn.3202 (2012). [PubMed: 23007187]
52. Rogers PJ Food and drug addictions: Similarities and differences. *Pharmacol Biochem Behav* 153, 182–190, doi:10.1016/j.pbb.2017.01.001 (2017). [PubMed: 28063947]
53. Ouellette AS et al. Establishing a food addiction diagnosis using the Yale Food Addiction Scale: A closer look at the clinically significant distress/functional impairment criterion. *Appetite* 129, 55–61, doi:10.1016/j.appet.2018.06.031 (2018). [PubMed: 29966727]
54. Davis C et al. Evidence that ‘food addiction’ is a valid phenotype of obesity. *Appetite* 57, 711–717, doi:10.1016/j.appet.2011.08.017 (2011). [PubMed: 21907742]
55. Meule A How Prevalent is “Food Addiction”? *Front Psychiatry* 2, 61, doi:10.3389/fpsy.2011.00061 (2011). [PubMed: 22065960]
56. Avena NM, Gearhardt AN, Gold MS, Wang GJ & Potenza MN Tossing the baby out with the bathwater after a brief rinse? The potential downside of dismissing food addiction based on limited data. *Nat Rev Neurosci* 13, 514; author reply 514, doi:10.1038/nrn3212-c1 (2012). [PubMed: 22714023]
57. Ziauddeen H & Fletcher PC Is food addiction a valid and useful concept? *Obesity reviews : an official journal of the International Association for the Study of Obesity* 14, 19–28, doi:10.1111/j.1467-789X.2012.01046.x (2013). [PubMed: 23057499]
58. Ziauddeen H, Farooqi IS & Fletcher PC Obesity and the brain: how convincing is the addiction model? *Nat Rev Neurosci* 13, 279–286, doi:10.1038/nrn3212 (2012). [PubMed: 22414944]
59. Muller A et al. Food addiction and other addictive behaviours in bariatric surgery candidates. *Eur Eat Disord Rev* 26, 585–596, doi:10.1002/erv.2629 (2018). [PubMed: 30094889]
60. Sevincer GM, Konuk N, Bozkurt S & Coskun H Food addiction and the outcome of bariatric surgery at 1-year: Prospective observational study. *Psychiatry Res* 244, 159–164, doi:10.1016/j.psychres.2016.07.022 (2016). [PubMed: 27479107]

61. Gearhardt AN, Boswell RG & White MA The association of “food addiction” with disordered eating and body mass index. *Eat Behav* 15, 427–433, doi:10.1016/j.eatbeh.2014.05.001 (2014). [PubMed: 25064294]
62. Gearhardt AN, White MA & Potenza MN Binge eating disorder and food addiction. *Curr Drug Abuse Rev* 4, 201–207 (2011). [PubMed: 21999695]
63. Nakazato M et al. A role for ghrelin in the central regulation of feeding. *Nature* 409, 194–198, doi:10.1038/35051587 (2001). [PubMed: 11196643]
64. Wren AM et al. Ghrelin causes hyperphagia and obesity in rats. *Diabetes* 50, 2540–2547, doi:10.2337/diabetes.50.11.2540 (2001). [PubMed: 11679432]
65. Jiang H, Betancourt L & Smith RG Ghrelin amplifies dopamine signaling by cross talk involving formation of growth hormone secretagogue receptor/dopamine receptor subtype 1 heterodimers. *Mol Endocrinol* 20, 1772–1785, doi:10.1210/me.2005-0084 (2006). [PubMed: 16601073]
66. Shah M & Vella A Effects of GLP-1 on appetite and weight. *Rev Endocr Metab Disord* 15, 181–187, doi:10.1007/s11154-014-9289-5 (2014). [PubMed: 24811133]
67. Karra E, Chandarana K & Batterham RL The role of peptide YY in appetite regulation and obesity. *J Physiol* 587, 19–25, doi:10.1113/jphysiol.2008.164269 (2009). [PubMed: 19064614]
68. Tolhurst G et al. Short-chain fatty acids stimulate glucagon-like peptide-1 secretion via the G-protein-coupled receptor FFAR2. *Diabetes* 61, 364–371, doi:10.2337/db11-1019 (2012). [PubMed: 22190648]
69. Cani PD et al. Gut microbiota fermentation of prebiotics increases satietogenic and incretin gut peptide production with consequences for appetite sensation and glucose response after a meal. *Am J Clin Nutr* 90, 1236–1243, doi:10.3945/ajcn.2009.28095 (2009). [PubMed: 19776140]
70. Parnell JA & Reimer RA Weight loss during oligofructose supplementation is associated with decreased ghrelin and increased peptide YY in overweight and obese adults. *Am J Clin Nutr* 89, 1751–1759, doi:10.3945/ajcn.2009.27465 (2009). [PubMed: 19386741]
71. Rodin J, Wack J, Ferrannini E & DeFronzo RA Effect of insulin and glucose on feeding behavior. *Metabolism* 34, 826–831 (1985). [PubMed: 3897769]
72. Jiao N et al. Gut microbiome may contribute to insulin resistance and systemic inflammation in obese rodents: a meta-analysis. *Physiol Genomics* 50, 244–254, doi:10.1152/physiolgenomics.00114.2017 (2018). [PubMed: 29373083]
73. Perry RJ et al. Acetate mediates a microbiome-brain-beta-cell axis to promote metabolic syndrome. *Nature* 534, 213–217, doi:10.1038/nature18309 (2016). [PubMed: 27279214]
74. Fang S et al. Intestinal FXR agonism promotes adipose tissue browning and reduces obesity and insulin resistance. *Nat Med* 21, 159–165, doi:10.1038/nm.3760 (2015). [PubMed: 25559344]
75. Pathak P et al. Intestine farnesoid X receptor agonist and the gut microbiota activate G-protein bile acid receptor-1 signaling to improve metabolism. *Hepatology* 68, 1574–1588, doi:10.1002/hep.29857 (2018). [PubMed: 29486523]
76. Vrieze A et al. Impact of oral vancomycin on gut microbiota, bile acid metabolism, and insulin sensitivity. *J Hepatol* 60, 824–831, doi:10.1016/j.jhep.2013.11.034 (2014). [PubMed: 24316517]
77. Leitao-Goncalves R et al. Commensal bacteria and essential amino acids control food choice behavior and reproduction. *PLoS Biol* 15, e2000862, doi:10.1371/journal.pbio.2000862 (2017). [PubMed: 28441450]
78. Ribeiro C & Dickson BJ Sex peptide receptor and neuronal TOR/S6K signaling modulate nutrient balancing in *Drosophila*. *Curr Biol* 20, 1000–1005, doi:10.1016/j.cub.2010.03.061 (2010). [PubMed: 20471268]
79. Storelli G et al. *Lactobacillus plantarum* promotes *Drosophila* systemic growth by modulating hormonal signals through TOR-dependent nutrient sensing. *Cell Metab* 14, 403–414, doi:10.1016/j.cmet.2011.07.012 (2011). [PubMed: 21907145]
80. Kahathuduwa CN, Boyd LA, Davis T, O’Boyle M & Binks M Brain regions involved in ingestive behavior and related psychological constructs in people undergoing calorie restriction. *Appetite* 107, 348–361, doi:10.1016/j.appet.2016.08.112 (2016). [PubMed: 27565377]
81. Berthoud HR Metabolic and hedonic drives in the neural control of appetite: who is the boss? *Curr Opin Neurobiol* 21, 888–896, doi:10.1016/j.conb.2011.09.004 (2011). [PubMed: 21981809]

82. Simon JJ et al. Integration of homeostatic signaling and food reward processing in the human brain. *JCI Insight* 2, doi:10.1172/jci.insight.92970 (2017).
83. Abizaid A, Gao Q & Horvath TL Thoughts for food: brain mechanisms and peripheral energy balance. *Neuron* 51, 691–702, doi:10.1016/j.neuron.2006.08.025 (2006). [PubMed: 16982416]
84. Gao Q & Horvath TL Neurobiology of feeding and energy expenditure. *Annual review of neuroscience* 30, 367–398, doi:10.1146/annurev.neuro.30.051606.094324 (2007).
85. Berthoud HR, Munzberg H & Morrison CD Blaming the Brain for Obesity: Integration of Hedonic and Homeostatic Mechanisms. *Gastroenterology* 152, 1728–1738, doi:10.1053/j.gastro.2016.12.050 (2017). [PubMed: 28192106]
86. Berthoud HR & Morrison C The brain, appetite, and obesity. *Annu Rev Psychol* 59, 55–92, doi:10.1146/annurev.psych.59.103006.093551 (2008). [PubMed: 18154499]
87. Harding IH et al. Brain substrates of unhealthy versus healthy food choices: influence of homeostatic status and body mass index. *International journal of obesity* 42, 448–454, doi:10.1038/ijo.2017.237 (2018). [PubMed: 29064475]
88. Gupta A et al. Patterns of brain structural connectivity differentiate normal weight from overweight subjects. *NeuroImage. Clinical* 7, 506–517, doi:10.1016/j.nicl.2015.01.005 (2015). [PubMed: 25737959]
89. Gupta A et al. Sex differences in the influence of body mass index on anatomical architecture of brain networks. *International journal of obesity* (2005) 41, 1185–1195, doi:10.1038/ijo.2017.86 (2017). [PubMed: 28360430]
90. Kenny PJ Reward mechanisms in obesity: new insights and future directions. *Neuron* 69, 664–679, doi:10.1016/j.neuron.2011.02.016 (2011). [PubMed: 21338878]
91. Volkow ND, Wang GJ & Baler RD Reward, dopamine and the control of food intake: implications for obesity. *Trends Cogn Sci* 15, 37–46, doi:10.1016/j.tics.2010.11.001 (2011). [PubMed: 21109477]
92. Bartholdy S, Dalton B, O'Daly OG, Campbell IC & Schmidt U A systematic review of the relationship between eating, weight and inhibitory control using the stop signal task. *Neurosci Biobehav Rev* 64, 35–62, doi:10.1016/j.neubiorev.2016.02.010 (2016). [PubMed: 26900651]
93. Gearhardt AN, Yokum S, Stice E, Harris JL & Brownell KD Relation of obesity to neural activation in response to food commercials. *Soc Cogn Affect Neurosci* 9, 932–938, doi:10.1093/scan/nst059 (2014). [PubMed: 23576811]
94. Steward T et al. Food addiction and impaired executive functions in women with obesity. *Eur Eat Disord Rev* 26, 574–584, doi:10.1002/erv.2636 (2018). [PubMed: 30159982]
95. Stice E, Yokum S, Burger KS, Epstein LH & Small DM Youth at risk for obesity show greater activation of striatal and somatosensory regions to food. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 31, 4360–4366, doi:10.1523/JNEUROSCI.6604-10.2011 (2011). [PubMed: 21430137]
96. Olivo G et al. Resting-State Brain and the FTO Obesity Risk Allele: Default Mode, Sensorimotor, and Salience Network Connectivity Underlying Different Somatosensory Integration and Reward Processing between Genotypes. *Front Hum Neurosci* 10, 52, doi:10.3389/fnhum.2016.00052 (2016). [PubMed: 26924971]
97. Morrow JD, Maren S & Robinson TE Individual variation in the propensity to attribute incentive salience to an appetitive cue predicts the propensity to attribute motivational salience to an aversive cue. *Behavioural brain research* 220, 238–243, doi:10.1016/j.bbr.2011.02.013 (2011). [PubMed: 21316397]
98. Garcia-Garcia I et al. Alterations of the salience network in obesity: a resting-state fMRI study. *Human brain mapping* 34, 2786–2797, doi:10.1002/hbm.22104 (2013). [PubMed: 22522963]
99. Volkow ND & Baler RD NOW vs LATER brain circuits: implications for obesity and addiction. *Trends in neurosciences* 38, 345–352, doi:10.1016/j.tins.2015.04.002 (2015). [PubMed: 25959611]
100. Wijngaarden MA et al. Obesity is marked by distinct functional connectivity in brain networks involved in food reward and salience. *Behav Brain Res* 287, 127–134, doi:10.1016/j.bbr.2015.03.016 (2015). [PubMed: 25779924]

101. Seeley WW et al. Dissociable intrinsic connectivity networks for salience processing and executive control. *J Neurosci* 27, 2349–2356, doi:10.1523/Jneurosci.5587-06.2007 (2007). [PubMed: 17329432]
102. Menon V & Uddin LQ Saliency, switching, attention and control: a network model of insula function. *Brain structure & function* 214, 655–667, doi:10.1007/s00429-010-0262-0 (2010). [PubMed: 20512370]
103. Zald DH The human amygdala and the emotional evaluation of sensory stimuli. *Brain research. Brain research reviews* 41, 88–123 (2003). [PubMed: 12505650]
104. Kilpatrick LA et al. Influence of sucrose ingestion on brainstem and hypothalamic intrinsic oscillations in lean and obese women. *Gastroenterology* 146, 1212–1221, doi:10.1053/j.gastro.2014.01.023 (2014). [PubMed: 24480616]
105. Park BY, Chung CS, Lee MJ & Park H Accurate neuroimaging biomarkers to predict body mass index in adolescents: a longitudinal study. *Brain Imaging Behav*, doi:10.1007/s11682-019-00101-y (2019).
106. Meng Q et al. Disrupted topological organization of the frontal-mesolimbic network in obese patients. *Brain Imaging Behav* 12, 1544–1555, doi:10.1007/s11682-017-9802-z (2018). [PubMed: 29318488]
107. Baek K, Morris LS, Kundu P & Voon V Disrupted resting-state brain network properties in obesity: decreased global and putaminal cortico-striatal network efficiency. *Psychol Med* 47, 585–596, doi:10.1017/S0033291716002646 (2017). [PubMed: 27804899]
108. Harrold JA & Halford JC The hypothalamus and obesity. *Recent Pat CNS Drug Discov* 1, 305–314 (2006). [PubMed: 18221212]
109. Munzberg H, Qualls-Creekmore E, Berthoud HR, Morrison CD & Yu S Neural Control of Energy Expenditure. *Handb Exp Pharmacol* 233, 173–194, doi:10.1007/164_2015_33 (2016). [PubMed: 26578523]
110. King BM The rise, fall, and resurrection of the ventromedial hypothalamus in the regulation of feeding behavior and body weight. *Physiol Behav* 87, 221–244, doi:10.1016/j.physbeh.2005.10.007 (2006). [PubMed: 16412483]
111. Purnell JQ, Lahna DL, Samuels MH, Rooney WD & Hoffman WF Loss of pons-to-hypothalamic white matter tracks in brainstem obesity. *Int J Obes (Lond)* 38, 1573–1577, doi:10.1038/ijo.2014.57 (2014). [PubMed: 24727578]
112. Carmo-Silva S & Cavadas C Hypothalamic Dysfunction in Obesity and Metabolic Disorders. *Adv Neurobiol* 19, 73–116, doi:10.1007/978-3-319-63260-5_4 (2017). [PubMed: 28933062]
113. Fu O et al. Hypothalamic neuronal circuits regulating hunger-induced taste modification. *Nat Commun* 10, 4560, doi:10.1038/s41467-019-12478-x (2019). [PubMed: 31594935]
114. Zagmutt S, Mera P, Soler-Vazquez MC, Herrero L & Serra D Targeting AgRP neurons to maintain energy balance: Lessons from animal models. *Biochem Pharmacol* 155, 224–232, doi:10.1016/j.bcp.2018.07.008 (2018). [PubMed: 30012460]
115. Morrison CD & Berthoud HR Neurobiology of nutrition and obesity. *Nutr Rev* 65, 517–534 (2007). [PubMed: 18236691]
116. Pedram P et al. Food addiction: its prevalence and significant association with obesity in the general population. *PLoS One* 8, e74832, doi:10.1371/journal.pone.0074832 (2013). [PubMed: 24023964]
117. Chao AM et al. Prevalence and psychosocial correlates of food addiction in persons with obesity seeking weight reduction. *Compr Psychiatry* 73, 97–104, doi:10.1016/j.comppsy.2016.11.009 (2017). [PubMed: 27930952]
118. Pursey KM, Stanwell P, Gearhardt AN, Collins CE & Burrows TL The prevalence of food addiction as assessed by the Yale Food Addiction Scale: a systematic review. *Nutrients* 6, 4552–4590, doi:10.3390/nu6104552 (2014). [PubMed: 25338274]
119. Eichen DM, Lent MR, Goldbacher E & Foster GD Exploration of “food addiction” in overweight and obese treatment-seeking adults. *Appetite* 67, 22–24, doi:10.1016/j.appet.2013.03.008 (2013). [PubMed: 23535004]
120. Michaelides M, Thanos PK, Volkow ND & Wang GJ Translational neuroimaging in drug addiction and obesity. *ILAR J* 53, 59–68 (2012). [PubMed: 23520600]

121. Hommel JD et al. Leptin receptor signaling in midbrain dopamine neurons regulates feeding. *Neuron* 51, 801–810, doi:10.1016/j.neuron.2006.08.023 (2006). [PubMed: 16982424]
122. Lutter M & Nestler EJ Homeostatic and hedonic signals interact in the regulation of food intake. *J Nutr* 139, 629–632, doi:10.3945/jn.108.097618 (2009). [PubMed: 19176746]
123. Morton GJ, Cummings DE, Baskin DG, Barsh GS & Schwartz MW Central nervous system control of food intake and body weight. *Nature* 443, 289–295, doi:10.1038/nature05026 (2006). [PubMed: 16988703]
124. Sinha R & Jastreboff AM Stress as a common risk factor for obesity and addiction. *Biol Psychiatry* 73, 827–835, doi:10.1016/j.biopsych.2013.01.032 (2013). [PubMed: 23541000]
125. Sinha R Role of addiction and stress neurobiology on food intake and obesity. *Biological psychology* 131, 5–13, doi:10.1016/j.biopsycho.2017.05.001 (2018). [PubMed: 28479142]
126. Onalapo AY & Onalapo OJ Food additives, food and the concept of ‘food addiction’: Is stimulation of the brain reward circuit by food sufficient to trigger addiction? *Pathophysiology* 25, 263–276, doi:10.1016/j.pathophys.2018.04.002 (2018). [PubMed: 29673924]
127. Stice E, Spoor S, Bohon C, Veldhuizen MG & Small DM Relation of reward from food intake and anticipated food intake to obesity: a functional magnetic resonance imaging study. *Journal of abnormal psychology* 117, 924–935, doi:10.1037/a0013600 (2008). [PubMed: 19025237]
128. Stoeckel LE et al. Effective connectivity of a reward network in obese women. *Brain Res Bull* 79, 388–395, doi:10.1016/j.brainresbull.2009.05.016 (2009). [PubMed: 19467298]
129. Stoeckel LE et al. Widespread reward-system activation in obese women in response to pictures of high-calorie foods. *NeuroImage* 41, 636–647, doi:10.1016/j.neuroimage.2008.02.031 (2008). [PubMed: 18413289]
130. Volkow ND et al. Cocaine cues and dopamine in dorsal striatum: mechanism of craving in cocaine addiction. *J Neurosci* 26, 6583–6588, doi:10.1523/JNEUROSCI.1544-06.2006 (2006). [PubMed: 16775146]
131. Blum K, Thanos PK & Gold MS Dopamine and glucose, obesity, and reward deficiency syndrome. *Front Psychol* 5, 919, doi:10.3389/fpsyg.2014.00919 (2014). [PubMed: 25278909]
132. Jastreboff AM et al. Neural correlates of stress- and food cue-induced food craving in obesity: association with insulin levels. *Diabetes care* 36, 394–402, doi:10.2337/dc12-1112 (2013). [PubMed: 23069840]
133. Loeber S et al. Impairment of inhibitory control in response to food-associated cues and attentional bias of obese participants and normal-weight controls. *Int J Obes (Lond)* 36, 1334–1339, doi:10.1038/ijo.2011.184 (2012). [PubMed: 21986703]
134. Martin LE et al. Neural mechanisms associated with food motivation in obese and healthy weight adults. *Obesity (Silver Spring, Md.)* 18, 254–260, doi:10.1038/oby.2009.220 (2010).
135. Blum K, Oscar-Berman M, Barh D, Giordano J & Gold M Dopamine Genetics and Function in Food and Substance Abuse. *J Genet Syndr Gene Ther* 4, doi:10.4172/2157-7412.1000121 (2013).
136. Hardman CA, Herbert VM, Brunstrom JM, Munafo MR & Rogers PJ Dopamine and food reward: effects of acute tyrosine/phenylalanine depletion on appetite. *Physiol Behav* 105, 1202–1207, doi:10.1016/j.physbeh.2011.12.022 (2012). [PubMed: 22230253]
137. Volkow ND et al. Low dopamine striatal D2 receptors are associated with prefrontal metabolism in obese subjects: possible contributing factors. *NeuroImage* 42, 1537–1543, doi:10.1016/j.neuroimage.2008.06.002 (2008). [PubMed: 18598772]
138. Gaiser EC et al. Elevated Dopamine D2/3 Receptor Availability in Obese Individuals: A PET Imaging Study with [(11)C](+)-PHNO. *Neuropsychopharmacology* 41, 3042–3050, doi:10.1038/npp.2016.115 (2016). [PubMed: 27374277]
139. Volkow ND, Wang GJ, Fowler JS & Telang F Overlapping neuronal circuits in addiction and obesity: evidence of systems pathology. *Philos T R Soc B* 363, 3191–3200, doi:10.1098/Rstb.2008.0107 (2008).
140. Volkow ND, Wang GJ, Fowler JS & Telang F Overlapping neuronal circuits in addiction and obesity: evidence of systems pathology. *Philos Trans R Soc Lond B Biol Sci* 363, 3191–3200, doi:10.1098/rstb.2008.0107 (2008). [PubMed: 18640912]

141. Sinha R, Gu P, Hart R & Guarnaccia JB Food craving, cortisol and ghrelin responses in modeling highly palatable snack intake in the laboratory. *Physiology & behavior* 208, 112563, doi:10.1016/j.physbeh.2019.112563 (2019). [PubMed: 31145919]
142. Johnson PM & Kenny PJ Dopamine D2 receptors in addiction-like reward dysfunction and compulsive eating in obese rats. *Nat Neurosci* 13, 635–641, doi:10.1038/nn.2519 (2010). [PubMed: 20348917]
143. Pepino MY et al. Sweet Dopamine: Sucrose Preferences Relate Differentially to Striatal D2 Receptor Binding and Age in Obesity. *Diabetes* 65, 2618–2623, doi:10.2337/db16-0407 (2016). [PubMed: 27307220]
144. Molteni R, Barnard RJ, Ying Z, Roberts CK & Gomez-Pinilla F A high-fat, refined sugar diet reduces hippocampal brain-derived neurotrophic factor, neuronal plasticity, and learning. *Neuroscience* 112, 803–814, doi:10.1016/s0306-4522(02)00123-9 (2002). [PubMed: 12088740]
145. Roessmann U & Gambetti P Astrocytes in the developing human brain. An immunohistochemical study. *Acta Neuropathol* 70, 308–313 (1986). [PubMed: 3766128]
146. Zhang SC Defining glial cells during CNS development. *Nat Rev Neurosci* 2, 840–843, doi:10.1038/35097593 (2001). [PubMed: 11715061]
147. Hoban AE et al. Regulation of prefrontal cortex myelination by the microbiota. *Transl Psychiatry* 6, e774, doi:10.1038/tp.2016.42 (2016). [PubMed: 27045844]
148. Diaz Heijtz R et al. Normal gut microbiota modulates brain development and behavior. *Proc Natl Acad Sci U S A* 108, 3047–3052, doi:10.1073/pnas.1010529108 (2011). [PubMed: 21282636]
149. Sudo N et al. Postnatal microbial colonization programs the hypothalamic-pituitary-adrenal system for stress response in mice. *J Physiol* 558, 263–275, doi:10.1113/jphysiol.2004.063388 (2004). [PubMed: 15133062]
150. Zellner DA et al. Food selection changes under stress. *Physiol Behav* 87, 789–793, doi:10.1016/j.physbeh.2006.01.014 (2006). [PubMed: 16519909]
151. Oliver G, Wardle J & Gibson EL Stress and food choice: a laboratory study. *Psychosom Med* 62, 853–865 (2000). [PubMed: 11139006]
152. Epel E, Lapidus R, McEwen B & Brownell K Stress may add bite to appetite in women: a laboratory study of stress-induced cortisol and eating behavior. *Psychoneuroendocrinology* 26, 37–49, doi:10.1016/s0306-4530(00)00035-4 (2001). [PubMed: 11070333]
153. Bose M, Olivan B & Laferrere B Stress and obesity: the role of the hypothalamic-pituitary-adrenal axis in metabolic disease. *Curr Opin Endocrinol Diabetes Obes* 16, 340–346, doi:10.1097/MED.0b013e32832fa137 (2009). [PubMed: 19584720]
154. Lee MJ, Pramyothin P, Karastergiou K & Fried SK Deconstructing the roles of glucocorticoids in adipose tissue biology and the development of central obesity. *Biochim Biophys Acta* 1842, 473–481, doi:10.1016/j.bbdis.2013.05.029 (2014). [PubMed: 23735216]
155. Cong X, Henderson WA, Graf J & McGrath JM Early Life Experience and Gut Microbiome: The Brain-Gut-Microbiota Signaling System. *Adv Neonatal Care* 15, 314–323; quiz E311–312, doi:10.1097/ANC.000000000000191 (2015). [PubMed: 26240939]
156. Neuman H, Forsythe P, Uzan A, Avni O & Koren O Antibiotics in early life: dysbiosis and the damage done. *FEMS Microbiol Rev* 42, 489–499, doi:10.1093/femsre/fuy018 (2018). [PubMed: 29945240]
157. Lundgren SN et al. Maternal diet during pregnancy is related with the infant stool microbiome in a delivery mode-dependent manner. *Microbiome* 6, 109, doi:10.1186/s40168-018-0490-8 (2018). [PubMed: 29973274]
158. Chu DM et al. The early infant gut microbiome varies in association with a maternal high-fat diet. *Genome Med* 8, 77, doi:10.1186/s13073-016-0330-z (2016). [PubMed: 27503374]
159. Bhagavata Srinivasan SP, Raipuria M, Bahari H, Kaakoush NO & Morris MJ Impacts of Diet and Exercise on Maternal Gut Microbiota Are Transferred to Offspring. *Front Endocrinol (Lausanne)* 9, 716, doi:10.3389/fendo.2018.00716 (2018). [PubMed: 30559716]
160. Hohwu L, Li J, Olsen J, Sorensen TI & Obel C Severe maternal stress exposure due to bereavement before, during and after pregnancy and risk of overweight and obesity in young adult men: a Danish National Cohort Study. *PLoS One* 9, e97490, doi:10.1371/journal.pone.0097490 (2014). [PubMed: 24828434]

161. Jasarevic E et al. The maternal vaginal microbiome partially mediates the effects of prenatal stress on offspring gut and hypothalamus. *Nat Neurosci*, doi:10.1038/s41593-018-0182-5 (2018).
162. Mueller NT et al. Prenatal exposure to antibiotics, cesarean section and risk of childhood obesity. *Int J Obes (Lond)* 39, 665–670, doi:10.1038/ijo.2014.180 (2015). [PubMed: 25298276]
163. Dominguez-Bello MG et al. Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc Natl Acad Sci U S A* 107, 11971–11975, doi:10.1073/pnas.1002601107 (2010). [PubMed: 20566857]
164. Marcobal A & Sonnenburg JL Human milk oligosaccharide consumption by intestinal microbiota. *Clin Microbiol Infect* 18 Suppl 4, 12–15, doi:10.1111/j.1469-0691.2012.03863.x (2012). [PubMed: 22647041]
165. Roger LC, Costabile A, Holland DT, Hoyles L & McCartney AL Examination of faecal *Bifidobacterium* populations in breast- and formula-fed infants during the first 18 months of life. *Microbiology* 156, 3329–3341, doi:10.1099/mic.0.043224-0 (2010). [PubMed: 20864478]
166. Tamburini S, Shen N, Wu HC & Clemente JC The microbiome in early life: implications for health outcomes. *Nat Med* 22, 713–722, doi:10.1038/nm.4142 (2016). [PubMed: 27387886]
167. Hart AL et al. Modulation of human dendritic cell phenotype and function by probiotic bacteria. *Gut* 53, 1602–1609, doi:10.1136/gut.2003.037325 (2004). [PubMed: 15479680]
168. O’Sullivan A, Farver M & Smilowitz JT The Influence of Early Infant-Feeding Practices on the Intestinal Microbiome and Body Composition in Infants. *Nutr Metab Insights* 8, 1–9, doi:10.4137/NMI.S29530 (2015).
169. Bogen DL, Hanusa BH & Whitaker RC The effect of breast-feeding with and without formula use on the risk of obesity at 4 years of age. *Obes Res* 12, 1527–1535, doi:10.1038/oby.2004.190 (2004). [PubMed: 15483218]
170. Koenig JE et al. Succession of microbial consortia in the developing infant gut microbiome. *Proc Natl Acad Sci U S A* 108 Suppl 1, 4578–4585, doi:10.1073/pnas.1000081107 (2011). [PubMed: 20668239]
171. Forbes JD et al. Association of Exposure to Formula in the Hospital and Subsequent Infant Feeding Practices With Gut Microbiota and Risk of Overweight in the First Year of Life. *JAMA Pediatr* 172, e181161, doi:10.1001/jamapediatrics.2018.1161 (2018). [PubMed: 29868719]
172. Yan J, Liu L, Zhu Y, Huang G & Wang PP The association between breastfeeding and childhood obesity: a meta-analysis. *BMC Public Health* 14, 1267, doi:10.1186/1471-2458-14-1267 (2014). [PubMed: 25495402]
173. Monteiro CA et al. Household availability of ultra-processed foods and obesity in nineteen European countries. *Public Health Nutr* 21, 18–26, doi:10.1017/S1368980017001379 (2018). [PubMed: 28714422]
174. Martinez Steele E et al. Ultra-processed foods and added sugars in the US diet: evidence from a nationally representative cross-sectional study. *BMJ Open* 6, e009892, doi:10.1136/bmjopen-2015-009892 (2016).
175. Hall KD Did the Food Environment Cause the Obesity Epidemic? *Obesity* 26, 11–13, doi:10.1002/oby.22073 (2018). [PubMed: 29265772]
176. Sadeghirad B, Duhaney T, Motaghipisheh S, Campbell NR & Johnston BC Influence of unhealthy food and beverage marketing on children’s dietary intake and preference: a systematic review and meta-analysis of randomized trials. *Obesity reviews : an official journal of the International Association for the Study of Obesity* 17, 945–959, doi:10.1111/obr.12445 (2016). [PubMed: 27427474]
177. Uribe R & Fuentes-Garcia A The effects of TV unhealthy food brand placement on children. Its separate and joint effect with advertising. *Appetite* 91, 165–172, doi:10.1016/j.appet.2015.03.030 (2015). [PubMed: 25839731]
178. Hicks LA, Taylor TH Jr. & Hunkler RJ US outpatient antibiotic prescribing, 2010. *N Engl J Med* 368, 1461–1462, doi:10.1056/NEJMc1212055 (2013). [PubMed: 23574140]
179. Stark CM, Susi A, Emerick J & Nylund CM Antibiotic and acid-suppression medications during early childhood are associated with obesity. *Gut* 68, 62–69, doi:10.1136/gutjnl-2017-314971 (2019). [PubMed: 30377188]

180. Yassour M et al. Natural history of the infant gut microbiome and impact of antibiotic treatment on bacterial strain diversity and stability. *Sci Transl Med* 8, 343ra381, doi:10.1126/scitranslmed.aad0917 (2016).
181. Ajslev TA, Andersen CS, Gamborg M, Sorensen TI & Jess T Childhood overweight after establishment of the gut microbiota: the role of delivery mode, pre-pregnancy weight and early administration of antibiotics. *Int J Obes (Lond)* 35, 522–529, doi:10.1038/ijo.2011.27 (2011). [PubMed: 21386800]
182. Cox LM et al. Altering the intestinal microbiota during a critical developmental window has lasting metabolic consequences. *Cell* 158, 705–721, doi:10.1016/j.cell.2014.05.052 (2014). [PubMed: 25126780]
183. Candon S et al. Antibiotics in early life alter the gut microbiome and increase disease incidence in a spontaneous mouse model of autoimmune insulin-dependent diabetes. *PLoS One* 10, e0125448, doi:10.1371/journal.pone.0125448 (2015). [PubMed: 25970503]
184. Gaskins HR, Collier CT & Anderson DB Antibiotics as growth promotants: mode of action. *Anim Biotechnol* 13, 29–42, doi:10.1081/ABIO-120005768 (2002). [PubMed: 12212942]
185. Cho I et al. Antibiotics in early life alter the murine colonic microbiome and adiposity. *Nature* 488, 621–626, doi:10.1038/nature11400 (2012). [PubMed: 22914093]
186. Ruiz VE et al. A single early-in-life macrolide course has lasting effects on murine microbial network topology and immunity. *Nat Commun* 8, 518, doi:10.1038/s41467-017-00531-6 (2017). [PubMed: 28894149]
187. Hemmingsson E Early Childhood Obesity Risk Factors: Socioeconomic Adversity, Family Dysfunction, Offspring Distress, and Junk Food Self-Medication. *Curr Obes Rep* 7, 204–209, doi:10.1007/s13679-018-0310-2 (2018). [PubMed: 29704182]
188. Farr OM et al. Posttraumatic stress disorder, alone or additively with early life adversity, is associated with obesity and cardiometabolic risk. *Nutr Metab Cardiovasc Dis* 25, 479–488, doi:10.1016/j.numecd.2015.01.007 (2015). [PubMed: 25770759]
189. Hemmingsson E, Johansson K & Reynisdottir S Effects of childhood abuse on adult obesity: a systematic review and meta-analysis. *Obesity reviews : an official journal of the International Association for the Study of Obesity* 15, 882–893, doi:10.1111/obr.12216 (2014). [PubMed: 25123205]
190. Forster GL, Anderson EM, Scholl JL, Lukkes JL & Watt MJ Negative consequences of early-life adversity on substance use as mediated by corticotropin-releasing factor modulation of serotonin activity. *Neurobiol Stress* 9, 29–39, doi:10.1016/j.ynstr.2018.08.001 (2018). [PubMed: 30151419]
191. Whitesell NR et al. Childhood exposure to adversity and risk of substance-use disorder in two American Indian populations: the meditational role of early substance-use initiation. *J Stud Alcohol Drugs* 70, 971–981 (2009). [PubMed: 19895776]
192. Moffett MC et al. Maternal separation alters drug intake patterns in adulthood in rats. *Biochem Pharmacol* 73, 321–330, doi:10.1016/j.bcp.2006.08.003 (2007). [PubMed: 16962564]
193. Moussaoui N et al. Chronic Early-life Stress in Rat Pups Alters Basal Corticosterone, Intestinal Permeability, and Fecal Microbiota at Weaning: Influence of Sex. *J Neurogastroenterol Motil* 23, 135–143, doi:10.5056/jnm16105 (2017). [PubMed: 27829577]
194. Isohookana R, Marttunen M, Hakko H, Riipinen P & Riala K The impact of adverse childhood experiences on obesity and unhealthy weight control behaviors among adolescents. *Compr Psychiatry* 71, 17–24, doi:10.1016/j.comppsy.2016.08.002 (2016). [PubMed: 27580313]
195. Windle M et al. A multivariate analysis of adverse childhood experiences and health behaviors and outcomes among college students. *J Am Coll Health* 66, 246–251, doi:10.1080/07448481.2018.1431892 (2018). [PubMed: 29405856]
196. Campbell JA, Farmer GC, Nguyen-Rodriguez S, Walker RJ & Egede LE Using path analysis to examine the relationship between sexual abuse in childhood and diabetes in adulthood in a sample of US adults. *Prev Med* 108, 1–7, doi:10.1016/j.ypmed.2017.12.013 (2018). [PubMed: 29277408]

197. Van Niel C, Pachter LM, Wade R Jr., Felitti VJ & Stein MT Adverse events in children: predictors of adult physical and mental conditions. *J Dev Behav Pediatr* 35, 549–551, doi:10.1097/DBP.000000000000102 (2014). [PubMed: 25225793]
198. Osadchiy V et al. History of early life adversity is associated with increased food addiction and sex-specific alterations in reward network connectivity in obesity. *Obes Sci Pract* 5, 416–436, doi:10.1002/osp4.362 (2019). [PubMed: 31687167]
199. Martin B et al. Sex-dependent metabolic, neuroendocrine, and cognitive responses to dietary energy restriction and excess. *Endocrinology* 148, 4318–4333, doi:10.1210/en.2007-0161 (2007). [PubMed: 17569758]
200. Inam QU, Ikram H, Shireen E & Haleem DJ Effects of sugar rich diet on brain serotonin, hyperphagia and anxiety in animal model of both genders. *Pak J Pharm Sci* 29, 757–763 (2016). [PubMed: 27166525]
201. Osadchiy V et al. Correlation of tryptophan metabolites with connectivity of extended central reward network in healthy subjects. *PloS one* 13, e0201772, doi:10.1371/journal.pone.0201772 (2018). [PubMed: 30080865]
202. Leigh SJ, Lee F & Morris MJ Hyperpalatability and the Generation of Obesity: Roles of Environment, Stress Exposure and Individual Difference. *Curr Obes Rep* 7, 6–18, doi:10.1007/s13679-018-0292-0 (2018). [PubMed: 29435959]
203. Chao A, Grilo CM, White MA & Sinha R Food cravings, food intake, and weight status in a community-based sample. *Eat Behav* 15, 478–482, doi:10.1016/j.eatbeh.2014.06.003 (2014). [PubMed: 25064302]
204. Mozaffarian D, Hao T, Rimm EB, Willett WC & Hu FB Changes in diet and lifestyle and long-term weight gain in women and men. *N Engl J Med* 364, 2392–2404, doi:10.1056/NEJMoal014296 (2011). [PubMed: 21696306]
205. Schulte EM, Joyner MA, Potenza MN, Grilo CM & Gearhardt AN Current considerations regarding food addiction. *Curr Psychiatry Rep* 17, 563, doi:10.1007/s11920-015-0563-3 (2015). [PubMed: 25749750]
206. Nunes-Neto PR et al. Food addiction: Prevalence, psychopathological correlates and associations with quality of life in a large sample. *J Psychiatr Res* 96, 145–152, doi:10.1016/j.jpsychires.2017.10.003 (2018). [PubMed: 29049971]
207. Wu GD et al. Linking long-term dietary patterns with gut microbial enterotypes. *Science* 334, 105–108, doi:10.1126/science.1208344 (2011). [PubMed: 21885731]
208. Xu Z & Knight R Dietary effects on human gut microbiome diversity. *Br J Nutr* 113 Suppl, S1–5, doi:10.1017/S0007114514004127 (2015).
209. English L, Lasschuijt M & Keller KL Mechanisms of the portion size effect. What is known and where do we go from here? *Appetite* 88, 39–49, doi:10.1016/j.appet.2014.11.004 (2015). [PubMed: 25447010]
210. Kerr MA, McCann MT & Livingstone MB Food and the consumer: could labelling be the answer? *Proc Nutr Soc* 74, 158–163, doi:10.1017/S0029665115001676 (2015). [PubMed: 25998678]
211. Lucan SC, Maroko AR, Sanon OC & Schechter CB Unhealthful Food-and-Beverage Advertising in Subway Stations: Targeted Marketing, Vulnerable Groups, Dietary Intake, and Poor Health. *J Urban Health* 94, 220–232, doi:10.1007/s11524-016-0127-9 (2017). [PubMed: 28271237]
212. Coates AE, Hardman CA, Halford JCG, Christiansen P & Boyland EJ Social Media Influencer Marketing and Children's Food Intake: A Randomized Trial. *Pediatrics* 143, doi:10.1542/peds.2018-2554 (2019).
213. Norman J et al. Sustained impact of energy-dense TV and online food advertising on children's dietary intake: a within-subject, randomised, crossover, counter-balanced trial. *Int J Behav Nutr Phys Act* 15, 37, doi:10.1186/s12966-018-0672-6 (2018). [PubMed: 29650023]
214. Folkvord F, Anschutz DJ, Wiers RW & Buijzen M The role of attentional bias in the effect of food advertising on actual food intake among children. *Appetite* 84, 251–258, doi:10.1016/j.appet.2014.10.016 (2015). [PubMed: 25451582]

215. Deglaire A et al. Associations between weight status and liking scores for sweet, salt and fat according to the gender in adults (The Nutrinet-Sante study). *Eur J Clin Nutr* 69, 40–46, doi:10.1038/ejcn.2014.139 (2015). [PubMed: 25074389]
216. Geiker NRW et al. Does stress influence sleep patterns, food intake, weight gain, abdominal obesity and weight loss interventions and vice versa? *Obesity reviews : an official journal of the International Association for the Study of Obesity* 19, 81–97, doi:10.1111/obr.12603 (2018). [PubMed: 28849612]
217. Chao A, Grilo CM, White MA & Sinha R Food cravings mediate the relationship between chronic stress and body mass index. *J Health Psychol* 20, 721–729, doi:10.1177/1359105315573448 (2015). [PubMed: 26032789]
218. Sinha R Chronic stress, drug use, and vulnerability to addiction. *Ann N Y Acad Sci* 1141, 105–130, doi:10.1196/annals.1441.030 (2008). [PubMed: 18991954]
219. Chao AM, Jastreboff AM, White MA, Grilo CM & Sinha R Stress, cortisol, and other appetite-related hormones: Prospective prediction of 6-month changes in food cravings and weight. *Obesity (Silver Spring, Md.)* 25, 713–720, doi:10.1002/oby.21790 (2017).
220. Dallman MF Stress-induced obesity and the emotional nervous system. *Trends Endocrinol Metab* 21, 159–165, doi:10.1016/j.tem.2009.10.004 (2010). [PubMed: 19926299]
221. Christiansen AM, Dekloet AD, Ulrich-Lai YM & Herman JP “Snacking” causes long term attenuation of HPA axis stress responses and enhancement of brain FosB/deltaFosB expression in rats. *Physiol Behav* 103, 111–116, doi:10.1016/j.physbeh.2011.01.015 (2011). [PubMed: 21262247]
222. Ulrich-Lai YM et al. Pleasurable behaviors reduce stress via brain reward pathways. *Proc Natl Acad Sci U S A* 107, 20529–20534, doi:10.1073/pnas.1007740107 (2010). [PubMed: 21059919]
223. Bharwani A et al. Structural & functional consequences of chronic psychosocial stress on the microbiome & host. *Psychoneuroendocrinology* 63, 217–227, doi:10.1016/j.psyneuen.2015.10.001 (2016). [PubMed: 26479188]
224. Depommier C et al. Supplementation with Akkermansia muciniphila in overweight and obese human volunteers: a proof-of-concept exploratory study. *Nat Med* 25, 1096–1103, doi:10.1038/s41591-019-0495-2 (2019). [PubMed: 31263284]
225. Gheorghe CE et al. Focus on the essentials: tryptophan metabolism and the microbiome-gut-brain axis. *Curr Opin Pharmacol* 48, 137–145, doi:10.1016/j.coph.2019.08.004 (2019). [PubMed: 31610413]
226. Osadchiy V, Martin CR & Mayer EA Gut Microbiome and Modulation of CNS Function. *Compr Physiol* 10, 57–72, doi:10.1002/cphy.c180031 (2019). [PubMed: 31853944]
227. O’Mahony SM, Clarke G, Borre YE, Dinan TG & Cryan JF Serotonin, tryptophan metabolism and the brain-gut-microbiome axis. *Behav Brain Res* 277, 32–48, doi:10.1016/j.bbr.2014.07.027 (2015). [PubMed: 25078296]
228. Wikoff WR et al. Metabolomics analysis reveals large effects of gut microflora on mammalian blood metabolites. *Proc Natl Acad Sci U S A* 106, 3698–3703, doi:10.1073/pnas.0812874106 (2009). [PubMed: 19234110]
229. Yano JM et al. Indigenous bacteria from the gut microbiota regulate host serotonin biosynthesis. *Cell* 161, 264–276, doi:10.1016/j.cell.2015.02.047 (2015). [PubMed: 25860609]
230. Kim DY & Camilleri M Serotonin: a mediator of the brain-gut connection. *Am J Gastroenterol* 95, 2698–2709, doi:10.1111/j.1572-0241.2000.03177.x (2000). [PubMed: 11051338]
231. Hood SD, Bell CJ & Nutt DJ Acute tryptophan depletion. Part I: rationale and methodology. *Aust N Z J Psychiatry* 39, 558–564, doi:10.1080/j.1440-1614.2005.01627.x (2005). [PubMed: 15996136]
232. Pagoto SL et al. Acute tryptophan depletion and sweet food consumption by overweight adults. *Eat Behav* 10, 36–41, doi:10.1016/j.eatbeh.2008.10.010 (2009). [PubMed: 19171315]
233. Roager HM & Licht TR Microbial tryptophan catabolites in health and disease. *Nat Commun* 9, 3294, doi:10.1038/s41467-018-05470-4 (2018). [PubMed: 30120222]
234. Bender DA Biochemistry of tryptophan in health and disease. *Mol Aspects Med* 6, 101–197 (1983). [PubMed: 6371429]

235. Schwarcz R, Bruno JP, Muchowski PJ & Wu HQ Kynurenines in the mammalian brain: when physiology meets pathology. *Nat Rev Neurosci* 13, 465–477, doi:10.1038/nrn3257 (2012). [PubMed: 22678511]
236. Marin IA et al. Microbiota alteration is associated with the development of stress-induced despair behavior. *Sci Rep* 7, 43859, doi:10.1038/srep43859 (2017). [PubMed: 28266612]
237. Stavrum AK, Heiland I, Schuster S, Puntervoll P & Ziegler M Model of tryptophan metabolism, readily scalable using tissue-specific gene expression data. *J Biol Chem* 288, 34555–34566, doi:10.1074/jbc.M113.474908 (2013). [PubMed: 24129579]
238. Favennec M et al. The kynurenine pathway is activated in human obesity and shifted toward kynurenine monooxygenase activation. *Obesity* 23, 2066–2074, doi:10.1002/oby.21199 (2015). [PubMed: 26347385]
239. Kennedy PJ, Cryan JF, Dinan TG & Clarke G Kynurenine pathway metabolism and the microbiota-gut-brain axis. *Neuropharmacology* 112, 399–412, doi:10.1016/j.neuropharm.2016.07.002 (2017). [PubMed: 27392632]
240. Chimere C et al. Bacterial metabolite indole modulates incretin secretion from intestinal enteroendocrine L cells. *Cell Rep* 9, 1202–1208, doi:10.1016/j.celrep.2014.10.032 (2014). [PubMed: 25456122]
241. Mangge H et al. Obesity-related dysregulation of the tryptophan-kynurenine metabolism: role of age and parameters of the metabolic syndrome. *Obesity* 22, 195–201, doi:10.1002/oby.20491 (2014). [PubMed: 23625535]
242. Buckman LB et al. Obesity induced by a high-fat diet is associated with increased immune cell entry into the central nervous system. *Brain Behav Immun* 35, 33–42, doi:10.1016/j.bbi.2013.06.007 (2014). [PubMed: 23831150]
243. Teixeira TF, Collado MC, Ferreira CL, Bressan J & Peluzio Mdo C Potential mechanisms for the emerging link between obesity and increased intestinal permeability. *Nutr Res* 32, 637–647, doi:10.1016/j.nutres.2012.07.003 (2012). [PubMed: 23084636]
244. Sonnenburg ED et al. Diet-induced extinctions in the gut microbiota compound over generations. *Nature* 529, 212–215, doi:10.1038/nature16504 (2016). [PubMed: 26762459]
245. Rasoamanana R, Even PC, Darcel N, Tome D & Fromentin G Dietary fibers reduce food intake by satiation without conditioned taste aversion in mice. *Physiol Behav* 110–111, 13–19, doi:10.1016/j.physbeh.2012.12.008 (2013).
246. Tailford LE, Crost EH, Kavanaugh D & Juge N Mucin glycan foraging in the human gut microbiome. *Front Genet* 6, 81, doi:10.3389/fgene.2015.00081 (2015). [PubMed: 25852737]
247. Shen W et al. Intestinal and systemic inflammatory responses are positively associated with sulfidogenic bacteria abundance in high-fat-fed male C57BL/6J mice. *J Nutr* 144, 1181–1187, doi:10.3945/jn.114.194332 (2014). [PubMed: 24919690]
248. Ding S et al. High-fat diet: bacteria interactions promote intestinal inflammation which precedes and correlates with obesity and insulin resistance in mouse. *PLoS One* 5, e12191, doi:10.1371/journal.pone.0012191 (2010). [PubMed: 20808947]
249. Cani PD et al. Metabolic endotoxemia initiates obesity and insulin resistance. *Diabetes* 56, 1761–1772, doi:10.2337/db06-1491 (2007). [PubMed: 17456850]
250. Chang M, Alsaigh T, Kistler EB & Schmid-Schonbein GW Breakdown of mucin as barrier to digestive enzymes in the ischemic rat small intestine. *PLoS One* 7, e40087, doi:10.1371/journal.pone.0040087 (2012). [PubMed: 22768227]
251. Hotamisligil GS Inflammation and metabolic disorders. *Nature* 444, 860–867, doi:10.1038/nature05485 (2006). [PubMed: 17167474]
252. Wang Q, Liu D, Song P & Zou MH Tryptophan-kynurenine pathway is dysregulated in inflammation, and immune activation. *Front Biosci (Landmark Ed)* 20, 1116–1143 (2015). [PubMed: 25961549]
253. Larraufie P, Dore J, Lapaque N & Blottiere HM TLR ligands and butyrate increase Pyy expression through two distinct but inter-regulated pathways. *Cell Microbiol* 19, doi:10.1111/cmi.12648 (2017).
254. Palazzo M et al. Activation of enteroendocrine cells via TLRs induces hormone, chemokine, and defensin secretion. *J Immunol* 178, 4296–4303 (2007). [PubMed: 17371986]

255. Kidd M, Gustafsson BI, Drozdov I & Modlin IM IL1 β - and LPS-induced serotonin secretion is increased in EC cells derived from Crohn's disease. *Neurogastroenterol Motil* 21, 439–450, doi:10.1111/j.1365-2982.2008.01210.x (2009). [PubMed: 19019013]
256. de Lartigue G, Ronveaux CC & Raybould HE Vagal plasticity the key to obesity. *Mol Metab* 3, 855–856, doi:10.1016/j.molmet.2014.09.009 (2014). [PubMed: 25506551]
257. de Lartigue G, Barbier de la Serre C, Espero E, Lee J & Raybould HE Diet-induced obesity leads to the development of leptin resistance in vagal afferent neurons. *Am J Physiol Endocrinol Metab* 301, E187–195, doi:10.1152/ajpendo.00056.2011 (2011). [PubMed: 21521717]
258. de Lartigue G, de La Serre CB & Raybould HE Vagal afferent neurons in high fat diet-induced obesity; intestinal microflora, gut inflammation and cholecystokinin. *Physiol Behav* 105, 100–105, doi:10.1016/j.physbeh.2011.02.040 (2011). [PubMed: 21376066]
259. Qin Y et al. An obesity-associated gut microbiome reprograms the intestinal epigenome and leads to altered colonic gene expression. *Genome Biol* 19, 7, doi:10.1186/s13059-018-1389-1 (2018). [PubMed: 29361968]
260. Cani PD et al. Selective increases of bifidobacteria in gut microflora improve high-fat-diet-induced diabetes in mice through a mechanism associated with endotoxaemia. *Diabetologia* 50, 2374–2383, doi:10.1007/s00125-007-0791-0 (2007). [PubMed: 17823788]
261. Peterli R et al. Effect of Laparoscopic Sleeve Gastrectomy vs Laparoscopic Roux-en-Y Gastric Bypass on Weight Loss in Patients With Morbid Obesity: The SM-BOSS Randomized Clinical Trial. *JAMA* 319, 255–265, doi:10.1001/jama.2017.20897 (2018). [PubMed: 29340679]
262. Chang SH et al. The effectiveness and risks of bariatric surgery: an updated systematic review and meta-analysis, 2003–2012. *JAMA Surg* 149, 275–287, doi:10.1001/jamasurg.2013.3654 (2014). [PubMed: 24352617]
263. Scholtz S et al. Obese patients after gastric bypass surgery have lower brain-hedonic responses to food than after gastric banding. *Gut* 63, 891–902, doi:10.1136/gutjnl-2013-305008 (2014). [PubMed: 23964100]
264. Pepino MY et al. Changes in taste perception and eating behavior after bariatric surgery-induced weight loss in women. *Obesity (Silver Spring)* 22, E13–20, doi:10.1002/oby.20649 (2014). [PubMed: 24167016]
265. Sanmiguel C et al. Bariatric surgery is associated with changes in the brain's reward system architecture and eating behaviors. *Gastroenterology* 150, S824 (2016).
266. Kanerva N, Larsson I, Peltonen M, Lindroos AK & Carlsson LM Changes in total energy intake and macronutrient composition after bariatric surgery predict long-term weight outcome: findings from the Swedish Obese Subjects (SOS) study. *Am J Clin Nutr* 106, 136–145, doi:10.3945/ajcn.116.149112 (2017). [PubMed: 28515062]
267. Konttinen H, Peltonen M, Sjostrom L, Carlsson L & Karlsson J Psychological aspects of eating behavior as predictors of 10-y weight changes after surgical and conventional treatment of severe obesity: results from the Swedish Obese Subjects intervention study. *The American journal of clinical nutrition* 101, 16–24, doi:10.3945/ajcn.114.095182 (2015). [PubMed: 25527746]
268. Makaronidis JM et al. Reported appetite, taste and smell changes following Roux-en-Y gastric bypass and sleeve gastrectomy: Effect of gender, type 2 diabetes and relationship to post-operative weight loss. *Appetite* 107, 93–105, doi:10.1016/j.appet.2016.07.029 (2016). [PubMed: 27453553]
269. Pepino MY, Stein RI, Eagon JC & Klein S Bariatric surgery-induced weight loss causes remission of food addiction in extreme obesity. *Obesity (Silver Spring)* 22, 1792–1798, doi:10.1002/oby.20797 (2014). [PubMed: 24852693]
270. Basso N et al. First-phase insulin secretion, insulin sensitivity, ghrelin, GLP-1, and PYY changes 72 h after sleeve gastrectomy in obese diabetic patients: the gastric hypothesis. *Surg Endosc* 25, 3540–3550, doi:10.1007/s00464-011-1755-5 (2011). [PubMed: 21638183]
271. le Roux CW et al. Gut hormones as mediators of appetite and weight loss after Roux-en-Y gastric bypass. *Ann Surg* 246, 780–785, doi:10.1097/SLA.0b013e3180caa3e3 (2007). [PubMed: 17968169]

272. Faulconbridge LF et al. Changes in neural responsivity to highly palatable foods following roux-en-Y gastric bypass, sleeve gastrectomy, or weight stability: An fMRI study. *Obesity* (Silver Spring) 24, 1054–1060, doi:10.1002/oby.21464 (2016). [PubMed: 27112067]
273. Li JV et al. Metabolic surgery profoundly influences gut microbial-host metabolic cross-talk. *Gut* 60, 1214–1223, doi:10.1136/gut.2010.234708 (2011). [PubMed: 21572120]
274. Li M et al. Symbiotic gut microbes modulate human metabolic phenotypes. *Proc Natl Acad Sci U S A* 105, 2117–2122, doi:10.1073/pnas.0712038105 (2008). [PubMed: 18252821]
275. Arora T et al. Roux-en-Y Gastric Bypass Surgery Induces Early Plasma Metabolomic and Lipidomic Alterations in Humans Associated with Diabetes Remission. *PLoS One* 10, e0126401, doi:10.1371/journal.pone.0126401 (2015). [PubMed: 25946120]
276. Gralka E et al. Metabolomic fingerprint of severe obesity is dynamically affected by bariatric surgery in a procedure-dependent manner. *Am J Clin Nutr* 102, 1313–1322, doi:10.3945/ajcn.115.110536 (2015). [PubMed: 26581381]
277. Ryan KK et al. FXR is a molecular target for the effects of vertical sleeve gastrectomy. *Nature* 509, 183–188, doi:10.1038/nature13135 (2014). [PubMed: 24670636]
278. Coveleskie K et al. The effect of the GLP-1 analogue Exenatide on functional connectivity within an NTS-based network in women with and without obesity. *Obes Sci Pract* 3, 434–445, doi:10.1002/osp4.124 (2017). [PubMed: 29259802]
279. Braas D et al. Dynamic Changes in Gut Microbial Derived Indole and Phenol Products after Bariatric Surgery and its Relationship to Weight Loss. *Gastroenterology* 154, S158 (2018).
280. Jacobs J et al. Glutamate and Hedonic Eating: Role of the Brain-Gut-Microbiome Axis on Changes on Hedonic Eating after Bariatric Surgery. *Gastroenterology* 154, s201 (2018).
281. Lee CJ et al. Changes in Gut Microbiome after Bariatric Surgery Versus Medical Weight Loss in a Pilot Randomized Trial. *Obes Surg*, doi:10.1007/s11695-019-03976-4 (2019).
282. Guo Y et al. Modulation of the gut microbiome: a systematic review of the effect of bariatric surgery. *Eur J Endocrinol* 178, 43–56, doi:10.1530/EJE-17-0403 (2018). [PubMed: 28916564]
283. Luijten J, Vugts G, Nieuwenhuijzen GAP & Luyer MDP The Importance of the Microbiome in Bariatric Surgery: a Systematic Review. *Obes Surg* 29, 2338–2349, doi:10.1007/s11695-019-03863-y (2019). [PubMed: 30982169]
284. Monte SV et al. Reduction in endotoxemia, oxidative and inflammatory stress, and insulin resistance after Roux-en-Y gastric bypass surgery in patients with morbid obesity and type 2 diabetes mellitus. *Surgery* 151, 587–593, doi:10.1016/j.surg.2011.09.038 (2012). [PubMed: 22088821]
285. Iannelli A, Anty R, Schneck AS, Tran A & Gugenheim J Inflammation, insulin resistance, lipid disturbances, anthropometrics, and metabolic syndrome in morbidly obese patients: a case control study comparing laparoscopic Roux-en-Y gastric bypass and laparoscopic sleeve gastrectomy. *Surgery* 149, 364–370, doi:10.1016/j.surg.2010.08.013 (2011). [PubMed: 20932542]
286. Iannelli A et al. Body composition, anthropometrics, energy expenditure, systemic inflammation, in premenopausal women 1 year after laparoscopic Roux-en-Y gastric bypass. *Surg Endosc* 28, 500–507, doi:10.1007/s00464-013-3191-1 (2014). [PubMed: 24013471]
287. Yadav R et al. Effect of Roux-en-Y Bariatric Surgery on Lipoproteins, Insulin Resistance, and Systemic and Vascular Inflammation in Obesity and Diabetes. *Front Immunol* 8, 1512, doi:10.3389/fimmu.2017.01512 (2017). [PubMed: 29187850]
288. Peng Y, Li JZ, You M & Murr MM Roux-en-Y gastric bypass improves glucose homeostasis, reduces oxidative stress and inflammation in livers of obese rats and in Kupffer cells via an AMPK-dependent pathway. *Surgery* 162, 59–67, doi:10.1016/j.surg.2017.01.012 (2017). [PubMed: 28291540]
289. Lindegaard KK, Jorgensen NB, Just R, Heegaard PM & Madsbad S Effects of Roux-en-Y gastric bypass on fasting and postprandial inflammation-related parameters in obese subjects with normal glucose tolerance and in obese subjects with type 2 diabetes. *Diabetol Metab Syndr* 7, 12, doi:10.1186/s13098-015-0012-9 (2015). [PubMed: 25763111]

290. van de Sande-Lee S et al. Partial reversibility of hypothalamic dysfunction and changes in brain activity after body mass reduction in obese subjects. *Diabetes* 60, 1699–1704, doi:10.2337/db10-1614 (2011). [PubMed: 21515852]
291. Blackburn AN, Hajnal A & Leggio L The gut in the brain: the effects of bariatric surgery on alcohol consumption. *Addict Biol* 22, 1540–1553, doi:10.1111/adb.12436 (2017). [PubMed: 27578259]
292. Vrieze A et al. Transfer of intestinal microbiota from lean donors increases insulin sensitivity in individuals with metabolic syndrome. *Gastroenterology* 143, 913–916 e917, doi:10.1053/j.gastro.2012.06.031 (2012). [PubMed: 22728514]
293. Kootte RS et al. Improvement of Insulin Sensitivity after Lean Donor Feces in Metabolic Syndrome Is Driven by Baseline Intestinal Microbiota Composition. *Cell Metab* 26, 611–619 e616, doi:10.1016/j.cmet.2017.09.008 (2017). [PubMed: 28978426]
294. Dailey FE, Turse EP, Daglilar E & Tahan V The dirty aspects of fecal microbiota transplantation: a review of its adverse effects and complications. *Curr Opin Pharmacol* 49, 29–33, doi:10.1016/j.coph.2019.04.008 (2019). [PubMed: 31103793]
295. Longo VD & Panda S Fasting, Circadian Rhythms, and Time-Restricted Feeding in Healthy Lifespan. *Cell Metab* 23, 1048–1059, doi:10.1016/j.cmet.2016.06.001 (2016). [PubMed: 27304506]
296. Melkani GC & Panda S Time-restricted feeding for prevention and treatment of cardiometabolic disorders. *J Physiol* 595, 3691–3700, doi:10.1113/JP273094 (2017). [PubMed: 28295377]
297. Di Francesco A, Di Germanio C, Bernier M & de Cabo R A time to fast. *Science* 362, 770–775, doi:10.1126/science.aau2095 (2018). [PubMed: 30442801]
298. Kohsaka A et al. High-fat diet disrupts behavioral and molecular circadian rhythms in mice. *Cell Metab* 6, 414–421, doi:10.1016/j.cmet.2007.09.006 (2007). [PubMed: 17983587]
299. Hatori M et al. Time-restricted feeding without reducing caloric intake prevents metabolic diseases in mice fed a high-fat diet. *Cell Metab* 15, 848–860, doi:10.1016/j.cmet.2012.04.019 (2012). [PubMed: 22608008]
300. Gill S & Panda S A Smartphone App Reveals Erratic Diurnal Eating Patterns in Humans that Can Be Modulated for Health Benefits. *Cell Metab* 22, 789–798, doi:10.1016/j.cmet.2015.09.005 (2015). [PubMed: 26411343]
301. Thaïss CA et al. Transkingdom control of microbiota diurnal oscillations promotes metabolic homeostasis. *Cell* 159, 514–529, doi:10.1016/j.cell.2014.09.048 (2014). [PubMed: 25417104]
302. Racz B, Duskova M, Starka L, Hainer V & Kunesova M Links between the circadian rhythm, obesity and the microbiome. *Physiol Res* 67, S409–S420, doi:10.33549/physiolres.934020 (2018). [PubMed: 30484668]
303. Ara R et al. What is the clinical effectiveness and cost-effectiveness of using drugs in treating obese patients in primary care? A systematic review. *Health Technol Assess* 16, iii–xiv, 1–195, doi:10.3310/hta16050 (2012).
304. Shin JH & Gadde KM Clinical utility of phentermine/topiramate (Qsymia) combination for the treatment of obesity. *Diabetes Metab Syndr Obes* 6, 131–139, doi:10.2147/DMSO.S43403 (2013). [PubMed: 23630428]
305. Hainer V & Aldhoon-Hainerova I Tolerability and safety of the new anti-obesity medications. *Drug Saf* 37, 693–702, doi:10.1007/s40264-014-0206-3 (2014). [PubMed: 25096956]
306. Billes SK, Sinnayah P & Cowley MA Naltrexone/bupropion for obesity: an investigational combination pharmacotherapy for weight loss. *Pharmacol Res* 84, 1–11, doi:10.1016/j.phrs.2014.04.004 (2014). [PubMed: 24754973]
307. Wellman PJ & Maher TJ Synergistic interactions between fenfluramine and phentermine. *Int J Obes Relat Metab Disord* 23, 723–732 (1999). [PubMed: 10454106]
308. Lam DD et al. Serotonin 5-HT_{2C} receptor agonist promotes hypophagia via downstream activation of melanocortin 4 receptors. *Endocrinology* 149, 1323–1328, doi:10.1210/en.2007-1321 (2008). [PubMed: 18039773]
309. McElroy SL et al. Topiramate for the treatment of binge eating disorder associated with obesity: a placebo-controlled study. *Biol Psychiatry* 61, 1039–1048, doi:10.1016/j.biopsych.2006.08.008 (2007). [PubMed: 17258690]

310. Anderberg RH et al. Glucagon-Like Peptide 1 and Its Analogs Act in the Dorsal Raphe and Modulate Central Serotonin to Reduce Appetite and Body Weight. *Diabetes* 66, 1062–1073, doi:10.2337/db16-0755 (2017). [PubMed: 28057699]
311. Wang Z et al. Gut microbiome differences between metformin- and liraglutide-treated T2DM subjects. *Endocrinol Diabetes Metab* 1, e00009, doi:10.1002/edm2.9 (2018). [PubMed: 30815546]
312. Foster D, Sanchez-Collins S & Cheskin LJ Multidisciplinary Team-Based Obesity Treatment in Patients With Diabetes: Current Practices and the State of the Science. *Diabetes Spectr* 30, 244–249, doi:10.2337/ds17-0045 (2017). [PubMed: 29151714]
313. Cooper Z & Fairburn CG A new cognitive behavioural approach to the treatment of obesity. *Behav Res Ther* 39, 499–511 (2001). [PubMed: 11341247]
314. Klumpp H, Fitzgerald DA, Angstadt M, Post D & Phan KL Neural response during attentional control and emotion processing predicts improvement after cognitive behavioral therapy in generalized social anxiety disorder. *Psychol Med* 44, 3109–3121, doi:10.1017/S0033291714000567 (2014). [PubMed: 25066308]
315. Jensen KB et al. Cognitive Behavioral Therapy increases pain-evoked activation of the prefrontal cortex in patients with fibromyalgia. *Pain* 153, 1495–1503, doi:10.1016/j.pain.2012.04.010 (2012). [PubMed: 22617632]
316. Seminowicz DA et al. Cognitive-behavioral therapy increases prefrontal cortex gray matter in patients with chronic pain. *J Pain* 14, 1573–1584, doi:10.1016/j.jpain.2013.07.020 (2013). [PubMed: 24135432]
317. Brunoni AR et al. Cognitive control therapy and transcranial direct current stimulation for depression: a randomized, double-blinded, controlled trial. *J Affect Disord* 162, 43–49, doi:10.1016/j.jad.2014.03.026 (2014). [PubMed: 24767004]
318. Clay SW, Allen J & Parran T A review of addiction. *Postgrad Med* 120, E01–07, doi:10.3810/pgm.2008.07.1802 (2008).
319. Labus J et al. Randomised clinical trial: symptoms of the irritable bowel syndrome are improved by a psycho-education group intervention. *Aliment Pharmacol Ther* 37, 304–315, doi:10.1111/apt.12171 (2013). [PubMed: 23205588]
320. An H, He RH, Zheng YR & Tao R Cognitive-Behavioral Therapy. *Adv Exp Med Biol* 1010, 321–329, doi:10.1007/978-981-10-5562-1_16 (2017). [PubMed: 29098680]
321. Sawamoto R et al. Predictors of successful long-term weight loss maintenance: a two-year follow-up. *Biopsychosoc Med* 11, 14, doi:10.1186/s13030-017-0099-3 (2017). [PubMed: 28592990]
322. O'reilly GA, Cook L, Spruijt-Metz D & Black DS Mindfulness-based interventions for obesity-related eating behaviours: a literature review. *Obesity Reviews* 15, 453–461, doi:10.1111/Obr.12156 (2014). [PubMed: 24636206]
323. Lappalainen R et al. The effectiveness and applicability of different lifestyle interventions for enhancing wellbeing: the study design for a randomized controlled trial for persons with metabolic syndrome risk factors and psychological distress. *Bmc Public Health* 14, doi:10.1186/1471-2458-14-310 (2014).
324. Makki K, Deehan EC, Walter J & Backhed F The Impact of Dietary Fiber on Gut Microbiota in Host Health and Disease. *Cell Host Microbe* 23, 705–715, doi:10.1016/j.chom.2018.05.012 (2018). [PubMed: 29902436]
325. Desai MS et al. A Dietary Fiber-Deprived Gut Microbiota Degrades the Colonic Mucus Barrier and Enhances Pathogen Susceptibility. *Cell* 167, 1339–1353 e1321, doi:10.1016/j.cell.2016.10.043 (2016). [PubMed: 27863247]

Box 1 |**Therapeutic targets within the brain gut microbiome axis for obesity**

Brain Targets	Approaches for Obesity Treatment
Cognitive Control	<ul style="list-style-type: none"> • Cognitive behavioral therapy
Reward Processes	<ul style="list-style-type: none"> • Topiramate + Phentermine • Bupropion + Naltrexone • GLP-1 agonists (e.g. Liraglutide)
Homeostatic Control	<ul style="list-style-type: none"> • Serotonin modulators (e.g. Lorcaserin) • Amphetamine class (e.g. Phentermine) • GLP-1 agonists (e.g. Liraglutide) • Leptin agonists • Vagus nerve electrical modulation
Gut Remodeling	<ul style="list-style-type: none"> • Bariatric surgery (e.g. RYGB, LSG) • Bariatric endoscopy (e.g. gastric balloons, gastric plication, gastric content aspiration)
Gut Absorption	<ul style="list-style-type: none"> • Lipase inhibitors (e.g. Orlistat) • Mucosal barriers/ablation (e.g. duodenal sleeve)
Gut Microbiota	<ul style="list-style-type: none"> • Prebiotics (e.g. oligofructose, oligosacharides) • Probiotics (e.g. Lactobacillus) • Bariatric surgery • Fecal transplant

Key Points

- Food addiction refers to maladaptive ingestive behaviors resulting from a shift from primarily homeostatic to hedonic regulatory mechanisms of food intake; this shift reflects alterations at all levels of the brain–gut–microbiome (BGM) axis.
- Normal ingestive behavior is the result of the tightly regulated interplay between orexogenic and anorexogenic gut hormones, leptin signaling from adipose tissue, hypothalamic nuclei, the dopaminergic reward system and prefrontal inhibitory influences.
- In food addiction, a disinhibition of reward and anorexogenic mechanisms at all levels of the BGM axis results in unrestrained craving for food.
- Several adverse early life events, including nutrition, stress and antibiotic intake can influence the development of BGM interactions and of ingestive behavior.
- Lifelong dietary choices can modulate BGM Interactions and eating behaviors; for example, chronic ingestion of a typical Western diet can result in systemic low-grade immune system activation that can reduce feedback inhibitory mechanisms restraining food intake.
- Current pharmacological treatment options for food addiction are limited, bariatric surgery is the only therapy providing long-term benefits, but novel treatment approaches (including time restricted eating and cognitive behavioral interventions) are being evaluated.

Figure 1. Model of brain–gut–microbiome interactions in ingestive behavior.

In the periphery, gut-generated and vagally transmitted orexogenic and anorexogenic signals interact with specific nuclei in the hypothalamus in the homeostatic regulation of food intake. Food-related factors interact with gut microorganisms and gut microbial metabolites modulate the release of orexogenic and anorexogenic peptides from enteroendocrine cells in the distal small intestine, shifting the balance between anorexogenic and orexogenic signaling in the hypothalamus. In addition, gut microorganisms can signal to the brain via inflammatory mediators (such as lipopolysaccharides) and neuroactive metabolites (such as tryptophan metabolites). Centrally, interactions between several brain networks, including the prefrontal cortex, the dopaminergic reward system and the sensorimotor system underlie the hedonic regulation of food intake. Several environmental influences such as food advertisements, food cues engage the extended reward system which can override the homeostatic control mechanisms. Exposure to visual and sensory cues, as well as psychosocial stress play important role in this process. Blue boxes on the left represent different parts of the BGM axis. Light green boxes in the center show mechanisms involved in altered BGM interactions in food addiction. Upward arrows show upregulation,

downward arrows show downregulation. Modified with permission from Volkow et al.
Biological Psychiatry 2013 EM: DONE

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 2. Model of altered brain network interactions in food addiction.

Several brain networks interact in the regulation of ingestive behavior. In food addiction, increased engagement of the salience network by food cues engages the executive control network leading to increased attention to food, and the emotional arousal network. Insufficient inhibitory control of the reward and of the emotional arousal networks by the executive control network plays a key part in shifting the balance from predominantly homeostatic to hedonic and regulation of food intake. The **salience network** (aMCC, anterior mid cingulate cortex; aINS, anterior insula) responds according to the subjective salience of any interoceptive or exteroceptive stimulus reaching the brain, or to the expectation of such stimuli, and coordinates appropriate attentional, behavioral, affective and visceral autonomic responses to such stimuli. The **executive control network** (dlPFC, dorsolateral prefrontal cortex; vlPFC, ventrolateral prefrontal cortex; mPFC, medial prefrontal cortex; OFG, orbitofrontal gyrus) is activated during tasks involving executive functions such as attention, working memory, planning and response selection. Under normal circumstances, it exerts an inhibitory influence on the emotional arousal and the reward networks. The **reward network** (Nacc, nucleus accumbens; VTA-SN, ventral tegmental area – substantia nigra; CaN, caudate nucleus; Pal, pallidum) is a group of neural structures responsible for motivation, ‘wanting’ desire or craving for a reward. It is under inhibitory control by the executive control network. Its main neurotransmitter is dopamine. The **sensorimotor network** (Thal, thalamus; Put, putamen; pINS, posterior insula; S1/S2, primary/secondary sensory cortex; M1/M2, primary/secondary motor cortex) receives sensory input from the body, is important for awareness of body sensations and the generation of appropriate motor responses and behaviors.

The **Emotional arousal network** (ACC, anterior cingulate cortex; Hipp, hippocampus; Amyg, amygdala; sgACC, subgenual anterior cingulate cortex) is activated by perceived or real perturbation of the organism's homeostasis. Bidirectional arrows between brains depict reported bidirectional network interactions. Up and down arrows next to brains illustrate reported up and down regulation of the individual networks

Figure 3. Mechanisms in the homeostatic and hedonic systems leading to food addiction.

a | Diet-induced disinhibition of vagal and hypothalamic satiety mechanisms. A high fat, low fibre diet reduces the release of satiety hormones (GLP1, PYY, CCK) from enteroendocrine cells (EECs) in the gut by dietary fibre-derived short-chain fatty acids (SCFAs), leads to downregulation of receptors for satiety hormones molecules on vagal afferents innervating the EECs, and to a downregulation of the vagally-mediated satiety signaling to the arcuate nucleus of the hypothalamus (ARC). Hypothalamic receptors mediating the effect of other anorexigenic signals (leptin) reaching the ARC via the systemic circulation are also downregulated, resulting in an unrestrained effect of orexigenic signals (ghrelin, insulin, cortisol). Thus, chronic exposure to a high fat, low fibre diet downregulates the inhibitory mechanisms of homeostatic regulation of ingestive behaviors. **b** | Diet-induced changes in the extended reward system. According to the dopamine deficiency hypothesis, a reduction of dopaminergic stimulation of neurons in the NAc as a result of reduced dopamine release from the VTA, and a downregulation of dopamine receptors on NAc neurons, reduces the rewarding effects of ingested foods, and leads to craving and overconsumption of unhealthy food in an attempt to compensate for the reduced dopamine signaling. Chronic stress-induced CRF release and glucocorticoid levels also have an inhibitory effect on dopamine signaling. Up and downward arrows inside boxes illustrate reported up and downregulation of respective mechanisms.

Figure 4. Interactions between food, gut microbiota and intestinal permeability in the regulation of ingestive behavior.

Left panel: A healthy diet (high in fibre, low in fat and sugar) is associated with a high diversity of the gut microbiota, including an abundance of taxa involved in stimulating mucus production in humans and animal models.^{323,324} The combination of an intact mucus layer and tight intestinal epithelium results in an intact gut barrier. **Right panel:** An unhealthy diet (high in fat, sugar and low in fiber) is associated with a reduced microbial diversity, reduction of mucus stimulating microorganisms, reduction in mucus layer thickness and an increase in epithelial leakiness. This process results in reduced intestinal barrier function (leaky gut) and activation of the gut associated immune system by microbial products such as lipopolysaccharide (LPS). The combination of a leaky gut and an overabundance of inflammatory bacterial products is thought to result in elevated plasma levels of LPS and proinflammatory cytokines. This state of metabolic endotoxaemia has been shown to reduce central satiety mechanisms by influencing enteroendocrine secretion of the satiety hormones PYY, cholecystokinin and serotonin (5-HT), and by reducing the expression of anorexigenic peptide receptors and leptin receptors on vagal afferents and in the hypothalamus, respectively, leading to a disinhibition of satiety mechanisms. Up and downward arrows inside boxes represent reported up and downregulation of mechanisms and of microbiome measures. Modified with permission from Cani & Everard. *Mol. Nutr Food Res* 60:58-66; 2016 **OK, but would replace the last one with reduction of net mucus production**

Figure 5. Circular model of brain gut microbiome interactions in obesity and targets for intervention.

The interaction of genetic and epigenetic factors influences the balance between hedonic and homeostatic control of ingestive behavior, and the risk for the development of hedonic dominance. When exposed to ubiquitous food of high caloric density (fat, sugar) and low in fiber, predisposed individual will overconsume such foods, resulting in changes in the gut and the microbiome as shown in Fig. 3. The resulting change in gut to brain signaling can further compromises homeostatic regulation of food intake and reinforces the disinhibition of the reward system. Targets for intervention and therapeutic modalities include altered ingestive behavior (cognitive behavioral therapy, CBT; time restricted eating; dietary counseling), alterations of gut and microbiome (bariatric endoscopic and surgical treatment; faecal microbota transplantation (FMT); prebiotics and probiotics); altered gut to brain feedback (postbiotics such as butyrate, tryptophan derived compounds, including indoles, and other amino acid metabolites.) and altered reward system (centrally acting medications).