Images in Cardiovascular Disease

Anomalous Takeoff of Left Internal Mammary Artery from Thyrocervical Trunk Arising off the Aortic Arch

Ryan Malek , DO¹, Vasim Lala , DO¹, Adam Bykowski , DO¹, and Hiroshi Yamasaki , MD²

¹Department of Cardiology, Michigan State University, McLaren Macomb/Oakland Cardiovascular Disease Fellowship, Clinton Township, MI, USA

²Department of Cardiology, Wayne State University, Ascension St. John Hospital, Detroit, MI, USA

OPEN ACCESS

Received: Apr 22, 2020

Revised: Jun 5, 2020

Accepted: Jun 9, 2020

Address for Correspondence:

Ryan Malek, DO

Department of Cardiology, Michigan State University, McLaren Macomb/Oakland Cardiovascular Disease Fellowship, 41340 Fortuna Drive East, Clinton Township, MI 48038, USA.

E-mail: Ryan.Malek1@mclarenmeded.org

Copyright © 2021 Korean Society of

Echocardiography

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Ryan Malek

<https://orcid.org/0000-0003-0177-4393>

Vasim Lala

<https://orcid.org/0000-0002-4881-6709>

Adam Bykowski

<https://orcid.org/0000-0001-7563-6749>

Hiroshi Yamasaki

<https://orcid.org/0000-0001-7903-4200>

Conflict of Interest

The authors have no financial conflicts of interest.

The patient is 72 years old with a past medical history of atherosclerotic coronary disease and coronary artery bypass grafting with left internal mammary artery (LIMA) to the left anterior descending (LAD) presented with chest pain.

Coronary angiography was pursued to elucidate the etiology of the patient's symptoms. This revealed an angiographically patent right internal mammary artery. The LIMA graft to the LAD was unable to be selected for angiography at the expected proximal third of the subclavian artery (**Figure 1A**). Further angiography performed of distal segments of subclavian artery (**Figure 1B**). This was followed by selective thyrocervical trunk angiography (**Figure 1C**). This revealed an anomalous origin of the LIMA, which was anastomosed to the mid LAD (**Figure 1D**) and appeared to arise from the thyrocervical trunk.

Anatomically, the subclavian artery is divided into three segments. From the cadaver study by Henriquez-Pino and Prates,¹⁾ it was determined that the LIMA arises from the first segment of the subclavian artery in 92% of subjects. LIMA was found to arise from the second portion of the subclavian in 7% and from the third segment in 1%. Two reports described a LIMA takeoff directly from the aortic arch.²⁾³⁾ A case study described a dual anomalous origin of the LIMA off the thyrocervical trunk which arises from the third segment of the subclavian artery.⁴⁾ To our knowledge, there has been only one additional case describing a LIMA takeoff from the thyrocervical trunk arising from the aortic arch.⁵⁾

Figure 1. (A) Left subclavian artery angiography. (B) Distal left subclavian and axillary angiography. (C) Engagement of TCT with aortic arch origin. LIMA takeoff from TCT. (D) LIMA anastomosis to mid LAD. LAD: left anterior descending, LIMA: left internal mammary artery, TCT: thyrocervical trunk.

SUPPLEMENTARY MATERIAL

Movie 1

[Click here to view](#)

REFERENCES

1. Henriquez-Pino JA, Prates JC. An anatomical study of the proximal portion of the internal thoracic artery. *Arq Bras Cardiol* 1992;59:177-80.
[PUBMED](#)
2. Hailan A, Obaid D, Zaidi A, Smith D. Anomalous origin of left internal mammary artery arising directly from the aortic arch. *BMJ Case Rep* 2014;2014:bcr2014206583.
[PUBMED](#) | [CROSSREF](#)
3. Incani A, Lee JC, Poon KK, Crowhurst JA, Pincus M, Walters DL. Anomalous origin of the left internal mammary artery from the aortic arch. *JACC Cardiovasc Interv* 2012;5:e27-8.
[PUBMED](#) | [CROSSREF](#)

4. Kang K, Ferraro R, Kang G. Anomalous left internal mammary artery off the thyrocervical trunk and off the third segment of subclavian artery. *Indian Heart J* 2015;67:600-1.
[PUBMED](#) | [CROSSREF](#)
5. Nuqali A, Shafiq Q, Sheikh M. Dual anomalous origins of the thyrocervical trunk and left internal mammary artery. *JACC Cardiovasc Interv* 2016;9:e105-6.
[PUBMED](#) | [CROSSREF](#)