Published in final edited form as:

Ann Diagn Pathol. 2020 October ; 48: 151576. doi:10.1016/j.anndiagpath.2020.151576.

Clinicopathologic Features of Breast Cancer Reclassified as *HER2*-Amplified by Fluorescence In Situ Hybridization with Alternative Chromosome 17 Probes

Kristen C Blanton, B.S.¹, Allison M. Deal, Ph.D.², Kathleen A. Kaiser-Rogers, Ph.D.³, Carey K. Anders, M.D.⁴, Siobhan M. O'Connor, M.D.³, Johann D. Hertel, M.D.³, Benjamin C. Calhoun, M.D. Ph.D.^{2,3,*}

¹School of Medicine, University of North Carolina, Chapel Hill, NC, USA

²Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC, USA

³Department of Pathology and Laboratory Medicine, University of North Carolina, Chapel Hill, NC, USA

⁴Department of Medicine, Division of Medical Oncology, Duke Cancer Institute, Durham, NC, USA

Abstract

Objective: Dual probe fluorescence in situ hybridization (FISH) assays for determination of human epidermal growth factor receptor 2 (*HER2*) gene amplification in breast cancer provide a ratio of *HER2* to chromosome 17. The ratio may be skewed by copy number alterations (CNA) in the control locus for chromosome 17 (CEP17). We analyzed the impact of alternative chromosome 17 control probes on *HER2* status in a series of breast cancers with an emphasis on patients reclassified as amplified.

Methods: Breast cancer patients with equivocal *HER2* immunohistochemistry (2+) and equivocal FISH with CEP17 were included. Reclassification of *HER2* status was assessed with alternative chromosome 17 control probes (*LIS1* and *RARA*).

Results: A total of 40 unique patients with 46 specimens reflexed to alternative chromosome 17 probe testing were identified. The majority (>80%) of patients had pT1–2, hormone receptor-positive tumors with an intermediate or high combined histologic grade. There were 34/46 (73.9%) specimens reclassified as amplified with alternative probes, corresponding to 29/40 (72.5%) patients. Of the patients reclassified as amplified with alternative probes, 34.5% (10/29) received *HER2*-targeted therapy.

Conclusion: In this series, the majority of breast cancers tested with alternative chromosome 17 control probes under the 2013 ASCO/CAP Guidelines were converted to *HER2*-amplified. The treatment data and the clinicopathologic profile of the tumors suggest that most of these patients will neither receive nor benefit from *HER2*-targeted therapy. The findings support the

*Corresponding author: Benjamin C. Calhoun MD PhD, Department of Pathology and Laboratory Medicine, University of North Carolina - Chapel Hill, 160 North Medical Drive, Campus Box #7525, Chapel Hill, NC 27599-7525.

Prior Presentation: Preliminary data were presented in abstract form at the 2018 annual poster review for the Medical Student Research Program at the University of North Carolina at Chapel Hill.

recommendation in the 2018 ASCO/CAP HER2 Guidelines to discontinue the use of alternative chromosome 17 probes.

Keywords

Breast; HER2; FISH; ASCO/CAP Guidelines; chromosome 17

1. Introduction

Approximately 15–20% of invasive breast cancers are positive for human epidermal growth factor receptor 2 (*HER2*; *ERBB2*) gene amplification and protein over-expression.^{1–3} The humanized anti-HER2 monoclonal antibody trastuzumab is a highly effective treatment with a favorable side effect profile.⁴ Newer anti-HER2 agents, including pertuzumab, TDM1, and tucatinib, have also been shown to be effective in HER2-positive breast cancer.^{5–7} Reliable assessment of patients' eligibility for HER2-targeted therapies requires accurate pathologic evaluation of HER2 status.

The Guidelines for the interpretation and reporting of HER2 testing results in breast cancer were first published in 2007 by the American Society of Clinical Oncology and the College of American Pathologists (ASCO/CAP).⁸ The ASCO/CAP Guidelines were updated in 2013⁹ and again in 2018¹⁰. In all three versions, the Expert Panel did not make a specific recommendation for the use of immunohistochemistry (IHC) or fluorescence in situ hybridization (FISH) for first-line HER2 testing. However, most laboratories use IHC for initial HER2 testing with FISH as the reflex testing method for cases with equivocal (2+) IHC. FISH is used for first-line testing in some large academic and reference laboratories.

The most commonly used FISH assays are the US Food and Drug Administration (FDA)-approved dual probe assays. These assays employ fluorochrome-labeled probes for *HER2*, located on the long arm of chromosome 17, and a separate probe for chromosome 17 that hybridizes to alpha satellite sequences at the centromere of chromosome 17 (CEP17; *D17Z1*). *HER2* and CEP17 signals are counted in a subset of tumor cells to generate a *HER2*:CEP17 ratio. *HER2*:CEP17 ratios ≥ 2.0 are classified as amplified.^{9, 10} The chromosome 17 centromere probe/control probes are used to control for aneusomy (especially polysomy) for chromosome 17. However, several studies have shown that whole chromosome polysomy for chromosome 17 is uncommon in breast tumors.^{11, 12} Copy number alterations (CNA) in the DNA that encompass the hybridization site for the control probe may explain the increased number of chromosome 17 signals in many cases.¹³ The increased CEP17 copy number seen in some cases may skew the *HER2*:CEP17 ratio toward normal, while loss of one or more copies of this locus can produce a falsely positive *HER2*:CEP17 ratio. In both situations, the *HER2* FISH results may not appear to correlate with *HER2* mRNA or protein expression levels.^{13, 14}

The 2013 ASCO/CAP Guidelines⁹ recommended testing with an alternative chromosome 17 probe in cases with equivocal FISH results (i.e., ISH Group 4 with *HER2*:CEP17 ratio <2.0 and mean *HER2* copy number ≥ 4.0 and <6.0)^{10, 15}. However, the 2018 ASCO/CAP HER2 Guidelines essentially eliminated the equivocal FISH category and the use of alternative chromosome 17 probes.¹⁰ The recommendation for the use of alternative

chromosome probes in the 2013 Guidelines was based on the idea that these probes may “correct” the estimation of chromosome 17 copy number, but some studies have shown CNA in the hybridization sites for alternative probes as well as CEP17.^{16, 17} The goal of this study was to evaluate the impact of alternative chromosome 17 probes [*LIS1* (*PAFAH1B1*) and *RARA*] on the assessment of *HER2* gene amplification by FISH. The classification of these cases based on the updated 2018 ASCO/CAP Guidelines¹⁰ is emphasized and data on treatment with HER2-targeted therapies are also reported.

2. Materials and Methods

This study was approved by the University of North Carolina Institutional Review Board. The study population was selected from primary and metastatic breast cancer cases undergoing dual probe FISH for HER2 gene status at the UNC Cytogenetics Laboratory from 2013 to 2018. Cases with equivocal IHC and FISH results were reflexed to a second FISH assay with an alternative chromosome 17 probe, including *LIS1* and/or *RARA*. The distribution of cases assigned to each HER2 category by *LIS1* and/or *RARA* with the 2013 ASCP/CAP Guidelines was recorded.

During the study period, the HER2 status of breast carcinomas was determined by FISH using the PathVysion *HER2* DNA Probe Kit, (Abbott Molecular, Inc., Des Plaines, IL). Tumors were tested with FISH using a dual probe assay with a *HER2* gene probe labeled in Spectrum Orange and a chromosome 17 centromeric enumeration probe (CEP17; *D17Z1*) labeled in Spectrum Green. To confirm invasive carcinoma, 4- μ m-thick sections stained with hematoxylin and eosin (H&E) were evaluated, and separate cut sections that remained unstained were analyzed by *HER2*:CEP17 FISH. For all FISH cases, two areas of 30 cells each in a region designated by a breast pathologist as invasive carcinoma, are routinely analyzed by two independent observers. If the results of the initial analysis were equivocal, FISH with probes for *LIS1* and/or *RARA* was performed.

Surgical pathology and *HER2* FISH data were retrieved from the Anatomic Pathology Laboratory Information System (Epic Beaker, Epic Systems Corporation, Verona, WI) and reviewed by a dedicated breast pathologist (BCC). Patient demographics and treatment data were obtained from the electronic medical record in Epic (Epic Systems Corporation, Verona, WI).

3. Results

3.1 HER2 FISH with alternative chromosome 17 probes

A total of 46 specimens from 40 unique patients were tested with *HER2*:CEP17 FISH and reflexed to *LIS1* and/or *RARA* FISH (Table 1). A schematic representation of the positions of the CEP17, *LIS1*, *RARA* and *HER2* probes on chromosome 17 is shown in Figure 1. The specimens tested included 25 (54.3%) core biopsies of the primary tumor in the breast, 9 (19.5%) surgical excision breast specimens, and 12 (26.1%) specimens collected from local or distant metastatic sites. There were 6 patients with alternative probe results on 2 specimens and hormone receptor studies were not repeated on the second specimen.

HER2 immunohistochemistry (IHC) results were available for 43 specimens: 40 were equivocal (2+) and the scores for the remaining 3 cases were 1+ (Patients 6 and 20), and 0 (Patient 23). The indication for *HER2* FISH in 40/46 (87.0%) specimens was an equivocal (2+) IHC result. In 6/46 (13.0%) the indication for *HER2* FISH testing was an equivocal FISH result from an outside laboratory, conflicting IHC and FISH results from an outside laboratory, or a prior equivocal FISH result at our institution. *HER2* FISH was performed on 1 excision specimen (Patient 6 in Table 1) due to conflicting IHC results: 3+ at an outside laboratory on a pretreatment breast core biopsy and 1+ at UNC on an ipsilateral skin punch biopsy. Testing with alternative chromosome 17 probes was performed on 3 specimens (Patients 7, 10 and 40 in Table 1) based on an equivocal FISH result using the 2007 ASCO/CAP Guidelines (i.e., just prior to implementation of the 2013 Guidelines).

A total of 34/46 (73.9%) specimens tested with alternative chromosome 17 probes were reclassified as amplified using the 2013 ASCO/CAP guidelines (Figure 2). Of these reclassified specimens, all 34 were amplified based on reflex testing with the *LIS1* probe and 3 of these 34 cases also were amplified with the *RARA* probe. In 1 of 46 specimens, reflex testing with alternative probes was performed to investigate an unusual hybridization pattern for CEP17. In this single case, there were few or no CEP17 signals in the tumor cells, but the expected disomic pattern of hybridization in adjacent normal tissue was preserved (patient 9 in Table 1). In 41 (91.1%) of the remaining 45 specimens, the chromosome 17 copy number obtained with *LIS1* was lower (closer to the disomic number 2.0) than the number obtained with CEP17. However as indicated above, in only 34 of these cases was the number of chromosome 17 control probe signals low enough to yield a *HER2*:control locus ratio of greater than or equal to 2.

3.2 Clinicopathological features

There was no significant difference in the clinicopathologic features of those patients whose tumors were reclassified from equivocal to amplified with alternative chromosome 17 probes (n=29) when compared to the rest of the cohort (n=11) (all $p > 0.09$). The median age was 57.4 years (range: 28–90). Of the 31/40 patients with available pT or pN data, 27/31 (87.1%) were pT1 or pT2, 15 (48.4%) were lymph node-negative (pN0), 7 (22.6%) had 1–3 positive lymph nodes (pN1) and 2 (6.5%) had 4–9 positive lymph nodes (pN2). Of the patients with alternative probe testing, 33/40 (82.5%) had hormone receptor-positive tumors and 35/40 (87.5%) had an intermediate or high combined histologic grade.

3.3 Treatment and follow-up

The results for each pathologic specimen described above correspond to re-classification of 29/40 (72.5%) patients as amplified. Of the 36/40 (90.0%) patients with available treatment data, 15/36 (41.7%) received HER2-targeted therapy, 10 of whom were reclassified as amplified with alternative chromosome 17 probes. Follow-up data was available for 37/40 patients at a median of 16 months (range: 0.3 – 62 months). At the time of last follow-up, 17 patients were alive with disease, 16 had no evidence of disease (NED) and 4 were deceased. All 4 of the patients who were deceased at last follow up had tumors that were reclassified as amplified with alternative chromosome 17 probes. Three (3) of the deceased patients were

referred to our institution for evaluation and treatment of metastatic disease and 2 had received HER2-targeted therapy.

4. Discussion

The 2013 ASCO/CAP HER2 Guidelines classified cases based on a combination of the *HER2*:CEP17 ratio and the mean *HER2* copy number, and allowed the use of alternative chromosome 17 probes.⁹ The emphasis on *HER2* signal number and alternative chromosome 17 probes represented attempts to address issues related to the potential for a skewed *HER2*:CEP17 ratio in cases with CEP17 CNA. One of the most significant changes in the 2013 guidelines was the classification of cases with a mean *HER2* copy number ≥ 4.0 and <6.0 and a *HER2*:CEP17 ratio <2.0 as equivocal.⁹ Several subsequent studies showed that the new equivocal category and the use of alternative chromosome 17 probes resulted in an increase in the proportion of equivocal and amplified *HER2* FISH results, particularly among cases with equivocal (2+) IHC results and CEP17 CNA.^{17–32} However, copy number variations also have been reported for the most commonly used alternative chromosome 17 control probes [i.e., *RAI1*, *LIS1* (*PAFAH1B1*), *RARA*, and *TP53*].^{16, 17} Therefore, it is not clear which probe provides the most accurate reflection of the chromosome 17 copy number and ultimately the most accurate *HER2*:control ratio.

The emphasis on *HER2* copy number in the 2013 Guidelines effectively created 5 categories of FISH results.¹⁵ The group of FISH cases defined as equivocal in the 2013 Guidelines⁹ corresponds to ISH Group 4 in the scheme developed by Press et al. and adopted in the most recent 2018 ASCO/CAP HER2 FISH Guidelines.^{10, 15, 33} Based on the data from patients screened for enrollment in three Breast Cancer International Research Group (BCIRG) clinical trials¹⁵ and the NSABP-B47 trial of “Trastuzumab for Women with HER2-low Breast Cancer”,³⁴ patients in ISH Group 4 are unlikely to benefit from HER2-targeted therapy. However, it should be noted that patients with any *HER2* FISH result with a mean *HER2* copy number ≥ 4.0 may have been excluded from NSABP-B47.²⁰

In the majority of cases in this series, the indication for *HER2* FISH with an alternative chromosome 17 probe was an equivocal result (as defined in the 2013 Guidelines) obtained from a dual probe assay using CEP17. When patients with known metastases are excluded, the majority in this series had pathological Stage I-II, intermediate to high combined histologic grade, hormone receptor-positive, lymph node-negative tumors. The clinicopathological characteristics of these tumors are not typical for *HER2*-amplified tumors but they are similar to the “excess” equivocal cases identified using the 2013 Guidelines.^{19, 25, 28, 35} These tumors also were unlikely to have positive (3+) IHC results, similar to the data from other studies of the equivocal category in the 2013 Guidelines.^{15, 18, 19, 21, 22} The clinicopathologic features and *HER2* IHC data may account for the fact that approximately 60% of patients in this series who were re-classified as amplified with alternative probes were not treated with HER2-targeted therapy. In studies using gene expression analysis^{36, 37} and surrogate immunohistochemistry markers²⁵ for intrinsic molecular subtypes, the majority of *HER2* FISH equivocal tumors (as defined in the 2013 Guidelines) are luminal-type, and not *HER2*-enriched. In the series reported by Desai et al.,

most of the tumors reclassified as amplified with alternative chromosome 17 probes were Luminal A or B subtype, indicating that these are not *HER2*-driven tumors.³⁶

Three specimens in this series were tested with alternative chromosome 17 probes based on an equivocal FISH result using the 2007 Guidelines, just prior to the implementation of the 2013 Guidelines (Patients 7, 10 and 40 in Table 1). The core biopsies for Patients 7 and 40, with mean *HER2* copy numbers of 6.4 and *HER2*:CEP17 ratios of 1.8, would be classified as amplified using the 2013 Guidelines and presumably would be classified as amplified (ISH Group 3) in the 2018 Guidelines. The excision specimen from Patient 10, with a mean *HER2* copy number of 5.8 and a *HER2*:CEP17 ratio of 2.0 would be classified as amplified in the 2013 Guidelines and amplified (ISH Group 1) in the 2018 Guidelines. It should be noted that Patient 10 also had a core biopsy that was converted to amplified with alternative chromosome 17 probes. That specimen, with a mean *HER2* copy number of 4.8 and a *HER2*:CEP17 ratio of 1.3, would have been classified as equivocal by the 2013 Guidelines. Using the 2018 Guidelines, it would correspond to ISH Group 4 and would be classified as not amplified in the absence of a 3+ *HER2* IHC result. Taking all of this into consideration, if the 3 specimens classified as equivocal under the 2007 Guidelines were excluded from analysis, the rate of conversion to amplified (with alternative probes and the 2013 Guidelines) would remain high: 31/43 (72.1%) specimens and 27/38 (71.5%) patients.

The use of alternative chromosome 17 probes in this series increased the number of patients potentially eligible for *HER2*-targeted therapy. However, there is no well-established benefit from *HER2*-targeted therapy in this patient population and many of these “amplified” cases may in fact represent false-positives.³⁸ In a study of over 10,000 patients screened for enrollment in three BCIRG trials, there was no significant difference in DFS or OS when the patients in ISH Group 4 were compared to those with clearly negative FISH results (i.e., ISH Group 5; *HER2*:CEP17 ratio <2.0 and a mean *HER2* copy number of <4.0).¹⁵ Based on the outcomes data and the absence of *HER2* protein overexpression in ISH Group 4, Press et al. concluded that tumors with a *HER2*:CEP17 ratio <2.0 and a mean *HER2* copy number 4.0 and <6.0 should be classified as not amplified.¹⁵ This recommendation was incorporated into the updated 2018 ASCO/CAP *HER2* Guidelines.¹⁰

Limitations of this study include the single-institution retrospective design and limited sample size. Strengths of the study include the focus on tumors reclassified as amplified with alternative probes, the detailed characterization of the FISH and IHC findings, and thorough clinicopathologic characterization with data on treatment with *HER2*-targeted therapy. Although the sample size is limited, the detailed characterization of this group of patients provides an accurate reflection of the impact of alternative chromosome 17 probes in complex breast cancer cases at a tertiary referral center. No prospective clinical trials using alternative chromosome 17 probes to select patients for *HER2*-targeted therapy have been reported.

In summary, *HER2* FISH with alternative chromosome 17 probes reclassified the tumors in 29/40 (72.5%) patients as amplified. The majority of these patients were not treated with *HER2*-targeted therapy. The best available retrospective data derived from large clinical trials indicate that this group of patients is unlikely to benefit from trastuzumab.¹⁵ Our

findings support the classification of these cases as not amplified in the 2018 ASCO/CAP HER2 Guidelines.¹⁰

Acknowledgments

Research Support: K. Blanton was supported by a short-term research training (STRT) grant from the National Institute of Diabetes, Digestive and Kidney diseases (NIDDK), grant number T35-DK007386 (University of North Carolina at Chapel Hill, School of Medicine, Office of Medical Student Research).

Conflict of Interest Disclosures: Dr. Anders receives research funding from PUMA, Lilly, Merck, Seattle Genetics, Nektar, Tesaro, G1-Therapeutics; compensation as a consultant for Genentech (1/2019-), Eisai (1/2019-), IPSEN (2/2019 -), Seattle Genetics (11/15/2019 – 11/15/2020); and royalties from UpToDate and Jones and Bartlett. Dr. Calhoun is a member of the Oncology Advisory Board for Luminex Corporation. The other authors have no relevant financial disclosures.

Protection of Human and Animal Subjects: This study was approved by the Institutional Review Board of the University of North Carolina at Chapel Hill

References

1. Lal P, Salazar PA, Hudis CA, Ladanyi M, Chen B. HER-2 testing in breast cancer using immunohistochemical analysis and fluorescence in situ hybridization: a single-institution experience of 2,279 cases and comparison of dual-color and single-color scoring. *Am J Clin Pathol*. 2004;121:631–636. [PubMed: 15151202]
2. Varga Z, Noske A, Ramach C, Padberg B, Moch H. Assessment of HER2 status in breast cancer: overall positivity rate and accuracy by fluorescence in situ hybridization and immunohistochemistry in a single institution over 12 years: a quality control study. *BMC Cancer*. 2013;13:615. [PubMed: 24377754]
3. Tubbs RR, Hicks DG, Cook J, et al. Fluorescence in situ hybridization (FISH) as primary methodology for the assessment of HER2 Status in adenocarcinoma of the breast: a single institution experience. *Diagn Mol Pathol*. 2007;16:207–210. [PubMed: 18043283]
4. Pegram MD. Treating the HER2 pathway in early and advanced breast cancer. *Hematol Oncol Clin North Am*. 2013;27:751–765, viii. [PubMed: 23915743]
5. Saini KS, Azim HA, Metzger-Filho O, et al. Beyond trastuzumab: New treatment options for HER2-positive breast cancer. *The Breast*. 2011;20:S20–S27.
6. Verma S, Miles D, Gianni L, et al. Trastuzumab Emtansine for HER2-Positive Advanced Breast Cancer. *New England Journal of Medicine*. 2012;367:1783–1791.
7. Murthy RK, Loi S, Okines A, et al. Tucatinib, Trastuzumab, and Capecitabine for HER2-Positive Metastatic Breast Cancer. *New England Journal of Medicine*. 2019;382:597–609.
8. Wolff AC, Hammond ME, Schwartz JN, et al. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for human epidermal growth factor receptor 2 testing in breast cancer. *J Clin Oncol*. 2007;25:118–145. [PubMed: 17159189]
9. Wolff AC, Hammond ME, Hicks DG, et al. Recommendations for Human Epidermal Growth Factor Receptor 2 Testing in Breast Cancer: American Society of Clinical Oncology/College of American Pathologists Clinical Practice Guideline Update. *J Clin Oncol*. 2013;31:3997–4013. [PubMed: 24101045]
10. Wolff AC, Hammond MEH, Allison KH, et al. Human Epidermal Growth Factor Receptor 2 Testing in Breast Cancer: American Society of Clinical Oncology/College of American Pathologists Clinical Practice Guideline Focused Update. *J Clin Oncol*. 2018;36:2105–2122. [PubMed: 29846122]
11. Marchio C, Lambros MB, Gugliotta P, et al. Does chromosome 17 centromere copy number predict polysomy in breast cancer? A fluorescence in situ hybridization and microarray-based CGH analysis. *J Pathol*. 2009;219:16–24. [PubMed: 19670217]
12. Moelans CB, de Weger RA, van Diest PJ. Absence of chromosome 17 polysomy in breast cancer: analysis by CEP17 chromogenic in situ hybridization and multiplex ligation-dependent probe amplification. *Breast Cancer Res Treat*. 2010;120:1–7. [PubMed: 19760503]

13. Hanna WM, Ruschhoff J, Bilous M, et al. HER2 in situ hybridization in breast cancer: clinical implications of polysomy 17 and genetic heterogeneity. *Mod Pathol*. 2014;27:4–18.
14. Downs-Kelly E, Yoder BJ, Stoler M, et al. The influence of polysomy 17 on HER2 gene and protein expression in adenocarcinoma of the breast: a fluorescent in situ hybridization, immunohistochemical, and isotopic mRNA in situ hybridization study. *Am J Surg Pathol*. 2005;29:1221–1227. [PubMed: 16096413]
15. Press MF, Sauter G, Buyse M, et al. HER2 Gene Amplification Testing by Fluorescent In Situ Hybridization (FISH): Comparison of the ASCO-College of American Pathologists Guidelines With FISH Scores Used for Enrollment in Breast Cancer International Research Group Clinical Trials. *J Clin Oncol*. 2016.
16. Tse CH, Hwang HC, Goldstein LC, et al. Determining true HER2 gene status in breast cancers with polysomy by using alternative chromosome 17 reference genes: implications for anti-HER2 targeted therapy. *J Clin Oncol*. 2011;29:4168–4174. [PubMed: 21947821]
17. Jang MH, Kim EJ, Kim HJ, Chung YR, Park SY. Assessment of HER2 status in invasive breast cancers with increased centromere 17 copy number. *Breast Cancer Res Treat*. 2015;153:67–77. [PubMed: 26223814]
18. Shah MV, Wiktor AE, Meyer RG, et al. Change in Pattern of HER2 Fluorescent in Situ Hybridization (FISH) Results in Breast Cancers Submitted for FISH Testing: Experience of a Reference Laboratory Using US Food and Drug Administration Criteria and American Society of Clinical Oncology and College of American Pathologists Guidelines. *J Clin Oncol*. 2016.
19. Muller KE, Marotti JD, Memoli VA, Wells WA, Tafe LJ. Impact of the 2013 ASCO/CAP HER2 Guideline Updates at an Academic Medical Center That Performs Primary HER2 FISH Testing: Increase in Equivocal Results and Utility of Reflex Immunohistochemistry. *Am J Clin Pathol*. 2015;144:247–252. [PubMed: 26185309]
20. Long TH, Lawce H, Durum C, et al. The New Equivocal: Changes to HER2 FISH Results When Applying the 2013 ASCO/CAP Guidelines. *Am J Clin Pathol*. 2015;144:253–262. [PubMed: 26185310]
21. Varga Z, Noske A. Impact of Modified 2013 ASCO/CAP Guidelines on HER2 Testing in Breast Cancer. One Year Experience. *PLoS One*. 2015;10:e0140652. [PubMed: 26473483]
22. Stoss OC, Scheel A, Nagelmeier I, et al. Impact of updated HER2 testing guidelines in breast cancer--re-evaluation of HERA trial fluorescence in situ hybridization data. *Mod Pathol*. 2015;28:1528–1534. [PubMed: 26403781]
23. Singh K, Tantravahi U, Lomme MM, Pasquariello T, Steinhoff M, Sung CJ. Updated 2013 College of American Pathologists/American Society of Clinical Oncology (CAP/ASCO) guideline recommendations for human epidermal growth factor receptor 2 (HER2) fluorescent in situ hybridization (FISH) testing increase HER2 positive and HER2 equivocal breast cancer cases; retrospective study of HER2 FISH results of 836 invasive breast cancers. *Breast Cancer Res Treat*. 2016;157:405–411. [PubMed: 27180259]
24. Lim TH, Lim AS, Thike AA, Tien SL, Tan PH. Implications of the Updated 2013 American Society of Clinical Oncology/College of American Pathologists Guideline Recommendations on Human Epidermal Growth Factor Receptor 2 Gene Testing Using Immunohistochemistry and Fluorescence In Situ Hybridization for Breast Cancer. *Arch Pathol Lab Med*. 2016;140:140–147. [PubMed: 26910218]
25. Bethune GC, Veldhuijzen van Zanten D, MacIntosh RF, et al. Impact of the 2013 American Society of Clinical Oncology/College of American Pathologists guideline recommendations for human epidermal growth factor receptor 2 (HER2) testing of invasive breast carcinoma: a focus on tumours assessed as 'equivocal' for HER2 gene amplification by fluorescence in-situ hybridization. *Histopathology*. 2015;67:880–887. [PubMed: 25913507]
26. Fan YS, Casas CE, Peng J, et al. HER2 FISH classification of equivocal HER2 IHC breast cancers with use of the 2013 ASCO/CAP practice guideline. *Breast Cancer Res Treat*. 2016;155:457–462. [PubMed: 26895325]
27. Polonia A, Leita D, Schmitt F. Application of the 2013 ASCO/CAP guideline and the SISH technique for HER2 testing of breast cancer selects more patients for anti-HER2 treatment. *Virchows Arch*. 2016;468:417–423. [PubMed: 26754674]

28. Sapino A, Maletta F, Verdun di Cantogno L, et al. Gene status in HER2 equivocal breast carcinomas: impact of distinct recommendations and contribution of a polymerase chain reaction-based method. *Oncologist*. 2014;19:1118–1126. [PubMed: 25323485]
29. Griffin BB, Pincus JL, Siziopikou KP, Blanco LZ Jr. Double-Equivocal HER2 Invasive Breast Carcinomas: Institutional Experience and Review of Literature. *Arch Pathol Lab Med*. 2018;142:1511–1516. [PubMed: 29595316]
30. Pu X, Shi J, Li Z, Feng A, Ye Q. Comparison of the 2007 and 2013 ASCO/CAP evaluation systems for HER2 amplification in breast cancer. *Pathol Res Pract*. 2015;211:421–425. [PubMed: 25818873]
31. Donaldson AR, Shetty S, Wang Z, et al. Impact of an alternative chromosome 17 probe and the 2013 American Society of Clinical Oncology and College of American Pathologists guidelines on fluorescence in situ hybridization for the determination of HER2 gene amplification in breast cancer. *Cancer*. 2017;123:2230–2239. [PubMed: 28192599]
32. Hui L, Geiersbach KB, Downs-Kelly E, Gulbahce HE. RAI1 Alternate Probe Identifies Additional Breast Cancer Cases as Amplified Following Equivocal HER2 Fluorescence In Situ Hybridization Testing: Experience From a National Reference Laboratory. *Arch Pathol Lab Med*. 2017;141:274–278. [PubMed: 27959582]
33. Press MF, Villalobos I, Santiago A, et al. Assessing the New American Society of Clinical Oncology/College of American Pathologists Guidelines for HER2 Testing by Fluorescence In Situ Hybridization: Experience of an Academic Consultation Practice. *Arch Pathol Lab Med*. 2016;140:1250–1258. [PubMed: 27081877]
34. Fehrenbacher L, Cecchini RS, Geyer CE Jr., et al. NSABP B-47/NRG Oncology Phase III Randomized Trial Comparing Adjuvant Chemotherapy With or Without Trastuzumab in High-Risk Invasive Breast Cancer Negative for HER2 by FISH and With IHC 1+ or 2. *J Clin Oncol*. 2019;Jco1901455.
35. Ji H, Xuan Q, Nanding A, Zhang H, Zhang Q. The Clinicopathologic and Prognostic Value of Altered Chromosome 17 Centromere Copy Number in HER2 Fish Equivocal Breast Carcinomas. *PLoS One*. 2015;10:e0132824. [PubMed: 26161550]
36. Desai NV, Torous V, Parker J, et al. Intrinsic molecular subtypes of breast cancers categorized as HER2-positive using an alternative chromosome 17 probe assay. *Breast Cancer Res*. 2018;20:75. [PubMed: 29996866]
37. Marchio C, Dell’Orto P, Annaratone L, et al. The Dilemma of HER2 Double-equivocal Breast Carcinomas: Genomic Profiling and Implications for Treatment. *Am J Surg Pathol*. 2018;42:1190–1200. [PubMed: 29975246]
38. Press MF, Seoane JA, Curtis C, et al. Assessment of ERBB2/HER2 Status in HER2-Equivocal Breast Cancers by FISH and 2013/2014 ASCO-CAP Guidelines. *JAMA Oncol*. 2019;5:366–375. [PubMed: 30520947]

Highlights

- Most breast cancers with equivocal HER2 FISH results under the 2013 ASCO/CAP HER2 Guidelines are converted to HER2-amplified with alternative chromosome 17 probes
- The available treatment data and the clinicopathologic profile of these tumors suggest that most of these patients will not receive or benefit from HER2-targeted therapy
- The findings support the recommendation in the 2018 ASCO/CAP HER2 Guidelines to discontinue the use of alternative chromosome 17 probes

Figure 1:
Schematic of Chromosome 17 and Associated Probe Locations. Pictured are the probe locations for the human epidermal growth factor receptor-2 (*HER2*), chromosome 17 centromere enumerator (CEP17; D17Z1), lissencephaly (*LIS1*), and retinoic acid receptor- α (*RARA*) loci.

Figure 2:

Representative images of an equivocal *HER2* FISH result reclassified as amplified with alternative chromosome 17 probes and the 2013 ASCO/CAP Guidelines (patient 34 in Table 1). Dual-probe FISH yielded a mean *HER2* (red) copy number of 4.1, a mean CEP17 (green) copy number of 2.8 and a *HER2*:CEP17 ratio of 1.5 (A). Alternative chromosome 17 probes demonstrated a mean number of 3.9 *RARA* (green) and 1.6 *PAFAH1B1/LIS1* (red) signals per cell; CEP17 (aqua) (B). The *HER2*:*PAFAH1B1/LIS1* ratio was 2.5 (amplified) and the *HER2*:*RARA* ratio was 1.1 (not amplified).

Table 1. Clinicopathologic Features of Breast Cancer with HER2 FISH and Reflex testing with Alternative Chromosome 17 probes (n = 40 patients)

Pt	Specimen Type	IHC	CEP17 Probe				HER2 Status	Alternative Probes				HER2 Status		Hormone Receptor Status		HER2 Therapy
			Avg. copy number and ratio					Avg. copy number and ratio						δ		
			HER2	CEP17	Ratio			LIS1	Ratio	RARA	Ratio			ER	PR	
1	Met	2+	4.0	2.5	1.6	Equivocal	2.6	1.5	3.9	1.0	Equivocal	Positive	Positive	Positive		
2	Met	2+	4.9	2.6	1.9	Equivocal	1.7	2.9	4.6	1.1	Amplified	Positive	Negative	Negative	Yes	
3	Core	2+	5.7	4.0	1.4	Equivocal	2.3	2.5	7.2	0.8	Amplified	Positive	Positive	Positive		
4	Met	2+	5.5	3.9	1.4	Equivocal	1.7	3.2	3.4	1.6	Amplified	Negative	Negative	Negative		
5	Core	2+	4.8	2.9	1.6	Equivocal	2.1	2.2	4.4	1.1	Amplified	Positive	Positive	Positive	Yes	
6	Exc	1+	5.3	4.8	1.1	Equivocal	3.4	1.6	3.5	1.5	Equivocal	Negative	Negative	Negative	Yes	
7	Core	2+	6.4	3.5	1.8	Equivocal ²	3.0	2.1	NR	NR	Amplified	Positive	Positive	Positive		
8	Core	2+	4.3	2.3	1.8	Equivocal	3.6	1.2	3.8	1.1	Equivocal	Positive	Positive	Positive	Yes	
9	Met	2+	4.2	NR ³	NR ³	Equivocal	2.9	1.4	3.8	1.1	Equivocal	Negative	Negative	Negative	Yes	
10	Core	2+	4.8	3.5	1.3	Equivocal	1.8	2.6	NR	NR	Amplified	Positive	Positive	Negative		
10	Exc	2+	5.8	2.9	2.0	Equivocal ²	1.8	3.2	NR	NR	Amplified	NR	NR	NR		
11	Core	2+	5.6	3.9	1.4	Equivocal	3.6	1.6	5.9	1.0	Equivocal	Positive	Positive	Positive	Yes	
12	Core	2+	4.9	2.9	1.7	Equivocal	1.6	3.1	3.7	1.3	Amplified	Positive	Positive	Positive		
13	Core	2+	4.7	2.6	1.8	Equivocal	1.8	2.6	3.9	1.2	Amplified	Positive	Positive	Positive		
14	Core	2+	4.3	3.1	1.4	Equivocal	1.8	2.4	3.8	1.1	Amplified	Negative	Negative	Positive		
15	Core	2+	5.2	3.6	1.5	Equivocal	1.8	2.9	4.0	1.3	Amplified	Positive	Positive	Positive		
16	Met	2+	4.3	3.7	1.2	Equivocal	1.8	2.4	4.3	1.0	Amplified	Negative	Negative	Negative	Yes	
17	Met	2+	4.2	2.7	1.6	Equivocal	1.9	2.2	4.8	0.9	Amplified	Positive	Positive	Negative		
18	Core	2+	4.5	2.6	1.7	Equivocal	2.1	2.1	2.2	2.0	Amplified	Negative	Negative	Negative	Yes	
19	Core	2+	4.8	3.5	1.4	Equivocal	1.9	2.6	4.2	1.1	Amplified	Positive	Positive	Positive	Yes	
20	Core	1+ ⁴	4.0	2.7	1.5	Equivocal	2.1	1.9	4.2	0.9	Equivocal	Positive	Positive	Positive		
21	Met	2+	4.3	3.4	1.2	Equivocal	3.6	1.2	3.9	1.1	Equivocal	Positive	Positive	Positive		
22	Core	2+	4.4	3.9	1.1	Equivocal	2.0	2.2	4.1	1.1	Amplified	Positive	Positive	Positive	Yes	

Pt	Specimen Type	IHC	CEP17 Probe				HER2 Status	Alternative Probes				HER2 Status	Hormone Receptor Status		HER2 Therapy
			Avg. copy number and ratio					Avg. copy number and ratio					δ		
			HER2	CEP17	Ratio			LIS1	Ratio	RARA	Ratio		ER	PR	
23	Core	2+	4.1	4.2	1.0		Equivocal	4.0	1.0	3.8	1.1	Equivocal	Negative	Negative	
23	Exc	0	4.3	3.3	1.3		Equivocal	3.8	1.1	3.5	1.2	Equivocal	NR	NR	
24	Exc	NR ⁵	4.8	3.8	1.3		Equivocal	1.5	3.2	4.9	1.0	Amplified	Positive	Positive	
25	Core	2+	5.0	2.7	1.8		Equivocal	1.6	3.2	5.3	0.9	Amplified	Positive	Positive	Yes
26	Exc	2+	5.5	4.1	1.4		Equivocal	2.5	2.2	5.3	1.0	Amplified	Positive	Positive	
27	Core	2+	4.9	3.0	1.7		Equivocal	1.8	2.7	4.7	1.1	Amplified	Positive	Positive	
27	Exc	NR	4.6	2.6	1.8		Equivocal	1.7	2.7	5.4	0.8	Amplified	NR	NR	
28	Exc	2+	4.4	3.0	1.5		Equivocal	2.2	2.0	4.7	1.0	Amplified	Positive	Positive	
28	Met	2+	5.3	3.8	1.4		Equivocal	1.7	3.1	5.3	1.0	Amplified	NR	NR	
29	Met	2+	4.3	3.9	1.1		Equivocal	2.6	1.7	3.3	1.3	Equivocal	Positive	Positive	
30	Core	2+	4.8	3.3	1.5		Equivocal	1.8	2.7	4.1	1.2	Amplified	Positive	Positive	Yes
30	Exc	2+	5.1	3.5	1.4		Equivocal	1.6	3.1	4.7	1.1	Amplified	NR	NR	
31	Core	2+	5.0	3.2	1.6		Equivocal	1.8	2.9	4.8	1.0	Amplified	Positive	Negative	
32	Met	2+	5.0	2.8	1.8		Equivocal	2.7	1.9	4.7	1.1	Equivocal	Positive	Positive	Yes
33	Core	2+	5.1	3.1	1.6		Equivocal	1.5	3.3	4.2	1.2	Amplified	Positive	Positive	
33	Exc	NR	5.0	3.7	1.4		Equivocal	1.7	2.9	4.9	1.0	Amplified	NR	NR	
34	Core	2+	4.1	2.8	1.5		Equivocal	1.6	2.5	3.9	1.1	Amplified	Positive	Positive	
35	Core	2+	3.1	2.7	1.1		Equivocal	1.5	2.1	3.0	1.1	Amplified	Positive	Positive	
36	Core	2+	4.0	3.1	1.3		Equivocal	1.9	2.1	1.9	2.2	Amplified	Positive	Positive	Yes
37	Met	2+	4.5	3.9	1.2		Equivocal	3.2	1.4	4.0	1.1	Equivocal	Negative	Negative	
38	Core	2+	4.9	3.9	1.3		Equivocal	1.9	2.6	4.4	1.1	Amplified	Positive	Positive	
39	Met	2+	4.4	2.6	1.7		Equivocal	1.8	2.5	3.4	1.3	Amplified	Positive	Negative	Yes
40	Core	2+	6.4	3.5	1.8		Equivocal ²	1.9	3.5	NR	NR	Amplified	Positive	Positive	

¹Patient 6: Positive (3+) HER2 IHC on a breast core biopsy at an outside laboratory. Negative (1+) HER2 IHC on a skin punch biopsy at UNC. HER2 FISH on a post-treatment breast specimen was equivocal and reflexed to alternative probe testing.

²Patients 7, 10, 40: Original results were reported prior to implementation of 2013 Guidelines. Equivocal based on 2007 Guidelines and classified as amplified by 2013 Guidelines.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

³ Patient 9: Absent CEP17 signals in tumor cells; disomic hybridization pattern for CEP17 in adjacent normal tissue. CEP17 and HER2:CEP17 could not be reported (NR) for this specimen.

⁴ Patient 20: Prior negative (1+) IHC and equivocal HER2 FISH at an outside laboratory. Repeat HER2 FISH at UNC was equivocal, prompting alternative probe testing.

⁵ Patient 24: Amplified HER2 FISH at an outside institution based on HER2 copy number (mean HER2 6.4, mean CEP17 4.7) and an equivocal HERmark result. Alternative probe testing was performed at our institution to exclude co-amplification

⁶ For patients with more than one specimen tested with alternative chromosome 17 probes, hormone receptor studies were not repeated on the subsequent specimens (i.e., NR).

Abbreviations: Pt, patient; IHC, Immunohistochemistry; CEP17, chromosome 17 centromere enumerator probe; HER2, human epidermal growth factor receptor 2; ER, estrogen receptor; PR, progesterone receptor; Avg, Average; RARA, retinoic acid receptor- α ; LJS1, lissencephaly gene; Exc, Excision; Met, Metastasis; NR, Not performed/not reported.