Published in final edited form as:

Public Health Nutr. 2021 October ; 24(14): 4537–4545. doi:10.1017/S1368980020003936.

B-vitamins and metabolic syndrome in Mesoamerican children and their adult parents

Claudia R Villatoro-Santos¹, Manuel Ramirez-Zea², Eduardo Villamor^{1,*}, Nine Mesoamerican Countries Metabolic Syndrome (NiMeCoMeS) Study group

¹Department of Epidemiology, University of Michigan School of Public Health, Ann Arbor, MI, USA

²INCAP Research Center for the Prevention of Chronic Diseases, Institute of Nutrition of Central America and Panama, Guatemala City, Guatemala

Abstract

Objective.—To examine the associations between vitamins of the methionine-homocysteine cycle (B6, B12, and folate) and homocysteine (Hcys) with metabolic syndrome (MetS) among Mesoamerican children and their adult parents.

Design.—We conducted a cross-sectional study. Exposures were plasma vitamin B6 and B12 concentrations, erythrocyte folate, and plasma Hcys. In children, the outcome was a continuous metabolic risk score calculated through sex- and age-standardization of waist circumference, the homeostatic model assessment for insulin resistance, mean arterial pressure (MAP), serum high-density lipoprotein (HDL) cholesterol, and serum triglycerides. In parents, the outcome was the prevalence of MetS according to the Adult Treatment Panel III criteria. We estimated mean differences in the metabolic risk score and prevalence ratios (PR) of MetS between quartiles of the exposures using multivariable-adjusted linear and Poisson regression models, respectively.

Setting.—Capital cities of Belize, Guatemala, El Salvador, the Dominican Republic, Honduras, Nicaragua, Panama, Costa Rica, and Chiapas State in Mexico.

Participants.—237 school-age children and 524 parents.

Results.—Among children, vitamin B12 was inversely associated with the metabolic risk score (quartiles 4–1 adjusted difference = -0.13 ; 95% CI: -0.21 , -0.04 ; P , trend=0.008) through MAP, HDL cholesterol, and triglycerides. By contrast, folate was positively associated with the metabolic risk score (quartiles 4–1 adjusted difference = 0.11 ; 95% CI: 0.01 , 0.20 ; P , trend=0.02).

*Corresponding author. villamor@umich.edu.

Authorship: E.V. designed the research. M.R.Z. conducted the research. CV performed the statistical analysis. C.V and E.V wrote the manuscript and had the primary responsibility of the final content. All authors have read and approved the final version of the manuscript.

Publisher's Disclaimer: This is an Accepted Manuscript for Public Health Nutrition as part of the Cambridge Coronavirus Collection. This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process.

Conflict of Interest: None.

Ethical Standards Disclosure: This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving research study participants were approved by the Institutional Review Boards (IRB) of collaborating institutions in each country and by the University of Michigan Health and Behavioral Sciences IRB. Written parental informed consent and children's assent was obtained from all participants.

In adults, vitamin B6 was inversely associated with MetS prevalence whereas vitamin B12 and folate were positively related to this outcome.

Conclusions.—Vitamins of the methionine-homocysteine cycle are associated with metabolic syndrome in different directions. The associations differ between children and adults.

Keywords

Metabolic syndrome; vitamin B6; vitamin B12; erythrocyte folate; homocysteine

INTRODUCTION

Cardiovascular disease is the leading cause of death worldwide ⁽¹⁾. The metabolic syndrome (MetS) is a cluster of independent risk factors for cardiovascular disease, type 2 diabetes, and mortality that includes abdominal obesity, impaired glucose tolerance, hypertension, elevated triglycerides, and reduced high-density lipoprotein (HDL)-cholesterol levels ⁽²⁾. The prevalence of MetS is on the rise in many world regions; in Latin America, it ranges from 19 to 43% ^(3, 4), and is comparable to estimates from Europe ⁽⁵⁾ and the United States ⁽⁶⁾. MetS components can become apparent from childhood ⁽⁷⁾ and are highly prevalent in children and adolescents from various Latin American settings ^(3, 8–10) due, in part, to the growing childhood obesity epidemic in this region ⁽¹¹⁾. For example, at least one in every five children aged 6 to 12 y from the Chilean Araucania region had insulin resistance ⁽⁸⁾ and one-third of normal-weight Mexican children of the same age group had >1 MetS component ⁽¹⁰⁾. These early metabolic abnormalities track into adulthood, ⁽¹²⁾ increasing risk for cardiovascular disease. Because MetS is mostly preventable, identifying potentially modifiable risk factors is an important research priority.

Vitamins B6, B9 (folate), and B12 are cofactors in the methionine-homocysteine (Hcys) cycle which is essential in the synthesis of methyl donors. Methyl donors play critical roles in multiple physiologic pathways including DNA methylation and lipid and protein metabolism. B-vitamin deficiencies could impair the synthesis of methyl groups and induce hyperhomocysteinemia (H-Hcys), which increases risk of atherosclerosis and cardiovascular disease ⁽¹³⁾. The potential role of these vitamins on MetS is less certain. In adults, vitamin B6 serostatus ⁽¹⁴⁾ and intake ⁽¹⁵⁾, and vitamin B12 ^(16–18) and folate ⁽¹⁶⁾ biomarkers and supplementation ⁽¹⁹⁾ have been inversely related to MetS, whereas H-Hcys has been positively associated with this outcome ^(20, 21). In children, the evidence is scant. Vitamin B12 serostatus was inversely associated with MetS ⁽²²⁾ and obesity ^(22, 23) in cross-sectional studies of Turkish and Israeli school-aged children and adolescents. Maternal folate supplementation ⁽²⁴⁾ and serostatus ^(25, 26) were inversely associated with MetS ⁽²⁴⁾, blood pressure ⁽²⁶⁾, and body mass index (BMI)-for-age Z score ⁽²⁵⁾ at ages 6 to 9 years in longitudinal studies from Nepal and the United States. A cross-sectional study of Czech adolescents found an inverse association of serum folate with hypertension ⁽²⁷⁾, whereas H-Hcys was positively related to MetS prevalence in Nepalese school-aged children ⁽²⁸⁾.

Because the prevalence of B-vitamin deficiencies is high in Latinamerican children ⁽²⁹⁾ and interventions during childhood may modulate cardiovascular risk in the long term ⁽³⁰⁾ there is a need to clarify the role of the B-vitamins and Hcys on MetS in this understudied

population. This study aimed to examine the associations of vitamin B6, B9, and B12 biomarkers and Hcys with MetS and its components among school-aged Mesoamerican children and their parents. The family design offered an opportunity to account for measured parental confounders in the estimation of associations among children.

METHODS

Study population

The Nine Mesoamerican Countries Metabolic Syndrome (NiMeCoMes) Study was a cross-sectional investigation on nutrition and cardiovascular health conducted in the capital cities of Belize, Guatemala, El Salvador, the Dominican Republic, Honduras, Nicaragua, Panama, Costa Rica, and Chiapas State in Mexico. Details of the study have been described elsewhere ⁽³⁾. In brief, we recruited a convenience sample of 267 family groups, a number determined by the availability of funding, each consisting of a school-aged child and their two parents from public primary schools in periurban areas of the cities, using convenience sampling. Eligibility criteria included child's age between 7 and 12 years, living with both biological parents, not being pregnant or having a pregnant mother, and not having a sibling already recruited into the study. We chose only few eligibility criteria to enhance generalizability of the findings. The study protocol and procedures were approved by the Institutional Review Boards (IRB) of collaborating institutions in each country and by the University of Michigan Health and Behavioral Sciences IRB. Written parental informed consent and children's assent to participate were obtained before enrollment.

Data collection

Data collection took place at a single home or clinic visit after a minimum 6-hour fasting. At this visit, participants responded to a questionnaire on sociodemographic characteristics including age, education level, household assets and income, home ownership, and past and current health status. The mothers answered the Latin American and Caribbean Food Security Scale (ELCSA) ⁽³¹⁾, a 15 yes/no item survey about food security experiences during the past three months; seven questions only concern households with children. Each affirmative response corresponded to one point and a household's food insecurity status was classified as: none (sum=0), mild (sum=1–3 or 1–5 in households without or with children, respectively), moderate (sum=4–6 or 6–10), or severe (sum=7–8 or 11–15). Research assistants administered a 97-item semiquantitative food frequency questionnaire (FFQ) separately to mothers, fathers, and children to estimate average intake during the past 12 months. The FFQ was based on a previously validated instrument developed for Costa Rican adults ⁽³²⁾. The FFQ characteristics have been detailed elsewhere ⁽³³⁾. In brief, items included major sources of energy and micronutrients in food groups such as fruits, vegetables, animal protein (dairy, eggs, and meat), carbohydrates (breads, flours, and cereals), beverages (dairy, sugar-sweetened beverages, coffee, and alcohol), fast foods, desserts, sauces and dressings, cooking fats and oils, and dietary supplements. For each item, research assistants described reference portion sizes in natural units or standard measures for commonly consumed servings among this population. There were nine frequency of intake options: 6 times/d, 4–5 times/d, 2–3 times/d, 1 time/d, 5–6 times/wk, 2–4 times/wk, 1 time/wk, 1–3 times/mo, or <1 time/mo. Researchers measured height, weight, and waist

circumference with the use of standardized methods and calibrated instruments. Waist circumference was measured at the end of an unforced exhalation to the nearest millimeter, at the midpoint between the lower edge of the ribcage and the iliac crest in adults and above the uppermost lateral border of the right ilium in children. All anthropometric measures were obtained in triplicate, and the median of the three values was used⁽³⁴⁾. Blood pressure was measured while seated, using Omron HEM-712C digital blood pressure monitors (Omron Healthcare, Inc., Lake Forest, IL, USA). Three measurements were taken, separated by at least one minute, and the average of the second and third measures was the value of blood pressure used. At the end of the visit, phlebotomists obtained 7.5 ml of blood through antecubital venipuncture. The median (interquartile range) estimated fasting time at the time of sample collection was 12.9 (12.0–14.0) h. Samples were placed in refrigerated containers and transported on the day of collection to each country's collaborating laboratories where the serum, plasma, and red blood cells were separated, aliquoted, and cryostored at –20°C. Stored samples from all countries were transported to the Institute of Nutrition of Central America and Panama (INCAP) in Guatemala City.

Laboratory methods

Quantification of insulin, glucose, lipids, vitamin B12, erythrocyte folate, and Hcys was conducted at INCAP. Plasma insulin, vitamin B12, and Hcys were measured using a chemiluminescent immunoassay on an Immulite 1000 system (Siemens Healthcare Diagnostics Products GmbH, Marburg, Germany). Plasma glucose and serum lipids (total and HDL-cholesterol and triglycerides) were quantified on an automated chemistry analyzer (Cobas c111 system; Roche Diagnostics, Mannheim, Germany). Erythrocyte folate was measured in red blood cell lysates using competitive immunoassays on an Immulite 2000 system (Siemens Healthcare Diagnostics Products GmbH, Marburg, Germany). Vitamin B6 biologically active metabolite, pyridoxal 5' phosphate (PLP), was measured in plasma using Liquid Chromatography-Mass Spectrometry on a Thermo Fisher/Cohesive Technologies Aria TLX Series HTLC System at the Mayo Medical Laboratories (Minnesota, USA).

Definition of outcomes

Children.—Because the metabolic syndrome is undefined in children <10 years of age, we calculated a metabolic risk score using MetS components that correspond to the definition in adults: waist circumference, the homeostatic model assessment for insulin resistance (HOMA-IR)⁽³⁵⁾, mean arterial pressure (MAP) serum high-density lipoprotein (HDL)-cholesterol, and serum triglycerides. MAP was calculated as [(2 x diastolic blood pressure) + systolic blood pressure]/3. The score was created by regressing each log-transformed component on sex and log-transformed age using linear regression models to obtain standardized regression residuals. We then averaged the residuals of the components with the negative of HDL to create the score. Higher scores indicate worse metabolic profile.

Adults.—The presence of MetS was defined according to the Adult Treatment Panel III criteria⁽²⁾ as having any 3 of the following: 1) waist circumference 102 cm in men and 88 cm in women; 2) fasting blood glucose 100 mg/dL; 3) systolic blood pressure 130 mmHg or diastolic blood pressure 85 mmHg, or treatment with an antihypertensive drug; 4) serum HDL-cholesterol <40 mg/dL in men and <50 mg/dL in women, or drug treatment for low

HDL-cholesterol; and 5) serum triglycerides ≥ 150 mg/dL or drug treatment for elevated triglycerides.

Definition of exposures

Exposures were plasma concentrations of PLP and vitamin B12, erythrocyte folate, and plasma Hcys. Prevalences of B-vitamin deficiencies according to conventional cutpoints were low; thus, biomarker concentrations were categorized into quartiles.

Covariates

Children.—Height-for-age and BMI-for-age Z scores were calculated using the World Health Organization (WHO) reference ⁽³⁶⁾. Parental height was categorized into quartiles and parental BMI according to the WHO classification. Household education was the maximum number of years of schooling achieved by either parent. Household food insecurity was categorized by the number of affirmative responses in the ELCSA survey (no insecurity, 0; mild insecurity, 1–5; moderate insecurity, 6–10; severe insecurity, ≥ 11). The number of household assets was the sum of car, bicycle, refrigerator/freezer, gas stove, electric stove, blender, microwave, washing machine, color television, sound set, computer, or Internet. Maternal parity was categorized as 1, 2, 3, or ≥ 4 births. We also considered as covariates food sources that may be predictors of the outcome. Using information collected with the FFQ, we created five food intake frequency groups (dairy, meat, fish, green leafy vegetables, and fortified foods) by adding intake frequency weights of individual foods corresponding to each group. Total energy intake was estimated by multiplying the intake frequency of each food by the energy contents of the specific food portion using values from the USDA's Standard Reference food composition database.

Adults.—Height, BMI, education level, home ownership, number of household assets, and household food security were categorized as presented in Supplemental Table 3. Parental smoking was categorized as never, past, or current. Household income was categorized into country-specific quartiles. Dietary covariates were defined as described in children.

Data analysis

The final analytic sample comprised 237 children and 524 adult parents who had information on both the exposures and MetS.

Children.—In bivariate analysis, we compared the distribution of MetS and MetS components by quartiles of exposures using means and SD. We estimated tests for trend by introducing a variable representing the median value of each quartile as a continuous predictor in a linear regression model with MetS or each MetS component as the outcome. In multivariable analyses, we estimated adjusted differences with 95% CI in mean metabolic scores between quartiles of exposures using linear regression models with robust variances. Adjustment covariates were independent predictors of the outcome including height-for-age Z score, maternal height, parental education, household food security, number of household assets, country of origin, habitual intake of vitamin food sources, and total energy intake. Because the MetS is standardized for the age and sex distributions of the study population, the inclusion of these variables in multivariable models is unnecessary. Estimates for each

vitamin were adjusted for the others but not for Hcys as this could be an intermediate variable. Estimates for Hcys were further adjusted for all vitamins. Models for each component, except waist circumference, were further adjusted for the child's BMI-for-age Z score.

Adults.—In bivariate analysis, we compared unadjusted MetS prevalences by quartiles of the exposures. We conducted tests for linear trend by introducing a variable representing the median value of each exposure quartile as a continuous predictor into a generalized estimation equations (GEE) model with the Poisson distribution. In multivariable analysis, we estimated prevalence ratios with 95% CI between quartiles of the exposures from GEE models using an analogous approach to adjustment as described for children. Adjustment variables were independent predictors of the outcome that could be related to the exposures but not be their consequence. Robust variances were specified in all models to account for within-household correlations.

All analyses were conducted using Statistical Analysis Software version 9.4 (SAS Institute, Cary, NC, USA).

RESULTS

Children

Mean age ($\pm$ SD) of children was 9.9 ± 1.6 years; 52.3% were girls. Thirty-five percent of children were overweight (BMI Z > 1 SD) and 19.4 % were obese (BMI Z > 2 SD). The mean ($\pm$ SD) metabolic risk score was 0.00 ± 0.22 . Deficiency prevalence of vitamin B6 (<20 nmol/), vitamin B12 (<148 pmol/L) and folate (<305 nmol/L) ⁽³⁷⁾ was 0.9%, 4.7%, and 2.1%, respectively; whereas 4.6% had H-Hcys (Hcys $>95^{\text{th}}$ percentile for age) ⁽³⁸⁾.

Correlates of B-vitamin status biomarkers.—Plasma PLP was inversely associated with paternal height and food insecurity, and positively related to parental education, number of household assets, and household income (Supplemental Table 1). Plasma vitamin B12 was positively related to female sex, parental height and education, and number of household assets; and inversely to maternal parity and household food insecurity. It varied significantly between countries. Erythrocyte folate was inversely related to the child's height-for-age Z score and household food insecurity. Hcys was positively associated with male sex, child's age, and maternal height, BMI, and parity; and inversely with parental education. It differed significantly by country. Plasma PLP was positively related to dairy intake (Supplemental Table 2); plasma vitamin B12 was associated with dairy, red meat, and fish intake, and erythrocyte folate was positively related to dairy, green leafy vegetables, and fortified foods intake.

B-vitamin biomarkers and metabolic score.—PLP was not significantly associated with the metabolic score (Table 1) or its components (Supplemental Table 3). Plasma vitamin B12 was inversely related to the metabolic score (Table 1); compared with children at the lowest vitamin B12 quartile, those at the highest quartile had an adjusted 0.13 units lower metabolic score (P, trend=0.008). The metabolic score components associated with plasma vitamin B12 included MAP, serum HDL-cholesterol, and triglycerides

(Supplemental Table 3). Erythrocyte folate was positively associated with the metabolic risk score (Table 1); the mean adjusted difference between quartiles 4 and 1 was 0.11 units (P , trend=0.02). Folate was positively related to waist circumference and MAP and inversely to HDL-cholesterol (Supplemental Table 3). Plasma Hcys was not associated with the metabolic score overall but was positively related to the waist circumference and arterial pressure components.

Adults

Mean age ($\pm$ SD) of participants was 38.7 ± 7.5 years and 50.4% were women; 34.2% had obesity ($\text{BMI} > 30 \text{ kg/m}^2$). The prevalence of MetS was 36.6%. Deficiency prevalence of vitamin B6 ($< 20 \text{ nmol/L}$), vitamin B12 ($< 148 \text{ pmol/L}$), and folate ($< 305 \text{ nmol/L}$) ⁽³⁷⁾ was 4.7%, 10.4%, and 8.4%, respectively; whereas 16.4% had H-Hcys ($\text{Hcys} > 14 \text{ } \mu\text{mol/L}$) ⁽³⁹⁾.

Correlates of B-vitamin biomarkers.—Plasma PLP was inversely associated with female sex and BMI and positively with home ownership, number of assets, and income (Supplemental Table 4). Vitamin B12 was positively related to the number of household assets, and income; it was highest in Mexico and lowest in Honduras and the Dominican Republic. Erythrocyte folate was inversely associated with male sex and height, and positively to home ownership and food security. It was highest in Mexico and lowest in the Dominican Republic. Hcys was positively associated with male sex and smoking status, and was highest in the Dominican Republic and lowest in Mexico. Vitamin B12 was positively associated with red meat, fish, and fortified foods intake (Supplemental Table 5) whereas erythrocyte folate was positively associated with intake of green leafy vegetables and fortified foods.

B-vitamin biomarkers and metabolic syndrome.—Plasma PLP was inversely associated with MetS (Table 2); adults in the highest quartile of plasma PLP had a 33% lower adjusted prevalence of MetS compared with adults in the lowest quartile (P , trend=0.02). Of the MetS components, only abdominal obesity was related to PLP (Supplemental Table 6). Vitamin B12 was positively associated with MetS in a non-linear fashion (Table 2). Compared with parents at the lowest quartile of plasma vitamin B12, those at the highest three quartiles combined had a 49% higher prevalence of MetS (95% CI: 7%, 108%; $P = 0.02$). Vitamin B12 was positively associated with high fasting blood glucose and high blood pressure (Supplemental Table 6). Erythrocyte folate was positively associated with MetS prevalence (Table 2); compared with adults at the lowest quartile, those in the highest had twice the prevalence of MetS (P , trend=0.003). Erythrocyte folate was not related to any of the components (Supplemental Table 6). Hcys was not significantly associated with MetS (Table 2) or its components (Supplemental Table 6).

DISCUSSION

In this cross-sectional study, a metabolic risk score in children was inversely associated with plasma vitamin B12 and positively related to erythrocyte folate. In adults, plasma PLP was inversely associated with MetS prevalence, whereas plasma vitamin B12 and erythrocyte folate were positively related to this outcome.

Vitamin B6 was not associated with the metabolic risk score in children; however, we found an inverse association of plasma PLP with MetS prevalence and abdominal obesity in adults. These results are consistent with previous cross-sectional studies in Nigeria ⁽¹⁴⁾ and Japan ⁽¹⁵⁾. In a cross-sectional study, plasma PLP was lower in adults with morbid obesity than in healthy-weight controls ⁽⁴⁰⁾. Low plasma PLP has been associated with increased levels of inflammatory and oxidative stress biomarkers ⁽⁴¹⁾ which might contribute to explain an inverse association with MetS since inflammation and oxidative stress are considered underlying etiologic factors ⁽²⁾.

Vitamin B12 was inversely associated with the metabolic risk score in children; possibly through blood pressure and serum lipids. Previous studies of European adolescents ⁽⁴²⁾ and Japanese preschool children ⁽⁴³⁾ found inverse associations between vitamin B12 serostatus and blood pressure. Also, cross-sectional investigations of Turkish ⁽²²⁾ and Israeli ⁽⁴⁴⁾ children found inverse associations between vitamin B12 serostatus and body weight. Our findings in children are in contrast to those in adults since vitamin B12 was positively related to MetS prevalence, possibly through high fasting blood glucose and high blood pressure. This finding was also inconsistent with prior cross-sectional studies that showed inverse associations between vitamin B12 and MetS ^(16–18). Similarly, a randomized trial ⁽¹⁹⁾, a Mendelian randomization study ⁽⁴⁵⁾, and a case control investigation ⁽¹⁶⁾ found inverse associations between vitamin B12 and measures of insulin resistance in adults, whereas a cross-sectional study found an inverse association of serum vitamin B12 with hypertension ⁽⁴⁶⁾. There are some possible explanations for the discrepancies. Although we were able to adjust for potential foods like red and processed meats that could also be associated with increased risk of cardiometabolic disease, there is a possibility of residual confounding which could explain the positive association of vitamin B12 with MetS. The discrepancy between our findings in children compared to adults could be explained by the different measures used to assess MetS, and direct comparisons may not be accurate. Additionally, children had higher dairy consumption compared to adults and there is evidence of an inverse association between dairy consumption and MetS in other populations ⁽⁴⁷⁾.

Erythrocyte folate was positively associated with the metabolic risk score in children, probably through increased waist circumference and blood pressure, and lowered HDL-cholesterol. Similarly, erythrocyte folate was positively associated with MetS prevalence in adults. Direct evaluations of the role of folate on MetS in children are scarce, although some studies had reported inverse associations between maternal folate and MetS or some of its components. In a prenatal supplementation study conducted in rural Nepal, offspring from mothers who had received antenatal folic acid with vitamin A supplements had a lower risk of MetS at ages 6 to 8 years compared with vitamin A alone ⁽²⁴⁾. Likewise, results from the Boston Birth Cohort suggested an L-shaped association of maternal plasma folate levels at the end of pregnancy with childhood overweight or obesity at ages 3 to 9 years; the highest risk was among children from mothers with obesity and low folate compared to children from mothers with normal weight and folate ⁽²⁵⁾. In the same cohort, there was also an inverse association of maternal folate levels with systolic blood pressure among children from mothers with cardiometabolic risk factors ⁽²⁶⁾. Two cross-sectional investigations in children found an inverse association between folate and blood pressure ^(27, 43), but another suggested a positive relation between erythrocyte folate and systolic

blood pressure among adolescent girls ⁽⁴²⁾. Our findings on both children and adults seem to contradict existing evidence on folate's relation with MetS and its components. However, some previous evidence suggests a potential adverse effect of folate on MetS components; notably, cross-sectional evaluations of adults with obesity found a positive association between erythrocyte folate and HOMA-IR, especially among participants with low vitamin B12 status ⁽⁴⁸⁾. One proposed mechanism to explain a potential deleterious effect of folate on metabolism is the methyl-pool depletion caused by folic acid, the standard form of folate used for fortification ⁽⁴⁹⁾, which is mandatory for wheat and corn flour in Mesoamerica ⁽⁵⁰⁾. In our population, fortified foods intake was significantly associated with erythrocyte folate and folate concentrations were generally high. Whether folic acid from fortified foods could have a deleterious role on metabolic health warrants further investigation.

Hcys was positively associated with waist circumference and blood pressure in children, but there was no association with MetS or any of its components in adults. Previous cross-sectional studies in school-aged children found that H-Hcys was positively associated with MetS, high blood pressure ⁽²⁸⁾, and waist circumference >90th percentile ⁽⁵¹⁾. Nonetheless, one cohort study found no association between serum Hcys with MetS components or insulin sensitivity among healthy men from London ⁽⁵²⁾. A supplementation trial suggested that, while folate and vitamin B12 might reduce Hcys, this effect may not extend to cardiometabolic outcomes ⁽⁵³⁾.

Our study has several strengths. We used valid biomarkers of intake, which precludes recall bias in the ascertainment of exposure compared to methods that rely on memory. The prevalence of B-vitamin deficiencies was within range of other studies in comparable populations, which enhances external validity. For example, 4.7% of children in this study were vitamin B12-deficient, compared with 2.6% in Mexico in 2012 ⁽⁵⁴⁾ and 2.8% in Colombia in 2010 ⁽⁵⁵⁾; and only 2.1% of children were folate-deficient, consistent with reports from Mexico ⁽⁵⁴⁾ and Guatemala ⁽⁵⁶⁾, where prevalence was zero. Among adults, the prevalence of vitamin B12 deficiency, 10.4%, was also within range of reports among women from Costa Rica (4.8% in 2008) ⁽⁵⁷⁾, Mexico (8.5% in 2012) ⁽⁵⁸⁾, Argentina (11.9% in 2007) ⁽⁵⁹⁾, and Colombia (18.5% in 2010) ⁽⁵⁵⁾. Prevalence of folate deficiency in adults, 8.4%, was slightly higher than that reported among women from Mexico (1.9% in 2012) ⁽⁵⁸⁾, Costa Rica (3.8% in 2008) ⁽⁵⁷⁾, Guatemala (3.1% in 2008) ⁽⁶⁰⁾, and Argentina (1.3% in 2007) ⁽⁵⁷⁾. No previous studies in the region have reported on the prevalence of vitamin B6 deficiency according to PLP concentrations; the figure among adults, 4.7%, was lower than that reported in the United States for people 9 y-old, 11% ⁽⁶¹⁾. The prevalence of MetS in adults, 37%, was close to that in Mexico, 41% ⁽⁶²⁾, but lower than that in Guatemala, 64% ⁽⁶³⁾. The family design involving children with their parents allowed adjustment of the associations in children for parental confounders. Evaluating the associations of B-vitamins with MetS in Mesoamerican children is a novel endeavor in an understudied population.

Some limitations are also noteworthy. The cross-sectional design limits the making of causal inference. Comparisons of the absolute value of the children's metabolic risk score with other populations is not possible because it was standardized for the population-specific sex and age distributions. Since the study sample was not representative of the entire

Mesoamerican population, generalizability of the findings is limited. Finally, due to small sample size, country-specific analyses were not possible.

In conclusion, vitamin B12 was inversely associated with a metabolic risk score in Mesoamerican children but was positively associated with MetS prevalence in adults. Unexpectedly, erythrocyte folate was positively associated with MetS and some of its components in both children and adults. The potential role of mandatory folate fortification on these findings deserves careful consideration in future studies.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements:

Participants in the Nine Mesoamerican Countries Metabolic Syndrome Study (NiMeCoMeS) Group:

Mexico: Erika Lopez, Liz Peña, Alejandra Maldonado, Aldeni Vasquez, Aldrin Lopez

Belize: Lilly Mahung, Diomar Salazar

Guatemala: Ana Victoria Román, Fernanda Kroker, Maria Alejandra Cordova, Regina Garcia, Lilian Navas

El Salvador: Josefina Sibrian, Mauricio Flores, Noel Avalos

Honduras: Astarte Alegria, Jorge A. Sierra, Hector Murillo

Nicaragua: Ana María Gutierrez, Carmen María Flores, Mario Romero

Costa Rica: Emilce Ulate, Natalia Valverde, Andrea Fiatt, Juan Manuel Valverde

Panama: Flavia Fontes, Raísa Rodríguez, Emerita Pons, Lino Chue, Elka Gonzalez

Dominican Republic: Rafael Montero, Francisco Torres, Amarilis Then, Melvi Perez

Financial Support: This work was supported by the United States National Heart, Lung, and Blood Institute [grant number BAA-NHLBI-HV-09-12]. The funding sources had no role in the study design, collection or interpretation of data, writing of the report, and decision to submit the article for publication.

REFERENCES

1. Global Burden of Disease Causes of Death Collaborators (2017) Global, regional, and national age-sex specific mortality for 264 causes of death, 1980–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 390, 1151–1210. [PubMed: 28919116]
2. Grundy SM, Cleeman JI, Daniels SR et al. (2005) Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* 112, 2735–2752. [PubMed: 16157765]
3. Villamor E, Finan CC, Ramirez-Zea M et al. (2017) Prevalence and sociodemographic correlates of metabolic syndrome in school-aged children and their parents in nine Mesoamerican countries. *Public Health Nutr* 20, 255–265. [PubMed: 27609776]
4. Marquez-Sandoval F, Macedo-Ojeda G, Viramontes-Horner D et al. (2011) The prevalence of metabolic syndrome in Latin America: a systematic review. *Public Health Nutr* 14, 1702–1713. [PubMed: 21486521]
5. Balkau B, Charles MA, Drivsholm T et al. (2002) Frequency of the WHO metabolic syndrome in European cohorts, and an alternative definition of an insulin resistance syndrome. *Diabetes Metab* 28, 364–376. [PubMed: 12461473]

6. Moore JX, Chaudhary NAKinyemiju T (2017) Metabolic Syndrome Prevalence by Race/Ethnicity and Sex in the United States, National Health and Nutrition Examination Survey, 1988–2012. *Prev Chronic Dis* 14, E24. [PubMed: 28301314]
7. Zimmet P, Alberti KG, Kaufman Fet al. (2007) The metabolic syndrome in children and adolescents - an IDF consensus report. *Pediatr Diabetes* 8, 299–306. [PubMed: 17850473]
8. Sapunar J, Aguilar-Farias N, Navarro Jet al. (2018) [High prevalence of overweight, obesity, insulin resistance and metabolic syndrome in rural children and adolescents]. *Rev Med Chil* 146, 978–986. [PubMed: 30725017]
9. Burrows R, Correa-Burrows P, Reyes Met al. (2016) High cardiometabolic risk in healthy Chilean adolescents: associations with anthropometric, biological and lifestyle factors. *Public Health Nutr* 19, 486–493. [PubMed: 25990645]
10. Guzman-Guzman IP, Salgado-Bernabe AB, Munoz Valle JFet al. (2015) [Prevalence of metabolic syndrome in children with and without obesity]. *Med Clin (Barc)* 144, 198–203. [PubMed: 24721677]
11. Rivera JA, de Cossio TG, Pedraza LSet al. (2014) Childhood and adolescent overweight and obesity in Latin America: a systematic review. *Lancet Diabetes Endocrinol* 2, 321–332. [PubMed: 24703050]
12. Morrison JA, Friedman LA, Wang Pet al. (2008) Metabolic syndrome in childhood predicts adult metabolic syndrome and type 2 diabetes mellitus 25 to 30 years later. *J Pediatr* 152, 201–206. [PubMed: 18206689]
13. Strain JJ, Dowey L, Ward Met al. (2004) B-vitamins, homocysteine metabolism and CVD. *Proc Nutr Soc* 63, 597–603. [PubMed: 15831132]
14. Odum EP & Wakwe VC (2012) Plasma concentrations of water-soluble vitamins in metabolic syndrome subjects. *Niger J Clin Pract* 15, 442–447. [PubMed: 23238195]
15. Otsuka R, Imai T, Kato Yet al. (2010) Relationship between number of metabolic syndrome components and dietary factors in middle-aged and elderly Japanese subjects. *Hypertens Res* 33, 548–554. [PubMed: 20224573]
16. Narang M, Singh M, Dange S (2016) Serum Homocysteine, Vitamin B12 and Folic Acid Levels in Patients with Metabolic Syndrome. *J Assoc Physicians India* 64, 22–26.
17. Guven A, Inanc F, Kilinc Met al. (2005) Plasma homocysteine and lipoprotein (a) levels in Turkish patients with metabolic syndrome. *Heart Vessels* 20, 290–295. [PubMed: 16314912]
18. Onat A, Hergenc G, Kucukdurmaz Zet al. (2008) Serum folate is associated with coronary heart disease independently of homocysteine in Turkish men. *Clin Nutr* 27, 732–739. [PubMed: 18694616]
19. Setola E, Monti LD, Galluccio Eet al. (2004) Insulin resistance and endothelial function are improved after folate and vitamin B12 therapy in patients with metabolic syndrome: relationship between homocysteine levels and hyperinsulinemia. *Eur J Endocrinol* 151, 483–489. [PubMed: 15476449]
20. Oron-Herman M, Rosenthal TSela BA (2003) Hyperhomocysteinemia as a component of syndrome X. *Metabolism* 52, 1491–1495. [PubMed: 14624412]
21. Catena C, Colussi G, Nait Fet al. (2015) Elevated Homocysteine Levels Are Associated With the Metabolic Syndrome and Cardiovascular Events in Hypertensive Patients. *Am J Hypertens* 28, 943–950. [PubMed: 25498997]
22. Ozer S, Sonmezgoz EDemir O (2017) Negative correlation among vitamin B12 levels, obesity severity and metabolic syndrome in obese children: A case control study. *J Pak Med Assoc* 67, 1648–1653. [PubMed: 29171553]
23. Ho M, Halim JH, Gow MLet al. (2014) Vitamin B12 in obese adolescents with clinical features of insulin resistance. *Nutrients* 6, 5611–5618. [PubMed: 25486369]
24. Stewart CP, Christian P, Schulze KJet al. (2009) Antenatal micronutrient supplementation reduces metabolic syndrome in 6- to 8-year-old children in rural Nepal. *J Nutr* 139, 1575–1581. [PubMed: 19549749]
25. Wang G, Hu FB, Mistry KBet al. (2016) Association Between Maternal Prepregnancy Body Mass Index and Plasma Folate Concentrations With Child Metabolic Health. *JAMA Pediatr* 170, e160845. [PubMed: 27295011]

26. Wang H, Mueller NT, Li Jet al. (2017) Association of Maternal Plasma Folate and Cardiometabolic Risk Factors in Pregnancy with Elevated Blood Pressure of Offspring in Childhood. *Am J Hypertens* 30, 532–540. [PubMed: 28338750]
27. Kahleova R, Palyzova D, Zvara Ket al. (2002) Essential hypertension in adolescents: association with insulin resistance and with metabolism of homocysteine and vitamins. *Am J Hypertens* 15, 857–864. [PubMed: 12372672]
28. Yakub M, Schulze KJ, Khatry SKet al. (2014) High plasma homocysteine increases risk of metabolic syndrome in 6 to 8 year old children in rural Nepal. *Nutrients* 6, 1649–1661. [PubMed: 24763111]
29. Allen LH (2004) Folate and vitamin B12 status in the Americas. *Nutr Rev* 62, S29–33. [PubMed: 15298445]
30. Nupponen M, Pakkala K, Juonala Met al. (2015) Metabolic syndrome from adolescence to early adulthood: effect of infancy-onset dietary counseling of low saturated fat: the Special Turku Coronary Risk Factor Intervention Project (STRIP). *Circulation* 131, 605–613. [PubMed: 25605660]
31. Pérez-Escamilla R, Melgar-Quiñonez H, Nord Met al. (2007) Escala Latinoamericana y Caribeña de Seguridad Alimentaria (ELCSA). *Perspectivas en Nutricion Humana* (October Supplement), 117–134.
32. Campos H, Willett WC, Peterson RMet al. (1991) Nutrient intake comparisons between Framingham and rural and Urban Puriscal, Costa Rica. Associations with lipoproteins, apolipoproteins, and low density lipoprotein particle size. *Arterioscler Thromb* 11, 1089–1099. [PubMed: 2065030]
33. Robinson SL, Ramirez-Zea M, Roman AVet al. (2017) Correlates and family aggregation of vitamin D concentrations in school-aged children and their parents in nine Mesoamerican countries. *Public Health Nutr* 20, 2754–2765. [PubMed: 28735597]
34. Villamor E & Bosch RJ (2015) Optimal treatment of replicate measurements in anthropometric studies. *Ann Hum Biol* 42, 507–510. [PubMed: 25426678]
35. Matthews DR, Hosker JP, Rudenski ASet al. (1985) Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28, 412–419. [PubMed: 3899825]
36. de Onis M, Onyango AW, Borghi Eet al. (2007) Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* 85, 660–667. [PubMed: 18026621]
37. Institute of Medicine (1998). In *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline*. Washington, DC: National Academic Press.
38. Papandreou D, Mavromichalis I, Makedou Aet al. (2006) Reference range of total serum homocysteine level and dietary indexes in healthy Greek schoolchildren aged 6–15 years. *Br J Nutr* 96, 719–724. [PubMed: 17010232]
39. Selhub J, Jacques PF, Bostom AGet al. (1995) Association between plasma homocysteine concentrations and extracranial carotid-artery stenosis. *N Engl J Med* 332, 286–291. [PubMed: 7816063]
40. Aasheim ET, Hofso D, Hjelmestaeth Jet al. (2008) Vitamin status in morbidly obese patients: a cross-sectional study. *Am J Clin Nutr* 87, 362–369. [PubMed: 18258626]
41. Shen J, Lai CQ, Mattei Jet al. (2010) Association of vitamin B-6 status with inflammation, oxidative stress, and chronic inflammatory conditions: the Boston Puerto Rican Health Study. *Am J Clin Nutr* 91, 337–342. [PubMed: 19955400]
42. de Moraes AC, Gracia-Marco L, Iglesia Iet al. (2014) Vitamins and iron blood biomarkers are associated with blood pressure levels in European adolescents. The HELENA study. *Nutrition* 30, 1294–1300. [PubMed: 25194965]
43. Tamai Y, Wada K, Tsuji Met al. (2011) Dietary intake of vitamin B12 and folic acid is associated with lower blood pressure in Japanese preschool children. *Am J Hypertens* 24, 1215–1221. [PubMed: 21814291]

44. Pinhas-Hamiel O, Doron-Panush N, Reichman Bet al. (2006) Obese children and adolescents: a risk group for low vitamin B12 concentration. *Arch Pediatr Adolesc Med* 160, 933–936. [PubMed: 16953016]
45. Moen GH, Qvigstad E, Birkeland KI et al. (2018) Are serum concentrations of vitamin B-12 causally related to cardiometabolic risk factors and disease? A Mendelian randomization study. *Am J Clin Nutr* 108, 398–404. [PubMed: 29982347]
46. Mahalle N, Kulkarni MV, Garg MK et al. (2013) Vitamin B12 deficiency and hyperhomocysteinemia as correlates of cardiovascular risk factors in Indian subjects with coronary artery disease. *J Cardiol* 61, 289–294. [PubMed: 23473764]
47. Chen GC, Szeto IM, Chen LH et al. (2015) Dairy products consumption and metabolic syndrome in adults: systematic review and meta-analysis of observational studies. *Sci Rep* 5, 14606. [PubMed: 26416233]
48. Li Z, Gueant-Rodriguez RM, Quilliot Det al. (2018) Folate and vitamin B12 status is associated with insulin resistance and metabolic syndrome in morbid obesity. *Clin Nutr* 37, 1700–1706. [PubMed: 28780990]
49. Bailey LB, Stover PJ, McNulty Het al. (2015) Biomarkers of Nutrition for Development-Folate Review. *J Nutr* 145, 1636S–1680S. [PubMed: 26451605]
50. Bonilla-Soto AI (2016) Fortificación de alimentos en Centroamérica y el Caribe. <http://ilsimesoamerica.org/wp-content/uploads/sites/14/2016/08/Monograf%C3%ADa-Fortificaci%C3%B3n-de-alimentos-para-Centroam%C3%A9rica-y-el-Caribe.pdf>. Guatemala City.
51. da Silva NP, de Souza FI, Pendergast AI et al. (2013) Homocysteine and cysteine levels in prepubertal children: association with waist circumference and lipid profile. *Nutrition* 29, 166–171. [PubMed: 23022121]
52. Godsland IF, Rosankiewicz JR, Proudler AJ et al. (2001) Plasma total homocysteine concentrations are unrelated to insulin sensitivity and components of the metabolic syndrome in healthy men. *J Clin Endocrinol Metab* 86, 719–723. [PubMed: 11158036]
53. Marti-Carvajal AJ, Sola IL et al. (2015) Homocysteine-lowering interventions for preventing cardiovascular events. *Cochrane Database Syst Rev* 1, CD006612.
54. Villalpando S, Cruz V de L, Shamah-Levy Tet al. (2015) Nutritional status of iron, vitamin B12, folate, retinol and anemia in children 1 to 11 years old: Results of the Ensanut 2012. *Salud Publica Mex* 57, 372–384. [PubMed: 26544998]
55. Herran OF, Ward JB, Villamor E (2015) Vitamin B12 serostatus in Colombian children and adult women: results from a nationally representative survey. *Public Health Nutr* 18, 836–843. [PubMed: 24969611]
56. Rogers LM, Boy E, Miller JW et al. (2003) High prevalence of cobalamin deficiency in Guatemalan schoolchildren: associations with low plasma holotranscobalamin II and elevated serum methylmalonic acid and plasma homocysteine concentrations. *Am J Clin Nutr* 77, 433–440. [PubMed: 12540405]
57. Brito A, Mujica-Coopman MF, Lopez de Romana Det al. (2015) Folate and Vitamin B12 Status in Latin America and the Caribbean: An Update. *Food Nutr Bull* 36, S109–118. [PubMed: 26125196]
58. Shamah-Levy T, Villalpando S, Mejia-Rodriguez Fet al. (2015) Prevalence of iron, folate, and vitamin B12 deficiencies in 20 to 49 years old women: Ensanut 2012. *Salud Publica Mex* 57, 385–393. [PubMed: 26544999]
59. Calvo EB & Biglieri A (2008) [Impact of folic acid fortification on women's nutritional status and on the prevalence of neural tube defects]. *Arch Argent Pediatr* 106, 492–498. [PubMed: 19107300]
60. Nguyen P, Grajeda R, Melgar Pet al. (2008) Weekly may be as efficacious as daily folic acid supplementation in improving folate status and lowering serum homocysteine concentrations in Guatemalan women. *J Nutr* 138, 1491–1498. [PubMed: 18641196]
61. Bird JK, Murphy RA, Ciappio ED et al. (2017) Risk of Deficiency in Multiple Concurrent Micronutrients in Children and Adults in the United States. *Nutrients* 9.
62. Gutierrez-Solis AL, Datta Banik SM, Mendez-Gonzalez RM (2018) Prevalence of Metabolic Syndrome in Mexico: A Systematic Review and Meta-Analysis. *Metab Syndr Relat Disord* 16, 395–405. [PubMed: 30063173]

63. Rivera-Andrade A, Kroker-Lobos MF, Lazo Met al. (2019) High prevalence of non-alcoholic fatty liver disease and metabolic risk factors in Guatemala: A population-based study. *Nutr Metab Cardiovasc Dis* 29, 191–200. [PubMed: 30573307]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1.

Metabolic syndrome score* according to plasma vitamins B6 and B12, erythrocyte folate, and plasma homocysteine concentrations among school-age children from Mesoamerica

Biomarker quartile (median)	n [†]	Mean	SD	Unadjusted difference 95% CI		Adjusted difference 95% CI			
						Model 1 [‡]		Model 2 [§]	
Plasma PLP (nmol/L)									
Q1 (36)	50	0.02	0.21		Reference	Reference		Reference	
Q2 (49)	52	−0.04	0.20	−0.05	−0.13, 0.02	−0.05	−0.12, 0.02	−0.07	−0.14, 0.00
Q3 (65)	67	0.03	0.24	0.01	−0.07, 0.09	0.02	−0.05, 0.09	0.00	−0.07, 0.07
Q4 (107)	60	0.01	0.24	0.00	−0.09, 0.08	−0.01	−0.09, 0.06	−0.02	−0.10, 0.06
P, trend ^{//}					0.69	0.88		0.87	
Plasma vitamin B12 (pmol/L)									
Q1 (191)	58	0.10	0.24		Reference	Reference		Reference	
Q2 (297)	58	−0.02	0.23	−0.12	−0.20, −0.03	−0.10	−0.18, −0.02	−0.07	−0.15, 0.01
Q3 (404)	60	−0.03	0.21	−0.13	−0.21, −0.05	−0.14	−0.22, −0.06	−0.12	−0.20, −0.05
Q4 (719)	59	−0.03	0.20	−0.12	−0.20, −0.04	−0.14	−0.22, −0.05	−0.13	−0.21, −0.04
P, trend					0.02	0.01		0.008	
Erythrocyte folate (nmol/L)									
Q1 (531)	58	0.02	0.23		Reference	Reference		Reference	
Q2 (735)	59	−0.01	0.22	−0.03	−0.11, 0.06	−0.03	−0.11, 0.04	0.01	−0.07, 0.08
Q3 (858)	59	−0.02	0.19	−0.03	−0.11, 0.04	0.00	−0.08, 0.07	0.04	−0.02, 0.11
Q4 (1169)	59	0.03	0.26	0.01	−0.08, 0.10	0.04	−0.05, 0.14	0.11	0.01, 0.20
P, trend					0.75	0.21		0.02	
Plasma Hcys (μmol/L)									
Q1 (3.8)	57	0.00	0.26		Reference	Reference		Reference	
Q2 (5.5)	60	0.02	0.23	0.02	−0.07, 0.11	0.05	−0.03, 0.13	0.02	−0.06, 0.10
Q3 (7.5)	59	−0.02	0.19	−0.03	−0.11, 0.05	0.05	−0.03, 0.14	0.03	−0.05, 0.12
Q4 (10.5)	59	0.02	0.23	0.01	−0.08, 0.10	0.07	−0.02, 0.17	0.06	−0.04, 0.16
P, trend					0.99	0.18		0.22	

PLP, pyridoxal phosphate; Hcys, homocysteine

* Component scores (waist circumference, HOMA-IR, MAP, serum HDL-cholesterol, and serum triglycerides) were computed by regressing each log-transformed component on sex and log-transformed age in linear regression models to obtain standardized residuals. After the HDL-cholesterol score was multiplied by -1, the overall score was calculated as the average of the five component scores.

[†] N may be less than 237 due to missing values.

[‡] From linear regression models adjusted for height-for-age Z score, maternal height, highest parental education level, household food security, number of household assets, and country of origin. Estimates for the vitamins were adjusted for each other but not for Hcys; estimates for Hcys were adjusted for all vitamins.

[§] Adjusted for covariates in model 1 plus log-transformed total energy intake, and intake of dairy, meat, fish, and green leafy vegetables.

^{//} From linear regression models with the metabolic score as the outcome and a variable representing medians of ordinal categories of the predictor introduced as a continuous covariate. Robust estimates of variance were specified in all models.

Table 2.

Metabolic syndrome^{*} according to plasma vitamins B6 and B12, erythrocyte folate, and plasma homocysteine concentrations among adults from Mesoamerica

Biomarker quartile (median)	n [†]	Prevalence %	Unadjusted PR	95% CI	Adjusted PR 95% CI			
					Model 1 [‡]		Model 2 [§]	
Plasma PLP (nmol/L)								
Q1 (24)	110	44.6	Reference		Reference		Reference	
Q2 (36)	119	44.5	1.00	0.74, 1.36	0.97	0.71, 1.34	0.95	0.69, 1.31
Q3 (53)	142	28.9	0.65	0.46, 0.91	0.61	0.43, 0.86	0.59	0.41, 0.83
Q4 (89)	141	30.5	0.68	0.49, 0.95	0.66	0.46, 0.93	0.67	0.47, 0.95
P, trend ^{//}			0.009		0.01		0.02	
Plasma vitamin B12 (pmol/L)								
Q1 (155)	130	25.4	Reference		Reference		Reference	
Q2 (232)	130	43.9	1.73	1.20, 2.49	1.61	1.12, 2.31	1.67	1.14, 2.43
Q3 (345)	130	37.7	1.48	1.01, 2.18	1.40	0.96, 2.06	1.47	0.99, 2.17
Q4 (619)	130	39.2	1.55	1.06, 2.24	1.42	0.97, 2.10	1.50	1.00, 2.26
P, trend			0.15		0.39		0.28	
Erythrocyte folate (nmol/L)								
Q1 (413)	130	32.3	Reference		Reference		Reference	
Q2 (699)	130	30.0	0.93	0.65, 1.34	1.19	0.77, 1.84	1.29	0.83, 2.02
Q3 (891)	130	38.5	1.19	0.84, 1.68	1.51	0.96, 2.37	1.62	1.02, 2.57
Q4 (1218)	132	45.5	1.41	1.02, 1.94	1.83	1.18, 2.84	1.97	1.22, 3.17
P, trend			0.02		0.003		0.003	
Plasma Hcys (μmol/L)								
Q1 (5.2)	129	42.6	Reference		Reference		Reference	
Q2 (7.8)	131	32.8	0.77	0.56, 1.05	0.71	0.51, 0.98	0.71	0.50, 1.00
Q3 (10.2)	130	36.9	0.87	0.65, 1.16	0.82	0.58, 1.14	0.75	0.52, 1.07
Q4 (15.2)	130	33.9	0.79	0.57, 1.10	0.78	0.54, 1.12	0.72	0.48, 1.09
P, trend			0.30		0.39		0.24	

PR, prevalence ratios; PLP, pyridoxal phosphate; Hcys, homocysteine

^{*} According to the National Cholesterol Education Program's Adult Treatment Panel III definition as having three or more of the following components: abdominal obesity, high fasting glucose, high blood pressure, low serum HDL-cholesterol, and high serum triglycerides.

[†] N may be less than 524 due to missing values.

[‡] From Poisson regression models adjusted for age, sex, smoking and education level, number of household assets, household food security, and country of origin. Estimates for the vitamins were adjusted for each other but not for Hcys; Hcys estimates were adjusted for all vitamins.

[§] Adjusted for covariates in model 1 plus log-transformed total energy intake, and intake of meat, fish, fortified foods, and green leafy vegetables.

^{//} From Poisson regression models with metabolic syndrome as the outcome and a variable representing medians of ordinal categories of the predictor introduced as a continuous covariate. Robust estimates of variance were specified in all models to account for clustering by family membership.